

Large Deviations for Brownian Intersection Measures

WOLFGANG KÖNIG

*Technische Universität Berlin
Weierstrass Institute Berlin*

CHIRANJIB MUKHERJEE

Max-Planck Institute for Mathematics in the Sciences in Leipzig

Abstract

We consider p independent Brownian motions in \mathbb{R}^d . We assume that $p \geq 2$ and $p(d-2) < d$. Let ℓ_t denote the intersection measure of the p paths by time t , i.e., the random measure on \mathbb{R}^d that assigns to any measurable set $A \subset \mathbb{R}^d$ the amount of intersection local time of the motions spent in A by time t . Earlier results of X. Chen derived the logarithmic asymptotics of the upper tails of the total mass $\ell_t(\mathbb{R}^d)$ as $t \rightarrow \infty$. In this paper, we derive a large-deviation principle for the normalized intersection measure $t^{-p}\ell_t$ on the set of positive measures on some open bounded set $B \subset \mathbb{R}^d$ as $t \rightarrow \infty$ before exiting B . The rate function is explicit and gives some rigorous meaning, in this asymptotic regime, to the understanding that the intersection measure is the pointwise product of the densities of the normalized occupation times measures of the p motions. Our proof makes the classical Donsker-Varadhan principle for the latter applicable to the intersection measure.

A second version of our principle is proved for the motions observed until the individual exit times from B , conditional on a large total mass in some compact set $U \subset B$. This extends earlier studies on the intersection measure by König and Mörters. © 2012 Wiley Periodicals, Inc.

1 Introduction and Main Results

1.1 Brownian Intersection Local Time

Let $W^{(1)}, \dots, W^{(p)}$ be p independent Brownian motions in \mathbb{R}^d . We assume throughout this paper that $p \geq 2$ and $d < \frac{2p}{p-1}$, which comprises the following cases:

$$p \begin{cases} \geq 2 & \text{arbitrary in } d = 2, \\ = 2 & \text{in } d = 3. \end{cases}$$

In the 1950s Dvoretzky, Erdős, Kakutani, and Taylor [10, 11, 12] showed that, almost surely, the intersection set of the p paths on individual time horizons,

$$S_b = \bigcap_{i=1}^p W_{[0, b_i]}^{(i)}, \quad b = (b_1, \dots, b_p) \in (0, \infty)^p,$$

is nonempty. Further results [14, 24] showed S_b has measure 0 in $d \geq 2$ and Hausdorff dimension 2 in $d = 2$ and one in $d = 3$. Hence S_b is a rather peculiar and interesting random set.

There is a natural measure ℓ_b supported on S_b counting the intensity of path intersections. This measure can be formally defined by

$$(1.1) \quad \ell_b(A) = \int_A dy \prod_{i=1}^p \int_0^{b_i} ds \delta_y(W_s^{(i)}) \quad \text{for every measurable } A \subset \mathbb{R}^d.$$

Hence, informally, ℓ_b is the pointwise product of the densities of the p occupation measures on the individual time horizons. This definition is rigorous in dimension $d = 1$, as the occupation measures of the motions have almost surely a density that is jointly continuous in the space and time variables. However, in $d \geq 2$, the occupation measures fail to have a density. Therefore, the above heuristic formula for ℓ_b needs a rigorous explanation. Geman, Horowitz, and Rosen [16] constructed ℓ_b as the local time of the confluent Brownian motion at 0, Le Gall [20] identified it as a renormalized limit of the Lebesgue measure on the intersection of Wiener sausages, and a third identification is in terms of a Hausdorff measure on S_b with explicit identification of the gauge function [21, 22]. These three rigorous constructions of ℓ_b are summarized in [4] and briefly surveyed in [18, sec. 2.1]. As a byproduct of the present paper, we will implicitly give a fourth construction in terms of a rescaled limit of pointwise products of smoothed occupation times; see Proposition 2.3. Some of the preceding results have been derived for b_1, \dots, b_p replaced by certain random times (independent exponential times or exit times from domains), but the proofs easily carry over to ℓ_b .

The measure ℓ_b is, with probability 1, positive and locally finite on \mathbb{R}^d . It is usually called *intersection local time (ISLT)* in the literature. However, its total mass, $\ell_b(\mathbb{R}^d)$, also enjoys this name, as it registers the total amount of intersections of the motions. Since the difference between these two objects will be significant in this paper, we will stick to the name *intersection measure* for ℓ_b and keep the name ISLT for its total mass $\ell_b(\mathbb{R}^d)$.

1.2 Asymptotics for Large Total Mass

The large- t behavior of the ISLT $\ell_{t\mathbf{1}}(\mathbb{R}^d)$ (where $\mathbf{1} = (1, \dots, 1)$) has been studied by X. Chen in a series of papers; see his monograph [4] for a comprehensive summary of these results, concepts of the proofs, and much more related material. The main result [4, theorem 3.3.2] is

$$(1.2) \quad \lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}(\ell_{t\mathbf{1}}(\mathbb{R}^d) > at^p) = -a^{2/d(p-1)} \chi, \quad a > 0,$$

where

$$(1.3) \quad \chi = \inf \left\{ \frac{p}{2} \|\nabla \psi\|_2^2 : \psi \in H^1(\mathbb{R}^d), \|\psi\|_{2p} = 1 = \|\psi\|_2 \right\}.$$

As we will explain in more detail in Section 1.4, the term ψ^2 informally plays the role of the normalized occupation measure density of any of the p motions, and ψ^{2p} the one of the intersection measure $t^{-p}\ell_{t1}$. This is one of the main features of intersection measures: How much rigorous meaning can be given to the intersection measure as a pointwise product of p occupation measures? The above result indicates that some heuristic sense can be given in terms of a large- t limit based on the understanding of the characteristic variational formula.

One of the main goals of this paper is to give a more rigorous meaning to this interpretation in terms of a large-deviation principle (LDP), at least for the case that the motions do not leave a given bounded set. Fix a bounded open set $B \subset \mathbb{R}^d$ with smooth boundary and compact closure \bar{B} and denote by $\tau_i = \inf\{t > 0 : W_t^{(i)} \notin B\}$ the exit time of the i^{th} motion from B . By $\ell = \ell_B$ we denote the intersection measure for the motions running up to their individual exit times from B ; i.e., we replace the time horizon $[0, b_1] \times \cdots \times [0, b_p]$ in (1.1) by $[0, \tau_1] \times \cdots \times [0, \tau_p]$. Then ℓ is a finite positive measure on B . Fix some compact subset U of B such that the boundary of U is a Lebesgue null set. The upper tails of $\ell(U)$ have been analyzed by König and Mörters [18], resulting in the asymptotics

$$(1.4) \quad \lim_{a \rightarrow \infty} a^{-1/p} \log \mathbb{P}(\ell(U) > a) = -\Theta_B(U)$$

for

$$(1.5) \quad \Theta_B(U) = \inf \left\{ \frac{p}{2} \|\nabla \phi\|_2^2 : \phi \in H_0^1(B), \|1_U \phi\|_{2p} = 1 \right\}.$$

This result is in the same spirit as the one above by Chen. Again, ϕ^2 and ϕ^{2p} can be informally interpreted as the densities of the individual occupation measures and the intersection measure, respectively. Denote by M the set of minimizing functions ϕ^{2p} ; then M is nonempty [18, theorem 1.3], and the elements of M carry some rigorous sense in terms of a law of large masses. Indeed, under the conditional measure $\mathbb{P}(\cdot \mid \ell(U) > a)$, it is shown in [19] that the distance of the normalized measure $\ell/\ell(U)$ (with harmonic extension to B) from M (where the elements of M are seen as probability measures on U) tends to 0 as $a \rightarrow \infty$. However, [19] failed to show that this convergence is exponential in $a^{1/p}$, and their proof was not a consequence of a large-deviation principle. It was the goal of [19] to get full control on the shape of $\ell/\ell(U)$ under $\mathbb{P}(\cdot \mid \ell(U) > a)$ in terms of asymptotics for integrals against many test functions, but the method used there (asymptotics for the k^{th} moments) did not produce a large-deviation principle; the technique precluded functions that assume negative values.

1.3 Main Results: Large Deviations

Our first main result is a large-deviation principle for large time for the motions before exiting the set B (defined as in Section 1.2). Assume that the p motions $W^{(1)}, \dots, W^{(p)}$ have some arbitrary starting distribution on B , possibly dependent on each other, which we suppress from the notation. Their occupation times

measures are denoted by

$$(1.6) \quad \ell_t^{(i)} = \int_0^t \delta_{W_s^{(i)}} ds, \quad i = 1, \dots, p, \quad t > 0.$$

We fix $b = (b_1, \dots, b_p) \in (0, \infty)^p$ and consider the time horizon $[0, tb_i]$ for the i^{th} motion. By

$$\mathbb{P}^{(tb)}(\cdot) = \mathbb{P}\left(\cdot \cap \bigcap_{i=1}^p \{tb_i < \tau_i\}\right)$$

we denote the subprobability measure under which the i^{th} motion does not exit B before time tb_i . Recall the heuristic definition of ℓ_{tb} from (1.1). Then ℓ_{tb} is a random element of the set $\mathcal{M}(B)$ of positive measures on B . We equip it with the weak topology induced by test integrals with respect to continuous bounded functions $B \rightarrow \mathbb{R}$. By $\mathcal{M}_1(B)$ we denote the set of probability measures on B , and by $H_0^1(B)$ the usual Sobolev space with zero boundary condition in B .

THEOREM 1.1 (LDP at Diverging Time). *The tuple*

$$\left(\frac{1}{t^p \prod_{i=1}^p b_i} \ell_{tb}; \frac{1}{tb_1} \ell_{tb_1}^{(1)}, \dots, \frac{1}{tb_p} \ell_{tb_p}^{(p)} \right)$$

satisfies, as $t \rightarrow \infty$, a large deviation principle in the space $\mathcal{M}(B) \times \mathcal{M}_1(B)^p$ under $\mathbb{P}^{(tb)}$ with speed t and rate function

$$(1.7) \quad I(\mu; \mu_1, \dots, \mu_p) = \frac{1}{2} \sum_{i=1}^p b_i \|\nabla \psi_i\|_2^2,$$

if μ, μ_1, \dots, μ_p each have densities ψ^{2p} and $\psi_1^2, \dots, \psi_p^2$ with $\|\psi_i\|_2 = 1$ for $i = 1, \dots, p$ such that $\psi, \psi_1, \dots, \psi_p \in H_0^1(B)$ and $\psi^{2p} = \prod_{i=1}^p \psi_i^2$; otherwise the rate function is ∞ . The level sets of the rate function I in (1.7) are compact.

To be more explicit in the special case $b = 1$, Theorem 1.1 says that, for any continuous and bounded test functions $f, f_1, \dots, f_p : B \rightarrow \mathbb{R}$,

$$(1.8) \quad \begin{aligned} & \lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}^{(t^1)} \left[\exp \left\{ t \left(\langle t^{-p} \ell_{t^1}, f \rangle + \sum_{i=1}^p \left\langle \frac{1}{t} \ell_t^{(i)}, f_i \right\rangle \right) \right\} \right] \\ &= \sup \left\{ \left\langle \prod_{i=1}^p \psi_i^2, f \right\rangle + \sum_{i=1}^p \langle \psi_i^2, f_i \rangle - \frac{1}{2} \sum_{i=1}^p \|\nabla \psi_i\|_2^2 : \right. \\ & \quad \left. \psi_i \in H_0^1(B) \text{ and } \|\psi_i\|_2 = 1 \text{ for } i = 1, \dots, p \right\}. \end{aligned}$$

Theorem 1.1 is an extension of the well-known Donsker-Varadhan LDP (see [6, 7, 8, 9] and [15]) for the occupation measure of a single Brownian motion in compacts to the intersection measure. It gives a rigorous meaning to the heuristic formula in (1.1) in the limit $t \rightarrow \infty$. Since B is bounded, ℓ_{tb} is a finite measure.

However, there is no natural normalization of ℓ_{tb} that turns it into a probability measure. Our result shows that $t^{-p}\ell_{tb}$ is asymptotically of finite order. A heuristic derivation of Theorem 1.1 in terms of the Donsker-Varadhan LDP is given in Section 1.4, and the proof in Sections 2 and 3.

Specializing to the first entry of the tuple, we get the following principle from the contraction principle [5, theorem 4.2.1]:

COROLLARY 1.2. *Fix $b = (b_1, \dots, b_p) \in (0, \infty)^p$. Then the family of measures*

$$\left(\left(t^p \prod_{i=1}^p b_i \right)^{-1} \ell_{tb} \right)_{t>0}$$

satisfies, as $t \rightarrow \infty$, a large-deviation principle in the space $\mathcal{M}(B)$ under $\mathbb{P}^{(tb)}$ with speed t and rate function

$$(1.9) \quad I(\mu) = \inf \left\{ \frac{1}{2} \sum_{i=1}^p b_i \|\nabla \psi_i\|_2^2 : \psi_i \in H_0^1(B), \|\psi_i\|_2 = 1 \forall i = 1, \dots, p, \right. \\ \left. \text{and } \prod_{i=1}^p \psi_i^2 = \frac{d\mu}{dx} \right\},$$

if μ has a density, and $I(\mu) = \infty$ otherwise. The level sets of the rate function I in (1.9) are compact.

To be more explicit in the special case $b = 1$, Corollary 1.2 says that, for any open set $G \subset \mathcal{M}(B)$ and every closed set $F \subset \mathcal{M}(B)$,

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}(t^{-p} \ell_t \in F, t < \tau_1 \wedge \dots \wedge \tau_p) \leq - \inf_{\mu \in F} I(\mu), \\ \liminf_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}(t^{-p} \ell_t \in G, t < \tau_1 \wedge \dots \wedge \tau_p) \geq - \inf_{\mu \in G} I(\mu).$$

In the special case $b = 1 = (1, \dots, 1)$, it is tempting to conjecture that, for (ψ_1, \dots, ψ_p) a minimizing tuple in (1.9), all the ψ_i should be identical. This would simplify the formula to $I(\mu) = \frac{p}{2} \|\nabla \psi\|_2^2$ if ψ^{2p} is a density of μ with $\psi \in H_0^1(B)$. However, we found no evidence for that and indeed conjecture that this is not true for general μ . But note that the result by Chen in (1.2)–(1.3), after replacing $\ell_t(\mathbb{R}^d)$ by $\ell_t(B)$ and $H^1(\mathbb{R}^d)$ by $H_0^1(B)$ for $a = 1$ suggests that, at least for the minimizer μ of $I(\mu)$, all the ψ_i should be identical, since the minimizer in (1.3) is just some ψ^{2p} .

As a corollary of Theorem 1.1, we now give a related LDP for the normalized intersection local time for the motions stopped at their first exit from B under conditioning on $\{\ell(U) > a\}$ as $a \rightarrow \infty$, where we recall that $U \subset B$ is a compact set whose boundary is a Lebesgue null set. This solves a problem left open in [19]; see Section 1.2. That is, instead of diverging deterministic time, we now consider a random time horizon and diverging ISLT. The measure $\ell/\ell(U)$ is a

positive measure on B , which is a probability measure on U . At the end of Section 1.2, we mentioned that the normalized probability measure $\ell/\ell(U)$ satisfies a law of large masses under the conditional law $\mathbb{P}(\cdot \mid \ell(U) > a)$. Here we in particular identify the precise rate of the exponential convergence. By $\mathcal{M}_U(B)$ we denote the set of positive finite measures on B whose restriction to U is a probability measure. Our second main result is the following:

THEOREM 1.3 (Large Deviations at Diverging Mass). *The normalized probability measures $\ell/\ell(U)$ under $\mathbb{P}(\cdot \mid \ell(U) > a)$ satisfy, as $a \rightarrow \infty$, a large-deviation principle in the space $\mathcal{M}_U(B)$, with speed $a^{1/p}$ and rate function $J - \Theta_B(U)$, where*

$$(1.10) \quad J(\mu) = \inf \left\{ \frac{1}{2} \sum_{i=1}^p \|\nabla \phi_i\|_2^2 : \phi_1, \dots, \phi_p \in H_0^1(B), \prod_{i=1}^p \phi_i^2 = \frac{d\mu}{dx} \right\},$$

if μ has a density and $J(\mu) = \infty$ otherwise, where $\Theta_B(U)$ is the number appearing in (1.5). The level sets of J are compact.

The proof of Theorem 1.3 is in Section 4; a heuristic derivation from Theorem 1.1 is in Section 1.4.

It follows from the compactness of the level sets and the lower semicontinuity that J possesses minimizers, and it follows from the LDP that $\min J = \Theta_B(U)$. Even more, the set of minimizers of J is equal to the set M of minimizers of the variational problem for $\Theta_B(U)$ in (1.5), where we consider the elements ϕ^{2p} of M as elements of $\mathcal{M}_U(B)$; see the discussion below (1.5). This is seen as follows: Let \tilde{M} be the set of minimizers of J . Then it is easily seen that $M \subset \tilde{M}$ by restricting the infimum in (1.10) to identical ϕ_1, \dots, ϕ_p . Suppose $\tilde{M} \setminus M \neq \emptyset$. Then for some $\mu \in \tilde{M} \setminus M$ with positive distance from M , we can find $\varepsilon > 0$ such that $B_\varepsilon(\mu)$, the ε -neighborhood of μ , is disjoint from M . Then it follows from the proof of [19, theorem 1.4] that

$$\limsup_{a \uparrow \infty} \frac{1}{a^{1/p}} \log \mathbb{P}(L \in B_\varepsilon(\mu) \mid \ell(U) > a) < 0.$$

But it follows from the LDP in Theorem 1.3 that the left-hand side is not smaller than $\Theta_B(U) - \inf_{\mu \in B_\varepsilon(\mu)} J(\mu)$, which implies that $J(\mu) - \Theta_B(U) > 0$. Hence, $\mu \notin \tilde{M}$. This is a contradiction. Hence \tilde{M} is equal to M . However, for fixed $\mu \in \mathcal{M}_U(B)$, it remains an open problem to give a sufficient condition on μ for having a minimizing tuple of p identical functions ϕ_1, \dots, ϕ_p in (1.10).

For Theorems 1.1 and 1.3 and Corollary 1.2, there are analogues for random walks on \mathbb{Z}^d instead of Brownian motions on \mathbb{R}^d . These are much easier to formulate and to prove since the heuristic formula in (1.1) can be taken as a definition without problems.

1.4 Heuristic Derivation of the Main Results

In this section we sketch heuristics that lead to Theorems 1.1 and 1.3, starting from the Donsker-Varadhan theory of large deviations. For simplicity, we drop compactness issues and formulate the principle on \mathbb{R}^d rather than on some bounded domain B . We also put $b = 1$ and write ℓ_t instead of ℓ_{t1} .

Recall the occupation measure of the i^{th} Brownian motion defined in (1.6). That is, $\ell_t^{(i)}(A)$ is the amount of time that $W^{(i)}$ spends in $A \subset \mathbb{R}^d$ by time t . The famous Donsker-Varadhan LDP (see [6, 8, 9] and [15]) states that

$$(1.11) \quad \mathbb{P}\left(\frac{1}{t} \ell_t^{(i)} \approx \mu\right) = \exp\left[-t \frac{1}{2} \left\| \nabla \sqrt{\frac{d\mu}{dx}} \right\|_2^2 + o(t)\right], \quad t \rightarrow \infty.$$

This is a simplified version of the statement that, under $\mathbb{P}(\cdot \cap \{W_{[0,t]}^{(i)} \subset B\})$, the distributions of $\frac{1}{t} \ell_t^{(i)}$ satisfy an LDP with speed t and rate function $\mu \mapsto \frac{1}{2} \left\| \nabla \sqrt{\frac{d\mu}{dx}} \right\|_2^2$ if the square root of the density of μ exists in $H^1(\mathbb{R}^d)$ and $\mu \mapsto \infty$ otherwise.

The heuristic formula in (1.1) states that

$$(1.12) \quad t^{-p} \frac{\ell_t(dy)}{dy} = \prod_{i=1}^p \frac{1}{t} \frac{\ell_t^{(i)}(dy)}{dy}.$$

Hence $t^{-p} \ell_t$ is a function of the tuple $(\frac{1}{t} \ell_t^{(1)}, \dots, \frac{1}{t} \ell_t^{(p)})$. Let us ignore that this map is far from continuous. Now the LDP in Theorem 1.1 follows from a formal application of the contraction principle.

Let us now give a heuristic derivation of the LDP in Theorem 1.3. The heuristic formula in (1.1) implies that

$$(1.13) \quad \frac{\ell(dy)}{\ell(U)} = \frac{1}{\int_U dx \prod_{i=1}^p \frac{\ell_{\tau_i}^{(i)}(dx)}{dx}} \left(\prod_{i=1}^p \frac{\ell_{\tau_i}^{(i)}(dy)}{dy} \right) dy.$$

Pick some $\mu \in \mathcal{M}_U(B)$ with density ϕ^{2p} . We make the ansatz that the event $\{\ell/\ell(U) \approx \mu, \ell(U) > a\}$ is realized by the event $\bigcap_{i=1}^p A(b_i, \psi_i)$, where

$$A(b_i, \psi_i) = \left\{ \tau_i > b_i a^{1/p}, \frac{1}{b_i a^{1/p}} \ell_{b_i a^{1/p}}^{(i)} \approx \psi_i^2(x) dx \text{ on } B \right\},$$

where $\psi_1, \dots, \psi_p \in H_0^1(B)$ are $L^2(B)$ -normalized and $b_1, \dots, b_p \in (0, \infty)$. Later we optimize over ψ_1, \dots, ψ_p and b_1, \dots, b_p . In other words, the i^{th} motion spends an amount of $\tau_i \approx b_i a^{1/p}$ time units in B until it leaves the set B , and its normalized occupation times measure resembles ψ_i^2 on B . We approximate $\ell(U) > a$ by $\ell(U) \approx a$ and therefore have the following condition for b_1, \dots, b_p :

$$(1.14) \quad 1 \approx \frac{1}{a} \ell(U) = \prod_{i=1}^p b_i \int_U dx \prod_{i=1}^p \psi_i^2(x).$$

Furthermore, from (1.13), we get the condition

$$(1.15) \quad \phi^{2p} = \frac{\ell}{\ell(U)} = \frac{\prod_{i=1}^p \psi_i^2}{\int_U dx \prod_{i=1}^p \psi_i^2(x)} = \prod_{i=1}^p (b_i \psi_i^2).$$

Hence we get, also using (1.11) with $t = b_i a^{1/p}$,

$$(1.16) \quad \begin{aligned} & \lim_{a \rightarrow \infty} a^{-1/p} \log \mathbb{P} \left(\frac{\ell}{\ell(U)} \approx \phi^{2p}, \ell(U) > a \right) \\ &= - \inf_{b_1, \dots, b_p, \psi_1, \dots, \psi_p} \lim_{a \rightarrow \infty} a^{-1/p} \log \mathbb{P} \left(\prod_{i=1}^p A(b_i, \psi_i) \right) \\ &= - \inf_{b_1, \dots, b_p, \psi_1, \dots, \psi_p} \sum_{i=1}^p b_i \frac{1}{2} \|\nabla \psi_i\|_2^2, \end{aligned}$$

where the infimum runs under the above-mentioned conditions, in particular (1.14) and (1.15). Now substituting $\phi_i^2 = b_i \psi_i^2$ for $i = 1, \dots, p$, we see that the right-hand side of (1.16) is indeed equal to $-J(\mu)$. This ends the heuristic derivation of Theorem 1.3.

2 Proof of Theorem 1.1: Large Deviations for Diverging Time

In this section, we prove our first main result, the LDP in Theorem 1.1. A summary of our proof is as follows. In Section 2.2 we introduce an approximation of the normalized intersection measure in terms of the pointwise product of smoothed versions of the normalized occupation times measures of the p motions and prove an LDP for the tuple built from them. This is quite easy, as this tuple is a continuous function of the normalized occupation times measures, for which we can apply the classical Donsker-Varadhan LDP. Furthermore, in Section 2.3 we show that the corresponding rate function converges to the rate function I of the LDP of Theorem 1.1 as the smoothing parameter vanishes. The convergence is in the sense of Γ -convergence, and its proof relies on standard analysis. In Section 2.4 we finish the proof of Theorem 1.1, subject to the fact that the smoothed versions of the intersection measure is indeed an exponentially good approximation of the (nonsmoothed) intersection measure. This fact is formulated as a proposition; its proof is deferred to Section 3. In the Section 2.1 we give some remarks on the relation to other proofs in this field in the literature.

2.1 Literature Remarks on the Proof

In the recent decades, with especially much success in this millennium, people have developed many techniques to derive the large-time or the large-mass asymptotics for the total mass of mutual intersections of several independent paths; we mentioned two important ones in Section 1.2. With the exception of the work in [19], these results concern only the total mass, but not integrals against test functions, as we consider in the present paper. Hence, the question arises which of

the existing proof strategies are also amenable to the refined problem about test integrals. In our setting of large deviations in a bounded set B , we do not have the (technically very nasty) additional problem of compactifying the space, which cannot be overcome by the well-known periodization technique, but was solved by Chen using an abstract compactness criterion by de Acosta. We are also not using the technique of comparing deterministic time t to random independent exponential time, as this works only in connection with the Brownian scaling property, which we cannot use for our refined problem.

The technique of finding the asymptotics of high polynomial moments and using them for the logarithmic asymptotics of probabilities was first carried out in [18] in the context of mutual Brownian intersection local times in a bounded set B ; see Section 1.2 and a thorough presentation in [4]. This has the advantage of avoiding a smoothing approximation; these are always technically involved. In [19], this technique was extended to the analysis of test integrals against a large class of measurable and bounded test functions. However, this technique was not able to yield an LDP, since it could only be applied to nonnegative test functions. Hence we believe that this technique will not be helpful for deriving LDPs.

Another possibility would be to use Le Gall's [20] approximation technique with the help of renormalized Lebesgue measure on the intersection of the Wiener sausages. The main task here would be to strengthen the L^p -convergence of test integrals of these measures to exponential convergence. However, we found no way to do this.

Chen developed a strategy of smoothing by convolution of the Dirac measure in the proof of [4, theorem 2.2.3] for finding the logarithmic asymptotics for the upper tails of the total mass of the intersection. However, the strategy of proving the exponentially good approximation was tailored there for the total mass and does not seem to be amenable to the study of test integrals against test functions that may take arbitrary, positive and negative, values.

On the other side, another technique developed in [3] seems to be amenable to prove an exponentially good approximation of the intersection measure for $p = 2$ using Fourier inversion. However, for $p > 2$, the mollifier used in [3] does not seem to admit an LDP, at least not without substantial work, and we did not see how.

Therefore, we chose to work with mollifying each occupation time and to approximate the intersection measure with their pointwise product, which itself is easily seen to satisfy an LDP. Our proof of the exponential approximation in Section 3 with this object requires combinatorial and analytical work.

2.2 Large Deviations for Smoothed Intersection Local Times

Recall from (1.6) the occupation measure $\ell_t^{(i)}$ of the i^{th} motion. Let $\varphi = \varphi_1$ be a nonnegative, C^∞ -function on \mathbb{R}^d with compact support, normalized such that

$\int_{\mathbb{R}^d} \varphi_1(y) dy = 1$. Now we define the approximation of the Dirac δ -function at 0 by

$$\varphi_\varepsilon(x) = \varepsilon^{-d} \varphi_1(x/\varepsilon).$$

Let us consider the convolution of the above occupation measures with φ_ε :

$$\ell_{\varepsilon,t}^{(i)}(y) = \varphi_\varepsilon \star \ell_t^{(i)}(y) = \int_0^t ds \varphi_\varepsilon(W_s^{(i)} - y).$$

Then $\ell_{\varepsilon,t}^{(i)}$ is a bounded C^∞ -function. As $\varepsilon \downarrow 0$, the measure with density $\ell_{\varepsilon,t}^{(i)}$ converges weakly towards the occupation measure $\ell_t^{(i)}$.

Consider the pointwise product of the above densities,

$$\ell_{\varepsilon,t}(y) = \prod_{i=1}^p \ell_{\varepsilon,t}^{(i)}(y).$$

We will write $\ell_{\varepsilon,t}(y)dy$ for the measure with density $\ell_{\varepsilon,t}$. It should come as no surprise that these measures are, for any fixed t , an approximation of the intersection local time ℓ_t as $\varepsilon \downarrow 0$, even though we could not find this statement in the literature. Actually, we will go much further and show that they are even an exponentially good approximation of the intersection local time ℓ_t in the sense of [5]; see below.

First we state a large-deviation principle for the measures with density $\ell_{\varepsilon,t}$ as $t \rightarrow \infty$ for fixed $\varepsilon > 0$. It is known by classical work by Donsker and Varadhan [6, 8, 9, 15] that each $\frac{1}{t} \ell_t^{(i)}$ satisfies, as $t \rightarrow \infty$, a large-deviation principle. In the proof of Lemma 2.1 below we will see that $\ell_{\varepsilon,t}(y)dy$ is a continuous functional of the tuple $(\ell_t^{(1)}, \dots, \ell_t^{(p)})$. Hence, by the contraction principle, $\ell_{\varepsilon,t}(y)dy$ itself satisfies an LDP with some (ε -dependent) rate function.

Recall that we equip $\mathcal{M}(\mathbb{R}^d)$, the space of finite measures on \mathbb{R}^d , with the weak topology induced by test integrals against continuous bounded functions. For a measure $\mu \in \mathcal{M}(\mathbb{R}^d)$ and a function $f : \mathbb{R}^d \rightarrow \mathbb{R}$, we denote by $\langle \mu, f \rangle$ the integral $\int f d\mu$.

LEMMA 2.1 (LDP for Smoothed Measures). *Fix $\varepsilon > 0$ and $b = (b_1, \dots, b_p) \in (0, \infty)^p$. Then the tuple of random measures*

$$\left(\frac{1}{t^p \prod_{i=1}^p b_i} \ell_{\varepsilon,tb}; \frac{1}{tb_1} \ell_{\varepsilon,tb_1}^{(1)}, \dots, \frac{1}{tb_p} \ell_{\varepsilon,tb_p}^{(p)} \right)$$

satisfies, as $t \rightarrow \infty$, a large-deviation principle in $\mathcal{M}(B) \times \mathcal{M}_1(B)^p$ under $\mathbb{P}^{(tb)}$ with speed t and rate function

$$(2.1) \quad \begin{aligned} & I_\varepsilon(\mu; \mu_1, \dots, \mu_p) \\ &= \inf \left\{ \frac{1}{2} \sum_{i=1}^p b_i \|\nabla \psi_i\|_2^2 : \psi_i \in H_0^1(B), \|\psi_i\|_2 = 1, \psi_i^2 \star \varphi_\varepsilon = \frac{d\mu_i}{dx} \right. \\ & \quad \left. \forall i = 1, \dots, p \text{ and } \prod_{i=1}^p \psi_i^2 \star \varphi_\varepsilon = \frac{d\mu}{dx} \right\}, \end{aligned}$$

if μ has a density, and $I_\varepsilon(\mu) = \infty$ otherwise. The level sets of I_ε are compact.

PROOF. First observe that the mapping

$$(2.2) \quad (\mathcal{M}_1(\mathbb{R}^d))^p \longrightarrow \mathcal{M}(\mathbb{R}^d), \quad (\mu_1, \dots, \mu_p) \mapsto \left(\prod_{i=1}^p \mu_i \star \varphi_\varepsilon(x) \right) dx,$$

is weakly continuous. Indeed, first note that the map $(\mu_1, \dots, \mu_p) \mapsto \mu_1 \otimes \dots \otimes \mu_p$ is continuous from $\mathcal{M}_1(\mathbb{R}^d)^p$ to $\mathcal{M}_1((\mathbb{R}^d)^p)$ since $\mathcal{M}_1(\mathbb{R}^d)$ is a Polish space. Furthermore, for every continuous bounded test function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ and any $\mu_1, \dots, \mu_p \in \mathcal{M}_1(\mathbb{R}^d)$, we have

$$\begin{aligned} \left\langle f, \left(\prod_{i=1}^p \mu_i \star \varphi_\varepsilon(x) \right) dx \right\rangle &= \int_{\mathbb{R}^d} dx f(x) \int_{(\mathbb{R}^d)^p} \prod_{i=1}^p (\mu_i(dy_i) \varphi_\varepsilon(x - y_i)) \\ &= \langle A_f, \mu_1 \otimes \dots \otimes \mu_p \rangle, \end{aligned}$$

where

$$A_f(y_1, \dots, y_p) = \int_{\mathbb{R}^d} dx f(x) \varphi_\varepsilon(x - y_1) \cdots \varphi_\varepsilon(x - y_p).$$

As φ_ε is smooth and compactly supported in \mathbb{R}^d , the function A_f is continuous and bounded in $(\mathbb{R}^d)^p$. This shows the continuity of the map in (2.2). Now the claimed LDP follows from the contraction principle [5, theorem 4.2.1]. \square

2.3 Gamma-Convergence of the Rate Function

In this section, we pass to the limit $\varepsilon \downarrow 0$ in the variational formula (2.1). The sense of convergence is the Γ -convergence, as will be required in the proof of Theorem 1.1. The proof of this convergence is based on standard analytic tools. By $B_\delta(\mu) = \{v \in \mathcal{M}(B) : d(v, \mu) < \delta\}$ we denote the open ball of radius δ around μ , where d is a metric that induces the weak topology in $\mathcal{M}(B)$. By d we also denote the product metric on $\mathcal{M}(B) \times \mathcal{M}_1(B)^p$, and by $B_\delta(\mu; \mu_1, \dots, \mu_p)$ the open δ -ball around $(\mu, \mu_1, \dots, \mu_p)$ in this space.

PROPOSITION 2.2. *For every $\mu \in \mathcal{M}(B)$, we have*

$$(2.3) \quad \sup_{\delta > 0} \liminf_{\varepsilon \downarrow 0} \inf_{B_\delta(\mu; \mu_1, \dots, \mu_p)} I_\varepsilon = I(\mu; \mu_1, \dots, \mu_p),$$

where I is the rate function defined in (1.7). Furthermore, the level sets of I are compact.

PROOF. We write $f(x)\mu(dx)$ for the measure with density f with respect to μ . We denote the Lebesgue measure by dx .

First we prove ‘ \leq ’. Let μ, μ_1, \dots, μ_p be given. Without loss of generality, we may assume that $\psi_i^2 = \frac{d\mu_i}{dx}$ exists, and $\frac{d\mu}{dx} = \prod_{i=1}^p \psi_i^2$. Fix $\delta > 0$ and take $\varepsilon > 0$ so small that $\psi_i^2 \star \varphi_\varepsilon(x)dx \in B_{\delta/2p}(\mu_i)$ for $i = 1, \dots, p$ and $(\prod_{i=1}^p \psi_i^2 \star \varphi_\varepsilon(x))dx \in B_{\delta/2p}(\mu)$. Hence the tuple

$$\left(\left(\prod_{i=1}^p \psi_i^2 \star \varphi_\varepsilon(x) \right) dx; \psi_1^2 \star \varphi_\varepsilon(x) dx, \dots, \psi_p^2 \star \varphi_\varepsilon(x) dx \right)$$

lies in $B_\delta(\mu; \mu_1, \dots, \mu_p)$. Hence

$$\begin{aligned} \inf_{B_\delta(\mu; \mu_1, \dots, \mu_p)} I_\varepsilon &\leq I_\varepsilon \left(\left(\prod_{i=1}^p \psi_i^2 \star \varphi_\varepsilon(x) \right) dx; \psi_1^2 \star \varphi_\varepsilon(x) dx, \dots, \psi_p^2 \star \varphi_\varepsilon(x) dx \right) \\ &\leq \frac{1}{2} \sum_{i=1}^p \|\nabla \psi_i\|_2^2, \end{aligned}$$

where in the last step we used the definition of I_ε .

Now we prove ‘ \geq ’. Let μ, μ_1, \dots, μ_p be given and let $I(\mu; \mu_1, \dots, \mu_p)$ be finite. Without loss of generality, the left-hand side of (2.3) is also finite. For $\delta, \varepsilon > 0$, we pick $(\mu^{(\delta, \varepsilon)}, \mu_1^{(\delta, \varepsilon)}, \dots, \mu_p^{(\delta, \varepsilon)})$ in $B_\delta(\mu; \mu_1, \dots, \mu_p)$ such that

$$\inf_{B_\delta(\mu; \mu_1, \dots, \mu_p)} I_\varepsilon \geq I_\varepsilon(\mu^{(\delta, \varepsilon)}; \mu_1^{(\delta, \varepsilon)}, \dots, \mu_p^{(\delta, \varepsilon)}) - \delta.$$

By definition of I_ε , there are L^2 -normalized $\psi_i^{(\delta, \varepsilon)} \in H_0^1(B)$ for $i = 1, \dots, p$ such that $\mu_i^{(\delta, \varepsilon)}(dx) = \psi_i^2 \star \varphi_\varepsilon(x)dx$ and $\mu^{(\delta, \varepsilon)}(dx) = (\prod_{i=1}^p \psi_i^2 \star \varphi_\varepsilon(x))dx$ and

$$I_\varepsilon(\mu^{(\delta, \varepsilon)}; \mu_1^{(\delta, \varepsilon)}, \dots, \mu_p^{(\delta, \varepsilon)}) \geq \frac{1}{2} \sum_{i=1}^p \|\nabla \psi_i^{(\delta, \varepsilon)}\|_2^2 - \varepsilon.$$

Then, by well-known analysis [23, chap. 8], along some subsequences, we may assume that $\psi_i^{(\delta, \varepsilon)} \rightarrow \psi_i^{(\delta)}$ as $\varepsilon \downarrow 0$ for some L^2 -normalized $\psi_i^{(\delta)} \in H_0^1(B)$ for $i = 1, \dots, p$ such that $\|\nabla \psi_i^{(\delta)}\|_2^2 \leq \liminf_{\varepsilon \downarrow 0} \|\nabla \psi_i^{(\delta, \varepsilon)}\|_2^2$. This convergence is true strongly in L^q for any $q > 1$ in $d = 2$ and $1 < q < 6$ in $d = 3$, and we have

$$(2.4) \quad \liminf_{\varepsilon \downarrow 0} \inf_{B_\delta(\mu; \mu_1, \dots, \mu_p)} I_\varepsilon \geq \frac{1}{2} \sum_{i=1}^p \|\nabla \psi_i^{(\delta)}\|_2^2 - \delta.$$

In particular, we have $\mu_i^{(\delta, \varepsilon)} \Rightarrow \psi_i^{(\delta)}(x)^2 dx =: \mu_i^{(\delta)}(dx)$ in the weak topology. It is elementary (using Hölder's inequality) to see that $(\psi_i^{(\delta, \varepsilon)})^2 \star \varphi_\varepsilon(x) dx \Rightarrow \mu_i^{(\delta)}(dx)$ in the weak topology. Hence, $\mu_i^{(\delta)} \in B_{\delta/2p}(\mu_i)$. Now we let $\delta \downarrow 0$ and take a subsequence of $\psi_i^{(\delta)}$ that converges to some ψ_i strongly in L^q for any $q > 1$ in $d = 2$ and $1 < q < 6$ in $d = 3$ and

$$\liminf_{\delta \downarrow 0} \sum_{i=1}^p \|\nabla \psi_i^{(\delta)}\|_2^2 \geq \sum_{i=1}^p \|\nabla \psi_i\|_2^2.$$

Since $\mu_i^{(\delta)} \in B_{\delta/2p}(\mu_i)$, ψ_i^2 must be a density of μ_i . Therefore, the right-hand side of the last display is $2I(\mu; \mu_1, \dots, \mu_p)$. Sending $\delta \downarrow 0$ in (2.4), the proof is finished for the case when $I(\mu; \mu_1, \dots, \mu_p)$ is finite.

Now we consider the case $I(\mu; \mu_1, \dots, \mu_p) = \infty$. First, we consider the case that all μ_1, \dots, μ_p have densities $\psi_1^2, \dots, \psi_p^2$ such that $\psi_i \in H_0^1(B)$, but μ fails either to have a density or to be the pointwise product of the ψ_i^2 . By way of contradiction, assume that the left-hand side of (2.3) is finite. Now we follow the same line of argument as above and define $\mu^{(\delta)} = (\prod_{i=1}^p (\psi_i^{(\delta)})^2(x)) dx$ and note that $\mu^{(\delta, \varepsilon)} \Rightarrow \mu^{(\delta)}$ as $\varepsilon \downarrow 0$. Indeed $\psi_i^{(\delta, \varepsilon)}$ converges as $\varepsilon \downarrow 0$ (strongly in L^q for $q > 1$ in $d = 2$ and $1 < q < 6$ in $d = 3$) to $\psi_i^{(\delta)}$, and taking the pointwise product of the densities is a weakly continuous operation. Hence $\mu^{(\delta)}$ lies in $B_{\delta/2p}(\mu)$. Now we send $\delta \downarrow 0$ and use the same argument to infer that $\mu^{(\delta)} \Rightarrow \mu = (\prod_{i=1}^p \psi_i^2(x)) dx$. This is a contradiction.

Furthermore, also in the case that one of the μ_i 's does not have a density or its square root is not in $H_0^1(B)$, the same arguments above (by contradiction) show

$$\liminf_{\delta \downarrow 0} \sum_{i=1}^p \|\nabla \psi_i^{(\delta)}\|_2^2 \geq +\infty = I(\mu; \mu_1, \dots, \mu_p). \quad \square$$

2.4 Completion of Proof of Theorem 1.1

The main step in the remaining part of the proof of Theorem 1.1 is to show that the intersection measure $t^{-p} \ell_{tb}$ is exponentially well approximated by $t^{-p} \ell_{\varepsilon, tb}$. This we formulate here as a result of its own interest.

PROPOSITION 2.3 (Exponential Approximation). *Fix $b = (b_1, \dots, b_p) \in (0, \infty)^p$ and a measurable and bounded function $f : B \rightarrow \mathbb{R}$. Then, for any $\varepsilon > 0$, there is $C(\varepsilon) > 0$ such that*

$$(2.5) \quad \mathbb{E}^{(tb)} [|\langle \ell_{tb} - \ell_{\varepsilon, tb}, f \rangle|^k] \leq (k!)^p C(\varepsilon)^k, \quad t \in (0, \infty), k \in \mathbb{N},$$

and $\lim_{\varepsilon \downarrow 0} C(\varepsilon) = 0$.

Note that this result implicitly shows that ℓ_t is indeed approximated by $\ell_{\varepsilon,t}$ in the L^k -topology for any k , as we announced in Section 1.1. The proof of Proposition 2.3 is given in Section 3. Now we finish the proof of our main result.

PROOF OF THEOREM 1.1. Recall that we have an LDP for the ε -depending tuple in Lemma 2.1. We now use Proposition 2.3 to see that this tuple is an exponentially good approximation of the tuple in Theorem 1.1. Recall that d is a metric on $\mathcal{M}(B)$ that induces the weak topology. We also denote by d a metric on $\mathcal{M}(B) \times \mathcal{M}_1(B)^p$ that induces the product topology of this topology. Then we have to show that the probability that the d -distance of the two tuples in Lemma 2.1 and Theorem 1.1 being larger than any $\delta > 0$ has an exponential rate as $t \rightarrow \infty$ that tends to $-\infty$ as $\varepsilon \downarrow 0$. Since the topology on $\mathcal{M}(B)$ is induced by test integrals against continuous bounded functions, it is enough to show that, for any such test functions $f, f_1, \dots, f_p : B \rightarrow \mathbb{R}$,

$$\lim_{\varepsilon \downarrow 0} \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}^{(tb)} \left(\left\{ \left| \left\langle \frac{1}{t^p \prod_{i=1}^p b_i} (\ell_{tb} - \ell_{\varepsilon,tb}), f \right\rangle \right| > \delta \right\} \cup \bigcup_{i=1}^p \left\{ \left| \left\langle \frac{1}{t b_i} (\ell_{tb_i}^{(i)} - \ell_{\varepsilon,tb_i}^{(i)}), f_i \right\rangle \right| > \delta \right\} \right) = -\infty.$$

This indeed follows from Proposition 2.3, together with a version of this for $p = 1$, which is indeed much simpler and also follows from [1, lemma 3.1], for example. Indeed, we have from Proposition 2.3 that

$$(2.6) \quad \lim_{\varepsilon \downarrow 0} \limsup_{t \uparrow \infty} \frac{1}{t} \log \mathbb{P}^{(tb)} \left(\left\{ \left| \left\langle \frac{1}{t^p \prod_{i=1}^p b_i} (\ell_{tb} - \ell_{\varepsilon,tb}), f \right\rangle \right| > \delta \right\} \right) = -\infty,$$

which follows from the Markov inequality applied to the function $x \mapsto x^k$ with $k = \lceil t \rceil$ as follows:

$$\begin{aligned} & \mathbb{P}^{(tb)} \left(\left\{ \left| \left\langle \frac{1}{t^p \prod_{i=1}^p b_i} (\ell_{tb} - \ell_{\varepsilon,tb}), f \right\rangle \right| > \delta \right\} \right) \\ & \leq \delta^{-k} t^{-pk} C^k \mathbb{E}^{(tb)} [|\langle \ell_{tb} - \ell_{\varepsilon,tb}, f \rangle|^k] \\ & \leq \delta^{-k} t^{-pk} C^k (k!)^p C(\varepsilon)^k \leq \tilde{C}(\varepsilon)^t, \end{aligned}$$

for any $t > 0$, where C , $C(\varepsilon)$, and $\tilde{C}(\varepsilon)$ depend on b , B , d , f , and δ (but not on t) and satisfy $\lim_{\varepsilon \downarrow 0} C(\varepsilon) = 0 = \lim_{\varepsilon \downarrow 0} \tilde{C}(\varepsilon)$, and $C(\varepsilon)$ is the constant from Proposition 2.3. Since $k = \lceil t \rceil$ and $\lim_{\varepsilon \downarrow 0} \tilde{C}(\varepsilon) = 0$, (2.6) follows.

Hence, according to [5, theorem 4.2.16], the LDP of Theorem 1.1 is true with the rate function on the left-hand side of (2.3). But Proposition 2.2 identifies this as I given in (1.9).

Note that by (2.3) and [5, theorem 4.2.16], I is a lower-semicontinuous functional. Hence its level sets are closed in $\mathcal{M}(B) \times \mathcal{M}_1(B)^p$. Since the infimum in (1.7) extends only over functions in $H_0^1(B)$ (i.e., with zero boundary conditions),

I can also be seen as a lower-semicontinuous functional on $\mathcal{M}(\overline{B}) \times \mathcal{M}_1(\overline{B})^p$, which is weakly compact by Prohorov's theorem. Hence the levels sets of I are also compact. That is, the proof of Theorem 1.1 is finished. \square

3 Proof of Proposition 2.3: Exponential Approximation

We turn to the proof of Proposition 2.3. We will do this only for $b = 1$ and write $\mathbb{E}^{(t)}$ instead of $\mathbb{E}^{(t^1)}$, etc. Fix a measurable bounded function f on B . Then our task is to prove that, for any $\varepsilon > 0$,

$$(3.1) \quad |\mathbb{E}^{(t)}[(\langle \ell_t, f \rangle - \langle \ell_{\varepsilon,t}, f \rangle)^k]| \leq k!^p C(\varepsilon)^k, \quad t \in (0, \infty), k \in \mathbb{N},$$

and $\lim_{\varepsilon \downarrow 0} C(\varepsilon) = 0$.

Note that we have now the absolute value signs outside the expectation, in contrast to (2.5). This is sufficient for proving (2.5), since, for k even, we can drop the absolute value signs anyway, and for k odd, we use Jensen's inequality to go from the power k to $k + 1$ and use that $((k + 1)!^p)^{k/(k+1)} \leq k!^p C^k$ for some $C \in (0, \infty)$ and all $k \in \mathbb{N}$.

Our proof of (3.1) is bulky and also technical; we divide it into several steps. In Section 3.1 we present a formula for the moments of integrals against $\ell_t - \ell_{\varepsilon,t}$ in terms of k -step transition densities, some of which are convolved. In Section 3.2 we present a heuristic proof for the regime $k \ll t$, which is meant to be a guiding philosophy that leads the actual proof strategy, though we do not use this section later. The second main tool of our proof, a standard expansion of the transition density in terms of eigenfunctions and eigenvalues of $-\frac{1}{2}\Delta$, is employed in Section 3.3. The latent ε presence also manifests here as some of the eigenfunctions are convolved (and the rest remain ε -free). Furthermore, we also estimate away some contributions (popping up from some singularities) to the main term. These are relatively easy to handle. The main term is attacked in Section 3.4, where we use an intricate counting technique that finally makes it possible to trace back our way using the binomial theorem and to extract the k^{th} power of some term that is small if ε is small.

3.1 Moment Formula

We begin with a moment formula for the left-hand side of (3.1), which is an adaptation of Le Gall's formula for the moments of $\ell(U)$ for compact subsets U of B [20, 21, 22].

Let us write $\mathbb{P}_{x,y}^{(t)}$ and $\mathbb{E}_{x,y}^{(t)}$ for the Brownian bridge subprobability measure

$$\bigotimes_{l=1}^p \mathbb{P}_{x^{(l)}}(\cdot, t < \tau; W_t \in dy^{(l)})/dy^{(l)}$$

(where $x = (x^{(1)}, \dots, x^{(p)})$, $y = (y^{(1)}, \dots, y^{(p)}) \in B^p$) and the corresponding expectation. In other words, under $\mathbb{P}_{x,y}^{(t)}$, we consider p independent Brownian bridges in B with time interval $[0, t]$ from $x^{(l)}$ to $y^{(l)}$ for $l = 1, \dots, p$. Later we

integrate over $x, y \in B^p$ with respect to $\nu(dx)dy$, where ν is the joint starting distribution of the p motions and hence $\mathbb{P}^{(t)} = \int_{B^p} \nu(dx) \int_{B^p} dy \mathbb{P}_{x,y}^{(t)}$.

Furthermore, we denote by $p_s^{(B)}(x, y) = \mathbb{P}_x(W_s \in dy; \tau > s)/dy$ the density of the distribution of a single Brownian motion at time s before the exit time τ from B when started at $x \in B$. By \mathfrak{S}_k we denote the set of permutations of $1, \dots, k$.

LEMMA 3.1 (Moment Formula). *For any continuous function $f : B \rightarrow \mathbb{R}$, any $k \in \mathbb{N}$, any $t > 0$, and any $x_0 = (x_0^{(1)}, \dots, x_0^{(p)})$ and $x_{k+1} = (x_{k+1}^{(1)}, \dots, x_{k+1}^{(p)}) \in B^p$,*

$$\begin{aligned}
& \mathbb{E}_{x_0, x_{k+1}}^{(t)} [(\langle f, \ell_t \rangle - \langle f, \ell_{\varepsilon, t} \rangle)^k] \\
&= \sum_{m=0}^k (-1)^m \binom{k}{m} \int_{B^k} \prod_{l=1}^k (f(y_l) dy_l) \\
& \quad \times \prod_{i=1}^p \left[\sum_{\sigma \in \mathfrak{S}_k[0, t]^k} \int dr_k \dots dr_1 \mathbf{1}_{\{\sum_{i=1}^k r_i \leq t\}} \right. \\
& \quad \quad \int_{B^{k-m}} \prod_{j=m+1}^k (\varphi_\varepsilon(y_j - z_j) dz_j) \\
& \quad \quad \left. \times \prod_{j=1}^{k+1} p_{r_j}^{(B)}(x_{j-1}^{(i)}, x_j^{(i)}) \right], \tag{3.2}
\end{aligned}$$

where we abbreviate $r_{k+1} = t - \sum_{i=1}^k r_i$ and, for $j = 1, \dots, k$,

$$x_j = x_j^{(i)} = \begin{cases} y_{\sigma^{-1}(j)} & \text{if } \sigma^{-1}(j) \leq m, \\ z_{\sigma^{-1}(j)} & \text{if } \sigma^{-1}(j) > m. \end{cases} \tag{3.3}$$

PROOF. We use the binomial theorem to split the k^{th} moment as follows:

$$\begin{aligned}
(3.4) \quad \mathbb{E}_{x_0, x_{k+1}}^{(t)} [(\langle f, \ell_t \rangle - \langle f, \ell_{\varepsilon, t} \rangle)^k] &= \\
& \sum_{m=0}^k (-1)^m \binom{k}{m} \mathbb{E}_{x_0, x_{k+1}}^{(t)} [\langle f, \ell_t \rangle^m \langle f, \ell_{\varepsilon, t} \rangle^{k-m}].
\end{aligned}$$

Now we handle the mixed moments above. We formulate the proof in a somewhat loose way, a mathematically correct way is described in [20]. For any $m \in \{0, \dots, k\}$,

$$(3.5) \quad \mathbb{E}_{x_0, x_{k+1}}^{(t)} [(f, \ell_t)^m (f, \ell_{\varepsilon, t})^{k-m}] = \int_{B^k} \prod_{l=1}^k f(y_l) \mathbb{E}_{x_0, x_{k+1}}^{(t)} \left[\bigotimes_{j=1}^m \ell_t(dy_j) \bigotimes_{j=m+1}^k \ell_{\varepsilon, t}(y_j) dy_j \right],$$

where we recall that ℓ_t does not have a density, but $\ell_{\varepsilon, t}$ is a smooth function. By definition of $\ell_{\varepsilon, t}$ and independence of paths, the expectation on the right-hand side of (3.5) can be written as

$$\prod_{i=1}^p \left[\int_{[0, t]^k} ds_k \cdots ds_1 \int_{B^{k-m}} \prod_{j=m+1}^k (\varphi_\varepsilon(y_j - z_j)) \times \mathbb{P}_{x_0^{(i)}, x_{k+1}^{(i)}}^{(t)} \left(\begin{cases} W_{s_j} \in dy_j & \text{if } j \leq m \\ W_{s_j} \in dz_j & \text{if } j > m. \end{cases} \right) \right],$$

where we remark that the integral over B^{k-m} refers to $dz_{m+1} \cdots dz_k$. Now we time-order the k -dimensional cube $[0, t]^k$ and write the last expression as

$$(3.6) \quad \prod_{i=1}^p \left[\sum_{\sigma \in \mathfrak{S}_k} \int_{0 \leq s_1 \leq \dots \leq s_k \leq t} ds_k \cdots ds_1 \int_{B^{k-m}} \prod_{j=m+1}^k (\varphi_\varepsilon(y_j - z_j)) \times \mathbb{P}_{x_0^{(i)}, x_{k+1}^{(i)}}^{(t)} \left(\begin{cases} W_{s_{\sigma(j)}} \in dy_j & \text{if } j \leq m \\ W_{s_{\sigma(j)}} \in dz_j & \text{if } j > m \end{cases} \right) \right].$$

The time-ordering allows us to invoke the Markov property at the consecutive times $s_1 < \dots < s_k$ and to split the path into k pieces. Each of the pieces is a Brownian motion before leaving B . Therefore the joint probability distribution above also splits into the corresponding k -step transition probability densities:

$$(3.7) \quad \begin{aligned} & \mathbb{P}_{x_0^{(i)}, x_{k+1}^{(i)}}^{(t)} \left(\begin{cases} W_{s_{\sigma(j)}} \in dy_j & \text{if } j \leq m \\ W_{s_{\sigma(j)}} \in dz_j & \text{if } j > m \end{cases} \right) \\ &= \mathbb{P}_{x_0^{(i)}, x_{k+1}^{(i)}}^{(t)} \left(\begin{cases} W_{s_j} \in dy_{\sigma^{-1}(j)} & \text{if } \sigma^{-1}(j) \leq m \\ W_{s_j} \in dz_{\sigma^{-1}(j)} & \text{if } \sigma^{-1}(j) > m \end{cases} \right) \\ &= \mathbb{P}_{x_0^{(i)}, x_{k+1}^{(i)}}^{(t)} (W_{s_j} \in dx_j^{(i)}, j = 1, \dots, k) \\ &= \left(\prod_{j=1}^{k+1} p_{s_j - s_{j-1}}^{(B)}(x_{j-1}^{(i)}, x_j^{(i)}) \right) dy_1 \cdots dy_m dz_{m+1} \cdots dz_k. \end{aligned}$$

Substituting $r_j = s_j - s_{j-1}$ and putting all the material together proves the lemma. \square

3.2 A Heuristic Proof for $k \ll t$

In order to give some guidance to the reader, let us briefly describe heuristically in which way we will succeed to estimate the bulky expression on the right of (3.2) in terms of $(k!)^p C(\varepsilon)^k$ with a small $C(\varepsilon)$. We do this only for the regime $k \ll t$, which we actually do not consider in Proposition 2.3, but this is only meant as a demonstration of the philosophy of our proof. Apart from the formulation of Lemma 3.2 below, the material of this section will not be used later in the proof of Proposition 2.3.

The problem is to extract an extinction coming from a difference of two close (for small ε) terms with a power of order k by use of the binomial theorem. Since this works only if certain powers of these close terms appear, one has to expand the probability terms on the right of (3.2) into sums of powers.

Our second main ingredient is a standard eigenvalue expansion with respect to the spectrum of the Laplace operator in B with zero boundary condition, which follows from the well-known spectral theorem for compact, self-adjoint operators [2, theorem 4.13]:

LEMMA 3.2 (Eigenvalue Expansion). *There exist eigenvalues $0 < \lambda_1 \leq \lambda_2 \leq \dots$ and an $L^2(B)$ -orthonormal basis of corresponding eigenfunctions ψ_1, ψ_2, \dots in B of $-\frac{1}{2}\Delta$ with zero boundary condition in B ; that is, $-\frac{1}{2}\Delta\psi_n = \lambda_n\psi_n$ for any $n \in \mathbb{N}$. Furthermore,*

$$(3.8) \quad p_s^{(B)}(x, y) = \sum_{n=1}^{\infty} e^{-s\lambda_n} \psi_n(x) \psi_n(y), \quad s > 0,$$

and the convergence is absolute and uniform in $x, y \in B$.

In the regime $k \ll t$, we use that r_j is large for any j and use the approximation

$$(3.9) \quad p_r^{(B)}(x, y) = e^{-r\lambda_1} (\psi_1(x)\psi_1(y) + o(1)), \quad r \rightarrow \infty.$$

That is, instead of plugging in the full eigenvalue expansion (3.8), we just pick the leading term of the expansion (3.9) in the last line of (3.2). This gives, for any $i = 1, \dots, p$,

$$(3.10) \quad \prod_{j=1}^{k+1} p_{r_j}^{(B)}(x_{j-1}^{(i)}, x_j^{(i)}) \\ \approx \prod_{j=1}^{k+1} (e^{-r_j\lambda_1} \psi_1(x_{j-1}^{(i)}) \psi_1(x_j^{(i)})) =$$

$$\begin{aligned}
&= e^{-t\lambda_1} \psi_1(x_0^{(i)}) \psi_1(x_{k+1}^{(i)}) \prod_{j=1}^k \psi_1^2(x_j) \\
&= e^{-t\lambda_1} \psi_1(x_0^{(i)}) \psi_1(x_{k+1}^{(i)}) \left(\prod_{j=1}^m \psi_1^2(y_j) \right) \left(\prod_{j=m+1}^k \psi_1^2(z_j) \right).
\end{aligned}$$

Note that the last term does not depend on $\sigma \in \mathfrak{S}_k$ or any $r_1, \dots, r_k \in [0, t]$. Also note that $|\mathfrak{S}_k| = k!$ and $\int_{[0,t]^k} dr_1 \cdots dr_k \mathbf{1}_{\{\sum_{i=1}^k r_i \leq t\}} = t^k/k!$. Substituting the last term of (3.10) in (3.2), we can integrate out the convolution integrals over z_{m+1}, \dots, z_k and afterwards the integrals over y_1, \dots, y_k and see that

$$\begin{aligned}
&\mathbb{E}_{x_0, x_{k+1}}^{(i)} [(\langle f, \ell_t \rangle - \langle f, \ell_{\varepsilon, t} \rangle)^k] \\
&\approx e^{-tp\lambda_1} t^{kp} \left(\prod_{i=1}^p \psi_1(x_0^{(i)}) \psi_1(x_{k+1}^{(i)}) \right) \sum_{m=0}^k (-1)^m \binom{k}{m} \\
&\quad \times \int_{B^k} dy_1 \cdots dy_k \left(\prod_{j=1}^k f(y_j) \right) \left(\prod_{j=1}^m \psi_1^{2p}(y_j) \right) \left(\prod_{j=m+1}^k (\varphi_\varepsilon \star \psi_1^2)^p(y_j) \right) \\
(3.11) \quad &= e^{-tp\lambda_1} t^{kp} \left(\prod_{i=1}^p \psi_1(x_0^{(i)}) \psi_1(x_{k+1}^{(i)}) \right) \\
&\quad \times \sum_{m=0}^k (-1)^m \binom{k}{m} \langle f, \psi_1^{2p} \rangle^m \langle f, (\varphi_\varepsilon \star \psi_1^2)^p \rangle^{k-m} \\
&= e^{-tp\lambda_1} t^{kp} \left(\prod_{i=1}^p \psi_1(x_0^{(i)}) \psi_1(x_{k+1}^{(i)}) \right) (\langle f, \psi_1^{2p} \rangle - \langle f, (\varphi_\varepsilon \star \psi_1^2)^p \rangle)^k,
\end{aligned}$$

according to the binomial theorem. Since φ_ε is an approximation of the Dirac delta measure at 0, it is clear that $\langle f, \psi_1^{2p} \rangle - \langle f, (\varphi_\varepsilon \star \psi_1^2)^p \rangle$ tends to 0 as $\varepsilon \downarrow 0$. Hence we have derived an upper bound as claimed in (2.5).

The above heuristic is the guiding philosophy of our proof. However, when we expand the transition densities $p_r^{(B)}(x, y)$ into a full eigenvalue expansion, we encounter two singularities: (1) the time parameters r_j getting small and (2) the indices n_j attached to the corresponding eigenfunction ψ_{n_j} getting large. These two singularities hinder us from integrating $\int_{[0,t]} dr_j$ along with the infinite sum $\sum_{n_j \in \mathbb{N}}$. Hence we expand only those transition densities $p_{r_j}^{(B)}(x, y)$ for which $r_j > \delta$. For this part, large n_j indices can easily be summed out, thanks to the factors $\exp\{-\lambda_{n_j} r_j\}$. The rest of the transition densities (for which $r_j \leq \delta$) stay over and are finally integrated out in terms of the Green's function. We spell out the details.

3.3 Eigenvalue Expansion

Recall that we have to show (3.1). We start from (3.2). For brevity, we set forth the following notation. We abbreviate, with a slight abuse of notation,

$$\begin{aligned} \int dy \prod f &= \int_B dy_1 \cdots \int_B dy_k \prod_{j=1}^k f(y_j), \\ \int_{<} dr &= \int_{[0,t]^k} dr_k \cdots dr_1 \mathbf{1}_{\{\sum_{i=1}^k r_i \leq t\}} \quad \left(r_{k+1} = t - \sum_{i=1}^k r_i \right), \\ \int dz \varphi_\varepsilon &= \int_B dz_{m+1} \cdots \int_B dz_k \prod_{j=m+1}^k \varphi_\varepsilon(y_j - z_j). \end{aligned}$$

Our next main step is to expand the transition density terms $p_{r_i}^{(B)}(x_{i-1}, x_i)$ in a standard Fourier series with respect to all the eigenvalues and eigenfunctions of $-\frac{1}{2}\Delta$ in B with zero boundary condition; see Lemma 3.2. However, this series then has good convergence properties only if the time parameter r_i is bounded away from 0. Therefore we introduce a new small parameter $\delta \in (0, \infty)$ and distinguish, for each integration variable r_i , if $r_i \leq \delta$ or $r_i > \delta$. Introducing another small parameter $\eta \in (0, \infty)$, we isolate the contribution from those multi-indices (r_1, \dots, r_k) such that fewer than ηk of the indices i satisfy $r_i \leq \delta$. In other words, we write

$$\int_{<} dr = \sum_{D \subset \{1, \dots, k+1\}} \int_{<} dr \prod_{j \in D} \mathbf{1}_{r_j \leq \delta} \prod_{j \notin D} \mathbf{1}_{r_j > \delta}$$

and see from (3.2) that

$$(3.12) \quad \mathbb{E}_{x_0, x_{k+1}}^{(t)} [(\langle f, \ell_t \rangle - \langle f, \ell_{\varepsilon, t} \rangle)^k] = (I)_{t,k}(\eta, \delta, \varepsilon) + (II)_{t,k}(\eta, \delta, \varepsilon),$$

where

$$(3.13) \quad \begin{aligned} &(I)_{t,k}(\eta, \delta, \varepsilon) \\ &= \sum_{m=0}^k (-1)^m \binom{k}{m} \int dy \prod f \sum_{\substack{\forall i=1, \dots, p: D_i \subset \{1, \dots, k+1\} \\ \#D_i \leq \eta k}} \\ &\quad \times \prod_{i=1}^p \left[\sum_{\sigma \in \mathfrak{S}_k} \int dr \prod_{j \in D_i} \mathbf{1}_{r_j \leq \delta} \prod_{j \in D_i^c} \mathbf{1}_{r_j > \delta} \int dz \varphi_\varepsilon \prod_{j=1}^{k+1} p_{r_j}^{(B)}(x_{j-1}, x_j) \right], \end{aligned}$$

and $(II)_{t,k}(\eta, \delta, \varepsilon)$ is defined accordingly, that is, with the sum on the D_i replaced by the sum on $D_1, \dots, D_p \subset \{1, \dots, k+1\}$ satisfying $\#D_i > \eta k$ for at least one $i \in \{1, \dots, p\}$. This last term has a small exponential rate for fixed η if δ is small, since there are at least ηk integrations $r_i \in [0, \delta]$.

LEMMA 3.3 (Riddance of Small δ). *For every $\eta, \delta > 0$, there is $C(\eta, \delta) > 0$ such that, for any $\varepsilon \in (0, 1]$,*

$$(3.14) \quad |(II)_{t,k}(\eta, \delta, \varepsilon)| \leq k!^p C(\eta, \delta)^k, \quad t \in (0, \infty), k \in \mathbb{N},$$

where $C(\eta, \delta) \downarrow 0$ as $\delta \downarrow 0$.

PROOF. Note that the only i -dependence of the factors in the last line of (3.13) sits in the starting and ending points, $x_0^{(i)}$ and $x_{k+1}^{(i)}$. We neglect the changing signs $(-1)^m$ and estimate $\binom{k}{m} \leq 2^k$ and estimate against the supremum over all $x_0^{(i)} \in B$ and all $x_{k+1}^{(i)}$ for each $i = 1, \dots, p$. Hence, the sum on D_1, \dots, D_p satisfying $\#D_i > \eta k$ for at least one i is equal to p times the sum on those D_1, \dots, D_p satisfying $\#D_1 > \eta k$. Estimating also $|f| \leq C$ and dropping the indicator on $\{\sum_{j=1}^k r_j \leq t\}$ and carrying out the integration on r_j , we obtain

$$\begin{aligned} |(II)| &\leq p(2C)^k \sup_{x_0, x_{k+1} \in B^k} \sum_{m=0}^k \int_{B^k} dy_1 \cdots dy_k \prod_{i=2}^p \left[\sum_{\sigma_i \in \mathfrak{S}_k} \int \varphi_\varepsilon \prod_{j=1}^{k+1} G(x_{j-1}, x_j) \right] \\ &\times \sum_{D_1: \#D_1 > \eta k} \sum_{\sigma_1 \in \mathfrak{S}_k} \int \varphi_\varepsilon \prod_{j \in D_1} G_\delta(x_{j-1}, x_j) \prod_{j \in D_1^c} G(x_{j-1}, x_j), \end{aligned}$$

where G is the Green's function in B and $G_\delta(v, w) = \int_0^\delta ds p_s^{(B)}(v, w)$ is the truncated Green's function. Over $dz_{m+1} \cdots dz_k$ we now carry out the convolution integrals, which turns some of the (truncated) Green's functions into convolved (truncated) Green's functions, each of which can be estimated against $G^{(\star\varepsilon)}$ and $G_\delta^{(\star\varepsilon)}$, respectively, where

$$(3.15) \quad G^{(\star\varepsilon)}(x, y) = \max\{G(x, y), (G(x, \cdot) \star \varphi_\varepsilon)(y)\},$$

and an analogous notation for G replaced by G_δ .

Now we interchange the integration over y_1, \dots, y_k and the sum on σ_1 , such that, after some elementary substitutions involving all the permutations, this sum on σ_1 is turned into $k!$ times the term with σ_1 equal to the identical permutation. This gives

$$\begin{aligned} |(II)| &\leq k! p(2C)^k \sup_{x_0, x_{k+1} \in B^k} \sum_{m=0}^k \int_{B^k} dy_1 \cdots dy_k \prod_{i=2}^p \left[\sum_{\sigma_i \in \mathfrak{S}_k} \prod_{j=1}^{k+1} G^{(\star\varepsilon)}(x_{j-1}, x_j) \right] \\ &\times \sum_{D_1: \#D_1 > \eta k} \prod_{j \in D_1} G_\delta^{(\star\varepsilon)}(y_{j-1}, y_j) \prod_{j \in D_1^c} G^{(\star\varepsilon)}(y_{j-1}, y_j). \end{aligned}$$

Note that, for any $\tilde{\delta} > 0$,

$$(3.16) \quad \begin{aligned} & \limsup_{\delta \downarrow 0} \sup_{\varepsilon \in (0,1]} \sup_{\substack{v, w \in B: \\ |v-w| \geq \tilde{\delta}}} G_{\delta}^{(\star\varepsilon)}(v, w) = 0 \quad \text{and} \\ & \limsup_{\tilde{\delta} \downarrow 0} \sup_{\varepsilon \in (0,1]} \sup_{x \in B} \int_{|x-y| \leq \tilde{\delta}} G^{(\star\varepsilon)}(x, y)^p dy = 0. \end{aligned}$$

In order to employ these two facts, we separate the product over $i = 2, \dots, p$ from the last line with the help of Hölder's inequality and distinguish in the latter term those integrals over $dy_1 \cdots dy_k$ that satisfy $\#\{j \in D_1 : |y_{j-1} - y_j| \leq \tilde{\delta}\} > \tilde{\eta}k$ and the remainder, where $\tilde{\delta} > 0$ and $\tilde{\eta} > 0$ are new small auxiliary parameters. The first contribution gives at least $\tilde{\eta}k$ integrals over $G_{\delta}^{(\star\varepsilon)}(y_{j-1}, y_j)^p dy_j$ with $|y_{j-1} - y_j| \leq \tilde{\delta}$ (and therefore a small number), and in the second, we have at least $\tilde{\eta}k$ indices j with $|y_{j-1} - y_j| > \tilde{\delta}$, which makes it possible to estimate $G_{\delta}^{(\star\varepsilon)}(y_{j-1}, y_j)$ against a small number. Hence the contribution from the last line is bounded by $k! \tilde{C}(\delta, \eta)^k$ for some suitable $\tilde{C}(\delta, \eta) \in (0, \infty)$ satisfying $\lim_{\delta \downarrow 0} \tilde{C}(\delta, \eta) = 0$. The other terms (that is, those that stem from the product over $i = 2, \dots, p$) can be bounded against $(k!)^{p-1} C^k$ for some constant C that does not depend on k . Summarizing, we obtain the estimate in (3.14) with some suitable $C(\delta, \eta)$. The details are pretty standard, and we refer the reader to the proof of [18, lemma 3.3]. \square

Now we go on with the term (I) defined in (3.13) and use the eigenvalue expansion of Lemma 3.2 for all times that are $\geq \delta$. For any $i = 1, \dots, p$ and each $j \in D_i^c$, i.e., for any time duration $r_j \geq \delta$, we expand $p_{r_j}^{(B)}(x_{j-1}, x_j)$ into an eigenvalue series as in Lemma 3.2, introducing a sum on $\mathcal{N}^{(i)} = (n_j^{(i)})_{j \in D_i^c} \in \mathbb{N}^{D_i^c}$. Because $r_j \geq \delta$ and the appearance of the factor $\exp\{-r_j \lambda_{n_j^{(i)}}\}$, the sum on $n_j^{(i)}$ converges exponentially fast.

The eigenfunctions $\psi_{n_j^{(i)}}$ will later be used for an application of the binomial theorem, but this will turn out to be helpful only if all indices $n_j^{(i)}$ appearing are taken from some bounded set. Therefore we truncate this infinite sum at a large cutoff level $R \in \mathbb{N}$. We write $\mathcal{R} = \{1, \dots, R\}$ and split each sum on $n_j^{(i)}$ into the two sums on $n_j^{(i)} \in \mathcal{R}$ and $n_j^{(i)} \in \mathcal{R}^c$. This gives, for every i , sums of the form

$$\prod_{j \in D_i^c} \left(\sum_{n_j^{(i)} \in \mathcal{R}} + \sum_{n_j^{(i)} \in \mathcal{R}^c} \right) = \sum_{E_i \subset D_i^c} \sum_{\mathcal{N}^{(i)} \in \mathcal{R}^{E_i}} \sum_{\mathcal{N}^{(i)} \in (\mathcal{R}^c)^{D_i^c \setminus E_i}},$$

with the understanding that $\mathcal{N}^{(i)} \in \mathcal{R}^{E_i}$ and $\mathcal{N}^{(i)} \in (\mathcal{R}^c)^{D_i^c \setminus E_i}$ may be concatenated to some map $\mathcal{N}^{(i)} : D_i^c \rightarrow \mathbb{N}$.

We now introduce another small parameter $\gamma \in (0, \infty)$ and distinguish the contribution coming from those multisums with sets E_i satisfying $\#(D_i^c \setminus E_i) \leq \gamma k$ for all i and the remainder. This implies the decomposition

$$(I)_{t,k}(\eta, \delta, \varepsilon) = (Ia)_{t,k}(\eta, \gamma, \delta, \varepsilon, R) + (Ib)_{t,k}(\eta, \gamma, \delta, \varepsilon, R),$$

where $(Ia) = (Ia)_{t,k}(\eta, \gamma, \delta, \varepsilon, R)$ is defined as

$$(3.17) \quad \begin{aligned} (Ia) &= \sum_{\substack{\forall i: D_i \subset \{1, \dots, k+1\} \\ \#D_i \leq \eta k}} \sum_{\substack{\forall i: E_i \subset D_i^c \\ \#(D_i^c \setminus E_i) \leq \gamma k}} \sum_{\forall i: \mathcal{N}^{(i)} \in \mathcal{R}^{E_i}} \\ &\times \sum_{\forall i: \mathcal{N}^{(i)} \in (\mathcal{R}^c)^{D_i^c \setminus E_i}} \sum_{m=0}^k (-1)^m \binom{k}{m} \\ &\times \int dy \prod_{i=1}^p f \prod_{\sigma \in \mathfrak{S}_k} \left[\sum_{\sigma} \int dr H_r(\mathcal{N}^{(i)}|_{D_i^c}; D_i) \int dz \varphi_\varepsilon \prod_{j \in D_i} p_{r_j}^{(B)}(x_{j-1}, x_j) \right. \\ &\quad \left. \times \prod_{j \in D_i^c} \psi_{n_j^{(i)}}(x_{j-1}) \psi_{n_j^{(i)}}(x_j) \right] \end{aligned}$$

where

$$(3.18) \quad H_r(\mathcal{N}^{(i)}; D_i) = \left(\prod_{j \in D_i} 1_{r_j \leq \delta} \right) \prod_{j \in D_i^c} (1_{r_j > \delta} \exp\{-r_j \lambda_{n_j^{(i)}}\}).$$

(Ib) is defined accordingly; i.e., for at least one $i \in \{1, \dots, p\}$, the set E_i satisfies $\#(D_i^c \setminus E_i) > \gamma k$. That is, for at least one i , the sum on $n_j^{(i)}$ runs over the remainder set \mathcal{R}^c for at least γk different j 's and gives therefore, for large R , a small factor with power at least γk . Let us first show that therefore $(Ib)_{t,k}(\eta, \gamma, \delta, \varepsilon, R)$ is a small error term if R is large for fixed γ :

LEMMA 3.4 (Riddance of Large \mathcal{N}). *For every $\eta, \gamma, \delta \in (0, 1)$ and $R \in \mathbb{N}$, there is $C^{(b)}(\eta, \gamma, \delta, R) > 0$ such that, for any $\varepsilon \in (0, 1)$,*

$$(3.19) \quad (Ib)_{t,k}(\eta, \gamma, \delta, \varepsilon, R) \leq k!^p C^{(b)}(\eta, \gamma, \delta, R)^k, \quad t \in (0, \infty), k \in \mathbb{N},$$

and $C^{(b)}(\eta, \gamma, \delta, \varepsilon, R) \downarrow 0$ as $R \uparrow \infty$.

PROOF. We use a generic constant C that does not depend on the parameters involved, but possibly only on f , B , and d . In (3.17) (with the necessary changes for (Ib)), we estimate $\sum_{m=0}^k (-1)^m \binom{k}{m} \leq 2^k$ and $\|f\|_\infty \leq C$ and $\int_{<} dr \leq \int_{[0, \infty)^k} dr_1 \cdots dr_k$ and

$$H_r(\mathcal{N}^{(i)}; D_i) \leq \left(\prod_{j \in D_i^c \setminus E_i} 1_{r_j > \delta} \exp\{-r_j \lambda_{n_j^{(i)}}\} \right) \prod_{j \in E_i} \exp\{-r_j \lambda_1\}.$$

Next, in (Ib) we estimate all the terms against their absolute value and then apply the uniform eigenfunction estimate [17]

$$(3.20) \quad \|\psi_n\|_\infty \leq C \lambda_n^{(d-1)/4}, \quad n \in \mathbb{N},$$

to the eigenfunction product $\prod_{j \in D_i^c} \psi_{n_j^{(i)}}(x_{j-1}) \psi_{n_j^{(i)}}(x_j)$ to see that (recall the notation in (3.15))

$$(3.21) \quad \begin{aligned} (Ib) &\leq C^k \sum_{\substack{\forall i: D_i \subset \{1, \dots, k+1\} \\ \#D_i \leq \eta k}} \sum_{\substack{\forall i: E_i \subset D_i^c \\ \exists j: \#(D_i^c \setminus E_j) > \gamma k}} \int dy \prod_{i=1}^p \left[\left(\sum_{\sigma \in \mathfrak{S}_k} \prod_{j \in D_i} G^{(\star \varepsilon)}(x_{j-1}, x_j) \right) \right. \\ &\quad \times \left(\prod_{j \in E_i} \sum_{n_j^{(i)} \in \mathcal{R}} \lambda_{n_j^{(i)}}^{(d-1)/2} \right) \left(\int_{[0, \infty)^{E_i}} dr \prod_{j \in E_i} e^{-r_j \lambda_1} \right) \\ &\quad \times \left. \left(\prod_{j \in D_i^c \setminus E_i} \sum_{n_j^{(i)} \in \mathcal{R}^c} \int_\delta^\infty dr e^{-r \lambda_{n_j^{(i)}}} \lambda_{n_j^{(i)}}^{(d-1)/2} \right) \right] \\ &\leq C^k C_\delta(R)^{\gamma k} C(R)^{pk} \\ &\quad \times \sum_{\substack{\forall i: D_i \\ \#D_i \leq \eta k}} \sum_{\substack{\forall i: E_i \subset D_i^c \\ \exists j: \#(D_i^c \setminus E_j) > \gamma k}} \int dy \prod_{i=1}^p \left(\sum_{\sigma \in \mathfrak{S}_k} \prod_{j \in D_i} G^{(\star \varepsilon)}(x_{j-1}, x_j) \right), \end{aligned}$$

where $C_\delta(R) = \sum_{n \in \mathcal{R}^c} \int_\delta^\infty dr e^{-r \lambda_n} \lambda_n^{(d-1)/2}$ and $C(R) = \sum_{n \in \mathcal{R}} \lambda_n^{(d-1)/2}$, and we have estimated $\int_0^\infty dr e^{-r \lambda_1} \leq C$ for some $C > 1$. We assumed that R is so large that $C_\delta(R) < 1$ and $C(R) \geq 1$. Use that

$$\sup_{\varepsilon \in (0, 1]} \sup_{x \in B} \int_B dy G^{(\star \varepsilon)}(x, y)^p \leq C$$

(see the second statement in (3.16)) to see that the sum on $\sigma \in \mathfrak{S}_k$ is not larger than $(k!)^p C^k$. The two sums on the sets D_i and E_i have no more than C^k terms.

By the well-known Weyl's lemma, λ_n tends to ∞ like $n^{2/d}$. Hence $C_\delta(R)$ decays stretched-exponentially fast to 0 as $R \uparrow \infty$ (the rate depends on δ only), and C_R tends to ∞ only polynomially; hence we may estimate $C^k C_\delta(R)^{\gamma k} C(R)^{pk} \leq C^{(b)}(\eta, \gamma, \delta, R)^k$ with some constant satisfying $C^{(b)}(\eta, \gamma, \delta, \varepsilon, R) \downarrow 0$ as $R \uparrow \infty$. This finishes the proof. \square

3.4 Estimating the Main Term

After the preparations in Lemma 3.3 and 3.4, we now estimate the main term (Ia) defined in (3.17), which is the heart of the proof. The proof of (3.1), and therefore the proof of Proposition 2.3, is finished by the two lemmas, together with the following proposition; see (3.12) and recall the decomposition $(I) = (Ia) + (Ib)$.

PROPOSITION 3.5 (Main Estimate). *For every $\eta, \gamma, \delta, \varepsilon \in (0, 1)$ with $\eta + \gamma < 1/2p$ and for every $R \in \mathbb{N}$, there is a constant $C^{(a)}(\eta, \gamma, \delta, \varepsilon, R) > 0$ such that*

$$(3.22) \quad |(Ia)_{t,k}(\eta, \gamma, \delta, \varepsilon, R)| \leq (k!)^p C^{(a)}(\eta, \gamma, \delta, \varepsilon, R)^k, \quad t \in (0, \infty), k \in \mathbb{N},$$

and $C^{(a)}(\eta, \gamma, \delta, \varepsilon, R) \downarrow 0$ as $\varepsilon \downarrow 0$.

PROOF.

Step 1. Rewriting of Eigenfunction Terms. First we unravel the last term involving the eigenfunctions appearing in the right-hand side of (3.17). Observe that $z_j = z_j^{(i)}$ and $x_j = x_j^{(i)}$ in the i^{th} factor both depend on i , and we write σ_i instead of σ . Recall from Lemma 3.1 that

$$(3.23) \quad x_j^{(i)} = \begin{cases} y_{\sigma_i^{-1}(j)} & \text{if } \sigma_i^{-1}(j) \leq m, \\ z_{\sigma_i^{-1}(j)}^{(i)} & \text{if } \sigma_i^{-1}(j) > m. \end{cases}$$

Therefore, the last term in the second line of (3.17) reads as follows:

$$\begin{aligned} & \prod_{j \in D_i^c} (\psi_{n_j^{(i)}}(x_{j-1}^{(i)}) \psi_{n_j^{(i)}}(x_j^{(i)})) \\ &= \left(\prod_{\substack{j \in \sigma_i^{-1}(D_i^c) \\ j \leq m}} \psi_{n_{\sigma_i(j)}^{(i)}}(y_j) \right) \left(\prod_{\substack{j \in \sigma_i^{-1}(D_i^c - 1) \\ j \leq m}} \psi_{n_{\sigma_i(j)+1}^{(i)}}(y_j) \right) \\ & \quad \times \left(\prod_{\substack{j \in \sigma_i^{-1}(D_i^c) \\ j > m}} \psi_{n_{\sigma_i(j)}^{(i)}}(z_j^{(i)}) \right) \left(\prod_{\substack{j \in \sigma_i^{-1}(D_i^c - 1) \\ j > m}} \psi_{n_{\sigma_i(j)+1}^{(i)}}(z_j^{(i)}) \right). \end{aligned}$$

We now carry out the φ_ε -convolution integration over all $z_j^{(i)}$ and the integration over all those y_j that satisfy the following: (1) they exclusively appear in the above product twice for every $i \in \{1, \dots, p\}$ (but not in the product over the $p_{r_j}^{(B)}$ -terms with $j \in D_i$ for any i); i.e., $\sigma_i(j)$ and $\sigma_i(j) + 1$ both lie in D_i^c , and (2) the indices $n_{\sigma_i(j)}^{(i)}$ and $n_{\sigma_i(j)+1}^{(i)}$, respectively, at the corresponding ψ lies in \mathcal{R} for every $i = 1, \dots, p$; i.e., both indices $\sigma_i(j)$ and $\sigma_i(j) + 1$ lie in E_i . Since $E_i \subset D_i^c$, these are precisely those j that satisfy $j \in S(\sigma)$, where we set, for each $\sigma = (\sigma_1, \dots, \sigma_p) \in \mathfrak{S}_k^p$,

$$S(\sigma) = \bigcap_{i=1}^p \sigma_i^{-1}(F_i) \quad \text{where } F_i = E_i \cap (E_i - 1).$$

Certainly, we have to obey that, for $j \leq m$, the integration is over y_j and for $j > m$ it is the convolution with φ_ε . To express this, we write, for every subset $S \subset \{1, \dots, k\}$,

$$S_{\leq} = S \cap \{1, \dots, m\} \quad \text{and} \quad S_{>} = S \cap \{m+1, \dots, k\}.$$

Each $j \in S(\sigma)$ appears only in the product over $\psi(\dots)$ or $\varphi_\varepsilon \star \psi(\dots)$, whereas for $j \in S(\sigma)^c = \{1, \dots, k\} \setminus S(\sigma)$, the eigenfunction products stay over and remain unconvolved. We write $\mathcal{N} = (\mathcal{N}^{(1)}, \dots, \mathcal{N}^{(p)})$ and $\mathcal{N}_j = (n_j^{(1)}, \dots, n_j^{(p)})$ and introduce, for $j \in S(\sigma)$,

$$(3.24) \quad a(\mathcal{N}_j, \mathcal{N}_{j+1}) = \left\langle f, \prod_{i=1}^p \psi_{n_j^{(i)}} \psi_{n_{j+1}^{(i)}} \right\rangle,$$

$$(3.25) \quad a_\varepsilon(\mathcal{N}_j, \mathcal{N}_{j+1}) = \left\langle f, \prod_{i=1}^p \varphi_\varepsilon \star (\psi_{n_j^{(i)}} \psi_{n_{j+1}^{(i)}}) \right\rangle.$$

Substituting this into (3.17), we conclude

$$(3.26) \quad \begin{aligned} (Ia) &= \sum_{\substack{\forall i: D_i \subset \{1, \dots, k+1\} \\ \#D_i \leq \eta k}} \sum_{\substack{\forall i: E_i \subset D_i^c \\ \#(D_i^c \setminus E_i) \leq \gamma k}} \sum_{\forall i: \mathcal{N}^{(i)} \in \mathcal{R}^{E_i}} \\ &\times \sum_{\forall i: \mathcal{N}^{(i)} \in (\mathcal{R}^c)^{D_i^c \setminus E_i}} \sum_{m=0}^k (-1)^m \binom{k}{m} \\ &\times \sum_{\sigma = (\sigma_1, \dots, \sigma_p) \in \mathfrak{S}_k^p} \left[\prod_{j \in S(\sigma) \leq} a(\mathcal{N}_{\sigma(j)}, \mathcal{N}_{\sigma(j)+1}) \right] \\ &\quad \times \left[\prod_{j \in S(\sigma) >} a_\varepsilon(\mathcal{N}_{\sigma(j)}, \mathcal{N}_{\sigma(j)+1}) \right] \\ &\times G_t(m, D, E, \sigma, \mathcal{N}), \end{aligned}$$

where we wrote $\mathcal{N}_{\sigma(j)} = (n_{\sigma_i(j)}^{(i)})_{i=1, \dots, p}$, $D = (D_1, \dots, D_p)$, and $E = (E_1, \dots, E_p)$, and the remainder term is given as

$$(3.27) \quad \begin{aligned} &G_t(m, D, E, \sigma, \mathcal{N}) \\ &= \int_{\mathcal{B}^{S(\sigma)^c}} dy \prod_{j \in S(\sigma)^c} f(y_j) \\ &\quad \prod_{i=1}^p \left[\int_{<} dr H_r(\mathcal{N}^{(i)}; D_i) \right. \\ &\quad \left. \int \prod_{j \in W_i: j > m} (dz_j^{(i)} \varphi_\varepsilon(y_j - z_j^{(i)})) \prod_{j \in D_i} p_{r_j}^{(B)}(x_{j-1}^{(i)}, x_j^{(i)}) \right. \\ &\quad \left. \times \left(\prod_{j \in \sigma_i^{-1}(D_i^c \setminus E_i): j \leq m} \psi_{n_{\sigma_i(j)}^{(i)}}(y_j) \right) \right] \times \end{aligned}$$

$$\begin{aligned}
& \times \left(\prod_{j \in \sigma_i^{-1}((D_i^c - 1) \setminus F_i): j \leq m} \psi_{n_{\sigma_i(j)+1}}^{(i)}(y_j) \right) \\
& \times \left(\prod_{j \in \sigma_i^{-1}(D_i^c \setminus F_i): j > m} \psi_{n_{\sigma_i(j)}}^{(i)}(z_j^{(i)}) \right) \\
& \times \left(\prod_{j \in \sigma_i^{-1}((D_i^c - 1) \setminus F_i): j > m} \psi_{n_{\sigma_i(j)+1}}^{(i)}(z_j^{(i)}) \right),
\end{aligned}$$

where we recall that $F_i = E_i \cap (E_i - 1)$. Note that G_t depends on $\mathcal{N}^{(i)}$ only via its restriction to D_i^c and on σ_i only via its restriction to

$$(3.28) \quad W_i^c = \sigma_i^{-1}((D_i^c \setminus F_i) \cup ((D_i^c - 1) \setminus F_i) \cup D_i \cup (D_i - 1)) = \sigma_i^{-1}(F_i^c),$$

where c denotes the complement in $\{1, \dots, k\}$.

Step 2. Splitting and Permutation Symmetry. We write $m = m_1 + m_2$ and $k - m = m_3 + m_4$, where $m_1 = \#S(\sigma)_\leq$ and $m_3 = \#S(\sigma)_>$. With $\sum_{m=0}^k (-1)^m \binom{k}{m}$ in front, the second line of (3.26) reads

$$\begin{aligned}
& \sum_{\substack{m_1, m_2, m_3, m_4 \in \mathbb{N}_0 \\ \sum_{l=1}^4 m_l = k}} (-1)^{m_2} \binom{k}{m_1 + m_2} \\
& \sum_{\substack{S_\leq \subset \{1, \dots, m_1 + m_2\} \\ \#S_\leq = m_1}} \sum_{\substack{S_\> \subset \{m_1 + m_2 + 1, \dots, k\} \\ \#S_\> = m_3}} \\
& \times \sum_{\sigma \in \mathfrak{S}_k^p} \mathbf{1}_{\left\{ \begin{array}{l} S_\leq = S(\sigma)_\leq \\ S_\> = S(\sigma)_\> \end{array} \right\}} \left[\prod_{j \in S_\leq} (-a(\mathcal{N}_{\sigma(j)}, \mathcal{N}_{\sigma(j)+1})) \right] \\
& \quad \times \left[\prod_{j \in S_\>} a_\varepsilon(\mathcal{N}_{\sigma(j)}, \mathcal{N}_{\sigma(j)+1}) \right] \\
& \times G_t(m_1 + m_2, D, E, \sigma, \mathcal{N}).
\end{aligned}$$

We claim that the term in the last two lines above is constant on the sets S_\leq and $S_\>$ and depends only on the cardinalities m_1 of S_\leq and m_3 of $S_\>$. More precisely, for $m = m_1 + m_2$ and any permutation $\tau \in \mathfrak{S}_k$ such that $\tau(\{1, \dots, m\}) = \{1, \dots, m\}$, we claim (putting $\sigma \circ \tau = (\sigma_1 \circ \tau, \dots, \sigma_p \circ \tau)$)

(1)

$$\tau^{-1}(S(\sigma)_\leq) = S(\sigma \circ \tau)_\leq \quad \text{and} \quad \tau^{-1}(S(\sigma)_\>) = S(\sigma \circ \tau)_\> ,$$

(2)

$$\begin{aligned} & \prod_{j \in S(\sigma)_{\leq}} a(\mathcal{N}_{\sigma(j)}, \mathcal{N}_{\sigma(j)+1}) \prod_{j \in S(\sigma)_{>}} a_{\varepsilon}(\mathcal{N}_{\sigma(j)}, \mathcal{N}_{\sigma(j)+1}) = \\ & \prod_{j \in S(\sigma \circ \tau)_{\leq}} a(\mathcal{N}_{(\sigma \circ \tau)(j)}, \mathcal{N}_{(\sigma \circ \tau)(j)+1}) \prod_{j \in S(\sigma \circ \tau)_{>}} a_{\varepsilon}(\mathcal{N}_{(\sigma \circ \tau)(j)}, \mathcal{N}_{(\sigma \circ \tau)(j)+1}), \end{aligned}$$

(3)

$$G_t(m_1 + m_2, D, E, \sigma, \mathcal{N}) = G_t(m_1 + m_2, D, E, \sigma \circ \tau, \mathcal{N}).$$

Proofs of these facts are rather easy and involve straightforward computations. Indeed, (i) is seen as follows:

$$\begin{aligned} \tau^{-1}(S(\sigma)_{\leq}) &= \tau^{-1}\left(\bigcap_{i=1}^p S_i(\sigma_i)\right) \cap \{1, \dots, m\} \\ &= \bigcap_{i=1}^p \tau^{-1}(\sigma_i^{-1}(F_i)) \cap \{1, \dots, m\} \\ &= \bigcap_{i=1}^p (\sigma_i \circ \tau)^{-1}(F_i) \cap \{1, \dots, m\} = S(\sigma \circ \tau)_{\leq}. \end{aligned}$$

This proves (i) and similarly one can prove (ii). For the third part, we substitute $\tilde{y}_j = y_{\tau(j)}$ and can perform a similar computation.

Therefore, the sums on S_{\leq} and $S_{>}$ may be replaced by the number of summands, which is $\binom{m_1+m_2}{m_1} \times \binom{k-m_1-m_2}{m_3}$ and the definite choices

$$S_{\leq}^* = \{1, \dots, m_1\} \quad \text{and} \quad S_{>}^* = \{m_1 + m_2 + 1, \dots, m_1 + m_2 + m_3\}.$$

Multiplied with the factor $\binom{k}{m_1+m_2}$, the number gives $\frac{k!}{m_1!m_2!m_3!m_4!}$.

Recall that G_t depends on any permutation σ_i only via its restriction to $W_i^c = \sigma_i^{-1}(F_i^c)$; see (3.28). Therefore, we split each permutation $\sigma_i \in \mathfrak{S}_k$ into two bijections $\sigma_i : W_i \rightarrow F_i$ and $\tau_i : W_i^c \rightarrow F_i^c$ and write

$$\sum_{\sigma \in \mathfrak{S}_k^p} = \sum_{\substack{\forall i: W_i \subset \{1, \dots, k\} \\ \#W_i = \#F_i}} \sum_{\forall i: \sigma_i: W_i \rightarrow F_i} \sum_{\forall i: \tau_i: W_i^c \rightarrow F_i^c},$$

where the two latter sums go over bijections σ_i and τ_i . Furthermore, from (3.24) we see that the a - and a_{ε} -terms depend on $\mathcal{N}^{(i)}$ via its restriction to $F_i = E_i \cap (E_i - 1)$. With this in mind, we decompose the sum on \mathcal{N} as

$$\sum_{\forall i: \mathcal{N}^{(i)} \in \mathcal{R}^{E_i}} = \sum_{\forall i: \mathcal{N}^{(i)} \in \mathcal{R}^{F_i}} \sum_{\forall i: \mathcal{N}^{(i)} \in \mathcal{R}^{E_i \setminus F_i}}.$$

Putting all the material together, we conclude

$$\begin{aligned}
(3.29) \quad (Ia) &= \sum_{\substack{\forall i: D_i \subset \{1, \dots, k\} \\ \#D_i \leq \eta k}} \sum_{\substack{\forall i: E_i \subset D_i^c \\ \#(D_i^c \setminus E_i) \leq \gamma k}} \sum_{\substack{\forall i: W_i \subset \{1, \dots, k\} \\ \#W_i = \#F_i}} \\
&\quad \sum_{\substack{m_1, m_2, m_3, m_4 \in \mathbb{N}_0 \\ \sum_{l=1}^4 m_l = k}} (-1)^{m_2} \frac{k!}{m_1! m_2! m_3! m_4!} \\
&\quad \times \sum_{\forall i: \mathcal{N}^{(i)} \in (\mathcal{R}^c)^{D_i^c \setminus E_i}} \sum_{\forall i: \mathcal{N}^{(i)} \in \mathcal{R}^{E_i \setminus F_i}} \sum_{\forall i: \tau_i: W_i^c \rightarrow F_i^c} \\
&\quad \times \sum_{\forall i: \mathcal{N}^{(i)} \in \mathcal{R}^{F_i}} G_t(m_1 + m_2, D, E, \tau, \mathcal{N}) \\
&\quad \times \sum_{\forall i: \sigma_i: W_i \rightarrow F_i} \left[\prod_{j \in S_{\leq}^*} (-a(\mathcal{N}_{\sigma(j)}, \mathcal{N}_{\sigma(j)+1})) \right] \\
&\quad \quad \times \left[\prod_{j \in S_{>}^*} a_\varepsilon(\mathcal{N}_{\sigma(j)}, \mathcal{N}_{\sigma(j)+1}) \right].
\end{aligned}$$

Step 3. Counting Permutations and Multi-indices. Our next goal is to simplify the terms starting from the sum on $\mathcal{N}^{(i)} \in \mathcal{R}^{F_i}$ on the right-hand side of (3.29) and to show that these terms contain the k^{th} power of a small number if ε is small, which lays the basis of an upper bound as in (3.22) with a small number to the power k . For doing this, we will count the number of $\mathcal{N}^{(1)}, \dots, \mathcal{N}^{(p)}$ and of $\sigma_1, \dots, \sigma_p$ that give precisely the same contribution and to apply the binomial theorem (incorporating the sum on m_1 and m_3) for a large power of terms of the form $a_\varepsilon(l) - a(l)$, which is uniformly small if ε is small. This is the point after which we are finally allowed to use more stable estimates like the triangle inequality for absolute signs.

The starting point is that many of the multi-indices $\mathcal{N}^{(i)} \in \mathcal{R}^{F_i}$ and of the permutations $\sigma_1, \dots, \sigma_p$, $i = 1, \dots, p$, give precisely the same contribution. Our task here is to identify what classes of such \mathcal{N} and σ do this and to evaluate their cardinality.

First we note that the two products in the third line do not depend on each value of $(\mathcal{N}_j, \mathcal{N}_{j+1})$ for $j \in S^*$, but only on their occupation numbers, i.e., on the number $A(l)$ of occurrences of a given vector $l \in (\mathcal{R}^2)^p$ in the vector $(\mathcal{N}_j, \mathcal{N}_{j+1})_{j \in S^*}$. Hence $A : (\mathcal{R}^2)^p \rightarrow \mathbb{N}_0$ is a map satisfying $\sum_{l \in (\mathcal{R}^2)^p} A(l) = m_1 + m_3$, and we will be summing on all such maps.

Furthermore, recall the definition of G_t from (3.27). Note that G_t depends on $\mathcal{N}^{(i)}|_{F_i}$ only via the appearance of these indices in the products

$$\prod_{j \in F_i} 1_{r_j > \delta} \exp\{-r_j \lambda_{n_j^{(i)}}\}$$

embedded in the H_r terms (recall (3.18)). Due to this product structure, for some suitable function \tilde{G}_t (which we do not make explicit here), we may write

$$G_t(m_1 + m_2, D, E, \tau, \mathcal{N}) = \tilde{G}_t(m_2 + m_4, D, E, \tau, A, (\mathcal{N}^{(i)}|_{D_i^c \setminus F_i})_{i=1, \dots, p})$$

if A is the occupation times vector of $(\mathcal{N}_j, \mathcal{N}_{j+1})_{j \in S^*}$.¹ Also we have used that $m_2 + m_4$ can be constructed from $m = m_1 + m_2$ and A .

However, in order to describe the last line on the right-hand side of (3.29), we also have to sum on all occupation numbers $r(l)$ of the vectors $(\mathcal{N}_j, \mathcal{N}_{j+1})$ in the first product and the occupation numbers (which are necessarily $A(l) - r(l)$) in the second product. This leads to a further sum on all maps $r : (\mathcal{R}^2)^p \rightarrow \mathbb{N}_0$ satisfying $\sum_{l \in (\mathcal{R}^2)^p} r(l) = m_1$ and $0 \leq r(l) \leq A(l)$ for any $l \in (\mathcal{R}^2)^p$. We denote by M_{m_1, m_3} the set of all pairs (A, r) of such maps and by $M_{m_1 + m_3}$ the set of all maps A as above. Our strategy is to write the right-hand side of (3.29) as a sum on $A \in M_{m_1 + m_3}$ and a sum on $(A, r) \in M_{k, m}$, express both the products over the a -terms as functions of A and r , and finally to count all the tuples $(\mathcal{N}^{(i)}|_{F_i}, \sigma_i)$, $i = 1, \dots, p$, such that (A, r) is the pair of occupation number vectors of the vectors $(\mathcal{N}_{\sigma(j)}, \mathcal{N}_{\sigma(j)+1})$ for $j \in S^*$. By the last we mean that $A(l)$ is equal to the number of $j \in S^*$ such that $l = (\mathcal{N}_{\sigma(j)}, \mathcal{N}_{\sigma(j)+1})$.

In view of this discussion, the terms starting from the sum on $\mathcal{N}^{(i)} \in \mathcal{R}^{F_i}$ on the right-hand side of (3.29) read as

$$(3.30) \quad \sum_{(A, r) \in M_{m_1, m_3}} \tilde{G}_t(m_2 + m_4, D, E, \tau, A, \mathcal{N}) \\ \times \prod_{l \in (\mathcal{R}^2)^p} [(-a(l))^{r(l)} a_\varepsilon(l)^{A(l) - r(l)}] \# \Psi(A, r),$$

where the set Ψ is given by

$$(3.31) \quad \Psi(A, r) = \left\{ (\mathcal{N}^{(i)}|_{F_i}, \sigma_i)_{i=1, \dots, p} : \forall l \in (\mathcal{R}^2)^p, \right. \\ \left. r(l) = \#\{j \in S_{\leq}^* : (\mathcal{N}_{\sigma(j)}, \mathcal{N}_{\sigma(j)+1}) = l\}, \right. \\ \left. A(l) - r(l) = \#\{j \in S_{>}^* : (\mathcal{N}_{\sigma(j)}, \mathcal{N}_{\sigma(j)+1}) = l\} \right\},$$

where the domains of the $\mathcal{N}^{(i)}|_{F_i}$ and the σ_i are as in (3.29).

Now we evaluate this counting term. We will decompose this in the two steps of counting first the multi-indices and afterwards the permutation. For every $i = 1, \dots, p$, we define the i^{th} marginal of $A \in M_{m_1 + m_3}$ by

$$(3.32) \quad A_i(l^{(i)}) = \sum_{(l^{(j)})_{j \neq i} \in (\mathcal{R}^2)^{p-1}} A(l^{(1)}, \dots, l^{(p)}), \quad l^{(i)} \in \mathcal{R}^2.$$

¹The definition of \tilde{G}_t is canonical: \tilde{G}_t as a function of A is equal to G_t as a function of some $(\mathcal{N}^{(i)}|_{F_i})_{i=1, \dots, p}$ such that the occupation times vector of $(\mathcal{N}_j, \mathcal{N}_{j+1})_{j \in S^*}$ is equal to A .

Now we consider the multi-indices \mathcal{N} that produce the occupation times vectors A_i :

$$(3.33) \quad \begin{aligned} \Phi(A) &= \Phi(A_1, \dots, A_p) \\ &= \left\{ (\mathcal{N}^{(i)}|_{F_i})_{i=1, \dots, p} : \forall i = 1, \dots, p, \forall l^{(i)} \in \mathcal{R}^2, \right. \\ &\quad \left. \#\{j \in S^* : (\mathcal{N}_j^{(i)}, \mathcal{N}_{j+1}^{(i)}) = l^{(i)}\} = A_i(l^{(i)}) \right\}. \end{aligned}$$

Given $\mathcal{N} \in \Phi(A)$, we denote

$$(3.34) \quad \Psi(A, r, \mathcal{N}) = \left\{ (\sigma_i)_{i=1, \dots, p} \in \bigotimes_{i=1}^p \mathcal{B}(W_i, F_i) : (\mathcal{N}, \sigma_1, \dots, \sigma_p) \in \Psi(A, r) \right\},$$

where we denote by $\mathcal{B}(W, F)$ the set of bijections $W \rightarrow F$. Then it is clear that $\#\Psi(A, r) = \sum_{\mathcal{N} \in \Phi(A)} \#\Psi(A, r, \mathcal{N})$. The cardinality of $\Psi(A, r, \mathcal{N})$ is given in the next lemma.

LEMMA 3.6 (Cardinality of $\Psi(A, r, \mathcal{N})$). *For any $m_1, m_3 \in \mathbb{N}_0$ and any $(A, r) \in M_{m_1, m_3}$ and any $\mathcal{N} \in \Phi(A)$,*

$$(3.35) \quad \#\Psi(A, r, \mathcal{N}) = m_1! m_3! \frac{\prod_{i=1}^p \prod_{l^{(i)} \in \mathcal{R}^2} A_i(l^{(i)})!}{\prod_{l \in (\mathcal{R}^2)^p} A(l)!} \prod_{l \in (\mathcal{R}^2)^p} \binom{A(l)}{r(l)}.$$

PROOF. We count the number of p independent bijections $\sigma_i : W_i \rightarrow F_i$ for $i = 1, \dots, p$ with the prescribed properties. Since $\#(\bigcap_{i=1}^p W_i) = \#(\bigcap_{i=1}^p F_i) = \#S^*$, clearly this task boils down to counting all permutations σ_i of $S^* = S_{\leq}^* \cup S_{>}^*$. From now on, therefore, we shall be counting permutations σ_i of S^* .

For $p = 1$, we want to find out the number of permutations σ of the numbers in S^* such that any $l \in \mathcal{R}^2$ appears $r(l)$ times as a pair $(n_{\sigma(j)}, n_{\sigma(j)+1})$ for $j \in S_{\leq}^*$ and $A(l) - r(l)$ times as a pair $(n_{\sigma(j)}, n_{\sigma(j)+1})$ for $j \in S_{>}^*$. We will now describe a two-step procedure that constructs all such σ . For each $l \in \mathcal{R}^2$, choose $r(l)$ out of $A(l)$ indices $j \in S^*$ such that $(n_j, n_{j+1}) = l$. Let D be the set of those j . Then D has precisely m_1 elements and there are $\prod_{l \in \mathcal{R}^2} \binom{A(l)}{r(l)}$ choices. Now any permutation σ that maps $\{1, \dots, m_1\}$ onto D has the above property. Obviously, for a given D , there are $m_1! m_3!$ such σ 's. This shows that there are at least as many as $m_1! m_3! \prod_{l \in \mathcal{R}^2} \binom{A(l)}{r(l)}$ such σ 's. In other words,

$$(3.36) \quad \#\Psi(A, r, \mathcal{N}) \geq \prod_{l \in \mathcal{R}^2} \binom{A(l)}{r(l)} m_1! m_3!.$$

To show the upper bound \leq , we pick a $\sigma \in \Psi$ and put $D = \{\sigma(1), \dots, \sigma(m_1)\}$. Then, by definition of Ψ , D contains, for any l , precisely $r(l)$ out of $A(l)$ indices j satisfying $(n_j, n_{j+1}) = l$. This means that the above construction also produces

the chosen σ . This shows that equality holds in (3.36). Hence we have proved (3.35) for $p = 1$.

For $p = 2$, we can go ahead similarly. For this, let us denote the left-hand side of (3.35) by $\#\Psi_2(A, r, \mathcal{N})$. Without loss of generality, we may assume that $\mathcal{N} \in \Phi(A)$. First we argue that

$$(3.37) \quad \{\sigma_1 \in \mathfrak{S}(S^*) : \exists \sigma_2 \in \mathfrak{S}(S^*) : (\sigma_1, \sigma_2) \in \Psi_2(A, r, \mathcal{N})\} = \Psi_1(A_1, r_1, \mathcal{N}^{(1)})$$

where $\Psi_1(A_1, r_1, \mathcal{N}^{(1)})$ is defined in (3.31) for $p = 1$ and A and r are replaced by their first marginals A_1 and r_1 , respectively. Indeed, let $\sigma_1, \sigma_2 \in \mathfrak{S}(S^*)$ be such that $r(\cdot)$ and $A(\cdot) - r(\cdot)$ are the occupation times vectors of $(n_{\sigma_i(j)}^{(i)}, n_{\sigma_i(j)+1}^{(i)})_{i=1,2}$ for $j = 1, \dots, m_1$ and of $(n_{\sigma_i(j)}^{(i)}, n_{\sigma_i(j)+1}^{(i)})_{i=1,2}$ for $j = m_1 + m_2 + 1, \dots, m_1 + m_2 + m_3$, respectively. Recall that for any $i \in \{1, \dots, p\}$, we have the i^{th} row vector $(n_j^{(i)})_{j \in S^*}$, which we denoted by $\mathcal{N}^{(i)}$. By projecting onto the first row, we see that r_1 and $A_1 - r_1$ are the occupation numbers of $(n_{\sigma_1(j)}^{(1)}, n_{\sigma_1(j)+1}^{(1)})$ for $j = 1, \dots, m_1$ and $(n_{\sigma_1(j)}^{(1)}, n_{\sigma_1(j)+1}^{(1)})$ for $j = m_1 + m_2 + 1, \dots, m_1 + m_2 + m_3$. This shows that $\sigma_1 \in \Psi_1(A_1, r_1, \mathcal{N}^{(1)})$.

Let us show that also \supset holds in (3.37). Pick $\sigma_1 \in \Psi_1(A_1, r_1, \mathcal{N}^{(1)})$. Since $\mathcal{N} \in \Phi(A)$, for each $l^{(2)} \in \mathcal{R}^2$, there are precisely $A_2(l^{(2)})$ indices j such that $(n_j^{(2)}, n_{j+1}^{(2)}) = l^{(2)}$. Therefore, there is an order (i.e., a permutation σ_2 of the second row) such that, for any $l^{(1)}$ and any $r(l^{(1)}, l^{(2)})$, the set $\{j \in S_{\leq}^* : (n_{\sigma_1(j)}^{(1)}, n_{\sigma_1(j)+1}^{(1)}) = l^{(1)}\}$ contains precisely as many as $r(l^{(1)}, l^{(2)})$ indices j satisfying $(n_{\sigma_2(j)}^{(2)}, n_{\sigma_2(j)+1}^{(2)}) = l^{(2)}$ for any $l^{(2)} \in \mathcal{R}^2$ and the set

$$\{j \in S_{>}^* : (n_{\sigma_1(j)}^{(1)}, n_{\sigma_1(j)+1}^{(1)}) = l^{(1)}\}$$

contains precisely as many as $A(l^{(1)}, l^{(2)}) - r(l^{(1)}, l^{(2)})$ indices j satisfying

$$(n_{\sigma_2(j)}^{(2)}, n_{\sigma_2(j)+1}^{(2)}) = l^{(2)}$$

for any $l^{(2)} \in \mathbb{N}^2$. Therefore, $(\sigma_1, \sigma_2) \in \Psi_2(A, r, \mathcal{N})$. This proves (3.37).

Hence we have

$$(3.38) \quad \#\Psi_2(A, r, \mathcal{N}) = \sum_{\sigma_1 \in \Psi_1(A_1, r_1, \mathcal{N}^{(1)})} \#\{\sigma_2 : (\sigma_1, \sigma_2) \in \Psi_2(A, r, \mathcal{N})\}.$$

Fix $\sigma_1 \in \Psi_1(A_1, r_1, \mathcal{N}^{(1)})$. We now give a two-step construction of all σ_2 satisfying $(\sigma_1, \sigma_2) \in \Psi_2(A, r, \mathcal{N})$. For each $l^{(1)}, l^{(2)} \in \mathcal{R}^2$, we decompose the set $\{j \in S_{\leq}^* : (n_{\sigma_1(j)}^{(1)}, n_{\sigma_1(j)+1}^{(1)}) = l^{(1)}\}$ into disjoint sets $D_{l^{(1)}, l^{(2)}}$ of cardinality

$r(l^{(1)}, l^{(2)})$ and the set $\{j \in S^* : (n_{\sigma_1(j)}^{(1)}, n_{\sigma_1(j)+1}^{(1)}) = l^{(1)}\}$ into sets $\bar{D}_{l^{(1)}, l^{(2)}}$ of cardinality $A(l^{(1)}, l^{(2)}) - r(l^{(1)}, l^{(2)})$. For doing this, we have

$$\prod_{l^{(1)} \in \mathcal{R}^2} \frac{r_1(l^{(1)})!(A_1 - r_1)(l^{(1)})!}{\prod_{l^{(2)} \in \mathcal{R}^2} (r(l^{(1)}, l^{(2)})!)((A - r)(l^{(1)}, l^{(2)})!)}$$

choices. Having fixed these sets, we see that every permutation σ_2 satisfying

$$\sigma_2(\{j \in S^* : (n_j^{(2)}, n_{j+1}^{(2)}) = l^{(1)}\}) = \bigcup_{l^{(1)} \in \mathcal{R}^2} (D_{l^{(1)}, l^{(2)}} \cup \bar{D}_{l^{(1)}, l^{(2)}}) \quad \forall l^{(2)} \in \mathcal{R}^2$$

has the property that each pair $(l^{(1)}, l^{(2)})$ appears precisely $r(l^{(1)}, l^{(2)})$ times in the vector $(n_{\sigma_i(j)}^{(i)}, n_{\sigma_i(j)+1}^{(i)})_{i=1,2}$ for $j = 1, \dots, m_1$ and precisely $(A - r)(l^{(1)}, l^{(2)})$ times in the vector $(n_{\sigma_i(j)}^{(i)}, n_{\sigma_i(j)+1}^{(i)})_{i=1,2}$ for $j = m_1 + m_2 + 1, \dots, m_1 + m_2 + m_3$. That is, $(\sigma_1, \sigma_2) \in \Psi_2(A, r, \mathcal{N})$. Obviously, there are $\prod_{l^{(2)}} A_2(l^{(2)})!$ such permutations σ_2 . Different choices of D and \bar{D} produce different choices of permutations σ_1, σ_2 . A little reflection shows that every σ_2 satisfying $(\sigma_1, \sigma_2) \in \Psi_2$ can be constructed in this way by putting

$$D_{(l^1, l^2)} = \{j \in S^* : (n_{\sigma_i(j)}^{(i)}, n_{\sigma_i(j)+1}^{(i)})_{i=1,2}\}$$

and

$$\bar{D}_{(l^1, l^2)} = \{j \in S^* : (n_{\sigma_i(j)}^{(i)}, n_{\sigma_i(j)+1}^{(i)})_{i=1,2}\}.$$

Therefore, we have

$$\begin{aligned} \# \Psi_2(A, r, \mathcal{N}) &= \# \Psi_1(A_1, r_1, \mathcal{N}^{(1)}) \prod_{l^{(2)} \in \mathcal{R}^2} A_2(l^{(2)})! \\ &\quad \times \prod_{l^{(1)} \in \mathcal{R}^2} \frac{r_1(l^{(1)})!(A_1 - r_1)(l^{(1)})!}{\prod_{l^{(2)} \in \mathcal{R}^2} r(l^{(1)}, l^{(2)})!(A - r)(l^{(1)}, l^{(2)})!} \\ (3.39) \quad &= m_1! m_3! \frac{\prod_{l^{(1)}} A_1(l^{(1)})! \prod_{l^{(2)}} A_2(l^{(2)})!}{\prod_{l^{(1)}, l^{(2)}} r(l^{(1)}, l^{(2)})!(A - r)(l^{(1)}, l^{(2)})!} \\ &= m_1! m_3! \frac{\prod_{i=1}^2 \prod_{l^{(i)} \in \mathcal{R}^2} A_i(l^{(i)})!}{\prod_{l \in (\mathcal{R}^2)^2} A(l)!} \prod_{l \in (\mathcal{R}^2)^2} \binom{A(l)}{r(l)}. \end{aligned}$$

This proves (3.35) for $p = 2$. We leave the proof for $p > 2$ to the reader, as it is similar and can be carried out in a recursive manner. \square

We recall that $\Psi(A, r) = \sum_{\mathcal{N} \in \Phi(A)} \Psi(A, r, \mathcal{N})$ and use (3.35) to plug in the expression for $\# \Psi(A, r, \mathcal{N})$ in (3.30). We now insert this new form of (3.30) in

(3.29). Replacing m_1 on the right-hand side of (3.29) by $\sum_l r(l)$, the only condition on r in the set $\bigcup_{m=0}^{m_1+m_3} M_{m_1, m_3}$ that is left is that $r(l) \in \{0, \dots, A(l)\}$ for any l . Therefore, we infer from (3.30) and (3.29) that

$$\begin{aligned}
(3.40) \quad (Ia) &= \sum_{\substack{\forall i: D_i \subset \{1, \dots, k\} \\ \#D_i \leq \eta k}} \sum_{\substack{\forall i: E_i \subset D_i^c \\ \#(D_i^c \setminus E_i) \leq \gamma k}} \sum_{\substack{\forall i: W_i \subset \{1, \dots, k\} \\ \#W_i = \#F_i}} \\
&\times \sum_{m_2+m_4 \leq k} (-1)^{m_2} \frac{k!}{m_2! m_4!} \\
&\times \sum_{\forall i: \mathcal{N}^{(i)} \in (\mathcal{R}^c)^{D_i^c \setminus E_i}} \sum_{\forall i: \mathcal{N}^{(i)} \in \mathcal{R}^{E_i \setminus F_i}} \sum_{\forall i: \tau_i: W_i^c \rightarrow F_i^c} \\
&\times \sum_{A \in M_{k-m_2-m_4}} \tilde{G}_t(m_2 + m_4, D, E, \tau, A, \mathcal{N}) \\
&\times \#\Phi(A) \frac{\prod_{i=1}^p \prod_{l^{(i)} \in \mathcal{R}^2} A_i(l^{(i)})!}{\prod_{l \in (\mathcal{R}^2)^p} A(l)!} \\
&\times \prod_{l \in (\mathcal{R}^2)^p} \left[\sum_{r(l)=0}^{A(l)} [(-a(l))^{r(l)} a_\varepsilon(l)^{A(l)-r(l)}] \binom{A(l)}{r(l)} \right].
\end{aligned}$$

By the binomial theorem, the last term in the brackets is equal to $(a(l) - a_\varepsilon(l))^{A(l)}$.

Step 4. Finishing: Some Estimates. In this step we shall prove (3.22) and finish the proof of Proposition 3.5. From now on, we will use that $|a(l) - a_\varepsilon(l)|$ is, for fixed R , small uniformly in $l \in \mathcal{R}^{2p}$ if $\varepsilon > 0$ is small, and we are allowed to use the triangle inequality to estimate all the other terms appearing in (3.40) in absolute value. We will use C to denote a generic positive constant that depends possibly only on f , B , and d and may change its value from appearance to appearance.

The main task now is to estimate the second line of (3.40) as follows: We claim that there is some $C_\delta \in (0, \infty)$ such that, for any $k, m_2, m_4 \in \mathbb{N}$ satisfying $m_2 + m_4 \leq k$ and for any $A \in M_{k-m_2-m_4}$ and for any $t \in (0, \infty)$,

$$\begin{aligned}
(3.41) \quad &\sum_{\forall i: \mathcal{N}^{(i)} \in (\mathcal{R}^c)^{D_i^c \setminus E_i}} \sum_{\forall i: \mathcal{N}^{(i)} \in \mathcal{R}^{E_i \setminus F_i}} \\
&\sum_{\forall i: \tau_i: W_i^c \rightarrow F_i^c} |\tilde{G}_t(m_2 + m_4, D, E, \tau, A, \mathcal{N})| \leq C_\delta^k \prod_{i=1}^p \#(F_i^c)!.
\end{aligned}$$

We defer the proof of (3.41) to the end of this step.

Next, it is a standard fact from combinatorics [13, II.2] that, for $A \in M_{k-m_2-m_4}$,

$$(3.42) \quad \#\Phi(A) \leq k^p \prod_{i=1}^p \frac{\prod_{l_1^{(i)} \in \mathcal{R}} \bar{A}_i(l_1^{(i)})!}{\prod_{l^{(i)} \in \mathcal{R}^2} A_i(l^{(i)})!}$$

where \overline{A}_i is the marginal of A_i on the first component, i.e.,

$$\overline{A}_i(l_1) = \sum_{l_2 \in \mathcal{R}} A_i(l_1, l_2) \quad \text{for every } l_1 \in \mathcal{R}.$$

We estimate the sum over W_i against $\binom{k}{\#F_i}$ and the sum over D_i and E_i against C^k . Combining everything, we conclude

$$(3.43) \quad \begin{aligned} (Ia) &\leq k^p C^k C_\delta^k \sum_{m_2+m_4 \leq k} \frac{k!}{m_2!m_4!} \prod_{i=1}^p \left[\binom{k}{\#F_i} \#F_i^{\circ!} \right] \\ &\quad \times \sum_{A \in M_{k-m_2-m_4}} \frac{\prod_{i=1}^p \prod_{l_1^{(i)} \in \mathcal{R}} \overline{A}_i(l_1^{(i)})!}{\prod_{l \in (\mathcal{R}^2)^p} A(l)!} \prod_{l \in (\mathcal{R}^2)^p} |a(l) - a_\varepsilon(l)|^{A(l)} \\ &\leq k^p C^k C_\delta^k k!^p \sum_{m_2+m_4 \leq k} \frac{k!}{m_2!m_4!(k-m_2-m_4)!} \\ &\quad \times \sum_{A \in M_{k-m_2-m_4}} \frac{(k-m_2-m_4)!}{\prod_{l \in (\mathcal{R}^2)^p} A(l)!} \prod_{l \in (\mathcal{R}^2)^p} |a(l) - a_\varepsilon(l)|^{A(l)}, \end{aligned}$$

where we estimated $\#F_i! \geq (k-m_2-m_4)!$, which is true for any i since $S^* \subset \sigma_i^{-1}(F_i)$, and $\prod_{i=1}^p \prod_{l_1^{(i)} \in \mathcal{R}} \overline{A}_i(l_1^{(i)})! \leq (k-m_2-m_4)!$, which is true since the numbers $\overline{A}_i(l_1^{(i)})$ sum up to $k-m_2-m_4$.

From the multinomial theorem we see that the last sum is equal to $C_{\varepsilon, R}^{k-m_2-m_4}$, where $C_{\varepsilon, R} = \sum_{l \in (\mathcal{R}^2)^p} |a(l) - a_\varepsilon(l)|$. Take ε so small that $C_{\varepsilon, R} < 1$; then we can estimate $C_{\varepsilon, R}^{k-m_2-m_4} \leq C_{\varepsilon, R}^{k(1-2p(\eta+\gamma))}$, since

$$\begin{aligned} k - m_2 - m_4 = \#S^* &= \# \bigcap_{i=1}^p W_i = \# \bigcap_{i=1}^p (E_i \cap (E_i - 1)) \\ &\geq k(1 - 2p(\eta + \gamma)), \end{aligned}$$

since $\#D_i^{\circ} \geq k(1-\eta)$ and $\#(D_i^{\circ} \setminus E_i) \leq \gamma k$ (and also $\#(D_i^{\circ} \setminus (E_i - 1)) \leq \gamma k$) and therefore $\#(E_i \cap (E_i - 1)) \geq k(1 - 2(\eta + \gamma))$.

The sum over $m_2 + m_4 \leq k$ on the right-hand side of (3.43) is equal to 3^k , which we absorb in the C^k . Hence we derive the estimate

$$(Ia) \leq (k!)^p k^p C^k C_\delta^k C_{\varepsilon, R}^{k(1-2p(\eta+\gamma))}.$$

Since $\lim_{\varepsilon \downarrow 0} C_{\varepsilon, R} = 0$ and $\eta + \gamma < 1/2p$, this estimate proves (3.22) and therefore finishes the proof of Proposition 3.5.

Now we owe the reader only the proof of (3.41). In (3.18), we estimate

$$\begin{aligned}
H_r(\mathcal{N}^{(i)}; D_i) &\leq \prod_{j \in D_i^c} \left(1_{r_j > \delta} \exp \left\{ -\frac{r_j}{2} \lambda_{n_j^{(i)}} \right\} \right) \\
&\quad \times \prod_{j \in (D_i^c - 1)} \left(1_{r_{j+1} > \delta} \exp \left\{ -\frac{r_{j+1}}{2} \lambda_{n_{j+1}^{(i)}} \right\} \right) \\
&\leq \prod_{j \in F_i} \exp\{-r_j \lambda_1\} \prod_{j \in D_i^c \setminus F_i} \left(1_{r_j > \delta} \exp \left\{ -\frac{r_j}{2} \lambda_{n_j^{(i)}} \right\} \right) \\
&\quad \times \prod_{j \in (D_i^c - 1) \setminus F_i} \left(1_{r_{j+1} > \delta} \exp \left\{ -\frac{r_{j+1}}{2} \lambda_{n_{j+1}^{(i)}} \right\} \right).
\end{aligned}$$

Furthermore, we drop the indicator on $\{\sum_{j=1}^{k+1} r_j \leq t\}$ such that all integrations on r_j can be executed freely (over $[\delta, \infty)$ for $j \notin F_i$ and over $[0, \infty)$ for $j \in F_i$) as an upper bound. In (3.27), we estimate the absolute value of G_t by using the triangle inequality and the uniform eigenfunction estimate from (3.20). Furthermore, we also summarize and estimate the sums over $\mathcal{N}^{(i)}|_{D_i^c \setminus E_i}$ and $\mathcal{N}^{(i)}|_{E_i \setminus F_i}$ as a sum over $\mathcal{N}^{(i)}|_{D_i^c \setminus F_i} \in \mathbb{N}^{D_i^c \setminus F_i}$ for $i = 1, \dots, p$. Hence, we obtain, also using the notation of (3.16),

$$\begin{aligned}
\text{LHS of (3.41)} &\leq C^k \int_{B(S^*)^c} dy \prod_{j \in (S^*)^c} \prod_{i=1}^p \left[\left(\sum_{\tau_i: W_i^c \rightarrow F_i^c} \prod_{j \in D_i} G^{(\star\epsilon)}(y_{\tau_i^{-1}(j-1)}, y_{\tau_i^{-1}(j)}) \right) \right. \\
&\quad \times \left(\prod_{j \in D_i^c \setminus F_i} \sum_{n_j^{(i)} \in \mathcal{R}^c} \int_{\delta}^{\infty} dr e^{-r \lambda_{n_j^{(i)}}} \lambda_{n_j^{(i)}}^{\frac{d-1}{2}} \right) \\
&\quad \times \left. \left(\int_{[0, \infty)^{F_i}} dr \prod_{j \in F_i} e^{-r_j \lambda_1} \right) \right] \\
&\leq C^k C_{\delta}^k \left(\prod_{i=1}^p \#F_i^c! \right) \int_{B(S^*)^c} dy \prod_{i=1}^p \prod_{j \in D_i} G^{(\star\epsilon)}(y_{j-1}, y_j)
\end{aligned}$$

where $C_{\delta} = \sum_{n \in \mathbb{N}} \int_{\delta}^{\infty} dr e^{-r \lambda_n} \lambda_n^{(d-1)/2} \vee 1$, and we absorbed the $\#F_i$ -fold power of $\int_0^{\infty} dr e^{-r \lambda_1} = 1/\lambda_1$ in the term C^k , and we used Jensen's inequality to the sum over τ_1, \dots, τ_p to get hold of the term $\prod_{i=1}^p (\#F_i^c)!$. The integrals over the y_j are now bounded by C^k , thanks to the classical fact $\sup_{x \in B} \int_B dy G^p(x, y) \leq C$ for $p < d/(d-2)$. Altering the value of C_{δ} suitably, we finish the proof of (3.41). \square

4 From Large Time to Large Mass: Proof of Theorem 1.3

In this section we prove Theorem 1.3. To do this, we carry over our LDP for ℓ_{tb} as the time t diverges (Theorem 1.1) to an LDP for $\ell = \ell_{(\tau_1, \dots, \tau_p)}$ with random time horizon $[0, \tau_1) \times \dots \times [0, \tau_p)$ as the mass $\ell(U)$ diverges. Recall that U is a compact subset of B whose boundary is a Lebesgue null set. We want large deviations for the probability measures $\ell/\ell(U)$ conditional on $\mathbb{P}(\cdot \mid \ell(U) > a)$, as $a \uparrow \infty$ with rate function J defined in (1.10). The basic idea is to replace ℓ with ℓ_{tb} where $t = a^{1/p}$ and to optimize over $b = (b_1, \dots, b_p)$. In other words, we cut each i^{th} Brownian path at some time tb_i smaller than τ_i for some $b_i > 0$ and control the cutoff part. Theorem 1.1 gives the large-deviation rate for ℓ_{tb} as $t \rightarrow \infty$. Optimizing over b_1, \dots, b_p gives us the desired asymptotics. Lemmas 4.1 and 4.2 below give the lower and upper bounds, respectively, in the LDP.

We pick a metric d on $\mathcal{M}(B)$ that induces the weak topology. Recall that $\mathcal{M}_U(B)$ is the subspace of positive measures on B whose restriction to U is a probability measure.

LEMMA 4.1 (Lower Bound). *For every open set $G \subset \mathcal{M}_U(B)$, we have*

$$(4.1) \quad \liminf_{a \uparrow \infty} \frac{1}{a^{1/p}} \log \mathbb{P} \left(\frac{\ell}{\ell(U)} \in G, \ell(U) > a \right) \geq - \inf_{\mu \in G} J(\mu).$$

PROOF. Set $t = a^{1/p}$ and fix $b = (b_1, \dots, b_p) \in (0, \infty)^p$. For any $\delta_1, \delta_2 > 0$, we have

$$\begin{aligned} \{\ell(U) > a\} &\supset \{a < \ell(U) < a(1 + \delta_1)\} \cap \bigcap_{i=1}^p \{tb_i < \tau_i < t(b_i + \delta_2)\} \\ &\supset \{a < \ell_{tb}(U) < a(1 + \delta_1) - (\ell_{t(b+\delta_2)}(U) - \ell_{tb}(U))\} \\ &\quad \cap \bigcap_{i=1}^p \{tb_i < \tau_i < t(b_i + \delta_2)\}. \end{aligned}$$

On the set on the right-hand side, we want to replace $\ell/\ell(U)$ by $\frac{1}{t^p} \ell_{tb} = \frac{1}{a} \ell_{tb}$. The difference is estimated as

$$(4.2) \quad \begin{aligned} \left| \frac{\ell}{\ell(U)} - \frac{\ell_{tb}}{a} \right| &= \left| \frac{\ell - \ell_{tb}}{\ell(U)} + \frac{1}{t^p} \ell_{tb} \left(\frac{a}{\ell(U)} - 1 \right) \right| \\ &\leq \frac{\ell_{t(b+\delta_2)} - \ell_{tb}}{t^p} + \frac{1}{t^p} \ell_{tb} \frac{\delta_1}{1 + \delta_1}. \end{aligned}$$

Pick some open set $\tilde{G} \subset \mathcal{M}(B)$ such that $G = \tilde{G} \cap \mathcal{M}(B)$. Fix $\varepsilon > 0$. Denote by $\tilde{G}_\varepsilon = \{\mu \in \tilde{G} : d(\mu, \tilde{G}^c) > \varepsilon\}$ the inner ε -neighborhood of \tilde{G} . Hence, for any $M > 0$, on the event $\{d(\frac{1}{t^p} \ell_{tb}, 0) < M\} \cap A$, where

$$(4.3) \quad A = \left\{ d \left(\frac{\ell_{t(b+\delta_2)} - \ell_{tb}}{t^p}, 0 \right) < \frac{\varepsilon}{2}, \ell_{t(b+\delta_2)}(U) - \ell_{tb}(U) \leq a \frac{\delta_1}{2} \right\},$$

we have, for sufficiently small $\delta_1, \delta_2 > 0$, that the event $\{\ell/\ell(U) \in G\}$ contains the event $\{\frac{1}{t^p}\ell_{tb} \in \tilde{G}_\varepsilon\}$. Thus we have the following lower bound:

$$\begin{aligned}
& \mathbb{P}\left(\frac{\ell}{\ell(U)} \in G, \ell(U) > a\right) \\
& \geq \mathbb{P}\left(\frac{1}{t^p}\ell_{tb} \in \tilde{G}_\varepsilon, a < \ell_{tb}(U) < a(1 + \frac{\delta_1}{2}), d(\frac{1}{t^p}\ell_{tb}, 0) < M, A, \right. \\
(4.4) \quad & \left. \forall i : tb_i < \tau_i < t(b_i + \delta_2)\right) \\
& = \mathbb{E}\left(1 \left\{ \frac{1}{t^p}\ell_{tb} \in \tilde{G}_\varepsilon, 1 < \frac{1}{t^p}\ell_{tb}(U) < 1 + \frac{\delta_1}{2}, d\left(\frac{1}{t^p}\ell_{tb}, 0\right) < M, \right. \right. \\
& \quad \left. \left. \forall i : tb_i < \tau_i \right\} F(W_{tb_1}^{(1)}, \dots, W_{tb_p}^{(p)})\right),
\end{aligned}$$

where we used the Markov property at times tb_1, \dots, tb_p and introduced

$$F(x) = \mathbb{P}_x\left(d\left(\frac{1}{t^p}\ell_{t\delta_2}, 0\right) < \frac{\varepsilon}{2}, \ell_{t\delta_2}(U) \leq t^p \frac{\delta_1}{2} \forall i : \tau_i < tb_i \delta_2\right);$$

we recall that \mathbb{P}_x denotes expectation with respect to the p motions starting in the sites x_1, \dots, x_p , respectively. It is easy to see, by choosing some appropriate joint strategy of the p motions, that $\liminf_{t \rightarrow \infty} \frac{1}{t} \log \inf_{x \in B^p} F(x) \geq 0$. To the remaining term on the right-hand side of (4.4), we can apply the lower bound in the LDP for $(t^p \prod_{i=1}^p b_i)^{-1} \ell_{tb}$ from Corollary 1.2 and obtain

$$\begin{aligned}
& \liminf_{a \rightarrow \infty} \frac{1}{a^{1/p}} \log \mathbb{P}\left(\frac{\ell}{\ell(U)} \in G, \ell(U) > a\right) \\
& \geq -\inf \left\{ \frac{1}{2} \sum_{i=1}^p b_i \|\nabla \psi_i\|_2^2 : \psi_i \in H_0^1(B), \|\psi_i\|_2 = 1 \forall i, \right. \\
& \quad \left. \prod_{i=1}^p (b_i \psi_i^2) \in \tilde{G}_\varepsilon, 1 < \int_U \prod_{i=1}^p (b_i \psi_i^2) < 1 + \frac{\delta_1}{2}, \right. \\
& \quad \left. d\left(\prod_{i=1}^p (b_i \psi_i^2), 0\right) < M \right\},
\end{aligned}$$

where we conceive the function $\prod_{i=1}^p (b_i \psi_i^2)$ as a measure on B .

Now let $M \rightarrow \infty$ to see that the last condition is immaterial, let $\delta_1 \downarrow 0$, substitute $\phi_i^2 = b_i \psi_i^2$, and take the supremum over b_1, \dots, b_p on the right-hand side

(i.e., drop the condition $\|\phi_i\|_2^2 = b_i$) to see that

$$\begin{aligned}
& \liminf_{a \rightarrow \infty} \frac{1}{a^{1/p}} \log \mathbb{P} \left(\frac{\ell}{\ell(U)} \in G, \ell(U) > a \right) \\
& \geq - \inf \left\{ \frac{1}{2} \sum_{i=1}^p \|\nabla \phi_i\|_2^2 : \phi_i \in H_0^1(B) \forall i, \prod_{i=1}^p \phi_i^2 \in \tilde{G}_\varepsilon, \right. \\
& \quad \left. 1 = \int_U \prod_{i=1}^p \phi_i^2 \right\} \\
& = - \inf_{\tilde{G}_\varepsilon} \tilde{J},
\end{aligned}$$

where \tilde{J} is the extension of J defined in (1.10) from $\mathcal{M}_U(B)$ to $\mathcal{M}(B)$ with $J(\mu) = \infty$ for $\mu \in \mathcal{M}(B) \setminus \mathcal{M}_U(B)$. Now let $\varepsilon \downarrow 0$ and use the lower semicontinuity of J to see that (4.1) holds. This concludes the proof of Lemma 4.1. \square

Now we handle the upper bound part.

LEMMA 4.2 (Upper Bound). *For every closed set $F \subset \mathcal{M}_U(B)$,*

$$(4.5) \quad \limsup_{a \uparrow \infty} \frac{1}{a^{1/p}} \log \mathbb{P} \left(\frac{\ell}{\ell(U)} \in F, \ell(U) > a \right) \leq - \inf_{\mu \in F} J(\mu).$$

PROOF. For any $R \in (0, \infty)$ and $\delta_1 \in (0, \infty)$, we have the following upper bound estimate:

$$\begin{aligned}
& \mathbb{P} \left(\frac{\ell}{\ell(U)} \in F, \ell(U) > a \right) \\
(4.6) \quad & \leq \sum_{j \in \mathbb{N} \cap [0, R/\delta_1]} \mathbb{P} \left(\frac{\ell}{\ell(U)} \in F, a(1 + (j-1)\delta_1) < \ell(U) \leq a(1 + j\delta_1) \right) \\
& \quad + \mathbb{P}(\ell(U) > aR).
\end{aligned}$$

The exponential rate of the second probability is known from [18] (see (1.4)):

$$(4.7) \quad \mathbb{P}(\ell(U) > aR) = \exp(-a^{1/p} R^{1/p} (\Theta_B(U) + o(1))),$$

where $\Theta_B(U) \in (0, \infty)$ is the variational formula appearing in (1.5).

With this in mind, let us now focus on one of the summands of the first term on the right-hand side of (4.6). By monotonicity in j , it is sufficient to consider the event for $j = 1$, as this gives the dominant term. Then, for any $\tilde{R} \in \mathbb{N}$ and

$\delta_2 \in (0, \infty)$,

$$\begin{aligned}
& \mathbb{P}\left(\frac{\ell}{\ell(U)} \in F, a < \ell(U) \leq a(1 + \delta_1)\right) \\
& \leq \sum_{b_1, \dots, b_p \in \delta_2 \mathbb{N} \cap [0, \tilde{R}]} \mathbb{P}\left(\frac{\ell}{\ell(U)} \in F, a < \ell(U) \leq a(1 + \delta_1), \right. \\
(4.8) \quad & \left. \forall i : a^{1/p} b_i < \tau_i \leq a^{1/p} (b_i + \delta_2)\right) \\
& + \sum_{i=1}^p \mathbb{P}(\tau_i > a^{1/p} \tilde{R}) + \sum_{i=1}^p \mathbb{P}(\ell(U) > a, \tau_i \leq a^{1/p} \delta_2).
\end{aligned}$$

The first probability on the last line has a strongly negative exponential rate for large \tilde{R} :

$$(4.9) \quad \mathbb{P}(\tau_i > a^{1/p} \tilde{R}) = \exp(-\tilde{R} a^{1/p} \lambda_1 + o(a^{1/p})), \quad a \uparrow \infty,$$

$\lambda_1 \in (0, \infty)$ being the principal eigenvalue of $-\frac{1}{2}\Delta$ in B with zero boundary condition. Furthermore, the last probability on the last line has a strongly negative exponential rate for small δ_2 , since

$$\begin{aligned}
(4.10) \quad & \lim_{\delta_2 \downarrow 0} \limsup_{a \uparrow \infty} \frac{1}{a^{1/p}} \log \mathbb{P}(\ell(U) > a, \tau_i \leq a^{1/p} \delta_2) = -\infty, \\
& i \in \{1, \dots, p\}.
\end{aligned}$$

This is shown as follows: For any $K \in (0, \infty)$, estimate

$$\begin{aligned}
& \mathbb{P}(\ell(U) > a, \tau_i \leq a^{1/p} \delta_2) \\
& \leq \mathbb{P}(\ell(U) > a, \tau_i \leq a^{1/p} \delta_2, \forall j \neq i : \tau_j \leq a^{1/p} K) \\
& + \sum_{j \neq i} \mathbb{P}(\tau_j > a^{1/p} K).
\end{aligned}$$

The last term has a very negative exponential rate for large K (see (4.9)), and for fixed K we estimate the first term on the right against $\mathbb{P}(\ell_{a^{1/p}v}(U) > a)$, where v is the vector in $(0, \infty)^p$ with δ_2 in the i^{th} component and K in all the other $p - 1$ components (we use the notation introduced in (1.1)). Now use the Markov inequality to estimate, for any $m \in \mathbb{N}$,

$$\begin{aligned}
\mathbb{P}(\ell_{a^{1/p}v}(U) > a) & \leq a^{-m} \mathbb{E}[\ell_{a^{1/p}v}(U)^m] \\
& \leq a^{-m} \mathbb{E}_0[\ell_{a^{1/p}v}(\mathbb{R}^d)^m] \\
& \leq a^{-m} \mathbb{E}_0[\ell_{a^{1/p}\delta_{21}}(\mathbb{R}^d)^m]^{1/p} \mathbb{E}_0[\ell_{a^{1/p}K_1}(\mathbb{R}^d)^m]^{(p-1)/p},
\end{aligned}$$

where we used the fact that the total mass of the intersection local time is stochastically larger if all the p motions start from the origin (see [4, (2.2.24)]) and Hölder's

inequality in the last step (see [4, (2.2.12)]); recall the notation $\mathbf{1} = (1, \dots, 1) \in \{1\}^p$. We have the bound

$$\begin{aligned} \mathbb{E}_0[\ell_{a^{1/p}\delta_2\mathbf{1}}(\mathbb{R}^d)^m] &= (a^{1/p}\delta_2)^{\frac{2p-d(p-1)}{2}m} \mathbb{E}_0[\ell_1(\mathbb{R}^d)^m] \\ &\leq m!^{\frac{d(p-1)}{2}} (a^{1/p}C_{\delta_2})^{\frac{2p-d(p-1)}{2}m} \end{aligned}$$

using Brownian scaling with some C_{δ_2} satisfying $\lim_{\delta_2 \downarrow 0} C_{\delta_2} = 0$. A similar bound holds for $\mathbb{E}_0[\ell_{a^{1/p}K_1}(\mathbb{R}^d)^m]$ (see [4, (2.2.22)] and the last display in the proof of [4, theorem 2.2.9]). Now pick $m \approx a^{1/p}$ and summarize the above facts to see that (4.10) holds.

Hence we focus on one of the summands of the first sum on the right-hand side of (4.8) for fixed $\delta_2, \tilde{R} \in (0, \infty)$. Set $t = a^{1/p}$ and $b = (b_1, \dots, b_p)$. We want to replace $\ell/\ell(U)$ by $\frac{1}{t^p}\ell_{tb}$. The difference is estimated as in (4.2) on the event $\{a < \ell(U) < a(1 + \delta_1)\} \cap \bigcap_{i=1}^p \{tb_i < \tau_i \leq t(b_i + \delta_2)\}$; this difference is small on the event $\{d(\frac{1}{t^p}\ell_{tb}, 0) \leq M\} \cap A$, with A as in (4.3), for any M and small δ_1 . Furthermore, note that, on the event $\bigcap_{i=1}^p \{tb_i < \tau_i \leq t(b_i + \delta_2)\}$,

$$(4.11) \quad \{a < \ell(U) < a(1 + \delta_1)\} \subset \{a - (\ell_{t(b+\delta_2\mathbf{1})}(U) - \ell_{tb}(U)) < \ell_{tb}(U) < a(1 + \delta_1)\}.$$

Fix $\varepsilon > 0$. Note that F is also closed in $\mathcal{M}(B)$. Denote the outer closed ε -neighborhood of F by $F_\varepsilon = \{\mu \in \mathcal{M}(B) : d(\mu, F) \leq \varepsilon\}$. Hence, for any $M > 0$, on the event $\{d(\frac{1}{t^p}\ell_{tb}, 0) \leq M\} \cap A$, we have, for sufficiently small $\delta_1 > 0$, that the event $\{\ell/\ell(U) \in F\}$ is contained in the event $\{\frac{1}{t^p}\ell_{tb} \in F_\varepsilon\}$, and furthermore we may estimate $\ell_{t(b+\delta_2\mathbf{1})}(U) - \ell_{tb}(U) \leq a\delta_1/2$ and use this on the right-hand side of (4.11). Thus,

$$(4.12) \quad \begin{aligned} &\mathbb{P}\left(\frac{\ell}{\ell(U)} \in F, a < \ell(U) \leq a(1 + \delta_1), \forall i : a^{1/p}b_i < \tau_i \leq a^{1/p}(b_i + \delta_2)\right) \\ &\leq \mathbb{P}\left(\frac{1}{t^p}\ell_{tb} \in F_\varepsilon, 1 - \frac{\delta_1}{2} < \frac{1}{t^p}\ell_{tb}(U) < 1 + \delta_1, \right. \\ &\quad \left. d\left(\frac{1}{t^p}\ell_{tb}, 0\right) \leq M, A, \forall i : \tau_i > tb_i\right) \\ &\quad + \mathbb{P}\left(d\left(\frac{1}{t^p}\ell_{tb}, 0\right) > M, \forall i : \tau_i > tb_i\right) + \mathbb{P}(A^c) \\ &\leq \mathbb{P}\left(\frac{1}{t^p}\ell_{tb} \in F_\varepsilon, 1 - \frac{\delta_1}{2} < \frac{1}{t^p}\ell_{tb}(U) < 1 + \delta_1, \forall i : \tau_i > tb_i\right) \\ &\quad + \mathbb{P}\left(d\left(\frac{1}{t^p}\ell_{tb}, 0\right) > M, \forall i : \tau_i > tb_i\right) \\ &\quad + \mathbb{P}\left(d\left(\frac{1}{t^p}(\ell_{t(b+\delta_2\mathbf{1})} - \ell_{tb}), 0\right) > \frac{\varepsilon}{2}\right) \\ &\quad + \mathbb{P}\left(\frac{1}{t^p}(\ell_{t(b+\delta_2\mathbf{1})}(U) - \ell_{tb}(U)) > \frac{\delta_1}{2}\right). \end{aligned}$$

Note that the exponential rates of the last three terms are strongly negative for large M and small δ_2 appearing in the respective terms. For the first of these this follows from an application of the LDP for $\frac{1}{\beta t^p} \ell_{tb}$ (with $\beta = \prod_{i=1}^p b_i$) from Corollary 1.2, where we note that large values of $d(\mu, 0)$ imply large values of $\mu(B)$. For the two latter terms, this follows from our proof of (4.10) (use the Markov property at times tb_1, \dots, tb_p , respectively).

For the first term on the right-hand side of (4.12), we put $\beta = \prod_{i=1}^p b_i$ and use the upper bound for the LDP of $\frac{1}{\beta t^p} \ell_{tb}$ from Corollary 1.2 and the continuity of the map $\mu \mapsto \mu(U)$ (recall that U is a Lebesgue-continuity set) to see that

$$\begin{aligned} & \limsup_{a \rightarrow \infty} \frac{1}{a^{1/p}} \log \mathbb{P} \left(\frac{1}{\beta t^p} \ell_{tb} \in \frac{F_\varepsilon}{\beta}, \frac{1 - \frac{\delta_1}{2}}{\beta} < \frac{1}{\beta t^p} \ell_{tb}(U) < \frac{1 + \delta_1}{\beta}, \right. \\ & \quad \left. \forall i : \tau_i > tb_i \right) \\ & \leq - \inf \left\{ \frac{1}{2} \sum_{i=1}^p b_i \|\nabla \psi_i\|_2^2 : \psi_i \in H_0^1(B), \|\psi_i\|_2 = 1 \forall i, \right. \\ & \quad \left. \prod_{i=1}^p \psi_i^2 \in \frac{F_\varepsilon}{\beta}, \frac{1 - \frac{\delta_1}{2}}{\beta} \leq \int_U \prod_{i=1}^p \psi_i^2 \leq \frac{1 + \delta_1}{\beta} \right\} \\ & \leq - \inf \left\{ \frac{1}{2} \sum_{i=1}^p \|\nabla \phi_i\|_2^2 : \phi_i \in H_0^1(B) \forall i, \prod_{i=1}^p \phi_i^2 \in F_\varepsilon, \right. \\ & \quad \left. 1 - \frac{\delta_1}{2} \leq \int_U \prod_{i=1}^p \phi_i^2 \leq 1 + \delta_1 \right\}, \end{aligned}$$

where we substituted $\phi_i^2 = b_i \psi_i^2$ and dropped the condition $\|\psi_i\|_2 = 1$. Now let $\delta_1 \downarrow 0$ and note that the right-hand side converges to $-\inf_{F_\varepsilon} \tilde{J}$, where \tilde{J} is the extension of J defined in (1.10) from $\mathcal{M}_U(B)$ to $\mathcal{M}(B)$ with $J(\mu) = \infty$ for $\mu \in \mathcal{M}(B) \setminus \mathcal{M}_U(B)$. By lower semicontinuity, this in turn tends to the right-hand side of (4.5). Collecting all preceding steps, this concludes the proof of Lemma 4.2. \square

Acknowledgment. The second author would like to thank the Courant Institute of Mathematical Sciences for its hospitality and Professor Varadhan for valuable discussions. He also thanks Prof. Peter Mörters for his help and encouragement as well as the support of the University of Bath, where part of this work was done.

Bibliography

- [1] Asselah, A.; Castell, F. Large deviations for Brownian motion in a random scenery. *Probab. Theory Related Fields* **126** (2003), no. 4, 497–527.

- [2] Bass, R. F. *Probabilistic techniques in analysis*. Probability and Its Applications (New York). Springer, New York, 1995.
- [3] Chen, X. Large deviations and laws of the iterated logarithm for the local times of additive stable processes. *Ann. Probab.* **35** (2007), no. 2, 602–648.
- [4] Chen, X. *Random walk intersections. Large deviations and related topics*. Mathematical Surveys and Monographs, 157. American Mathematical Society, Providence, R.I., 2010.
- [5] Dembo, A.; Zeitouni, O. *Large deviations techniques and applications*. 2nd ed. Applications of Mathematics (New York), 38. Springer, New York, 1998.
- [6] Donsker, M. D.; Varadhan, S. R. S. Asymptotic evaluation of certain Markov process expectations for large time.I. *Comm. Pure Appl. Math.* **28** (1975), 1–47.
- [7] Donsker, M. D.; Varadhan, S. R. S. Asymptotic evaluation of certain Markov process expectations for large time.II. *Comm. Pure Appl. Math.* **28** (1975), 279–301.
- [8] Donsker, M. D.; Varadhan, S. R. S. Asymptotic evaluation of certain Markov process expectations for large time. III. *Comm. Pure Appl. Math.* **29** (1976), no. 4, 389–461.
- [9] Donsker, M. D.; Varadhan, S. R. S. Asymptotic evaluation of certain Markov process expectations for large time. IV. *Comm. Pure Appl. Math.* **36** (1983), no. 2, 183–212.
- [10] Dvoretzky, A.; Erdős, P.; Kakutani, S. Double points of paths of Brownian motion in n -space. *Acta Sci. Math. Szeged* **12** (1950), 75–81.
- [11] Dvoretzky, A.; Erdős, P.; Kakutani, S. Multiple points of paths of Brownian motion in the plane. *Bull. Res. Council Israel* **3** (1954), 364–371.
- [12] Dvoretzky, A.; Erdős, P.; Kakutani, S.; Taylor, S. J. Triple points of Brownian paths in 3-space. *Proc. Cambridge Philos. Soc.* **53** (1957), 856–862.
- [13] den Hollander, F. Large deviations. Fields Institute Monographs, 14. American Mathematical Society, Providence, R.I., 2000.
- [14] Fristedt, B. An extension of a theorem of S. J. Taylor concerning the multiple points of the symmetric stable process. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **9** (1967), 62–64.
- [15] Gärtner, J. On large deviations from an invariant measure. *Teor. Veroyatnost. i Primenen.* **22** (1977), no. 1, 27–42; translation in *Theory Probab. Appl.* **22** (1977), 24–39. Available at: <http://epubs.siam.org/tvp/resource/1/tprbau/v22/i1>
- [16] Geman, D.; Horowitz, J.; Rosen, J. A local time analysis of intersections of Brownian motion in the plane. *Ann. Probab.* **12** (1984), no. 1, 86–107.
- [17] Grieser, D. Uniform bounds for eigenfunctions of the Laplacian on manifolds with boundary. *Comm. Partial Differential Equations* **27** (2002), no. 7-8, 1283–1299.
- [18] König, W.; Mörters, P. Brownian intersection local times: upper tail asymptotics and thick points. *Ann. Probab.* **30** (2002), no. 4, 1605–1656.
- [19] König, W.; Mörters, P. Brownian intersection local times: exponential moments and law of large masses. *Trans. Amer. Math. Soc.* **358** (2006), no. 3, 1223–1255 (electronic).
- [20] Le Gall, J.-F. Sur la saucisse de Wiener et les points multiples du mouvement brownien. *Ann. Probab.* **14** (1986), no. 4, 1219–1244.
- [21] Le Gall, J.-F. The exact Hausdorff measure of Brownian multiple points. *Seminar on stochastic processes, 1986 (Charlottesville, Va., 1986)*, 107–137. Progress in Probability and Statistics, 13. Birkhäuser Boston, Boston, 1987.
- [22] Le Gall, J.-F. The exact Hausdorff measure of Brownian multiple points. II. *Seminar on stochastic processes, 1988 (Gainesville, FL, 1988)*, 193–197. Progress in Probability, 17. Birkhäuser Boston, Boston, 1989.
- [23] Lieb, E. H.; Loss, M. *Analysis*. 2nd ed. Graduate Studies in Mathematics, 14. American Mathematical Society, Providence, R.I., 2001.
- [24] Taylor, S. J. The exact Hausdorff measure of the sample path for planar Brownian motion. *Proc. Cambridge Philos. Soc.* **60** (1964), 253–258.

WOLFGANG KÖNIG
Technische Universität Berlin
Str. des 17. Juni 136
10623 Berlin
and
Weierstrass Institute Berlin,
Mohrenstr. 39
10117 Berlin
GERMANY
E-mail: koenig@
wias-berlin.de

Received May 2011.

CHIRANJIB MUKHERJEE
Max-Planck Institute
for Mathematics in the Sciences
Inselstr. 22
04103 Leipzig
GERMANY
E-mail: mukherje@mis.mpg.de