

## Exercise Sheet 11

### Exercise 41. Elliptic and parabolic equation via Fourier transform in $\mathbb{R}^d$ .

(a) For  $A \in \mathbb{R}^{d \times d}$  symmetric and positive definite,  $b \in \mathbb{R}^d$ , and  $c \in ]0, \infty[$  show that the elliptic equation

$$-\sum_{i,j=1}^d A_{ij} \frac{\partial^2}{\partial x_i \partial x_j} u + \sum_{k=1}^d b_k \frac{\partial}{\partial x_k} u + c u = f \text{ auf } \mathbb{R}^d$$

has a unique solution  $u \in H^2(\mathbb{R}^d)$  for each  $f \in L^2(\mathbb{R}^d)$ .

(c) For  $c > 0$  consider the parabolic equation

$$\frac{\partial}{\partial t} u = \Delta u - c u + f \text{ for } (t, x) \in \mathbb{R}^{1+d}.$$

Show that for each  $f \in L^2(\mathbb{R}^{1+d})$  there exists a unique solution  $u \in L^2(\mathbb{R}^{1+d})$ , which satisfies  $\frac{\partial}{\partial t} u, \frac{\partial}{\partial x_j} u, \frac{\partial^2}{\partial x_i \partial x_j} u \in L^2(\mathbb{R}^{1+d})$ .

**Exercise 42 (in written form). Cea's Lemma** (JEAN CÉA 1962) Consider a symmetric, bounded and coercive bilinear form  $B : H \times H \rightarrow \mathbb{R}$  in a real Hilbert space  $H$ :

$$B(u, v) = B(v, u), \quad |B(u, v)| \leq C_1 \|u\| \|v\|, \quad B(u, u) \geq c_0 \|u\|^2.$$

(a) Let the approximation space  $H_N$  be a closed subspace of  $H$ . For  $F \in H^* = \text{Lin}(H, \mathbb{R})$  consider the solution  $u \in H$  and the approximation  $u_N \in H_N$  to be defined via

$$B(u, v) = F(v) \text{ for all } v \in H \quad \text{and} \quad B(u_N, v_N) = F(v_N) \text{ for all } v_N \in H_N,$$

respectively. Show that  $u_N$  is uniquely defined and satisfies the Cea estimate

$$\|u - u_N\| \leq \sqrt{C_1/c_0} \inf \{ \|u - w_N\| \mid w_N \in H_N \}, \quad (\text{Céa})$$

i.e.,  $u_N$  is, up to a constant, as good as the best approximant in  $H_N$ .

Hint: Show  $B(u - u_N, u - u_N) + B(w_N - u_N, w_N - u_N) = B(u - w_N, u - w_N)$ .

(b) Let  $H = H_0^1(]0, \pi[)$  and  $H_N = \text{span}\{\sin(k \cdot) \mid k = 1, \dots, N\}$ . For  $a \in C^1([0, \pi])$  with  $a(x) \in [1, 2]$  and  $f \in L^2(]0, \pi[)$  we set  $B(u, v) = \int_0^\pi a(x) u'(x) v'(x) dx$  and  $F(v) = \int_0^\pi f(x) v(x) dx$ . For  $u$  and  $u_N$  be as in (a) provide the estimates

$$\|u_N - u\|_{H^1} \leq \frac{1}{N} C_2 \|u\|_{H^2} \quad \text{and} \quad \|u\|_{H^2} \leq C_a \|f\|_{L^2}.$$

Hint: Combine Exercise 25 and (Céa).

please turn

**Exercise 43. Resolvents and eigenvalues of elliptic operators.** Consider a bounded Lipschitz domain  $\Omega \subset \mathbb{R}^d$ . Further,  $A \in L^\infty(\Omega; \mathbb{R}_{\text{sym}}^{d \times d})$ ,  $a \in L^\infty(\Omega; \mathbb{R}^d)$  and  $\alpha \in L^\infty(\Omega)$  are given such that for  $\delta > 0$  we have  $A(x)\xi \cdot \xi \geq \delta|\xi|^2$  for all  $x \in \Omega$  and all  $\xi \in \mathbb{R}^d$ . On the complex Hilbert space  $H_0^1(\Omega) \oplus iH_0^1(\Omega)$  we define the sesquilinear form

$$B(u, v) = \int_{\Omega} A(x) \nabla u(x) \cdot \overline{\nabla v(x)} + a(x) \cdot \nabla u(x) \overline{v(x)} + \alpha(x) u(x) \overline{v(x)} dx$$

and call  $\lambda \in \mathbb{C}$  an eigenvalue of  $B$ , if there exists  $u \in H_0^1(\Omega) \setminus \{0\}$  such that

$$B_\lambda(u, v) := B(u, v) - \lambda \int_{\Omega} u \bar{v} dx = 0 \quad \text{for all } v \in H_0^1(\Omega).$$

If for some  $\lambda \in \mathbb{C}$  there exists  $c_0 > 0$  such that  $|B_\lambda(u, u)| \geq c_0 \|u\|_{H^1}^2$ , then  $\lambda$  lies in the resolvent set of  $B$ . Our aim is to show that the resolvent set is large.

(a) Find  $C_1 > 0$ ,  $c_0 > 0$ , and  $\lambda_*$  such that  $|B(u, v)| \leq C_1 \|u\|_{H^1} \|v\|_{H^1}$  and  $B_\lambda(u, u) \geq c_0 \|u\|_{H^1}^2$  holds for all  $u, v \in H_0^1(\Omega)$  and  $\lambda \leq \lambda_*$ .

(b) Construct positive numbers  $\beta, \gamma$  such that the set  $\{\lambda \in \mathbb{R} \mid |\text{Im } \lambda| \geq \gamma(\text{Re } \lambda + \beta)\}$  is contained in the resolvent set. For this consider  $B_{\theta, \lambda}(u, v) = e^{i\theta} B_\lambda(u, v)$  and bound  $|B_{\theta, \lambda}(u, u)|$  from below via  $\text{Re}(B_{\theta, \lambda}(u, u))$ .

**Exercise 44. A bilinear form with Robin boundary conditions** Consider the two-dimensional domain  $\Omega = ]0, \pi[^2$ , the Dirichlet boundary  $\Gamma_D = \{(x_1, x_2) \mid x_1 \in \{0, \pi\}, x_2 \in ]0, \pi[ \}$  and the symmetric bilinear form

$$B(u, v) = \int_{\Omega} \nabla u(x) \cdot \nabla v(x) + \alpha u(x) v(x) dx + \int_{\partial\Omega \setminus \Gamma_D} \rho u(x) v(x) da \quad \text{with } \alpha, \rho \in \mathbb{R}$$

on the Hilbert space  $H_{\Gamma_D}^1(\Omega) = \{u \in H^1(\Omega) \mid u|_{\Gamma_D} = 0 \text{ in the trace sense}\}$ .

(a) Show that for  $\alpha > 0$  and  $\rho \geq 0$  the bilinear form is bounded and coercive. For the weak problem  $B(u, v) = \int_{\Omega} f v dx$  write the associated linear problem  $L_{\alpha, \rho} u = f$  in strong form.

(b) Let  $\alpha = \rho = 0$  and find all eigenfunctions  $u$  for  $L_{0,0}$  (i.e.  $L_{0,0} u = \lambda u$ ), which have the form  $u(x) = \psi(x_1) \phi(x_2)$ . Is the set of these eigenfunctions a complete ONS?