# Weierstraß–Institut für Angewandte Analysis und Stochastik

im Forschungsverbund Berlin e.V.

## On the existence of unbiased Monte Carlo estimators

## Peter Mathé

submitted: 10th August 1994

Weierstrass Institute for Applied Analysis and Stochastics Mohrenstraße 39 D - 10117 Berlin Germany

Preprint No. 113 Berlin 1994

1991 Mathematics Subject Classification. 65C05, Secondary 47B10. Key words and phrases. Unbiased Monte Carlo methods, operator ideals.

Edited by Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS) Mohrenstraße 39 D — 10117 Berlin Germany

Fax: + 49 30 2004975 e-mail (X.400): c=de;a=d400;p=iaas-berlin;s=preprint e-mail (Internet): preprint@iaas-berlin.d400.de

## ON THE EXISTENCE

#### OF

### UNBIASED MONTE CARLO ESTIMATORS

#### PETER MATHÉ

#### August 10, 1994

ABSTRACT. For many typical instances where Monte Carlo methods are applied attempts were made to find unbiased estimators, since for them the Monte Carlo error reduces to the statistical error. These problems usually take values in the scalar field. If we study vector valued Monte Carlo methods, then we are confronted with the question whether there can exist unbiased estimators. This problem is apparently new. Below it is settled precisely. Partial answers are given, indicating relations to several classes of linear operators in Banach spaces.

#### 1. INTRODUCTION, NOTATION

In many practical applications the program designer is confronted with the "curse of dimensionality", an exponential dependence on the dimension, which is inherent in most error estimates provided by classical numerical analysis, see e.g. [TWW88] for a sample of typical numerical problems and the respective error estimates.

Often this can be overcome by choosing Monte Carlo methods, i.e., numerical methods involving random parameters in the computational process, see [HH64] for an excellent, by now classical treatment on the applicability of Monte Carlo methods. Within the classical theory one prefers unbiased Monte Carlo estimators, since they are self-focusing if the numerical simulation is repeated. Such unbiased estimators are often hard to find, see [KW86, ENS89].

In statistics, the problem whether there are unbiased estimates (in statistical sense) has also been attractive, recently. While it has been known since 1956 that there cannot exist such unbiased estimates for density estimation, see [Ros56], advantage has been made by [LB93]. They developed a machinery which enables to prove that there are no unbiased *l*-informative estimates for singular problems. Within our framework, the notion of *l*-informativity corresponds to the requirement of finite errors. On the other hand, all problems studied below are not singular in their sense. This may express that our

<sup>1991</sup> Mathematics Subject Classification. 65 C 05, Secondary 47 B 10.

Key words and phrases. unbiased Monte Carlo methods, operator ideals.

arguments are completely different, indicating geometric properties of the target space, where the error is measured.

Below we are concerned with vector valued Monte Carlo methods for the randomized approximation of *linear* mappings. The classical results extend easily from real valued Monte Carlo methods to a finite number of them, if we gather unbiased estimates for each component. The problem is how far one can proceed by this: Are there infinite dimensional problems, which admit unbiased Monte Carlo estimators?

As an illustration let us briefly and informally introduce the following

Example 1. Let f be any periodic function defined on the interval  $[0, 2\pi]$ , which has a (generalized) derivative in  $\tilde{L}_2(0, 2\pi)$ , the space of all periodic square integrable functions. It is well known that any such function expands into a Fourier series (converging in  $\tilde{L}_2(0, 2\pi)$ ), which means, that we have

(1) 
$$f(x) = \sum_{k=1}^{\infty} \gamma_k(f) e^{-ikx}, \quad x \in [0, 2\pi],$$

with Fourier coefficients

(2) 
$$\gamma_k(f) := \frac{1}{2\pi} \int_0^{2\pi} f(\xi) e^{-ik\xi} d\xi.$$

For such functions an unbiased Monte Carlo estimator (using Fourier coefficients) can be obtained in the following (trivial) way by choosing a member of the Fourier series at random. Precisely, let  $p_k := \frac{6}{\pi^2 k^2}$ ,  $k \in \mathbb{N}$ , and choose  $\frac{\pi^2 k^2}{6} \gamma_k(f) e^{-ikx}$ ,  $k \in \mathbb{N}$ , with probability  $p_k$ , respectively. The average performance of this method yields the Fourier series of f, provided the random variable

$$k \longrightarrow \frac{\pi^2 k^2}{6} \gamma_k(f) e^{-ikx}$$

taking values in  $L_2(0, 2\pi)$  is integrable. But we even have the stronger square integrability from

(3) 
$$\sum_{k=1}^{\infty} p_k \frac{\pi^4 k^4}{36} |\gamma_k(f)|^2 ||e^{-ikx}||_2^2 = \frac{\pi^2}{6} \sum_{k=1}^{\infty} k^2 |\gamma_k(f)|^2 < \infty,$$

for functions with square integrable derivative, see [Pie87].

The above example can be considered as a special instance of approximating a diagonal mapping in the space  $l_2$  of square summable sequences. To see this we switch from the spaces of functions to the respective spaces of Fourier coefficients in the following way. Assign any function f the sequence  $x_k := k\gamma_k(f)$ ,  $k \in \mathbb{N}$ . Then the approximation of f is replaced by the approximation of  $(\gamma_k(f))_{k\in\mathbb{N}}$ . The differentiability assumption ensures that  $\sum_{k=1}^{\infty} |x_k|^2 < \infty$ . This means that we assign every sequence  $(x_k)_{k\in\mathbb{N}}$  the sequence  $(\frac{1}{k}x_k)_{k\in\mathbb{N}}$ , corresponding to a diagonal mapping  $D_{\sigma}: l_2 \to l_2$ .

Summarizing we switched from the random approximation of functions to the random approximation of a diagonal mapping between Hilbert spaces. Within this framework a theory of stochastic numerical methods is available, see [Mat91, Hei94]. The basic notion will be the notion of a (linear) Monte Carlo method. Throughout the paper we restrict

ourselves to linear Monte Carlo methods only. Let X and Y be Banach spaces. Denote by  $\mathfrak{L}(X,Y)$  the space of all bounded linear operators and by  $\mathfrak{F}(X,Y)$  the subspace of all operators of finite rank, cf. [Pie80, Pie87] for notation from the theory of operators in Banach spaces. By  $\mathfrak{F}^k(X,Y)$  we denote the subset of operators of rank at most k. Corresponding to [Mat91, Hei94, Mat94] we propose the following

Definition 1. A triple

$$\mathcal{P} := ([\Omega, \mathcal{F}, P], \mathbf{u}, \mathbf{k})$$

is called a *(linear)* Monte Carlo method, if

(1)  $[\Omega, \mathcal{F}, P]$  is a probability space.

(2) u:  $\Omega \to \mathfrak{F}(X,Y)$  is such that the mapping  $\Phi: X_0 \times \Omega \to Y$ , defined by

$$\Phi(x,\omega):=(\mathrm{u}(\omega))(x),\quad x\in X,\quad \omega\in\Omega,$$

is product measurable into Y and the set  $\{(u(\omega))(x), x \in X, \omega \in \Omega\}$  is a separable subset in Y.

(3) The cardinality function  $k: \Omega \to \mathbb{N}$  is a measurable natural number, for which

$$u_{\omega} := u(\omega) \in \mathfrak{F}^{k(\omega)}(X, Y), \quad \omega \in \Omega.$$

Remark 1. For linear Monte Carlo methods as introduced above we could directly assign  $k(\omega) := \operatorname{rank}(u(\omega))$ , since this would result in an appropriate measurable cardinality function. However, for general classes of methods such assignment would not be meaningful, such that we kept the definition with the more general choice of cardinality function, see [Hei94, Mat94].

For such Monte Carlo methods we can assign the cardinality

$$\operatorname{MC-card}(\mathcal{P}) := \int_{\Omega} \mathrm{k}(\omega) dP(\omega).$$

By definition, for every  $x \in X$  the mapping

$$\omega \longrightarrow \mathrm{u}_\omega(x)$$

is a Radon random variable in Y, see [LT91] for spaces of Banach space valued random variables. Therefore the error of any Monte Carlo method  $\mathcal{P} := ([\Omega, \mathcal{F}, P], u, k)$  for an operator  $S: X \to Y$  at input  $x \in X$  may be defined as

$$e(S,\mathcal{P},x):=\left(\int_{\Omega}\|S(x)-\mathrm{u}_{\omega}(x)\|_{Y}^{2}dP(\omega)
ight)^{1/2},$$

while the overall performance is given by

$$e(S,\mathcal{P}):=\sup_{||x||_X\leq 1}e(S,\mathcal{P},x).$$

We agree to denote by

$$a_n^{mc}(S) := \inf \left\{ \sup_{\|x\|_X \le 1} \left( \int_{\Omega} \|S(x) - u_{\omega}(x)\|_Y^2 dP(\omega) \right)^{1/2}, \quad \text{MC-card}(\mathcal{P}) < n \right\}$$

the *nth Monte Carlo approximation number* of the linear operator  $S: X \to Y$ , see [Mat91] for more information on that topic. We shall make use of the following submultiplicativity property

(4) 
$$a_n^{mc}(RST) \le ||R|| a_n^{mc}(S) ||T||,$$

whenever the product is correctly defined.

If a Monte Carlo method  $\mathcal{P}$  has a finite error for some linear operator S, i.e.,  $e(S, \mathcal{P}) < \infty$ , then the function

$$S_{\mathcal{P}}(x) := \int_{\Omega} \mathtt{u}_{\omega}(x) dP(\omega)$$

exists and denotes the respective expectation. From now on we make the assumption that

$$\sup_{\|x\|_{X}\leq 1}\int_{\Omega}\|u_{\omega}(x)\|_{Y}^{2}dP(\omega)<\infty,$$

ensuring that  $e(0, \mathcal{P}) < \infty$ , where 0 denotes the zero operator. It is the aim of this paper to study properties of  $S_{\mathcal{P}}$ .

Definition 2. An operator  $S \in \mathfrak{L}(X, Y)$  admits an unbiased Monte Carlo method if there is a Monte Carlo method  $\mathcal{P}$  with

(1) 
$$MC\operatorname{-card}(\mathcal{P}) < \infty,$$
(2) 
$$\sup_{\|x\|_X \le 1} \int_{\Omega} \|u_{\omega}(x)\|_Y^2 dP(\omega) < \infty$$

and

$$S(x) = S_{\mathcal{P}}(x), \quad x \in X.$$

If  $S \in \mathfrak{L}(X, Y)$  admits an unbiased Monte Carlo method then we let

(5) 
$$u(S) := \inf \left\{ \sup_{\|x\|_X \leq 1} \left( \int_{\Omega} \|u_{\omega}(x)\|_Y^2 dP(\omega) \right)^{1/2}, \quad \mathcal{P} \text{ is unbiased for } S \right\}.$$

This turns into a norm and we have  $||S: X \to Y|| \le u(S)$ . As explained above, property (2) is equivalent to the statement that  $\mathcal{P}$  has a finite error for some bounded operator acting between X and Y. Within this framework it is immediate that any finite rank operator L admits an unbiased Monte Carlo method by letting  $\mathcal{P}$  be choosing L with probability 1. However, there are different ways of representing a given finite dimensional mapping, leading to different Monte Carlo methods.

Example 2. Let  $\mathrm{Id}^m : \mathbb{R}^m \to \mathbb{R}^m$  be the identity and  $e_j, j = 1, \ldots, m$ , denote the unit vector basis in  $\mathbb{R}^m$ .

1. (trivial) representation:

$$\mathrm{Id}^{m}(x) = \frac{1}{m} \sum_{j=1}^{m} (mx_{j})e_{j}, \quad x = (x_{1}, \ldots, x_{m}).$$

#### 2. (nuclear) representation:

$$\mathrm{Id}^{m}(x) = \frac{1}{2^{m}} \sum_{e_{1}, \dots, e_{m} = \pm 1} \left( \sum_{j=1}^{m} \varepsilon_{j} x_{j} \right) \left( \sum_{j=1}^{m} \varepsilon_{j} e_{j} \right), \quad x = (x_{1}, \dots, x_{m}).$$

where  $\varepsilon_1, \ldots, \varepsilon_m$  denotes a Bernoulli sequence, see Remark 2 below. Observe that it is much more elaborate to find unbiased Monte Carlo methods with prescribed properties. We will not turn to that problem.

## 2. Diagonal mappings $D_{\sigma} \colon l_2 \to l_q, \quad 1 \leq q \leq \infty$

We are going to study diagonal mappings in sequence spaces. To make things precise we need to introduce the Lorentz sequence spaces, cf. [Pie87, 2.1]. For any sequence  $x = (\xi_j)_{j \in \mathbb{N}}$  of real numbers, which is convergent to 0 we assign by  $s_n(x)$  the non-increasing rearrangement (in modulus) of x. The (real) Lorentz space  $l_{r,w}$  consists of all sequences xfor which the sequence  $(n^{1/r-1/w}s_n(x))_{n \in \mathbb{N}}$  belongs to  $l_w$  equipped with the norm arising from this identification. Explicitly, if  $1 \leq w < \infty$  then we let

$$\|x\|_{r,w} := \left(\sum_{j=1}^{\infty} [n^{1/r-1/w}s_n(x)]^w\right)^{1/w}$$

while for  $w = \infty$  we put

$$||x||_{r,\infty} := \sup_{n\in\mathbb{N}} \left\{ n^{1/r} s_n(x) \right\}.$$

Observe that we can identify for any  $1 \leq p < \infty$  the classical sequence spaces  $l_p$  with  $l_{p,p}$ , while we denote by  $c_0$  the space  $l_{\infty,\infty}$  and by  $l_{\infty}$  the space of all bounded sequences equipped with the supremum norm.

We begin our study with the consideration of a specific class of diagonal operators. Given any sequence  $\sigma = (\sigma_j)_{j \in \mathbb{N}}$  of real numbers, let us consider the mapping  $x = (x_j)_{j \in \mathbb{N}} \to (\sigma_j x_j)_{j \in \mathbb{N}}$ . Denote this mapping by  $D_{\sigma}$ . For choices of  $1 \leq p, q \leq \infty$  the operator  $D_{\sigma}$  acts continuously from  $l_p$  to  $l_q$  if and only if the diagonal  $\sigma$  belongs to  $l_r$  with  $\frac{1}{r} = \max\left\{0, \frac{1}{\sigma} - \frac{1}{2}\right\}$ .

The simple construction outlined in the introductory section implies the existence of unbiased Monte Carlo methods for  $D_{\sigma}: l_2 \rightarrow l_2$  whenever  $\sigma \in l_2$ , which is more than required for continuity. This is typical and we shall derive necessary conditions later.

However, the class of operators admitting an unbiased Monte Carlo method can be enlarged for other spaces, employing a more involved construction. This is provided in the following

**Theorem 1.** A diagonal mapping  $D_{\sigma} \colon l_2 \to l_q$  admits an unbiased Monte Carlo method if  $\sigma \in l_q$ . Moreover we have  $u(D_{\sigma} \colon l_2 \to l_q) \leq ||\sigma||_q$ .

Before proving the theorem we need a preparatory lemma. Let  $\varepsilon^m$  denote the uniform distribution on the extreme points of  $B_{\infty}^m$ , i.e., all vectors  $\omega \in \{+1, -1\}^m$ . Thus the basic probability space is  $[\{+1, -1\}^m, \mathcal{F}^m, \varepsilon^m]$ . For computations involving  $\varepsilon^m$  the following lemma is useful.

Lemma 1. Let m > 1.

(1) For all  $1 \leq j, k \leq m$  we have

$$\int_{\{+1,-1\}^m} \langle \omega, e_j \rangle \langle \omega, e_k \rangle d\varepsilon^m(\omega) = \begin{cases} 1 & , \text{if } j = k \\ 0 & , \text{else} \end{cases}$$

(2) For all  $a \in \mathbb{R}^m$  we have

$$\left(\int_{\{+1,-1\}^m} |\langle \omega,a\rangle|^2 d\varepsilon^m(\omega)\right)^{1/2} = ||a||_2.$$

*Proof.* The second statement is a consequence of the first one. To prove the first topic, let us observe that for  $\omega \in \{\pm 1, -1\}^m$  and  $1 \leq j \leq m$  the number  $\langle \omega, e_j \rangle$  is  $\pm 1$ . This implies  $\int_{\{\pm 1, -1\}} \langle \omega, e_j \rangle^2 d\varepsilon^m(\omega) = 1$ . If  $j \neq k$  then

$$\begin{array}{l} \operatorname{card} \{\omega \in \{+1,-1\}^{m}, \quad \langle \omega, e_{j} \rangle \langle \omega, e_{k} \rangle = 1\} = \\ \\ \operatorname{card} \{\omega \in \{+1,-1\}^{m}, \langle \omega, e_{j} \rangle \langle \omega, e_{k} \rangle = -1\}, \end{array}$$

which is equal to  $2^{m-1}$ , and from which the proof can be completed.  $\Box$ 

Remark 2. The above lemma is well known. The reader familiar with probability theory will recognize that the distribution of  $\langle \omega, a \rangle$  is the distribution of the sum  $\sum_{j=1}^{m} \varepsilon_j \langle e_j, a \rangle$ , where the  $\varepsilon_j$  are independent numbers taking values +1 and -1 with equal probability, i.e., they form a Bernoulli sequence.

Proof of Theorem 1. The proof is constructive. We are going to design a concrete Monte Carlo method of cardinality 1. Let  $\Omega^m := \{+1, -1\}^m$  equipped with  $\varepsilon^m$ , introduced above as probability. Denote by  $\Omega := \bigcup_{m \in \mathbb{N}} \Omega^m$  the disjoint union, equipped with the  $\sigma$ -algebra  $\mathcal{F}$ , generated from the sequence  $\mathcal{F}^m$ . A probability P will be given as mixture of  $\varepsilon^m$  in the following way.

Observe, that for  $\varepsilon > 0$  we can find a decreasing sequence  $(\tau_m)_{m \in \mathbb{N}}$ , such that  $\tau_1 = 1$  and  $\sigma_m = \tau_m \beta_m$ ,  $m \in \mathbb{N}$ ,  $\lim_{m \to \infty} \tau_m = 0$  and  $\sum_{j=1}^{\infty} |\beta_j|^q \leq (1+\varepsilon)^q \sum_{j=1}^{\infty} |\sigma_j|^q$ . For a proof of this fact we refer to [Pie80, 8.6.4]. Let  $p_m := \tau_m - \tau_{m+1}$ ,  $m \in \mathbb{N}$ . Then we have  $p_m \geq 0$  and  $\sum_{m=1}^{\infty} p_m = \tau_1 = 1$ , which means that this sequence gives rise to a probability P by letting  $P := \sum_{m=1}^{\infty} p_m \varepsilon^m$ . (We implicitly extended the probabilities  $\varepsilon^m$  to all  $\Omega$ .) So far we have defined a probability space  $[\Omega, \mathcal{F}, P]$ . Now, define a mapping  $u: \Omega \to \mathfrak{F}(l_2, l_q)$  by

$$\mathrm{u}_\omega(x):=\left(\sum_{j=1}^m\omega_j\langle x,e_j
angle
ight)\left(\sum_{j=1}^m\omega_jeta_je_j
ight),\quad\omega=(\omega_1,\ldots,\omega_m),\quad x\in l_2.$$

It is readily checked that  $\mathcal{P} = ([\Omega, \mathcal{F}, P], u, 1)$  is a linear Monte Carlo method of cardina-

lity 1. Moreover, for  $x \in l_2$  we have by Lemma 1

$$\begin{split} \int_{\Omega} \| \mathbf{u}_{\omega}(x) \|_{q}^{2} dP(\omega) &= \sum_{m=1}^{\infty} p_{m} \int_{\Omega^{m}} \| \mathbf{u}_{\omega}(x) \|_{q}^{2} d\varepsilon^{m}(\omega) \\ &= \sum_{m=1}^{\infty} p_{m} \int_{\Omega^{m}} |\sum_{j=1}^{m} \omega_{j} \langle x, e_{j} \rangle |^{2} \| \sum_{j=1}^{m} \omega_{j} \beta_{j} e_{j} \|_{q}^{2} d\varepsilon^{m}(\omega) \\ &\leq \sum_{m=1}^{\infty} p_{m} \| x \|_{2}^{2} \left( \sum_{j=1}^{m} |\beta_{j}|^{q} \right)^{2/q} \\ &\leq (1+\varepsilon)^{2} \| x \|_{2}^{2} \left( \sum_{j=1}^{\infty} |\sigma_{j}|^{q} \right)^{2/q}, \end{split}$$

such that  $u(D_{\sigma}: l_2 \to l_q) \leq \left(\sum_{j=1}^{\infty} |\sigma_j|^q\right)^{1/q}$ , provided  $\mathcal{P}$  was unbiased for  $D_{\sigma}$ . But this is true, since

$$T(x):=\int_{\Omega}\mathrm{u}_{\omega}(x)dP(\omega),\quad x\in l_2,$$

is well-defined and equal to

$$T(x) = \sum_{m=1}^{\infty} p_m \int_{\Omega^m} \left( \sum_{j=1}^m \omega_j \langle x, e_j \rangle \right) \left( \sum_{j=1}^m \omega_j \beta_j e_j \right) d\varepsilon^m(\omega)$$
  
$$= \sum_{m=1}^{\infty} p_m \sum_{j=1}^m \beta_j \langle x, e_j \rangle e_j$$
  
$$= \sum_{j=1}^{\infty} \beta_j \left( \sum_{m=j}^{\infty} p_m \right) \langle x, e_j \rangle e_j$$
  
$$= \sum_{j=1}^{\infty} \beta_j \tau_j \langle x, e_j \rangle e_j$$
  
$$= D_{\sigma}(x),$$

again by Lemma 1. The proof of the theorem is complete.  $\Box$ 

Analogously we could prove

Corollary 1. A diagonal mapping  $D_{\sigma} : l_2 \to l_{\infty}$  admits an unbiased Monte Carlo method provided the diagonal tends to 0.

All operators considered so far have been compact. So the question arises, whether this is typical and leads to the following

**Open Problem.** Does the identity Id:  $l_2 \rightarrow l_{\infty}$  admit an unbiased Monte Carlo method?

Below we shall turn to the question whether there are necessary conditions for the existence of unbiased Monte Carlo methods. Such conditions will be expressed in terms of the Monte Carlo approximation numbers, introduced above. Therefore we are looking

for a reformulation of Theorem 1. To do this we need the following result, which points at the relation of the ordinary approximation numbers to the Monte Carlo counterpart, see [Mat91, Lemma 5]. Topics (2) and (3), below provide the typical instances where the Monte Carlo and the (ordinary) approximation numbers differ.

**Proposition 1.** There is a constant  $C < \infty$  such that for all  $m, n \in \mathbb{N}$  we have

(1)  $a_n^{mc}(\mathrm{Id}: l_2^m \to l_q^m) \leq C \frac{m^{1/q}}{n^{1/q'}},$ (2)  $a_n^{mc}(\mathrm{Id}: l_2^m \to l_q^m) \leq C \frac{m^{1/q}}{n^{1/2}},$ (3)  $a_n^{mc}(\mathrm{Id}: l_2^m \to l_\infty^m) \leq C \sqrt{\frac{\log(1+m)}{n}}.$ 

For the convenience of the reader we shall provide a proof, different from the one given previously in [Mat91, Hei94], thereby using the construction outlined above. We need a geometric property of Banach spaces.

Definition 3. A Banach space Y has type  $p, 1 \le p \le 2$ , if there is a constant  $C < \infty$ , such that for all probability spaces  $[\Omega, \mathcal{F}, P], k \in \mathbb{N}$  and independent random elements  $\Upsilon_1, \ldots, \Upsilon_k$  in  $L_2(\Omega, \mathcal{F}, P, Y)$ , for which  $\int_{\Omega} \Upsilon_j(\omega) dP(\omega) = 0$ ,  $j = 1, \ldots, k$ , we have

(7) 
$$\left(\int_{\Omega} \|\sum_{j=1}^{k} \Upsilon_{j}(\omega)\|_{Y}^{2} dP(\omega)\right)^{1/2} \leq C \left(\sum_{j=1}^{k} \int_{\Omega} \|\Upsilon_{j}(\omega)\|_{Y}^{p} dP(\omega)\right)^{1/2}$$

The smallest constant satisfying the above inequality shall be called the *type-p-constant* and is denoted by  $T_p(Y)$ .

Remark 3. The notion of a type of a Banach space was originally introduced in [MP76], where this definition was given in terms of Rademacher sequences. But, as can be seen easily, it can be extended to the above situation, albeit the type-p-constant is different by a factor of at most 2, see [LT91, Chapt. 9.2], but also [HJ74]. It can be seen that  $T_2(\mathbb{R}) = 1$ . Moreover, every finite-dimensional Banach space has type p,  $1 \le p \le 2$ , although the respective constant may depend on the dimension, cf. [TJ88, Ch. 1, §4]. However, we have

$$T_q(l_q^m) \le C, \quad \text{if } 1 \le q \le 2$$

and

$$T_2(l_q^m) \le C \begin{cases} 1 & \text{, if } 2 \le q < \infty \\ \sqrt{\log 1 + m} & \text{, if } q = \infty \end{cases}$$

for some universal constant  $C < \infty$ .

Now we are prepared to provide the proof of Proposition 1, following the one given in [Mat94].

Proof of Proposition 1. We shall carry out the estimates only for  $2 \leq q < \infty$ . The other cases follow the same lines. By Theorem 1 there is an unbiased Monte Carlo method  $\mathcal{P} = ([\Omega, \mathcal{F}, P], \mathbf{u}, 1)$  with  $u(\mathrm{Id}: l_2^m \to l_q^m) \leq m^{1/q}$ . The sample mean

 $\mathcal{P}^{n} = ([\Omega^{n}, \mathcal{F}^{n}, P^{n}], \mathbf{v}, n)$ 

of n independent copies of  $\mathcal{P}$ , which is defined by

$$\mathbf{v}_{\omega^n}(x) := \frac{1}{n} \sum_{j=1}^n \mathbf{u}_{\omega_j}(x), \quad x \in l_2^m, \quad \omega^n := (\omega_1, \dots, \omega_n) \in \Omega^n,$$

provides another unbiased Monte Carlo method, this time of cardinality n, which has an error at  $x \in l_2^m$ ,  $||x||_2 \leq 1$ , of

(8) 
$$e(\operatorname{Id}: l_2^m \to l_q^m, \mathcal{P}^n, x) = \left( \int_{\Omega} \|x - \frac{1}{n} \sum_{j=1}^n u_{\omega_j}(x)\|_q^2 dP^n(\omega^n) \right)^{1/2}$$

(9) 
$$= \frac{1}{n} \left( \int_{\Omega} \| \sum_{j=1}^{n} \left( x - \mathbf{u}_{\omega_j}(x) \right) \|_q^2 dP^n(\omega^n) \right)^{1/2}$$

(10) 
$$\leq \frac{1}{\sqrt{n}} T_2(l_q^m) e(\mathrm{Id} \colon l_2^m \to l_q^m, \mathcal{P}).$$

Since

$$e(\mathrm{Id}: l_2^m \to l_q^m, \mathcal{P}) \leq \|\mathrm{Id}: l_2^m \to l_q^m\| + u(\mathrm{Id}: l_2^m \to l_q^m),$$

the proof can be completed in case  $2 \leq q < \infty$ .  $\Box$ 

Now we are able to provide the reformulation of Theorem 1 in terms of the respective Monte Carlo approximation numbers  $a_n^{mc}(D_{\sigma}: l_2 \rightarrow l_q)$ , as promised above.

Corollary 2. Let  $1 \le q < \infty$  and put  $r := \max\{2, q'\}$ . If

$$(a_n^{mc}(D_{\sigma}\colon l_2\to l_q))_{n\in\mathbb{N}}\in l_{r,q}$$

then the mapping  $D_{\sigma} \colon l_2 \to l_q$  admits an unbiased Monte Carlo method.

*Proof.* To simplify the proof we shall assume that the diagonal is nonnegative and non-increasing.

Since the following diagram is commutative (where  $R^m$  assigns every vector in  $l_q$  the vector with the first m coordinates,  $J^m$  denotes the natural embedding and  $D_{\sigma}^m$  is the corresponding restriction),



and  $||J^{m}|| \leq 1$ ,  $||R^{m}|| \leq 1$ , we have by inequality (4) the estimate

 $a_n^{mc}(D_{\sigma}^m) \le a_n^{mc}(D_{\sigma}).$ 

Moreover, we have

(11) 
$$\sigma_m a_n^{mc} (\mathrm{Id}_{2,q}^m \colon l_2^m \to l_q^m) \le a_n (D_\sigma^m),$$

which can be seen as follows, cf. also [Pie87, 2.9.3]. Without loss of generality we may assume  $\sigma_m > 0$ . Since

$$\mathrm{Id}_{2,q}^m = (D_{\sigma}^m \colon l_2^m \to l_q^m) \cdot (D_{\sigma}^m \colon l_2^m \to l_2^m)^{-1}$$

and

$$||(D_{\sigma}^{m}: l_{2}^{m} \to l_{2}^{m})^{-1}|| = \sigma_{m}^{-1}$$

the estimate (11) follows immediately.

Letting m = 2n - 1 and inserting the results from Proposition 1 we see that the sequence  $(\sigma_{2n-1})_{n \in \mathbb{N}}$  belongs to  $l_q$  whenever the assumptions from Corollary 2 are fulfilled. But this is equivalent to  $(\sigma_n)_{n \in \mathbb{N}} \in l_q$ , see e.g. [Pie87, Prop. 2.1.9].  $\Box$ 

We turn to the question whether there are non-trivial necessary conditions to be imposed on an operator in order to admit an unbiased Monte Carlo method. In terms of the Monte Carlo approximation numbers a fairly general condition can be given, provided the target space has some type p. Indeed, the sample mean construction from the proof of Proposition 1 implies

**Theorem 2.** Suppose the Banach space Y has type p, p > 1. If an operator  $S: X \to Y$  admits an unbiased Monte Carlo method then necessarily  $(a_n^{mc}(S))_{n \in \mathbb{N}} \in l_{p',\infty}$ .

Taking into account the behavior of the Monte Carlo approximation numbers of the diagonal mappings and applying a technique similar to the one of Corollary 2 this transfers to

Corollary 3. Let  $1 \leq q < \infty$ . If  $D_{\sigma} : l_2 \to l_q$  admits an unbiased Monte Carlo method then  $\sigma \in l_{q,\infty}$ .

*Remark* 4. A look at the necessary and sufficient conditions proves that there is only a small gap left. Since the arguments to prove the necessary conditions are very rough we conjecture that the sufficient conditions are sharp.

On the class of operators acting between Hilbert spaces we immediately obtain

#### UNBIASED MONTE CARLO ESTIMATORS

**Theorem 3.** Any Hilbert-Schmidt operator (acting between Hilbert spaces) admits an unbiased Monte Carlo method. Conversely, if an operator between Hilbert spaces admits an unbiased Monte Carlo method then necessarily the sequence of singular numbers belongs to  $l_{2,\infty}$ 

Indeed, The norm u, as defined in (5), is easily seen to be unitarily invariant, such that the result for diagonal operators implies the respective result for arbitrary ones using the Schmidt - representation, see [Pie87, 2.11.4]. The class of Hilbert - Schmidt operators corresponds to the class of operators having square summable singular numbers, see [Pie80, 15.5.5], these corresponding to diagonal mappings having square summable diagonal.

#### 3. Some relations to operator ideals

As could be seen in Theorem 3, Hilbert - Schmidt operators admit an unbiased Monte Carlo method. The class of such operators is closely related to the theory of operator ideals as developed in [Pie80]. The first important (though simple) observation is

**Theorem 4.** The class of all operators admitting an unbiased Monte Carlo method turns into a normed operator ideal by letting  $S \to u(S)$  be the norm as defined in equation (5). This immediately implies

#### Corollary 4. Nuclear operators admit an unbiased Monte Carlo method.

We shall introduce this class of operators below. The assertion of the corollary follows from the fact that the ideal of the nuclear operators is the smallest normed ideal. However, a direct proof of the corollary could also be given using the trivial unbiased representation known from the introductory example.

Next we shall see that the class of operators admitting an unbiased Monte Carlo method can be enlarged considerably in many situations. We shall exemplify this by introducing the ideal of (r, p, q)-nuclear operators.

Let  $0 < r \leq \infty$ ,  $1 \leq p, q \leq \infty$  with  $1 + 1/r \geq \frac{1}{p} + \frac{1}{q}$ . Denote by  $\mathfrak{N}_{(r,p,q)}$  the ideal of (r, p, q)-nuclear operators, i.e., an operator  $S \in \mathfrak{L}(X, Y)$  is (r, p, q)-nuclear, if there is a representation

$$Sx = \sum_{j=1}^{\infty} \sigma_j \langle x, a_j \rangle y_j, \quad x \in X,$$

with  $a_j \in X'$  and  $y_j \in Y$  satisfying

$$\|\sigma\|_{r} < \infty$$
$$w_{q'}((a_{j})_{j \in \mathbb{N}}) := \sup_{\|x''\| \le 1} \left( \sum_{j=1}^{\infty} |\langle x'', a_{j} \rangle|^{q'} \right)^{1/q'} < \infty$$

and

$$w_{p'}((y_j)_{j\in\mathbb{N}}):=\sup_{||b||\leq 1}\left(\sum_{j=1}^{\infty}|\langle b,y_j\rangle|^{p'}\right)^{1/p'}<\infty.$$

Denote the respective quasi norm by

$$m_{(r,p,q)}(S) := \inf \|\sigma\|_{r} w_{q'}((a_j)_{j \in \mathbb{N}}) w_{p'}((y_j)_{j \in \mathbb{N}}),$$

where the infimum is taken over all possible representations, see [Pie80, 18.1] for more details. Especially we obtain as  $\mathfrak{N}_{(1,1,1)}$  the ideal of nuclear operators mentioned above. The ideal of (r, p, q)-nuclear operators can be characterized by a factorization property, see [Pie80, 18.1.3]. Especially, every (q, q, 2)-nuclear operator factors through a diagonal mapping from  $l_2$  to  $l_q$  with diagonal belonging to  $l_q$ . (In case  $q = \infty$  the diagonal converges to 0.)

Together with Theorem 1 and Corollary 1 this provides

Corollary 5. Let  $1 \leq q < \infty$ . Every (q, q, 2)-nuclear operator S admits an unbiased Monte Carlo method. Moreover we have

$$u(S) \leq n_{(q,q,2)}(S).$$

The largest ideal of this type which admits unbiased Monte Carlo methods is obtained as  $\mathfrak{N}_{(\infty,\infty,2)}$ .

For operators acting between Hilbert spaces H and K it is readily checked that

$$\mathfrak{N}_{(\infty,\infty,2)}(H,K) = \mathfrak{N}_{(2,2,2)}(H,K)$$

and we obtain the Hilbert - Schmidt operators once more, see [Pie80, 18.5.4].

#### References

- [ENS89] S.M. Ermakov, W.W. Nekrutkin, and A.S. Sipin. Random Processes for Classical Equations of Mathematical Physics. Mathematics and its Appl., Soviet Ser. 34. Kluwer Academic Publ., Dordrecht, 1989.
- [Hei94] S. Heinrich. Random approximation in numerical analysis. In K. D. Bierstedt, et al., editor, Functional Analysis: Proceedings of the Essen Conference, volume 150 of Lect. Notes in pure and appl. Math., pages 123 - 171, New York, Basel, Hong Kong, 1994. Marcel Dekker.
- [HH64] J. M. Hammersley and D. C. Handscomb. Monte Carlo methods. Methuen, London, 1964.
- [HJ74] J. Hoffmann-Jørgensen. Sums of independent Banach space valued random variables. Studia Math., 52:159 - 186, 1974.
- [KW86] M. H. Kalos and P. A. Whitlock. Monte Carlo methods, volume 1. Wiley, New York, 1986.
- [LB93] R. C. Liu and L. D. Brown. Nonexistence of informative unbiased estimators in singular problems. Ann. Statist., 21(1):1 - 13, 1993.
- [LT91] M. Ledoux and M. Talagrand. Probability in Banach spaces, volume 23 of Ergeb. Math. und ihrer Grenzgeb., 3. Folge. Springer, Berlin, Heidelberg, New York, 1991.
- [Mat91] P. Mathé. Random approximation of Sobolev embeddings. J. Complexity, 7:261 281, 1991.
- [Mat94] P. Mathé. Approximation theory of Monte Carlo methods. Habilitation thesis, 1994.
- [MP76] B. Maurey and G. Pisier. Séries de variables aléatoires vectorielles indépendantes et géométrie des espaces de Banach. *Studia Math.*, 58:45 90, 1976.
- [Pie80] A. Pietsch. Operator Ideals. North Holland, Amsterdam, New York, Oxford, 1980.
- [Pie87] A. Pietsch. Eigenvalues and s-Numbers, volume 43 of Math. und ihre Anw. in Phys. und Technik. Geest & Portig, Leipzig, 1987.
- [Ros56] M. Rosenblatt. Remarks on nonparametric estimates of a density function. Ann. Math. Statist., 27:832 - 837, 1956.

[TJ88] N. Tomczak-Jaegermann. Banach-Mazur Distances and Finite-dimensional Operator Ideals, volume 38 of Pitman Monographs and Surveys in pure and applied Math. Longman, Essex, 1988.

[TWW88] J.F. Traub, G.W. Wasilkowski, and H. Woźniakowski. Information-Based Complexity. Academic Press, New York, 1988.

WEIERSTRASS INSTITUTE FOR APPLIED ANALYSIS AND STOCHASTICS, D-10117 BERLIN, MOHREN-STRASSE 39

• • • -. · · · · · · · · · · · · ۰

## Recent publications of the Weierstraß–Institut für Angewandte Analysis und Stochastik

#### Preprints 1994

- 84. Karmeshu, Henri Schurz: Moment evolution of the outflow-rate from nonlinear conceptual reservoirs.
- 85. Wolfdietrich Müller, Klaus R. Schneider: Feedback stabilization of nonlinear discrete-time systems.
- 86. Gennadii A. Leonov: A method of constructing of dynamical systems with bounded nonperiodic trajectories.
- 87. Gennadii A. Leonov: Pendulum with positive and negative dry friction. Continuum of homoclinic orbits.
- 88. Reiner Lauterbach, Jan A. Sanders: Bifurcation analysis for spherically symmetric systems using invariant theory.
- 89. Milan Kučera: Stability of bifurcating periodic solutions of differential inequalities in  $\mathbb{R}^3$ .
- 90. Peter Knabner, Cornelius J. van Duijn, Sabine Hengst: An analysis of crystal dissolution fronts in flows through porous media Part I: Homogeneous charge distribution.
- 91. Werner Horn, Philippe Laurençot, Jürgen Sprekels: Global solutions to a Penrose-Fife phase-field model under flux boundary conditions for the inverse temperature.
- 92. Oleg V. Lepskii, Vladimir G. Spokoiny: Local adaptivity to inhomogeneous smoothness. 1. Resolution level.
- **93.** Wolfgang Wagner: A functional law of large numbers for Boltzmann type stochastic particle systems.
- 94. Hermann Haaf: Existence of periodic travelling waves to reaction-diffusion equations with excitable-oscillatory kinetics.
- 95. Anton Bovier, Véronique Gayrard, Pierre Picco: Large deviation principles for the Hopfield model and the Kac-Hopfield model.
- 96. Wolfgang Wagner: Approximation of the Boltzmann equation by discrete velocity models.
- 97. Anton Bovier, Véronique Gayrard, Pierre Picco: Gibbs states of the Hopfield model with extensively many patterns.

- 98. Lev D. Pustyl'nikov, Jörg Schmeling: On some estimations of Weyl sums.
- 99. Michael H. Neumann: Spectral density estimation via nonlinear wavelet methods for stationary non-Gaussian time series.
- 100. Karmeshu, Henri Schurz: Effects of distributed delays on the stability of structures under seismic excitation and multiplicative noise.
- 101. Jörg Schmeling: Estimates of Weyl sums over subsequences of natural numbers.
- 102. Grigori N. Milstein, Michael V. Tret'yakov: Mean-square approximation for stochastic differential equations with small noises.
- 103. Valentin Konakov: On convergence rates of suprema in the presence of nonnegligible trends.
- 104. Pierluigi Colli, Jürgen Sprekels: On a Penrose-Fife model with zero interfacial energy leading to a phase-field system of relaxed Stefan type.
- 105. Anton Bovier: Self-averaging in a class of generalized Hopfield models.
- 106. Andreas Rathsfeld: A wavelet algorithm for the solution of the double layer potential equation over polygonal boundaries.
- 107. Michael H. Neumann: Bootstrap confidence bands in nonparametric regression.
- 108. Henri Schurz: Asymptotical mean square stability of an equilibrium point of some linear numerical solutions with multiplicative noise.
- 109. Gottfried Bruckner: On the stabilization of trigonometric collocation methods for a class of ill-posed first kind equations.
- 110. Wolfdietrich Müller: Asymptotische Input-Output-Linearisierung und Störgrößenkompensation in nichtlinearen Reaktionssystemen.
- 111. Vladimir Maz'ya, Gunther Schmidt: On approximate approximations using Gaussian kernels.
- 112. Henri Schurz: A note on pathwise approximation of stationary Ornstein-Uhlenbeck processes with diagonalizable drift.