

Weierstraß-Institut für Angewandte Analysis und Stochastik

im Forschungsverbund Berlin e.V.

Preprint

ISSN 0946 – 8633

Linear non-autonomous Cauchy problems and evolution semigroups

Hagen Neidhardt¹, Valentin A. Zagrebnov²

submitted: November 07, 2007

¹ Weierstrass Institute for
Applied Analysis and Stochastics
Mohrenstr. 39
D-10117 Berlin
Germany
E-mail: neidhard@wias-berlin.de

² Université de la Méditerranée
(Aix-Marseille II) and
Centre de Physique Théorique
UMR 6207, Luminy - Case 907,
F-13288 Marseille Cedex 9,
France, **E-mail:** zagrebnov@cpt.univ-mrs.fr

No. 1266
Berlin 2007



2000 *Mathematics Subject Classification.* 35L90, 34G10, 47D06.

Key words and phrases. linear evolution equations, evolution semigroups, perturbation theory, time-dependent Schrödinger operators, moving potentials.

Edited by
Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS)
Mohrenstraße 39
10117 Berlin
Germany

Fax: + 49 30 2044975
E-Mail: preprint@wias-berlin.de
World Wide Web: <http://www.wias-berlin.de/>

Abstract

The paper is devoted to the problem of existence of propagators for an abstract linear non-autonomous evolution Cauchy problem of hyperbolic type in separable Banach spaces. The problem is solved using the so-called evolution semigroup approach which reduces the existence problem for propagators to a perturbation problem of semigroup generators. The results are specified to abstract linear non-autonomous evolution equations in Hilbert spaces where the assumption is made that the domains of the quadratic forms associated with the generators are independent of time. Finally, these results are applied to time-dependent Schrödinger operators with moving point interactions in 1D.

Contents

1	Introduction and setup of the Problem	2
2	Evolution generators	4
2.1	Forward generators	4
2.2	Backward generators	6
2.3	Bidirectional problems	7
2.4	Problems on \mathbb{R}	10
3	Semigroup perturbations	12
4	Solutions of evolution equations	17
4.1	Solutions of forward evolution equations	17
4.2	Backward and bidirectional evolution equations	19
5	Evolution equations in Hilbert spaces	21
5.1	Preliminaries	21
5.2	Auxiliary evolution equation	23
5.3	Back to the original problem	24
6	Examples	29
6.1	Point interactions with varying coupling constant	29
6.2	Moving potentials	32

1 Introduction and setup of the Problem

The aim of the present paper is to develop an approach to Cauchy problems for linear non-autonomous evolution equations of type

$$\frac{\partial}{\partial t}u(t) + A(t)u(t) = 0, \quad u(s) = u_s \in X, \quad t, s \in I, \quad (1.1)$$

where \mathcal{I} is a bounded open interval of \mathbb{R} and $\{A(t)\}_{t \in \mathcal{I}}$ is a family of closed linear operators in the separable Banach space X . Evolution equations of that type are called forward evolution equations if $s \leq t$, backward if $s \geq t$ and bidirectional evolution equations if s and t are arbitrary. The main question concerning the Cauchy problem (1.1) is to find a so-called “solution operator” or propagator $U(t, s)$ such that $u(t) := U(t, s)u_s$ is in some sense a solution of (1.1) satisfying the initial condition $u(s) = u_s$.

Usually it is assumed that either $\{A(t)\}_{t \in \mathcal{I}}$ or $\{-A(t)\}_{t \in \mathcal{I}}$ are families of generators of C_0 -semigroups in X . In order to distinct both cases we call an operator A a *generator* if it generates a C_0 -semigroup $\{e^{tA}\}_{t \geq 0}$. We call A an *anti-generator* if $-A$ generates a C_0 -semigroup $\{e^{-tA}\}_{t \geq 0}$, i.e., the operator $-A$ is the *generator* of a semigroup. If simultaneously A is an anti-generator and a generator, then A is called a group generator.

Very often the Cauchy problem (1.1) is attacked for a suitable dense subset of initial data u_s by solving it directly in the same manner as ordinary differential equation, which immediately implies the existence of the propagator, see e.g. [43]. For this purpose one assumes that $\{A(t)\}_{t \in \mathcal{I}}$ is a family of anti-generators of C_0 -semigroups such that they uniformly belong to the class of quasi-bounded semigroups $\mathcal{G}(M, \beta)$, cf. [21, Chapter IX]. If $\{A(t)\}_{t \in \mathcal{I}}$ is a family of anti-generators of class $\mathcal{G}(M, \beta)$ which are simultaneously anti-generators of *holomorphic* C_0 -semigroups, then the evolution equation is called of “parabolic” type. If it is not holomorphic, then it is called of “hyperbolic” type. In the following in this paper we are only interested in the “hyperbolic” case.

There is a rich literature on “hyperbolic” evolution equations problems. The first author who discussed these problems was Phillips [37]. A more general case was considered by Kato in [19, 20] and by Mizohata in [29]. These results were generalized in the sixties in [11, 16, 24, 51, 52, 27, 15, 13]. Kato has improved these results in two important papers [22, 23], where for the first time he introduced the assumptions of *stability* and *invariance*. In the seventies and eighties Kato’s result were generalized in [10, 18, 25, 49, 50]. For related results see also [26, 14, 9]. Recently several new results were obtained in [3, 36, 35, 44, 45, 46]. In the following we refer to these results as a “standard approach” or “standard methods”. Their common feature is that the propagator is constructed by using certain approximations of the family $\{A(t)\}_{t \in \mathcal{I}}$ for which the corresponding Cauchy problem can be easily solved. After that one has only to verify that the obtained sequence of propagators converges to the propagator of the original problem. Widely used approximations are a so-called *Yosida approximation* introduced in [52], *piecewise constant* approximations proposed by Kato, cf. [22, 23], as well as a combination of both, see [24].

In contrast to the *standard methods* another approach was developed in [12, 17, 31, 32, 33, 34]. It does not rely on any approximation, since it is based on the fact that the existence problem for the propagator in question is *equivalent* to an operator extension problem for a suitable defined operator in a vector-valued Banach space $L^p(\mathcal{I}, X)$ for some $p \in [1, \infty)$. More precisely, it turns out that any *forward propagator* $\{U(t, s)\}_{(t, s) \in \Delta_{\mathcal{I}}}$, $\Delta_{\mathcal{I}} := \{(t, s) \in \mathcal{I} \times \mathcal{I} : s \leq t\}$, (see Definition 2.1) defines a C_0 -semigroup in $L^p(\mathcal{I}, X)$ by

$$(\mathcal{U}(\sigma)f)(t) := U(t, t - \sigma)\chi_{\mathcal{I}}(t - \sigma)f(t - \sigma), \quad f \in L^p(\mathcal{I}, X), \quad \sigma \geq 0, \quad (1.2)$$

where $\chi_{\mathcal{I}}(\cdot)$ is the characteristic function of the open interval \mathcal{I} . C_0 -semigroups in $L^p(\mathcal{I}, X)$ admitting a forward propagator representation (1.2) are called *forward evolution semigroups*. The anti-generator K of the semigroup $\{\mathcal{U}(\sigma)\}_{\sigma \in \mathbb{R}_+}$, i.e. $\mathcal{U}(\sigma) = e^{-\sigma K}$, $\sigma \in \mathbb{R}_+$, is called the *forward generator*. Our approach is based on the important fact that the set of the forward generators can

be described explicitly, and that there is a *one-to-one correspondence* between forward propagators and forward generators, see [32].

Now, let us assume that the forward propagator $\{U(t, s)\}_{(t,s) \in \Delta_{\mathcal{I}}}$ is found by the standard approach and that it solves the forward evolution equation (1.1) in some sense. Then it turns out that the forward generator $K_{\mathcal{I}}$ defined by (1.2) is an extension of the so-called *evolution operator* $\tilde{K}_{\mathcal{I}}$ given by

$$(\tilde{K}_{\mathcal{I}}f)(t) = D_{\mathcal{I}}f + Af, \quad f \in \text{dom}(\tilde{K}_{\mathcal{I}}) = \text{dom}(D_{\mathcal{I}}) \cap \text{dom}(A), \quad (1.3)$$

in $L^p(\mathcal{I}, X)$ for some $p \in [1, \infty)$, where $D_{\mathcal{I}}$ is the anti-generator of the *right-shift semigroup* in $L^p(\mathcal{I}, X)$ and A is the *multiplication* operator in $L^p(\mathcal{I}, X)$ induced by the family $\{A(t)\}_{t \in \mathcal{I}}$, see Section 2.

This remark leads to the main idea of our approach: to solve the evolution equation (1.1) by extending the evolution operator $\tilde{K}_{\mathcal{I}}$ to an anti-generator of an (forward) evolution semigroup. Notice that in contrast to the standard approach now the focus has moved from the problem to construct a propagator to the problem to find a certain operator extension. This so-called “extension approach” or “extension method” has a lot of advantages, since it works in a very general setting, and it is quite flexible and transparent. The approach becomes very simple, if the closure of the evolution operator $\tilde{K}_{\mathcal{I}}$ is already an anti-generator, in other words, if $\tilde{K}_{\mathcal{I}}$ is *essentially* anti-generator. In this case one gets the forward generator by closing $\tilde{K}_{\mathcal{I}}$, see Theorem 2.4, which immediately implies the existence of a unique forward propagator for the non-autonomous Cauchy problem (1.1). Some recent results related to the extension method can be find in e.g. [28, 30, 35, 39, 40].

Below we exploit this approach extensively and we show how this method can be applied to evolution equations of type (1.1). We prove that under the stability and invariance assumptions of Kato [22, 23] the evolution operator $\tilde{K}_{\mathcal{I}}$ is already an *essential* anti-generator, which means that its closure $K_{\mathcal{I}}$ is a forward generator.

We apply also the extension method to *bidirectional* evolution equations of the type

$$i \frac{\partial}{\partial t} u(t) = H(t)u(t), \quad u(s) = u_s, \quad s, t \in \mathbb{R}, \quad (1.4)$$

on \mathbb{R} in Hilbert spaces, where $\{H(t)\}_{t \in \mathbb{R}}$ is a family of non-negative self-adjoint operators. Using the extension method we restore and obtain some generalizations of the Kiszyński result [24]. Moreover, we show that Kiszyński’s propagator is in fact the propagator of an auxiliary evolution equation problem closely related to (1.4). The solution of the auxiliary problem implies a solution for (1.4). The uniqueness of the auxiliary solution does not imply, however, uniqueness of the original problem (1.4), in general.

The paper is organized as follows. In Section 2 we recall some basic facts of the theory of evolution semigroups. Section 3 is devoted to a perturbation theorem for generators of these semigroups, which is used then in Section 4 to show that the closure $K_{\mathcal{I}}$ of the evolution operator (1.3) is an anti-generator. The results of Section 4 are specified in Section 5 to families $\{A(t)\}_{t \in \mathbb{R}}$ of the form $A(t) = iH(t)$ where $H(t)$ are semi-bounded self-adjoint operators with *time-independent* form domains in a Hilbert space. In Section 6, we apply these results of Section 5 to Schrödinger operators with time-dependent point interactions of the form:

$$H(t) := -\frac{1}{2} \frac{d}{dx} \frac{1}{m(x)} \frac{d}{dx} + V(x) + \sum_{j=1}^N \kappa_j(t) \delta(x - x_j) \quad x \in \mathbb{R}, \quad t \in \mathbb{R},$$

as well as to the case of moving point interactions of the form:

$$H(t) := -\frac{d^2}{dx^2} + \kappa_1(t) \delta(x - x_1(t)) + \kappa_2(t) \delta(t)(x - x_2(t))$$

where the coupling constants $\kappa_j(\cdot)$ are non-negative Lipschitz continuous functions in $t \in \mathbb{R}$ and $x_j(t)$ are C^2 -trajectories in \mathbb{R} . These kind of problems were the subject of publications [7, 6, 38, 42, 41, 48].

2 Evolution generators

In the following we are interested not only in the *forward* evolution equations but also in the *backward* ones as well as in the *bidirectional* evolution equations. The interest to these evolution equations rises from time reversible problems in quantum mechanics, which we consider in conclusion of this paper as applications. For this purpose we show in Section 2.2 how one has to modify the extension approach for backward evolution equations. Moreover, in application to quantum mechanics we are concerned with *infinite* time intervals, in particular, with $\mathcal{I} = \mathbb{R}$. In order to apply our approach to this situation it is useful to *localize* it in time, this means that instead to consider the Cauchy problem on \mathbb{R} we consider it on arbitrary finite subintervals of \mathbb{R} . In this case, however, one has to ensure that propagators for different time intervals are *compatible*.

2.1 Forward generators

We start with the definition of a forward propagator in a separable Banach space.

Definition 2.1 Let X be a separable Banach space. A strongly continuous operator-valued function $U(\cdot, \cdot) : \Delta_{\mathcal{I}} \rightarrow \mathcal{B}(X)$ is called a *forward propagator* on $\Delta_{\mathcal{I}} = \{(t, s) \in \mathcal{I} \times \mathcal{I} : s \leq t\}$, if

- (i) $U(t, t) = I_X$ for $t \in \mathcal{I}$,
- (ii) $U(t, r)U(r, s) = U(t, s)$ for $(t, r, s) \in \mathcal{I}^3$, $s \leq r \leq t$,
- (iii) $\|U\|_{\mathcal{B}(X)} := \sup_{(t, s) \in \Delta_{\mathcal{I}}} \|U(t, s)\|_{\mathcal{B}(X)} < \infty$.

We call a strongly continuous operator-valued function $U(\cdot, \cdot)$ defined on $\Delta_{\mathbb{R}} := \{(t, s) \in \mathbb{R} \times \mathbb{R} : s \leq t\}$ a forward propagator, if for any bounded interval \mathcal{I} the *restriction* of $U(\cdot, \cdot)$ to $\Delta_{\mathcal{I}}$ is a forward propagator.

Another important notion is the so-called *evolution operator*. To explain this notion we introduce the Banach space $L^p(\mathcal{I}, X)$, $p \in [1, \infty)$, where X is a separable Banach space. In $L^p(\mathcal{I}, X)$ we define the *multiplication operator*

$$(M(\phi)f)(t) := \phi(t)f(t), \quad \text{dom}(M(\phi)) = L^p(\mathcal{I}, X), \quad \phi \in L^\infty(\mathcal{I}). \quad (2.1)$$

Definition 2.2 A linear operator K in $L^p(\mathcal{I}, X)$, $p \in [1, \infty)$, is called an *evolution operator*, if

- (i) it satisfies the conditions:

$$\text{dom}(K) \subseteq C(\overline{\mathcal{I}}, X), \quad (2.2)$$

$$M(\phi)\text{dom}(K) \subseteq \text{dom}(K), \quad \phi \in H^{1, \infty}(\mathcal{I}), \quad (2.3)$$

and

$$KM(\phi)f - M(\phi)Kf = M(\dot{\phi})f, \quad f \in \text{dom}(K), \quad \phi \in H^{1, \infty}(\mathcal{I}), \quad (2.4)$$

where $\dot{\phi} := d\phi/dt$, and

- (ii) its domain $\text{dom}(K)$ has a dense *cross-section* in X , this means that

$$[\text{dom}(K)]_t := \{x \in X : \exists f \in \text{dom}(K) \text{ such that } f(t) = x\},$$

is dense in X for each $t \in \mathcal{I}$.

If in addition K is an anti-generator or a generator in $L^p(\mathcal{I}, X)$, then K is called a *forward* or *backward* generator, respectively.

The density of the cross-section is not a trivial condition. However, one has to mention that it is important to ensure the *continuity* of the propagator. Notice that if K is an evolution operator, then its domain $\text{dom}(K)$ is already dense in $L^p(\mathcal{I}, X)$, $1 \leq p < \infty$.

Further, by virtue of Theorem 4.12, [32], it turns out that there is a *one-to-one correspondence* between the set of forward propagators and the set of forward generators established by (1.2). This correspondence plays a crucial role in our arguments below.

Let $S_r(\sigma)$ be the right-shift semigroup in $L^p(\mathcal{I}, X)$, $1 \leq p < +\infty$, given by

$$(S_r(\sigma)f)(t) := f(t - \sigma)\chi_{\mathcal{I}}(t - \sigma), \quad f \in L^p(\mathcal{I}, X). \quad (2.5)$$

This is a C_0 -semigroup of class $\mathcal{G}(1, 0)$. Its generator is given by $-D_{\mathcal{I}}$, where

$$(D_{\mathcal{I}}f)(t) = \frac{\partial}{\partial t}f(t), \quad f \in \text{dom}(D_{\mathcal{I}}) := H_a^{1,p}(\mathcal{I}, X), \quad \mathcal{I} = (a, b).$$

According to our convention the operator $D_{\mathcal{I}}$ is an anti-generator. Here

$$H_a^{1,p}(\mathcal{I}, X) := \{f \in H^{1,p}(\mathcal{I}, X) : f(a) = 0\},$$

and $H_a^{1,p}(\mathcal{I}, X)$ is the Sobolev space of X -valued absolutely continuous functions on \mathcal{I} with p -summable derivative.

Notice that a family $\{A(t)\}_{t \in \mathcal{I}}$ of closed and densely defined linear operators is called *measurable*, if there is a $z \in \mathbb{C}$ such that z belongs to the *resolvent set* $\rho(A(t))$ of $A(t)$ for almost every (a.e.) $t \in \mathcal{I}$ and for each $x \in X$ the function

$$f(t) := (A(t) - z)^{-1}x, \quad t \in \mathcal{I},$$

is *strongly* measurable. If the family $\{A(t)\}_{t \in \mathcal{I}}$ is measurable, then one can show that the multiplication operator A ,

$$(Af)(t) := A(t)f(t), \quad f \in \text{dom}(A), \quad (2.6)$$

$$\text{dom}(A) := \left\{ f \in L^p(\mathcal{I}, X) : \begin{array}{l} f(t) \in \text{dom}(A(t)) \text{ for a.e. } t \in \mathcal{I}, \\ A(t)f(t) \in L^p(\mathcal{I}, X) \end{array} \right\} \quad (2.7)$$

is densely defined and closed in $L^p(\mathcal{I}, X)$.

Instead of solving the Cauchy problem (1.1) for a suitable set of initial data u_s we consider the operator

$$\tilde{K}_{\mathcal{I}}f := D_{\mathcal{I}}f + Af, \quad f \in \text{dom}(\tilde{K}_{\mathcal{I}}) := \text{dom}(D_{\mathcal{I}}) \cap \text{dom}(A), \quad (2.8)$$

in $L^p(\mathcal{I}, X)$, $p \in [1, \infty)$. If the domain $\text{dom}(\tilde{K}_{\mathcal{I}})$ has a dense cross-section, then by the definition above $\tilde{K}_{\mathcal{I}}$ is an evolution operator. This leads naturally to following definitions:

Definition 2.3 Let $\{A(t)\}_{t \in \mathcal{I}}$ be a measurable family of a closed and densely defined linear operators in the separable Banach space X .

(i) The forward evolution equation (1.1) is *well-posed* on \mathcal{I} for some $p \in [1, \infty)$ if $\tilde{K}_{\mathcal{I}}$ is an evolution operator.

(ii) A forward propagator $\{U(t, s)\}_{(t,s) \in \Delta_{\mathcal{I}}}$ is called a *solution* of the well-posed forward evolution equation (1.1) on \mathcal{I} if the corresponding forward generator $K_{\mathcal{I}}$, cf. (1.2), is an *extension* of $\tilde{K}_{\mathcal{I}}$.

(iii) The well-posed forward evolution equation (1.1) on \mathcal{I} has a unique solution if $\tilde{K}_{\mathcal{I}}$ admits only one extension which is a forward generator.

It is quite possible that the forward evolution equation (1.1) has several solutions, which means that the evolution operator $\tilde{K}_{\mathcal{I}}$ admits *several* extensions, and each of them is a forward generator. The dense cross-section property of the evolution operator is not sufficient to show that the evolution equation admits a unique solution.

In the following the next statement will be important for our reasoning.

Theorem 2.4 *Let $\{A(t)\}_{t \in \mathcal{I}}$ be a measurable family of closed and densely defined linear operators in the separable Banach space X . Assume that the forward evolution equation (1.1) is well-posed on \mathcal{I} for some $p \in [1, \infty)$. If the evolution operator $\tilde{K}_{\mathcal{I}}$ is closable in $L^p(\mathcal{I}, X)$ and its closure $K_{\mathcal{I}}$ is an anti-generator, then the forward evolution equation (1.1) on \mathcal{I} has a unique solution.*

Proof. Since the evolution equation is well-posed, the domain $\text{dom}(\tilde{K}_{\mathcal{I}})$ is densely defined in $L^p(\mathcal{I}, X)$. By assumptions the closure $K_{\mathcal{I}}$ is an anti-generator. Hence, it remains to show that the closure $K_{\mathcal{I}}$ satisfies the conditions (2.2)-(2.4). It is easy to verify that the closure $K_{\mathcal{I}}$ satisfies the conditions (2.3) and (2.4). To show (2.2) let us assume that $K_{\mathcal{I}}$ belongs to $\mathcal{G}(M, \beta)$. By Lemma 2.16 of [33] the closure $K_{\mathcal{I}}$ admits the estimate

$$\|f(t)\|_X \leq \frac{M}{(\xi - \beta)^{(p-1)/p}} \|(K_{\mathcal{I}} + \xi)f\|_{L^p(\mathcal{I}, X)}, \quad f \in \text{dom}(K_{\mathcal{I}}), \quad p \in [1, \infty),$$

for a.e. $t \in \bar{\mathcal{I}}$ and $\xi > \beta$. In particular, we have

$$\|f\|_{C(\bar{\mathcal{I}}, X)} \leq \frac{M}{(\xi - \beta)^{(p-1)/p}} \|(\tilde{K}_{\mathcal{I}} + \xi)f\|_{L^p(\mathcal{I}, X)}, \quad f \in \text{dom}(\tilde{K}_{\mathcal{I}}). \quad (2.9)$$

Since $\tilde{K}_{\mathcal{I}}$ has a closure $K_{\mathcal{I}}$, there is a sequence of elements $\{f_n\}_{n \in \mathbb{N}}$ for any $f \in \text{dom}(\tilde{K}_{\mathcal{I}})$ such that $f_n \in \text{dom}(\tilde{K}_{\mathcal{I}})$, $f_n \rightarrow f$ and $\tilde{K}_{\mathcal{I}}f_n \rightarrow K_{\mathcal{I}}f$ in the $L^p(\mathcal{I}, X)$ sense when $n \rightarrow \infty$. By (2.9) one gets that $\{f_n\}_{n \in \mathbb{N}}$ is a Cauchy sequence in $C(\bar{\mathcal{I}}, X)$. Hence $f \in C(\bar{\mathcal{I}}, X)$, that proves (2.2). Since $\text{dom}(\tilde{K}_{\mathcal{I}})$ has a dense cross-section for each $t \in \mathcal{I}$, one gets that its closure $K_{\mathcal{I}}$ has a dense cross-section for each $t \in \mathcal{I}$. Hence $K_{\mathcal{I}}$ is forward generator.

Let $K_{\mathcal{I}}$ and $K'_{\mathcal{I}}$ be two different extensions of $\tilde{K}_{\mathcal{I}}$, which are both forward generators. Since $K_{\mathcal{I}}$ is the closure of $\tilde{K}_{\mathcal{I}}$ one has $K_{\mathcal{I}} \subseteq K'_{\mathcal{I}}$. Since $K_{\mathcal{I}}$ and $K'_{\mathcal{I}}$ are generators of a C_0 -semigroup, one gets $K_{\mathcal{I}} = K'_{\mathcal{I}}$. Hence the evolution equation (1.1) is uniquely solvable. \square

2.2 Backward generators

In the following we are also interested in so-called *backward* evolution equation (1.1), $t \leq s$, $t, s \in \mathcal{I}$. Equations of that type require the introduction of the notion of *backward propagator*:

Definition 2.5 A strongly continuous operator-valued function $V(\cdot, \cdot) : \nabla_{\mathcal{I}} \rightarrow \mathcal{B}(X)$ is called a *backward propagator* on $\nabla_{\mathcal{I}} := \{(t, s) \in \mathcal{I} \times \mathcal{I} : t \leq s\}$, if

- (i) $V(t, t) = I_X$ for $t \in \mathcal{I}$,
- (ii) $V(t, r)V(r, s) = V(t, s)$ for $(t, r, s) \in \mathcal{I}^3$, $t \leq r \leq s$,
- (iii) $\sup_{(t, s) \in \nabla_{\mathcal{I}}} \|V(t, s)\|_{\mathcal{B}(X)} < \infty$.

We call a strongly continuous operator-valued function $V(\cdot, \cdot)$ defined on $\nabla_{\mathbb{R}} := \{(t, s) \in \mathbb{R} \times \mathbb{R} : s \leq t\}$ a *backward propagator* if for any bounded interval \mathcal{I} the restriction of $V(\cdot, \cdot)$ to $\nabla_{\mathcal{I}}$ is a backward propagator.

Similar to forward propagators there is a one-to-one correspondence between backward propagators and backward generators given by

$$(e^{\sigma K} f)(t) = V(t, t + \sigma) \chi_{\mathcal{I}}(t + \sigma) f(t + \sigma), \quad f \in L^p(\mathcal{I}, X), \quad \sigma \geq 0, \quad (2.10)$$

$p \in [1, \infty)$. With the backward evolution equation we associated the operator $\tilde{K}^{\mathcal{I}}$

$$\tilde{K}^{\mathcal{I}} f = D^{\mathcal{I}} f + A f, \quad f \in \text{dom}(\tilde{K}^{\mathcal{I}}) := \text{dom}(D^{\mathcal{I}}) \cap \text{dom}(A), \quad (2.11)$$

where

$$(D^{\mathcal{I}} f)(t) = \frac{\partial}{\partial t} f(t), \quad f \in \text{dom}(D^{\mathcal{I}}) := \{f \in H_b^{1,p}(\mathcal{I}, X) : f(b) = 0\}$$

is the generator of *left-shift* semigroup $S_l(\sigma) = e^{\sigma D^{\mathcal{I}}}$ on $L^p(\mathcal{I}, X)$, that is,

$$(S_l(\sigma) f)(t) = f(t + \sigma) \chi_{\mathcal{I}}(t + \sigma), \quad t \in \mathcal{I}, \quad f \in L^p(\mathcal{I}, X), \quad \sigma \geq 0.$$

Definition 2.6 Let $\{A(t)\}_{t \in \mathcal{I}}$ be a measurable family of closed and densely defined linear operators in the separable Banach space X .

(i) The backward evolution equation (1.1) is *well-posed* on \mathcal{I} for some $p \in [1, \infty)$, if $\tilde{K}^{\mathcal{I}}$ is an evolution operator.

(ii) A backward propagator $\{V(t, s)\}_{(t,s) \in \nabla_{\mathcal{I}}}$ is called a *solution* of the well-posed backward evolution equation (1.1) on \mathcal{I} if the corresponding backward generator $K^{\mathcal{I}}$, cf. (2.10), is an *extension* of $\tilde{K}^{\mathcal{I}}$.

(iii) The well-posed backward evolution equation (1.1) on \mathcal{I} has a solution if $\tilde{K}_{\mathcal{I}}$ admits only one extension which is a backward generator.

Now, following the same line of reasoning as in Theorem 2.4 we obtain a similar statement concerning the backward evolution equation (1.1):

Theorem 2.7 Let $\{A(t)\}_{t \in \mathcal{I}}$ be a measurable family of closed and densely defined linear operators in the separable Banach space X . Assume that the backward evolution equation (1.1) is well-posed on \mathcal{I} for some $p \in [1, \infty)$. If the evolution operator $\tilde{K}^{\mathcal{I}}$ is closable in $L^p(\mathcal{I}, X)$ and its closure $K^{\mathcal{I}}$ is a generator, then the backward evolution equation (1.1) on \mathcal{I} has a unique solution.

2.3 Bidirectional problems

Crucial for studying *bidirectional* evolution equations on bounded intervals is the following proposition.

Proposition 2.8 Let $\{U(t, s)\}_{(t,s) \in \Delta_{\mathcal{I}}}$ and $\{V(t, s)\}_{(t,s) \in \nabla_{\mathcal{I}}}$ be for- and backward propagators which correspond to the for- and backward generators $K_{\mathcal{I}}$ and $K^{\mathcal{I}}$, respectively. The relation

$$V(s, t)U(t, s) = U(t, s)V(s, t) = I_X, \quad (t, s) \in \Delta_{\mathcal{I}}, \quad (2.12)$$

holds if and only if for each $\phi \in H_a^{1,\infty}(\mathcal{I}) \cap H_b^{1,\infty}(\mathcal{I})$ the conditions

$$M(\phi) \text{dom}(K_{\mathcal{I}}) \subseteq \text{dom}(K^{\mathcal{I}}) \quad \text{and} \quad M(\phi) \text{dom}(K^{\mathcal{I}}) \subseteq \text{dom}(K_{\mathcal{I}}) \quad (2.13)$$

and

$$K^{\mathcal{I}} M(\phi) f = K_{\mathcal{I}} M(\phi) f, \quad f \in \text{dom}(K_{\mathcal{I}}) \quad \text{or} \quad f \in \text{dom}(K^{\mathcal{I}}), \quad (2.14)$$

are satisfied.

Proof. We set

$$g(\sigma) := e^{\sigma K^{\mathcal{I}}} M(\phi) e^{-\sigma K_{\mathcal{I}}} f, \quad f \in L^p(\mathcal{I}, X), \quad \phi \in H^{1,\infty}(\mathcal{I}).$$

Taking into account (1.2) and (2.10) we find

$$(g(\sigma))(t) = V(t, t + \sigma) \phi(t + \sigma) U(t + \sigma, t) \chi_{\mathcal{I}}(t + \sigma) \chi_{\mathcal{I}}(t) f(t), \quad t \in \mathcal{I}.$$

Using (2.12) we obtain

$$(g(\sigma))(t) = \phi(t + \sigma) \chi_{(a, b - \sigma)}(t) f(t), \quad t \in \mathcal{I}, \quad 0 \leq \sigma < b - a. \quad (2.15)$$

Since

$$(g(\sigma) - M(\phi))f = (e^{\sigma K^{\mathcal{I}}} - I)M(\phi)f + e^{\sigma K^{\mathcal{I}}} M(\phi)(e^{-\sigma K_{\mathcal{I}}} - I)f, \quad (2.16)$$

by (2.15) we get that

$$\lim_{\sigma \rightarrow +0} \frac{1}{\sigma} (g(\sigma) - M(\phi))f = 0, \quad f \in L^p(\mathcal{I}, X).$$

Assuming $f \in \text{dom}(K_{\mathcal{I}})$ we immediately find from (2.16) that $M(\phi)f \in \text{dom}(K^{\mathcal{I}})$ and (2.13). Interchanging $K^{\mathcal{I}}$ and $K_{\mathcal{I}}$ we prove $M(\phi)f \in \text{dom}(K_{\mathcal{I}})$ and (2.14).

Conversely, assuming (2.13) and (2.14) we get that the function $g(\sigma)$ is differentiable and

$$\frac{d}{d\sigma} g(\sigma) = e^{\sigma K^{\mathcal{I}}} (K^{\mathcal{I}} M(\phi) - M(\phi) K_{\mathcal{I}}) e^{-\sigma K_{\mathcal{I}}} f, \quad \sigma \geq 0.$$

By virtue of (2.4) we find

$$\frac{d}{d\sigma} g(\sigma) = e^{\sigma K^{\mathcal{I}}} M(\dot{\phi}) e^{-\sigma K_{\mathcal{I}}} f, \quad \sigma \geq 0$$

which yields

$$e^{\sigma K^{\mathcal{I}}} M(\phi) e^{-\sigma K_{\mathcal{I}}} f = M(\phi) f + \int_0^{\sigma} d\tau e^{\tau K^{\mathcal{I}}} M(\dot{\phi}) e^{-\tau K_{\mathcal{I}}} f, \quad \sigma \geq 0.$$

Therefore, using representations (1.2) and (2.10) we obtain

$$\begin{aligned} V(t, t + \sigma) U(t + \sigma, t) \phi(t + \sigma) \chi_{\mathcal{I}}(t + \sigma) f(t) = \\ \phi(t) f(t) + \int_0^{\sigma} d\tau V(t, t + \tau) U(t + \tau, t) \dot{\phi}(t + \tau) \chi_{\mathcal{I}}(t + \tau) f(t) \end{aligned}$$

for $t \in \mathcal{I}$ and $\sigma \geq 0$. Put $s := t + \sigma$. Then we get

$$\begin{aligned} V(t, s) U(s, t) \phi(s) \chi_{\mathcal{I}}(s) f(t) = \\ \phi(t) f(t) + \int_t^s dr V(t, r) U(r, t) \dot{\phi}(r) \chi_{\mathcal{I}}(r) f(t) \end{aligned}$$

for $(s, t) \in \Delta_{\mathcal{I}}$. Let $\bar{\mathcal{I}}_0 \subset \mathcal{I}$ be a closed subinterval such that restriction $\phi \upharpoonright \bar{\mathcal{I}}_0 = 1$. If $s, t \in \bar{\mathcal{I}}_0$, then

$$V(t, s) U(s, t) f(t) = f(t)$$

for $t \in \bar{\mathcal{I}}$. Since $[\text{dom}(K_{\mathcal{I}})]_t$ is dense in X for each $t \in \mathcal{I}$, we prove the first part of the equality (2.12). To prove the second part one has to interchange generators $K_{\mathcal{I}}$ and $K^{\mathcal{I}}$. \square

Corollary 2.9 Let $\tilde{K}_{\mathcal{I}}$ and $\tilde{K}^{\mathcal{I}}$, $p \in [1, \infty)$, be evolution operators in $L^p(\mathcal{I}, X)$. Assume that for each $\phi \in H_a^{1,\infty}(\mathcal{I}) \cap H_b^{1,\infty}(\mathcal{I})$ one has

$$M(\phi) \text{dom}(\tilde{K}_{\mathcal{I}}) \subseteq \text{dom}(\tilde{K}^{\mathcal{I}}) \quad \text{and} \quad M(\phi) \text{dom}(\tilde{K}^{\mathcal{I}}) \subseteq \text{dom}(\tilde{K}_{\mathcal{I}}) \quad (2.17)$$

and

$$\tilde{K}^{\mathcal{I}} M(\phi) f = \tilde{K}_{\mathcal{I}} M(\phi) f, \quad f \in \text{dom}(\tilde{K}_{\mathcal{I}}) \quad \text{or} \quad f \in \text{dom}(\tilde{K}^{\mathcal{I}}). \quad (2.18)$$

If the closures $K_{\mathcal{I}}$ and $K^{\mathcal{I}}$ of the evolution operators $\tilde{K}_{\mathcal{I}}$ and $\tilde{K}^{\mathcal{I}}$ exist and are (respectively) forward and backward generators, then the corresponding forward and backward propagators $\{U(t, s)\}_{(t,s) \in \Delta_{\mathcal{I}}}$ and $\{V(t, s)\}_{(t,s) \in \nabla_{\mathcal{I}}}$ verify the relation (2.12).

Proof. Let $f \in \text{dom}(\tilde{K}_{\mathcal{I}})$. Then from (2.18) and (2.4) we get

$$\tilde{K}^{\mathcal{I}}M(\phi)f = M(\phi)\tilde{K}_{\mathcal{I}}f - M(\dot{\phi})f.$$

Since $\tilde{K}_{\mathcal{I}}$ is closable, for each $f \in \text{dom}(K_{\mathcal{I}})$ there is a sequence $\{f_n\}_{n \in \mathbb{N}}$, $f_n \in \text{dom}(\tilde{K}_{\mathcal{I}})$, such that $\lim_{n \rightarrow \infty} f_n = f$ and $\lim_{n \rightarrow \infty} \tilde{K}_{\mathcal{I}}f_n = K_{\mathcal{I}}f$. Since

$$\tilde{K}^{\mathcal{I}}M(\phi)f_n = M(\phi)\tilde{K}_{\mathcal{I}}f_n - M(\dot{\phi})f_n, \quad n \in \mathbb{N},$$

we get $M(\phi)f \in \text{dom}(K^{\mathcal{I}})$ and

$$K^{\mathcal{I}}M(\phi)f = M(\phi)K_{\mathcal{I}}f - M(\dot{\phi})f$$

for $f \in \text{dom}(K_{\mathcal{I}})$. Using (2.4) we prove $M(\phi)\text{dom}(K_{\mathcal{I}}) \subseteq \text{dom}(K^{\mathcal{I}})$ and (2.14). Similarly, we prove also $M(\phi)\text{dom}(K^{\mathcal{I}}) \subseteq \text{dom}(K_{\mathcal{I}})$ and (2.14). Then application of Proposition 2.8 completes the proof. \square

Now it makes sense to introduce the following definition.

Definition 2.10 A strongly continuous operator-valued function $G(\cdot, \cdot) : \mathcal{I} \times \mathcal{I} \rightarrow \mathcal{B}(X)$ is called a *bidirectional propagator* on $\mathcal{I} \times \mathcal{I}$ if

- (i) $G(t, t) = I_X$ for $t \in \mathcal{I}$,
- (ii) $G(t, r)G(r, s) = G(t, s)$ for $(t, r, s) \in \mathcal{I}^3$,
- (iii) $\sup_{(t,s) \in \mathcal{I} \times \mathcal{I}} \|G(t, s)\|_{\mathcal{B}(X)} < \infty$.

A strongly continuous operator-valued function $G(\cdot, \cdot)$ defined on $\mathbb{R} \times \mathbb{R}$ is called a *bidirectional propagator* on $\mathbb{R} \times \mathbb{R}$, if for any bounded interval \mathcal{I} the restriction of $G(\cdot, \cdot)$ to $\mathcal{I} \times \mathcal{I}$ is a bidirectional propagator.

One can easily verify that if $G(\cdot, \cdot)$ is a bidirectional propagator, then $U(\cdot, \cdot) := G(\cdot, \cdot) \upharpoonright \Delta_{\mathcal{I}}$ and $V(\cdot, \cdot) := G(\cdot, \cdot) \upharpoonright \nabla_{\mathcal{I}}$ are, respectively, for- and backward propagators related by (2.12). Conversely, if $U(\cdot, \cdot)$ and $V(\cdot, \cdot)$ are, respectively, for- and backward propagators, which are related by (2.12), then

$$G(t, s) := \begin{cases} U(t, s), & (t, s) \in \Delta_{\mathcal{I}} \\ V(t, s), & (t, s) \in \nabla_{\mathcal{I}} \end{cases} \quad (2.19)$$

defines a bidirectional propagator.

Definition 2.11 Let $\{A(t)\}_{t \in \mathcal{I}}$ be a measurable family of closed and densely defined linear operators in the separable Banach space X .

- (i) The evolution equation (1.1) is *well-posed* on \mathcal{I} if the for- and backward evolution equations (1.1) are well-posed \mathcal{I} for some $p \in [1, \infty)$.
- (ii) The bidirectional propagator $\{G(t, s)\}_{(t,s) \in \mathcal{I} \times \mathcal{I}}$ is called a *solution* of the bidirectional evolution equation (1.1) on \mathcal{I} if the for- and backward propagators $\{U(t, s)\}_{(t,s) \in \Delta_{\mathcal{I}}}$, $U(\cdot, \cdot) := G(\cdot, \cdot) \upharpoonright \Delta_{\mathcal{I}}$, and $\{V(t, s)\}_{(t,s) \in \nabla_{\mathcal{I}}}$, $V(\cdot, \cdot) := G(\cdot, \cdot) \upharpoonright \nabla_{\mathcal{I}}$, are solutions of the for- and backward equations (1.1) on \mathcal{I} .
- (iii) The well-posed evolution equation (1.1) has a unique solution if the for- and backward evolution equation (1.1) has unique solutions.

Theorem 2.12 Let $\{A(t)\}_{t \in \mathcal{I}}$ be a measurable family of closed and densely defined linear operators in the separable Banach space X . Assume that the bidirectional evolution equation (1.1) is well-posed on \mathcal{I} for some $p \in [1, \infty)$. If the closures $K_{\mathcal{I}}$ and $K^{\mathcal{I}}$ of evolution operators $\tilde{K}_{\mathcal{I}}$ and $\tilde{K}^{\mathcal{I}}$ exist in $L^p(\mathcal{I}, X)$ and are anti-generators and generators, respectively, then the bidirectional evolution equation (1.1) has a unique solution on \mathcal{I} .

Proof. One easily verifies that the operators $\tilde{K}_{\mathcal{I}}$ and $\tilde{K}^{\mathcal{I}}$ defined by (2.8) and (2.11) satisfy the conditions (2.17), (2.18). Then application of Corollary 2.9 completes the proof. \square

2.4 Problems on \mathbb{R}

Let us consider the forward evolution equation (1.1) on \mathbb{R} . A natural way to study this problem is to consider the equation (1.1) on bounded open intervals $\mathcal{I} \subset \mathbb{R}$. In this case one gets a solution $\{U_{\mathcal{I}}(t, s)\}_{(t,s) \in \mathbb{R}}$ for any bounded interval \mathcal{I} . Then we have to guarantee that two solutions $\{U_{\mathcal{I}_1}(t, s)\}_{(t,s) \in \Delta_{\mathcal{I}_1}}$ and $\{U_{\mathcal{I}_2}(t, s)\}_{(t,s) \in \Delta_{\mathcal{I}_2}}$, which correspond to different bounded open intervals \mathcal{I}_1 and \mathcal{I}_2 , are *compatible*, i.e., one has

$$U_{\mathcal{I}_1}(t, s) = U_{\mathcal{I}_2}(t, s), \quad (t, s) \in \Delta_1 \subseteq \Delta_2, \quad (2.20)$$

for $\mathcal{I}_1 \subseteq \mathcal{I}_2$. Below we clarify this compatibility of propagators in terms of evolution generators.

If $\mathcal{I}_1 \subseteq \mathcal{I}_2$, then $L^p(\mathcal{I}_1, X)$ is a subspace of $L^p(\mathcal{I}_2, X)$. Let $Q_{\mathcal{I}_1}$ denote the projection from $L^p(\mathcal{I}_2, X)$ onto the subspace $L^p(\mathcal{I}_1, X)$ given by

$$(Q_{\mathcal{I}_1}f)(t) := \chi_{\mathcal{I}_1}(t)f(t), \quad f \in L^p(\mathcal{I}_2, X).$$

Let intervals $\mathcal{I}_1 = (a_1, b_1)$ and $\mathcal{I}_2 = (a_2, b_2)$ be related by $a_2 \leq a_1 < b_1 \leq b_2$. We set $\mathcal{I}' = (a_1, b_2)$.

Proposition 2.13 *Let $\mathcal{I}_1 = (a_1, b_1)$ and $\mathcal{I}_2 = (a_2, b_2)$ be two bounded intervals such that $\mathcal{I}_1 \subseteq \mathcal{I}_2$. Further, let $K_{\mathcal{I}_1}$ and $K_{\mathcal{I}_2}$ be forward generators, respectively, in $L^p(\mathcal{I}_1, X)$ and $L^p(\mathcal{I}_2, X)$. The corresponding propagators $\{U_{\mathcal{I}_1}(t, s)\}_{(t,s) \in \Delta_{\mathcal{I}_1}}$ and $\{U_{\mathcal{I}_2}(t, s)\}_{(t,s) \in \Delta_{\mathcal{I}_2}}$ are compatible if and only if for any $f \in L^p(\mathcal{I}_2, X)$ obeying $Q_{\mathcal{I}'}f \in \text{dom}(K_{\mathcal{I}_2})$ one has $Q_{\mathcal{I}_1}f \in \text{dom}(K_{\mathcal{I}_1})$ and relation*

$$K_{\mathcal{I}_1}Q_{\mathcal{I}_1}f = Q_{\mathcal{I}_1}K_{\mathcal{I}_2}Q_{\mathcal{I}'}f. \quad (2.21)$$

Proof. We put $K_j := K_{\mathcal{I}_j}$, $U_j(t, s) := U_{\mathcal{I}_j}$, $Q_j := Q_{\mathcal{I}_j}$, $j = 1, 2$, and $Q' := Q_{\mathcal{I}'}$. Assume that the propagators $U_1(t, s)$ and $U_2(t, s)$ are compatible. In this case one easily verifies that

$$e^{-\sigma K_1}Q_1f = Q_1e^{-\sigma K_2}Q'f, \quad \sigma \geq 0, \quad f \in L^p(\mathcal{I}_2, X).$$

Moreover, by

$$\frac{1}{\sigma}(I - e^{-\sigma K_1})Q_1f = \frac{1}{\sigma}Q_1(I - e^{-\sigma K_2})Q'f, \quad \sigma > 0,$$

one gets that $Q'f \in \text{dom}(K_2)$ yields $Q_1f \in \text{dom}(K_1)$ as well as (2.21).

To prove the converse we set

$$W(\sigma)f := e^{-(\tau-\sigma)K_1}Q_1e^{-\sigma K_2}Q'f, \quad 0 \leq \sigma \leq \tau.$$

If $g := Q'f \in \text{dom}(K_2)$, then $g(\sigma) := e^{-\sigma K_2}Q'f \in \text{dom}(K_2)$ for $\sigma \geq 0$. Since

$$Q'e^{-\sigma K_2}Q'f = e^{-\sigma K_2}Q'f, \quad f \in L^p(\mathcal{I}_2, X), \quad \sigma \geq 0, \quad (2.22)$$

we obtain $Q'g(\sigma) = Q'e^{-\sigma K_1}Q'f \in \text{dom}(K_2)$, which yields $Q_1e^{-\sigma K_2}Q'f \in \text{dom}(K_1)$. Hence

$$\frac{d}{d\sigma}W(\sigma)f = e^{-(\tau-\sigma)K_1}(K_1Q_1 - Q_1K_2)e^{-\sigma K_2}Q'f, \quad 0 \leq \sigma \leq \tau.$$

Applying to this equation the relation (2.21), we obtain $\partial_{\sigma}W(\sigma)f = 0$, which yields

$$W(\tau)f = W(0)f, \quad \tau \geq 0,$$

or

$$Q_1e^{-\sigma K_2}Q'f = e^{-\sigma K_1}Q_1f, \quad \sigma \geq 0, \quad (2.23)$$

for all those $f \in L^p(\mathcal{I}_2, X)$ that $Q'f \in \text{dom}(K_2)$.

Now we notice that the set

$$\mathcal{D}' := \{f \in L^p(\mathcal{I}, X) : Q'f \in \text{dom}(K_2)\}$$

is dense in $L^p(\mathcal{I}', X)$. Indeed, let $\phi \in H^{1,\infty}(\mathcal{I}_2)$ such that $\text{supp}(\phi) \subseteq \overline{\mathcal{I}'}$. By (2.3) we have $M(\phi)f \in \text{dom}(K_2)$ for $f \in \text{dom}(K_2)$. By virtue of $Q'M(\phi)f = M(\phi)f$, we get $M(\phi)\text{dom}(K_2) \subseteq \mathcal{D}'$. Since this holds for any $\phi \in H^{1,\infty}(\mathcal{I}_2)$ obeying $\text{supp}(\phi) \subseteq \overline{\mathcal{I}'}$, we immediately find that \mathcal{D}' is a dense in $L^p(\mathcal{I}', X)$.

Hence, the relation (2.21) holds for any $f \in L^p(\mathcal{I}_2, X)$. This implies the compatibility of the propagators $U_1(t, s)$ and $U_2(t, s)$. \square

Corollary 2.14 *Let $\tilde{K}_{\mathcal{I}_1}$ and $\tilde{K}_{\mathcal{I}_2}$, $\mathcal{I}_1 \subseteq \mathcal{I}_2$, be evolution operators such that $Q_{\mathcal{I}'}f \in \text{dom}(\tilde{K}_{a_2})$ yields $Q_{\mathcal{I}_1}f \in \text{dom}(K_{\mathcal{I}_1})$ and the relation*

$$\tilde{K}_{\mathcal{I}_1}Q_{\mathcal{I}_1}f = Q_{\mathcal{I}_1}\tilde{K}_{\mathcal{I}_2}Q_{\mathcal{I}'}f \quad (2.24)$$

holds. If the closures $K_{\mathcal{I}_1}$ and $K_{\mathcal{I}_2}$ of evolution operators $\tilde{K}_{\mathcal{I}_1}$ and $\tilde{K}_{\mathcal{I}_2}$ exists in $L^p(\mathcal{I}, X)$, $p \in [1, \infty)$, and they are forward generators, then the corresponding forward propagators are compatible.

Proof. Let $\mathcal{I}_1 = (a_1, b_1)$ and $\mathcal{I}_2 = (a_2, b_2)$, $\mathcal{I}_1 \subseteq \mathcal{I}_2$. As above we set $K_j := K_{\mathcal{I}_j}$, $U_j(t, s) := U_{\mathcal{I}_j}$, $Q_j := Q_{\mathcal{I}_j}$, $j = 1, 2$, and $Q' := Q_{\mathcal{I}'}$, where $\mathcal{I}' = (a_1, b_2)$, see above. Let $g := Q'f \in \text{dom}(K_{\mathcal{I}_2})$. Then there is a sequence $\{g_n\}_{n \in \mathbb{N}}$, $g_n \in \text{dom}(\tilde{K}_2)$, such that $g_n \rightarrow g$ and $\tilde{K}_2 g_n \rightarrow K_2 g$ as $n \rightarrow \infty$. Let $\phi \in H^{1,\infty}(\mathcal{I}_2)$ such that $\text{supp}(\phi) \subseteq \overline{\mathcal{I}'}$. By (2.3) we have $M(\phi)g_n \in \text{dom}(\tilde{K}_2)$ and $Q'M(\phi)g_n = M(\phi)g_n$, $n \in \mathbb{N}$. Then taking into account (2.24) we obtain

$$\tilde{K}_1 Q_1 M(\phi)g_n = Q_1 \tilde{K}_2 Q' M(\phi)g_n, \quad n \in \mathbb{N}.$$

Using (2.4) we find

$$Q_1 \tilde{K}_2 Q' M(\phi)g_n = Q_2 \tilde{K}_2 M(\phi)g_n = M(\phi)Q_1 \tilde{K}_2 g_n + M(\dot{\phi})Q_1 g_n, \quad n \in \mathbb{N},$$

which yields

$$Q_1 \tilde{K}_2 Q' M(\phi)g_n \rightarrow Q_1 K_2 M(\phi)Q'f \quad \text{as } n \rightarrow \infty.$$

Hence, we obtain

$$\tilde{K}_1 Q_1 M(\phi)g_n \rightarrow Q_1 K_2 M(\phi)Q'f \quad \text{as } n \rightarrow \infty,$$

which proves

$$\tilde{K}_1 Q_1 M(\phi)g_n \rightarrow K_1 Q_1 M(\phi)f \quad \text{as } n \rightarrow \infty$$

and

$$K_1 Q_1 M(\phi)f = Q_1 K_1 M(\phi)Q'f.$$

Using (2.4) we also get

$$K_1 Q_1 M(\phi)f = M(\phi)Q_1 K_2 Q'f + M(\dot{\phi})Q_1 f \quad (2.25)$$

for $\phi \in H^{1,\infty}(\mathcal{I}_2)$ obeying $\text{supp}(\phi) \subseteq \overline{\mathcal{I}'}$.

Let us put

$$\phi_\delta(t) := \begin{cases} 0 & t \in (a_2, a_1] \\ (t - a_1)/\delta & t \in (a_1, a_1 + \delta) \\ 1 & t \in [a_1 + \delta, b_2) \end{cases}$$

where $\delta > 0$. Then by (2.25) we obtain

$$K_1 Q_1 M(\phi_\delta)f = M(\phi_\delta)Q_1 K_2 Q'f + \frac{1}{\delta}M(\chi_{(a_1, a_1 + \delta)})Q_1 f \quad (2.26)$$

for any $\delta > 0$. If $g \in L^p(\mathcal{I}, X)$ is continuous at $t = a_1$ and $g(a_1) = 0$, then

$$s - \lim_{\tau \rightarrow 0} \frac{1}{\delta} M(\chi_{(a_1, a_1 + \delta)})g = 0.$$

Since $Q'f$ is continuous, one has $f(a_1) = 0$. Hence

$$s - \lim_{\tau \rightarrow 0} \frac{1}{\delta} M(\chi_{(a_1, a_1 + \delta)})Q'f = 0.$$

Since $s - \lim_{\delta \rightarrow 0} M(\phi_\delta) = Q'$, from (2.26) we obtain that

$$\lim_{\delta \rightarrow 0} K_1 Q_1 M(\phi_\delta) f = Q_1 K_2 Q' f$$

and

$$\lim_{\delta \rightarrow 0} Q_1 M(\phi_\delta) f = Q_1 f.$$

This yields $Q_1 f \in \text{dom}(K_1)$ and $K_1 Q_1 f = Q_1 K_2 Q' f$. Applying now Proposition 2.13 one completes the proof. \square

Definition 2.15 Let $\{A(t)\}_{t \in \mathbb{R}}$ be a measurable family of closed and densely defined linear operators in the separable Banach space X .

(i) The forward evolution equation (1.1) is *well-posed* on \mathbb{R} for some $p \in [1, \infty)$ if for any bounded open interval \mathcal{I} of \mathbb{R} the operator $\tilde{K}_{\mathcal{I}}$ is an evolution operator.

(ii) A forward propagator $\{U(t, s)\}_{(t, s) \in \Delta_{\mathbb{R}}}$ is called a *solution* of the well-posed forward evolution equation (1.1) on \mathbb{R} if $\{U_{\mathcal{I}}(t, s)\}_{(t, s) \in \Delta_{\mathcal{I}}}$, $U_{\mathcal{I}}(\cdot, \cdot) := U(\cdot, \cdot) \upharpoonright \Delta_{\mathcal{I}}$, is a solution of the forward evolution equation (1.1) for any bounded interval \mathcal{I} of \mathbb{R} .

(iii) The well-posed forward evolution equation (1.1) on \mathbb{R} has a unique solution if for any bounded interval $\mathcal{I} \subseteq \mathbb{R}$ the forward evolution equation (1.1) admits a unique solution.

This definition can be extended (*mutatis mutandis*) to backward and to bidirectional evolution equations on \mathbb{R} .

Theorem 2.16 Let $\{A(t)\}_{t \in \mathbb{R}}$ be a measurable family of closed and densely defined linear operators in the separable Banach space X . Assume that the forward evolution equation (1.1) is well-posed on \mathbb{R} for some $p \in [1, \infty)$. If for any bounded open interval \mathcal{I} of \mathbb{R} the closure $K_{\mathcal{I}}$ of the evolution operator $\tilde{K}_{\mathcal{I}}$ exists in $L^p(\mathcal{I}, X)$, $p \in [1, \infty)$, and it is an anti-generator, then the forward evolution equation (1.1) has a unique solution on \mathbb{R} .

Proof. Let $\mathcal{I}_1 \subseteq \mathcal{I}_2$. One can easily verify that the evolution operators $\tilde{K}_{\mathcal{I}_1}$ and $\tilde{K}_{\mathcal{I}_2}$, which are given by (2.8), satisfy the condition (2.24). Since the operators $\tilde{K}_{\mathcal{I}_1}$ and $\tilde{K}_{\mathcal{I}_2}$ are closable and their closures are already forward evolution generators, one gets from Corollary 2.14 that the corresponding forward propagators (they exist and are unique by Theorem 2.4) are compatible. \square

Proposition 2.13, Corollary 2.15 and Theorem 2.16 can be generalized (*mutatis mutandis*) to backward and bidirectional evolution equations.

3 Semigroup perturbations

Theorem 2.4 shows that the problem of the unique solution of the forward or backward evolution equation (1.1) can be transformed to the question: whether the evolution operators $\tilde{K}_{\mathcal{I}}$ or $\tilde{K}^{\mathcal{I}}$ is

are closable and their closures $K_{\mathcal{I}}$ or $K^{\mathcal{I}}$ are anti-generators or generators in $L^p(\mathcal{I}, X)$ for some $p \in [1, \infty)$? In applications $\{A(t)\}_{t \in \mathcal{I}}$ is often a measurable family of anti-generators or generators belonging uniformly to the class $\mathcal{G}(M, \beta)$, for some constants M and β . One can easily verify that in this case the induced multiplication operator A is an anti-generator or generator in $L^p(\mathcal{I}, X)$.

This reduces the problem to the following one: Let T and A be anti-generators or generators in some Banach space \mathfrak{X} , is it possible to find conditions ensuring that their operator sum \tilde{K} :

$$\tilde{K}f = Tf + Af, \quad \text{dom}(T) \cap \text{dom}(A), \quad (3.1)$$

is closable in \mathfrak{X} and its closure K is an anti-generator or generator? To prove such kind of result we rely on the following theorem.

Theorem 3.1 *Let in \mathfrak{X} the operators T and A be generators both belonging to the class $\mathcal{G}(1, 0)$. If $\text{dom}(T) \cap \text{dom}(A)$ is dense in \mathfrak{X} and $\text{ran}(T + A + \xi)$ is dense in \mathfrak{X} for some $\xi < 0$, then \tilde{K} is closable and its closure K is a generator from the class $\mathcal{G}(1, 0)$.*

This theorem was originally proved by Kato, see [21, Theorem IX.2.11], however, under the additional assumption that \tilde{K} is closable. This condition was dropped by Da Prato and Grisvard in [4, Theorem 5.6].

In general, the assumption $T, A \in \mathcal{G}(1, 0)$ is too restrictive for our purposes. So, we modify this assumption. It is known that in general it is possible to find in the Banach space \mathfrak{X} a new norm such that one of the operators: T or A , becomes a generator of the contraction semigroups on \mathfrak{X} . Indeed, since T is the generator of C_0 semigroup, i.e. $T \in \mathcal{G}(M, \beta_T)$, one has:

$$\|e^{\sigma T} f\| \leq M_T e^{\beta_T \sigma}. \quad (3.2)$$

Setting

$$|||f||| := \sup_{\sigma > 0} e^{-\beta_T \sigma} \|e^{\sigma T} f\|$$

one immediately gets that

$$|||e^{\tau T} f||| = e^{\beta_T \tau} \sup_{\sigma > 0} e^{-\beta_T \{\sigma + \tau\}} \|e^{\tau T} f\|.$$

This observation shows that in the Banach space \mathfrak{X} endowed with the norm $|||\cdot|||$ the semigroup $\{e^{\sigma T}\}_{\sigma}$ belongs to the class $\mathcal{G}(1, \beta_T)$ of *quasi-contractive* semigroups. Since

$$\|f\| \leq |||f||| \leq M_T \|f\|,$$

the norm $|||\cdot|||$ is equivalent to $\|\cdot\|$. The same reasoning can be applied to the semigroup $\{e^{\sigma A}\}_{\sigma}$, but in general it is *impossible* to find an equivalent norm such that *both* semigroups become quasi-contractive.

Definition 3.2 Let T and A be generators of C_0 -semigroups $e^{\sigma T}$ and $e^{\sigma A}$ in \mathfrak{X} . The pair $\{T, A\}$ is called *renormalizable* with constants β_A and β_T if for any sequences $\{\tau_k\}_{k=1}^N$, $\tau_k \geq 0$, and $\{\sigma_k\}_{k=1}^N$, $\sigma_k \geq 0$, $n \in \mathbb{N}$, one has

$$\sup_{\substack{\tau_1 \geq 0, \dots, \tau_n \geq 0 \\ \sigma_1 \geq 0, \dots, \sigma_n \geq 0 \\ n \in \mathbb{N}}} e^{-\beta_T \sum \tau_k} e^{-\beta_A \sum \sigma_k} \|e^{\tau_1 T} e^{\sigma_1 A} \dots e^{\tau_n T} e^{\sigma_n A} f\| < \infty \quad (3.3)$$

for each $f \in \mathfrak{X}$. In an obvious manner the definition carries over to pairs $\{T, A\}$ of anti-generators.

Remark 3.3 In the following we formulate the statements in terms of pairs of generators. However, it is easily to see that these statements remain true for pairs of anti-generators.

Lemma 3.4 (Lemma 5.1, [34]) *Let T and A be generators of C_0 -semigroups in \mathfrak{X} . There is an equivalent norm $||| \cdot |||$ on \mathfrak{X} and such that $T \in \mathcal{G}(1, \beta_T)$ and $A \in \mathcal{G}(1, \beta_A)$ if and only if the pair $\{T, A\}$ is renormalizable with constants β_T and β_A .*

Proof. Let the pair $\{T, A\}$ be renormalizable with constants β_T and β_A . On the space \mathfrak{X} we define a norm by

$$|||f||| := \sup_{\substack{\tau_1 \geq 0, \dots, \tau_n \geq 0 \\ \sigma_1 \geq 0, \dots, \sigma_n \geq 0 \\ n \in \mathbb{N}}} e^{-\beta_T \sum \tau_k} e^{-\beta_A \sum \sigma_k} \|e^{\tau_1 T} e^{\sigma_1 A} \dots e^{\tau_n T} e^{\sigma_n A} f\|.$$

Obviously, we have $\|f\| \leq |||f|||$, $f \in \mathfrak{X}$. On the other hand, by the *uniform boundedness principle*, see e.g. [21, Theorem I.1.29], we find that the value of

$$M := \sup_{\substack{\tau_1 \geq 0, \dots, \tau_n \geq 0 \\ \sigma_1 \geq 0, \dots, \sigma_n \geq 0 \\ n \in \mathbb{N}, \|f\| \leq 1}} e^{-\beta_T \sum \tau_k} e^{-\beta_A \sum \sigma_k} \|e^{\tau_1 T} e^{\sigma_1 A} \dots e^{\tau_n T} e^{\sigma_n A} f\|$$

is finite, which yields $|||f||| \leq M\|f\|$, $f \in \mathfrak{X}$. Hence, the norms $\|\cdot\|$ and $|||\cdot|||$ are equivalent. Moreover, it turns out that $T \in \mathcal{G}(M, \beta_T)$ and $A \in \mathcal{G}(M, \beta_A)$. Furthermore, a straightforward computation shows that

$$\begin{aligned} |||e^{\tau T} f||| &\leq e^{\beta_T \tau} |||f|||, & f \in \mathfrak{X}, \\ |||e^{\sigma A} f||| &\leq e^{\beta_A \sigma} |||f|||, & f \in \mathfrak{X}. \end{aligned}$$

Therefore, in the Banach space $\{\mathfrak{X}, |||\cdot|||\}$ the generators T and A belong, respectively, to $\mathcal{G}(1, \beta_T)$ and $\mathcal{G}(1, \beta_A)$.

Conversely, if there is an equivalent norm $|||\cdot|||$ in the Banach space \mathfrak{X} such that $T \in \mathcal{G}(1, \beta_T)$ and $A \in \mathcal{G}(1, \beta_A)$, then a straightforward computation yields (3.3), i.e., the pair $\{T, A\}$ is renormalizable with constants β_T and β_A . \square

Definition 3.5 Let \mathfrak{Y} be a Banach space which is densely and continuously embedded into the Banach space \mathfrak{X} , i.e. $\mathfrak{Y} \hookrightarrow \mathfrak{X}$, and let the operator T be the generator of a C_0 -semigroup in \mathfrak{X} . The Banach space \mathfrak{Y} is called *admissible* with respect to T , if the space \mathfrak{Y} is *invariant* with respect to the semigroup $e^{\sigma T}$, i.e.

$$e^{\sigma T} \mathfrak{Y} \subseteq \mathfrak{Y}, \quad \sigma \geq 0,$$

and restriction $e^{\sigma \hat{T}} := e^{\sigma T} \upharpoonright \mathfrak{Y}$, $\sigma \geq 0$, is a C_0 -semigroup on \mathfrak{Y} .

If $J : \mathfrak{Y} \rightarrow \mathfrak{X}$ is the *embedding operator* of \mathfrak{Y} into \mathfrak{X} , then we get

$$e^{\sigma T} Jf = J e^{\sigma \hat{T}} f, \quad f \in \mathfrak{Y},$$

which yields

$$TJf = J\hat{T}f, \quad f \in \text{dom}(\hat{T}).$$

Lemma 3.6 *Let \hat{T} and \hat{A} be generators of C_0 -semigroups of class $\mathcal{G}(1, 0)$ in the Banach space \mathfrak{Y} . If either $\text{dom}(\hat{T}^*)$ or $\text{dom}(\hat{A}^*)$ are dense in \mathfrak{Y}^* , then for any $\xi < 0$ one gets the inequality:*

$$|\xi| \|g\|_{\mathfrak{Y}^*} \leq \|\hat{T}^* g + \hat{A}^* g + \xi g\|_{\mathfrak{Y}^*}, \quad g \in \text{dom}(\hat{T}^*) \cap \text{dom}(\hat{A}^*). \quad (3.4)$$

Proof. Let $\text{dom}(\widehat{A}^*)$ be dense in \mathfrak{Y}^* . We define

$$\widehat{A}_\alpha := \widehat{A}(I + \alpha\widehat{A})^{-1}, \quad \alpha < 0.$$

Since $\widehat{A} \in \mathcal{G}(1, 0)$ we have $\widehat{A}_\alpha \in \mathcal{G}(1, 0)$ for $\alpha < 0$. Further, we set

$$\widehat{K}_\alpha f := \widehat{T}f + \widehat{A}_\alpha f, \quad f \in \text{dom}(\widehat{K}) := \text{dom}(\widehat{T}), \quad \alpha < 0.$$

Since $\widehat{T} \in \mathcal{G}(1, 0)$ and $\widehat{A}_\alpha \in \mathcal{G}(1, 0)$ we find that $\widehat{K}_\alpha \in \mathcal{G}(1, 0)$, $\alpha < 0$. This yields the estimate

$$\|(\widehat{K}_\alpha + \xi)^{-1}f\|_{\mathfrak{Y}} \leq \frac{1}{|\xi|} \|f\|_{\mathfrak{Y}}, \quad f \in \mathfrak{Y}, \quad \alpha < 0, \quad \xi < 0.$$

Hence, we obtain

$$\|(\widehat{K}_\alpha^* + \xi)^{-1}g\|_{\mathfrak{Y}^*} \leq \frac{1}{|\xi|} \|g\|_{\mathfrak{Y}^*}, \quad g \in \mathfrak{Y}^*, \quad \alpha < 0, \quad \xi < 0,$$

or

$$|\xi| \|g\|_{\mathfrak{Y}^*} \leq \|(\widehat{K}_\alpha^* + \xi)g\|_{\mathfrak{Y}^*}, \quad g \in \text{dom}(\widehat{K}_\alpha^*) = \text{dom}(\widehat{T}^*), \quad \alpha < 0, \quad \xi < 0. \quad (3.5)$$

Note that

$$\widehat{K}_\alpha^* g = \widehat{T}^* g + \widehat{A}_\alpha^* g, \quad g \in \text{dom}(\widehat{T}^*), \quad \alpha < 0, \quad \xi < 0.$$

Now, since $\text{dom}(\widehat{A}^*)$ is dense in \mathfrak{Y}^* , we get

$$s - \lim_{\alpha \rightarrow 0} (I + \alpha\widehat{A}^*)^{-1} = I, \quad \alpha < 0,$$

which yields

$$\lim_{\alpha \rightarrow 0} \widehat{K}_\alpha^* g = \widehat{T}^* g + \widehat{A}^* g, \quad \alpha < 0,$$

for $g \in \text{dom}(\widehat{T}^*) \cap \text{dom}(\widehat{A}^*)$. Then in the limit $\alpha \rightarrow 0$ the inequality (3.5) gives (3.4).

The proof is similar, if one supposes that $\text{dom}(\widehat{T}^*)$ is dense in \mathfrak{Y}^* . \square

Corollary 3.7 *Let T and A be generators of C_0 -semigroups of class $\mathcal{G}(1, 0)$ on \mathfrak{X} . Further, let $\mathfrak{Y} \hookrightarrow \mathfrak{X}$ be admissible with respect to T, A and let the operator A be such that*

$$\mathfrak{Y} \subseteq \text{dom}(A). \quad (3.6)$$

Assume that the induced generators \widehat{T} and \widehat{A} are of the class $\mathcal{G}(1, 0)$. If $\text{dom}(A^)$ is dense in \mathfrak{X}^* , then*

$$|\xi| \|g\|_{\mathfrak{Y}^*} \leq \|\widehat{T}^* g + \widehat{A}^* g + \xi g\|_{\mathfrak{Y}^*}, \quad g \in \text{dom}(\widehat{T}^*) \cap J^* \mathfrak{X}^*, \quad (3.7)$$

for $\xi < 0$ where $J : \mathfrak{Y} \rightarrow \mathfrak{X}$ is the embedding operator.

Proof. By condition (3.6) we get that $\text{dom}(\widehat{A}^*) \supseteq J^* \mathfrak{X}^*$. Let $g \in \text{dom}(\widehat{T}^*) \cap J^* \mathfrak{X}^*$. Then there is $h \in \mathfrak{X}^*$ such that $g = J^* h$. Hence

$$\widehat{K}_\alpha^* J^* h = \widehat{T}^* J^* h + \widehat{A}^* J^* (I + \alpha A^*)^{-1} h, \quad \alpha < 0.$$

By condition (3.6) the operator $B := AJ : \mathfrak{Y} \rightarrow \mathfrak{X}$ is bounded. This yields the representation

$$\widehat{K}_\alpha^* J^* h = \widehat{T}^* J^* h + B^* (I + \alpha A^*)^{-1} h.$$

Since $\text{dom}(A^*)$ is dense in \mathfrak{X}^* we have $s - \lim_{\alpha \rightarrow 0} (I + \alpha A^*)^{-1} = I$. Hence

$$\lim_{\alpha \rightarrow 0} \widehat{K}_\alpha^* J^* g = \widehat{T}^* J^* h + B^* h = \widehat{T}^* g + \widehat{A}^* g.$$

Using (3.5) we get (3.7). \square

Theorem 3.8 (Theorem 5.5, [34]) *Let $\{T, A\}$ be a renormalizable pair of generators of C_0 -semigroups on \mathfrak{X} . Further, let the Banach space $\mathfrak{Y} \hookrightarrow \mathfrak{X}$ be admissible with respect to operators T and A . Assume that A satisfies condition (3.6) and that the pair $\{\widehat{T}, \widehat{A}\}$ is renormalizable. If either one of the domains $\text{dom}(\widehat{T}^*)$, $\text{dom}(\widehat{A}^*)$ is dense in \mathfrak{Y}^* , or $\text{dom}(A^*)$ is dense in \mathfrak{X}^* , then the closure K of \widetilde{K} ,*

$$\widetilde{K}f := Tf + Af, \quad \text{dom}(\widetilde{K}_{\mathcal{I}}) = \text{dom}(T) \cap \text{dom}(A),$$

exists and K is the generator of a C_0 -semigroup.

Proof. Since the pairs $\{T, A\}$ and $\{\widehat{T}, \widehat{A}\}$ are renormalizable we can assume without loss of generality that $T, A \in \mathcal{G}(1, 0)$ as well as $\widehat{T}, \widehat{A} \in \mathcal{G}(1, 0)$. It is obvious that

$$TJf = J\widehat{T}f, \quad f \in \text{dom}(\widehat{T}),$$

and

$$AJf = J\widehat{A}f, \quad \text{dom}(\widehat{A}).$$

By condition (3.6) we get that $J^*\mathfrak{X}^* \subseteq \text{dom}(\widehat{A}^*)$. Since $\text{dom}(\widehat{T})$ is dense in \mathfrak{Y} and \mathfrak{Y} is densely embedded in \mathfrak{X} , we get that the operator \widetilde{K} is densely defined. In particular, we have

$$J \text{dom}(\widehat{T}) \subseteq \text{dom}(\widetilde{K}).$$

Let $g \in \text{dom}(\widetilde{K}^*) \subseteq \mathfrak{X}^*$. Then we have

$$\langle \widetilde{K}Jf, g \rangle = \langle TJf, g \rangle + \langle Bf, g \rangle = \langle J\widehat{T}f, g \rangle + \langle f, B^*g \rangle$$

for $f \in \text{dom}(\widehat{T})$. Hence

$$\langle \widehat{T}f, J^*g \rangle = \langle f, B^*g \rangle - \langle f, J^*\widetilde{K}^*g \rangle, \quad f \in \text{dom}(\widehat{T}),$$

which yields $J^*\text{dom}(\widetilde{K}^*) \subseteq \text{dom}(\widehat{T}^*)$. Since $J^*\mathfrak{X}^* \subseteq \text{dom}(\widehat{A}^*)$ we obtain

$$J^*\text{dom}(\widetilde{K}^*) \subseteq \text{dom}(\widehat{T}^*) \cap \text{dom}(\widehat{A}^*). \quad (3.8)$$

Now, assume that $\text{ran}(\widetilde{K} + \xi)$ is not dense in \mathfrak{X} for some $\xi < 0$. In this case there is a $g \in \mathfrak{X}^*$ such that

$$\langle (\widetilde{K} + \xi)f, g \rangle = 0, \quad f \in \text{dom}(\widetilde{K}).$$

Hence $g \in \text{dom}(\widetilde{K}^*)$ and $(\widetilde{K}^* + \xi)g = 0$. By (3.8) we obtain

$$J^*g \in \text{dom}(\widehat{T}^*) \cap \text{dom}(\widehat{A}^*).$$

If either $\text{dom}(\widehat{T}^*)$ or $\text{dom}(\widehat{A}^*)$ is dense in \mathfrak{Y}^* , then by Lemma 3.6 we get $J^*g = 0$, which yields $g = 0$. If $\text{dom}(A^*)$ is dense in \mathfrak{X}^* , then we apply Corollary 3.7 and find also $J^*g = 0$, which yields $g = 0$. Hence, the range $\text{ran}(\widetilde{K} + \xi)$ is dense in \mathfrak{X} . We note that by virtue of $T, A \in \mathcal{G}(1, 0)$ the operator \widetilde{K} is closable. Indeed, one has the estimate

$$|\xi| \|f\| \leq \|\widetilde{K}f + \xi f\|, \quad f \in \text{dom}(\widetilde{K}), \quad \xi < 0,$$

which yields the existence of the closure K . Applying now Theorem IX.2.11 of [21] one completes the proof. \square

Remark 3.9 Under the assumptions of Theorem 3.8 one can easily verify that the set $\mathfrak{D} := J\text{dom}(\widehat{T}) \subseteq \mathfrak{X}$ is a core of K , i.e., the closure of the restriction $K \upharpoonright \mathfrak{D}$ coincides with K . This follows from the observation that in fact we have proved the density of the set $(\widetilde{K} + \xi)\mathfrak{D}$, $\xi < 0$, in the space \mathfrak{X} .

Taking into account Theorem 3.8 and [47] one immediately obtains the following corollary.

Corollary 3.10 *Let the assumptions of Theorem 3.8 be satisfied. If either one of the domains $\text{dom}(\widehat{T}^*)$, $\text{dom}(\widehat{A}^*)$ is dense in \mathfrak{Y}^* , or $\text{dom}(A^*)$ is dense in \mathfrak{X}^* , then the Trotter product formula*

$$s - \lim_{n \rightarrow \infty} \left(e^{\sigma T/n} e^{\sigma A/n} \right)^n = e^{\sigma K}$$

holds uniformly in $\sigma \in [0, \sigma_0]$, for any $\sigma_0 > 0$.

4 Solutions of evolution equations

4.1 Solutions of forward evolution equations

Let $\{A(t)\}_{t \in \mathcal{I}}$ be a measurable family of anti-generators of class $\mathcal{G}(M, \beta)$, in the separable Banach space X . By A we denote the multiplication operator induced by (2.6) and (2.7) in the Banach space $\mathfrak{X} = L^p(\mathcal{I}, X)$, $1 \leq p < \infty$. Notice that A is an anti-generator of a C_0 -semigroup on $\mathfrak{X} = L^p(\mathcal{I}, X)$ of class $\mathcal{G}(M, \beta)$.

Definition 4.1 ([22, 23]) Let $\{A(t)\}_{t \in \mathcal{I}}$ be a measurable family of anti-generators of C_0 -semigroups in the separable Banach space X . The family is called *forward stable*, if there are constants $M > 0$ and $\beta \geq 0$ such that the estimate

$$\|e^{-\sigma_1 A(t_1)} e^{-\sigma_2 A(t_2)} \dots e^{-\sigma_n A(t_n)}\|_{\mathcal{B}(X)} \leq M e^{\beta \sum_{k=1}^n \sigma_k}$$

holds for each sequences $\{\sigma_k\}_{k=1}^n$, $\sigma_k \geq 0$, and a.e. $(t_1, t_2, \dots, t_n) \in \Delta_n := \{(t_1, t_2, \dots, t_n) \in \mathbb{R}^n : a < t_n \leq t_{n-1} \leq \dots \leq t_1 < b\}$ with respect of the \mathbb{R}^n -Lebesgue measure.

It is clear that if $\{A(t)\}_{t \in \mathcal{I}}$ is forward stable, then the anti-generators $A(t)$ belong to $\mathcal{G}(M, \beta)$ for a.e. $t \in \mathcal{I}$.

Lemma 4.2 (Lemma 5.9, [34]) *Let $\{A(t)\}_{t \in \mathcal{I}}$ be a measurable family of anti-generators of C_0 -semigroups in the separable Banach space X . The pair of anti-generators $\{D_{\mathcal{I}}, A\}$ is renormalizable on $\mathfrak{X} = L^p(\mathcal{I}, X)$, $1 \leq p < \infty$, if and only if the family of anti-generators $\{A(t)\}_{t \in \mathcal{I}}$ is forward stable.*

Definition 4.3 Let $\{A(t)\}_{t \in \mathcal{I}}$ be a measurable family of anti-generators (generators) of class $\mathcal{G}(M, \beta)$ in the separable Banach space X . Further, let Y be a separable Banach space which is densely and continuously embedded into X . The Banach space Y is called *admissible* with respect to the family $\{A(t)\}_{t \in \mathcal{I}}$ if:

- (i) for a.e. $t \in \mathcal{I}$, the Banach space Y is admissible with respect to $A(t)$,
- (ii) there are constants \widehat{M} and $\widehat{\beta}$ such that the anti-generators (generators) $\{\widehat{A}(t)\}_{t \in \mathcal{I}}$ of the induced semigroups belong to $\mathcal{G}(\widehat{M}, \widehat{\beta})$ for a.e. $t \in \mathcal{I}$,
- (iii) the family $\{\widehat{A}(t)\}_{t \in \mathcal{I}}$ is measurable in Y .

We note that the condition (iii) in Definition 4.3 is redundant if X^* is densely embedded into the Banach space Y^* .

Lemma 4.4 (Lemma 5.11, [34]) *Let $\{A(t)\}_{t \in \mathcal{I}}$ be a measurable family of anti-generators in the separable Banach space X belonging to $\mathcal{G}(M, \beta)$ and let the separable Banach space Y be densely and continuously embedded into X . The Banach space $\mathfrak{Y} = L^p(\mathcal{I}, Y)$, $1 \leq p < \infty$, is admissible with respect to the anti-generator A if and only if the family $\{A(t)\}_{t \in \mathcal{I}}$ is admissible with respect to Y .*

Summing up all those properties it is useful for further purposes to introduce the following definition:

Definition 4.5 Let $\{A(t)\}_{t \in \mathcal{I}}$ be a measurable family of anti-generators in the separable Banach space X . Further, let Y be a separable Banach space which is densely and continuously embedded into X . We say the family $\{A(t)\}_{t \in \mathcal{I}}$ satisfies the *forward Kato condition* if :

- (i) $\{A(t)\}_{t \in \mathcal{I}}$ is forward stable in X ,
- (ii) the Banach space Y is admissible with respect to the family $\{A(t)\}_{t \in \mathcal{I}}$,
- (iii) the induced family $\{\widehat{A}(t)\}_{t \in \mathcal{I}}$ is forward stable in Y ,
- (iv) $Y \subseteq \text{dom}(A(t))$ holds for a.e. $t \in \mathcal{I}$,
- (v) $A(\cdot) \upharpoonright Y \in L^\infty(\mathcal{I}, B(Y, X))$.

In the following we use a so-called *Radon-Nikodym property* of certain Banach spaces, see e.g. [8].

We recall that a scalar-valued measure $\mu(\cdot)$ defined on the Borel sets of \mathbb{R} satisfies the Radon-Nikodym property if, for instance, its continuity with respect to the Lebesgue measure implies the existence of a locally summable function $f(\cdot)$ such that $\mu(\delta) = \int_\delta f(x)dx$ for any bounded Borel set $\delta \subset \mathbb{R}$. In general, this property *does not* extend to measures taking their values in Banach spaces. However, there are classes of Banach spaces where this Radon-Nikodym property still holds. For example, *dual* spaces of *separable* Banach spaces admit this property if and only if they are itself separable. This, in particular, yields that the dual Banach space $L^p(\mathcal{I}, Y)^*$, $1 < p < \infty$, is isometric to $L^q(\mathcal{I}, Y^*)$, $\frac{1}{p} + \frac{1}{q} = 1$.

Theorem 4.6 Let $\{A(t)\}_{t \in \mathcal{I}}$ be a measurable family of anti-generators in a separable Banach space X . Further, let Y be a separable Banach space which is densely and continuously embedded into X . If $\{A(t)\}_{t \in \mathcal{I}}$ obeys the forward Kato condition and if, in addition, one of the following conditions :

- (A₁) Y^* satisfies the Radon-Nikodym property,
- (A₂) $\text{dom}(\widehat{A}^*(t))$ is dense in Y^* for a.e. $t \in \mathcal{I}$,
- (A₃) $\text{dom}(A(t)^*)$ is dense in X^* for a.e. $t \in \mathcal{I}$

holds, then the forward evolution equation (1.1) is well-posed on \mathcal{I} for some $p \in (1, \infty)$ and has a unique solution.

Proof. By Lemma 4.2 the pair $\{D_{\mathcal{I}}, A\}$ of anti-generators is renormalizable. Further, let us consider the Banach space $\mathfrak{Y} = L^p(\mathcal{I}, Y)$, $1 < p < \infty$. Since Y is densely and continuously embedded into X the Banach space \mathfrak{Y} is densely and continuously embedded in $\mathfrak{X} = L^p(\mathcal{I}, X)$. Since the family $\{A(t)\}_{t \in \mathcal{I}}$ is admissible with respect to Y , the operator A is admissible with \mathfrak{Y} , cf. Lemma 4.4. Then from conditions (iv) and (v) of Definition 4.5 we find that $\mathfrak{Y} \subseteq \text{dom}(A)$.

Let (see (A₁)) Y^* satisfy the Radon-Nikodym property. Then $\mathfrak{Y}^* = L^p(\mathcal{I}, Y)^* = L^q(\mathcal{I}, Y^*)$, $1/p + 1/q = 1$, which yields that $\text{dom}(\widehat{D}_{\mathcal{I}}^*)$ is dense in \mathfrak{Y}^* . Applying Theorem 3.8 we immediately get that $\widetilde{K}_{\mathcal{I}}$ is closable and its closure K generates a C_0 -semigroup. Taking into account Theorem 2.4 and Theorem 3.8 we complete the proof of the Theorem under condition (A₁).

If Y does not satisfy the Radon-Nikodym property, then the dual space \mathfrak{Y}^* can be identified with a space $L_w^q(\mathcal{I}, Y^*)$, cf. [2]. The space $L_w^q(\mathcal{I}, Y^*)$ consists of equivalence classes $[g]$ of w^* -measurable functions $g(\cdot) : \mathcal{I} \longrightarrow Y^*$ such that $\int_0^T \|g(t)\|_{Y^*}^q dt < \infty$. Two functions $g_1(\cdot) : \mathcal{I} \longrightarrow Y^*$ and $g_2(\cdot) : \mathcal{I} \longrightarrow Y^*$ are called *equivalent*, if $\langle x, g_1(t) \rangle = \langle x, g_2(t) \rangle$ holds for a.e. $t \in \mathcal{I}$ for each

$x \in Y$. Recall that a function $g(\cdot) : \mathcal{I} \longrightarrow Y^*$ is w^* -measurable, if $\langle x, g(\cdot) \rangle$ is measurable for each $x \in Y$. By a straightforward computation we obtain that $(\alpha \widehat{A}^* + \xi)^{-1}$, $\xi > \beta$, $\alpha > 0$, admits the representation

$$\left((\alpha \widehat{A}^* + \xi)^{-1} g \right) (t) = (\alpha \widehat{A}(t)^* + \xi)^{-1} g(t), \quad g \in L_w^q(\mathcal{I}, Y^*). \quad (4.1)$$

Hence,

$$\left\| \left((\alpha \widehat{A}^* + \xi)^{-1} g - g \right) \right\|_{\mathfrak{Y}^*}^q = \int_a^b \left\| \xi (\alpha \widehat{A}(t)^* + \xi)^{-1} g(t) - g(t) \right\|_{Y^*}^q dt.$$

Note that for a.e. $t \in \mathcal{I}$ we have the estimate:

$$\left\| \xi (\alpha \widehat{A}(t)^* + \xi)^{-1} g(t) \right\|_{Y^*} \leq \frac{\widehat{M}\xi}{\xi - \alpha\widehat{\beta}} \|g(t)\|_{Y^*},$$

which yields

$$\left\| \xi (\alpha \widehat{A}(t)^* + \xi)^{-1} g(t) - g(t) \right\|_{Y^*} \leq \left\{ 1 + \frac{\widehat{M}\xi}{\xi - \alpha\widehat{\beta}} \right\} \|g(t)\|_{Y^*}$$

for a.e. $t \in \mathcal{I}$. Since the domain $\text{dom}(\widehat{A}(t)^*)$ is dense in Y^* for a.e. $t \in \mathcal{I}$, by *assumption* (A₂) we get

$$\lim_{\alpha \rightarrow 0} \left\| \xi (\alpha \widehat{A}(t)^* + \xi)^{-1} g(t) - g(t) \right\|_{Y^*} = 0$$

for a.e. $t \in \mathcal{I}$. Hence, by the Lebesgue dominated convergence theorem we obtain

$$\lim_{\alpha \rightarrow 0} \left\| \left((\alpha \widehat{A}^* + \xi)^{-1} g - g \right) \right\| = 0,$$

which shows that $\text{dom}(\widehat{A}^*)$ is dense in \mathfrak{Y}^* . Taking into account Theorem 2.4 and Theorem 3.8 we again conclude that the forward evolution equation (1.1) is well-posed and uniquely solvable.

Finally, by the same reasoning we obtain that under the assumption (A₃) the domain $\text{dom}(A^*)$ is dense in \mathfrak{X}^* . Applying again Theorem 2.4 and Theorem 3.8 we deduce that the evolution equation is well-posed and uniquely solvable. \square

Notice that using (2.5) we get the following representation:

$$\begin{aligned} & \left(\left((e^{-\sigma D_{\mathcal{I}/n}} e^{-\sigma A/n})^n f \right) \right) (t) = \\ & e^{-\sigma A(t-\sigma/n)/n} e^{-\sigma A(t-2\sigma/n)/n} \dots e^{-\sigma A(t-\sigma)/n} \chi_{\mathcal{I}}(t-\sigma) f(t-\sigma) \end{aligned}$$

for a.e. $t \in \mathcal{I}$ and $\sigma \geq 0$.

Corollary 4.7 *If the assumptions of Theorem 4.6 are satisfied, then the propagator can be approximated as follows:*

$$\lim_{n \rightarrow \infty} \int_a^{b-\sigma} \left\| e^{-\frac{\sigma}{n} A(s+\frac{n-1}{n}\sigma)} e^{-\frac{\sigma}{n} A(s+\frac{n-2}{n}\sigma)} \dots e^{-\frac{\sigma}{n} A(s)} x - U(s+\sigma, s)x \right\|^p ds = 0$$

for each $x \in X$ and $0 \leq \sigma \leq b-a$, $1 < p < \infty$.

4.2 Backward and bidirectional evolution equations

To solve the backward evolution equation (1.1) we assume that $\{A(t)\}_{t \in \mathcal{I}}$ is a measurable family of generators of C_0 -semigroups of the class $\mathcal{G}(M, \beta)$. We note that the multiplication operator defined by (2.6) and (2.7) generates a C_0 -semigroup of class $\mathcal{G}(M, \beta)$.

Definition 4.8 Let $\{A(t)\}_{t \in \mathcal{I}}$ be a measurable family of generators of class $\mathcal{G}(M, \beta)$ in a separable Banach space X . The family $\{A(t)\}_{t \in \mathcal{I}}$ is called *backward stable* if

$$\|e^{\sigma_1 A(t_1)} e^{\sigma_2 A(t_2)} \dots e^{\sigma_n A(t_n)}\|_{\mathcal{B}(X)} \leq M e^{\beta \sum_{k=1}^n \sigma_k}$$

is valid for each sequence $\{\sigma_k\}_{k=1}^n$, $\sigma_k \geq 0$ and a.e. $t \in \nabla_n := \{(t_1, t_2, \dots, t_n) \in \mathbb{R}^n : a < t_1 \leq t_2 \leq \dots \leq t_n < b\}$.

Then Lemma 4.2 admits the following analogon.

Lemma 4.9 Let $\{A(t)\}_{t \in \mathcal{I}}$ be a measurable family of semigroup generators in the separable Banach space X , which is supposed to belong to $\mathcal{G}(M, \beta)$. Then the pair $\{D^{\mathcal{I}}, A\}$ is renormalizable on $\mathfrak{X} = L^p(\mathcal{I}, X)$, $1 \leq p < \infty$, if and only if the family of generators $\{A(t)\}_{t \in \mathcal{I}}$ is backward stable.

Proof. Let $\mathcal{I} = (a, b)$. We introduce the isometry $\mathcal{U} : L^p(\mathcal{I}, X) \longrightarrow L^p(\mathcal{I}, X)$, defined by

$$(\mathcal{U}f)(t) = f(a + b - t), \quad t \in \mathcal{I}, \quad f \in \text{dom}(\mathcal{U}) := L^p(\mathcal{I}, X). \quad (4.2)$$

Notice that $\mathcal{U}^2 = I$ which yields $\mathcal{U}^{-1} = \mathcal{U}$. A straightforward computation shows that $\mathcal{U}^{-1} D^{\mathcal{I}} \mathcal{U} = \mathcal{U} D^{\mathcal{I}} \mathcal{U} = -D_{\mathcal{I}}$. Introducing the family

$$A'(t) := -A(a + b - t), \quad t \in \mathcal{I},$$

and the multiplication operator A' in $L^p(\mathcal{I}, X)$ we get that $\mathcal{U}^{-1} A \mathcal{U} = \mathcal{U} A \mathcal{U} = A'$. Hence, $\mathcal{U}^{-1} \{D^{\mathcal{I}}, A\} \mathcal{U} = \mathcal{U} \{D^{\mathcal{I}}, A\} \mathcal{U} = \{-D_{\mathcal{I}}, -A'\}$. Thus, the generator pair $\{D^{\mathcal{I}}, A\}$ is renormalizable if and only if the corresponding anti-generator pair $\{D_{\mathcal{I}}, A'\}$ is renormalizable. From Lemma 4.2 we obtain that $\{D_{\mathcal{I}}, A'\}$ is renormalizable if and only if the family $\{A'(t)\}_{t \in \mathcal{I}}$ is forward stable. On the other hand, $\{A'(t)\}_{t \in \mathcal{I}}$ is forward stable if and only if $\{A(t)\}_{t \in \mathcal{I}}$ is backward stable, that finishes the proof. \square

Definition 4.10 Let $\{A(t)\}_{t \in \mathcal{I}}$ be a measurable family of generators on the separable Banach space X and let Y be a separable Banach space which is densely and continuously embedded into X . We say the family $\{A(t)\}_{t \in \mathcal{I}}$ satisfies the *backward Kato condition* if:

- (i) $\{A(t)\}_{t \in \mathcal{I}}$ is backward stable in X ,
- (ii) the Banach space Y is admissible with respect to the family $\{A(t)\}_{t \in \mathcal{I}}$,
- (iii) the induced family $\{\widehat{A}(t)\}_{t \in \mathcal{I}}$ (see Definition 4.3) is backward stable in Y ,

and, in addition, we assume that conditions (iv) and (v) of Definition 4.5 are valid.

Then, applying Theorem 3.8 we immediately obtain the following statement:

Theorem 4.11 Let $\{A(t)\}_{t \in \mathcal{I}}$ be a measurable family of generators in the separable Banach space X . Further, let Y be a separable Banach space which is densely and continuously embedded into X . If $\{A(t)\}_{t \in \mathcal{I}}$ obeys the backward Kato condition and if in addition one of the conditions (A₁)-(A₃) holds, then the backward evolution equation (1.1) is well-posed on \mathcal{I} for some $p \in (1, \infty)$ and has a unique solution.

Corollary 4.12 If the assumptions of Theorem 4.11 are satisfied, then for each $x \in X$ we obtain approximation of the propagator in the form:

$$\lim_{n \rightarrow \infty} \int_{a+\sigma}^b \left\| e^{\frac{\sigma}{n} A(s - \frac{n-1}{n} \sigma)} e^{\frac{\sigma}{n} A(s - \frac{n-2}{n} \sigma)} \dots e^{\frac{\sigma}{n} A(s)} x - U(s - \sigma, s) x \right\|^p ds = 0$$

for each $x \in X$ and $0 \leq \sigma \leq b - a$, $1 < p < \infty$.

The proofs of the Theorem 4.11 and Corollary 4.12 follow directly from Theorem 4.6 and Corollary 4.7 by using transformation (4.2).

Theorem 4.13 *Let $\{A(t)\}_{t \in \mathcal{I}}$ be a measurable family of group generators in the separable Banach space X and let Y be a separable Banach space, which is densely and continuously embedded into X . If the family $\{A(t)\}_{t \in \mathcal{I}}$ obeys the forward and backward Kato conditions and if one of the conditions (A_1) - (A_3) holds, then the bidirectional evolution equation (1.1) is well-posed on \mathcal{I} for some $p \in (1, \infty)$ and has a unique solution.*

The proof follows directly from Theorem 2.12, Theorem 4.6 and Theorem 4.11. Finally, let us consider bidirectional evolution equations (1.1) on \mathbb{R} .

Theorem 4.14 *Let $\{A(t)\}_{t \in \mathbb{R}}$ be a measurable family of group generators in the separable Banach space X . Further, let Y be a separable Banach space which is densely and continuously embedded into X . If for any bounded open interval of \mathbb{R} the family $\{A(t)\}_{t \in \mathcal{I}}$ obeys the forward and backward Kato conditions and if one of the conditions (A_1) - (A_3) holds, then the bidirectional equation (1.1) is well-posed on \mathbb{R} for some $p \in (1, \infty)$ and admits a unique solution.*

The proof follows from a bidirectional modification of Theorem 2.16 and from Theorem 4.13.

5 Evolution equations in Hilbert spaces

Our next aim is to apply the above results to evolution equations for families of semi-bounded self-adjoint operators $\{H(t)\}_{t \in \mathbb{R}}$ with *time independent form-domains*.

This case was studied by Kisyański in [24]. The main Theorem 8.1 of [24] states that if for all elements of the form-domain, the corresponding closed quadratic form is continuously differentiable for $t \in \mathbb{R}$, then one can associated with the bidirectional evolution equation

$$\frac{1}{i} \frac{\partial}{\partial t} u(t) + H(t)u(t) = 0, \quad u(s) = u_s, \quad s, t \in \mathbb{R}, \quad (5.1)$$

a unique propagator which is called the solution of (5.1). In the present section we elucidate and improve this result.

5.1 Preliminaries

Let $\{H(t)\}_{t \in \mathbb{R}}$ be a family of non-negative self-adjoint operators in a separable Hilbert space \mathfrak{H} . In the following we consider the non-autonomous Cauchy problem (1.4). As above we assume that the family of operators $\{H(t)\}_{t \in \mathbb{R}}$ is measurable. As in [24] we assume also that

$$\mathfrak{D}^+ = \text{dom}(H(t)^{1/2}) \subseteq \mathfrak{H}, \quad t \in \mathbb{R},$$

which means that the domain $\text{dom}(H(t)^{1/2})$ is independent of $t \in \mathbb{R}$. Introducing the scalar products

$$(f, g)_t^+ := (\sqrt{H(t)}f, \sqrt{H(t)}g) + (f, g), \quad t \in \mathbb{R}, \quad f, g \in \mathfrak{D}.$$

one defines a family of Hilbert spaces $\{\mathfrak{H}_t^+\}_{t \in \mathbb{R}}$, which is densely and continuously embedded, $\mathfrak{H}_t^+ \hookrightarrow \mathfrak{H}$, into \mathfrak{H} . The corresponding vector norm is denoted by $\|\cdot\|_t^+$. The natural embedding operator of \mathfrak{H}_t^+ into \mathfrak{H} is denoted by $J_t^+ : \mathfrak{H}_t^+ \longrightarrow \mathfrak{H}$.

By the *closed graph principle* it follows that for each $t, s \in \mathbb{R}$ the constants

$$c(t, s) := \left\| (H(t) + I)^{1/2} (H(s) + I)^{-1/2} \right\|_{\mathcal{B}(\mathfrak{H})}$$

are finite. Obviously, we have

$$\|f\|_t^+ \leq c(t, s)\|f\|_s^+, \quad f \in \mathfrak{D}, \quad t, s \in \mathbb{R},$$

which yields the estimates:

$$\frac{1}{c(t, s)}\|f\|_t^+ \leq \|f\|_s^+ \leq c(s, t)\|f\|_t^+, \quad f \in \mathfrak{D}, \quad t, s \in \mathbb{R}. \quad (5.2)$$

This means that the norms $\|\cdot\|_t^+$ are *mutually equivalent*.

We note that for each $t \in \mathbb{R}$ the Hilbert space \mathfrak{H}_t^+ is admissible with respect to $H(t)$. The corresponding *induced group* (see Definition 3.5) is denoted by $U_t^+(\sigma)$ and is unitary. Its generator is denoted by $H^+(t)$, i.e. $U_t^+(\sigma) = e^{-i\sigma H^+(t)}$. Using the embedding operator J_t^+ one gets that

$$U_t(\sigma)J_t^+f = J_t^+U_t^+(\sigma)f, \quad f \in \mathfrak{H}_t^+, \quad \sigma \in \mathbb{R}, \quad t \in \mathbb{R}, \quad (5.3)$$

Notice that

$$H^+(t)f = H(t)f, \quad f \in \text{dom}(H^+(t)) := \{f \in \text{dom}(H(t)) : H(t)f \in \mathfrak{H}_t^+\},$$

which gives

$$\text{dom}(H^+(t)) = \text{dom}(H(t)^{3/2}).$$

The dual space with respect to the scalar product (\cdot, \cdot) is denoted by \mathfrak{H}_t^- . We note that

$$\mathfrak{H}_t^+ \hookrightarrow \mathfrak{H} \hookrightarrow \mathfrak{H}_t^-, \quad t \in \mathbb{R}.$$

The dual space can be obtained as the completion of the Hilbert space \mathfrak{H} with respect to the norm

$$\|f\|_t^- := \|(H(t) + I)^{-1/2}f\|, \quad f \in \mathfrak{H}.$$

Then from (5.2) we get

$$\frac{1}{c(s, t)}\|f\|_t^- \leq \|f\|_s^- \leq c(t, s)\|f\|_t^-, \quad f \in \mathfrak{H}, \quad t, s \in \mathbb{R},$$

which shows that the set $\mathfrak{D}^- := \mathfrak{H}_t^-$ is independent of t and the norms $\|\cdot\|_t^-$, $t \in \mathbb{R}$, are mutually equivalent. The natural embedding operator of \mathfrak{H} into \mathfrak{H}_t^- is denoted by $J_t^- : \mathfrak{H} \rightarrow \mathfrak{H}_t^-$. Obviously, we have

$$J_t^- = (J_t^+)^* \quad \text{and} \quad J_t^+ = (J_t^-)^*, \quad t \in \mathbb{R}. \quad (5.4)$$

The group $U_t(\sigma)$, $\sigma \in \mathbb{R}$, $t \in \mathbb{R}$, admits a unitary extension to the Hilbert space \mathfrak{H}_t^- , which we denote by $U_t^-(\sigma)$, $\sigma \in \mathbb{R}$, $t \in \mathbb{R}$. The generator of this group is H_t^- , i.e. $U_t^-(\sigma) = e^{-i\sigma H_t^-}$, $\sigma \in \mathbb{R}$, $t \in \mathbb{R}$, and its domain is given by

$$\text{dom}(H^-(t)) = \text{dom}(H(t)^{1/2}) = \mathfrak{D}^+. \quad (5.5)$$

One can verify that the Hilbert space \mathfrak{H} is admissible with respect to H_t^- , $t \in \mathbb{R}$. The corresponding unitary group coincides with $U_t(\sigma)$. One also has

$$U_t^-(\sigma)J_t^-f = J_t^-U_t(\sigma)f, \quad f \in \mathfrak{H}, \quad \sigma \in \mathbb{R}, \quad t \in \mathbb{R}, \quad (5.6)$$

and

$$\text{dom}(H(t)) = \{f \in \text{dom}(H^-(t)) : H^-(t)f \in \mathfrak{H}\}.$$

Since \mathfrak{H}_t^+ is admissible with respect to $H(t)$, one gets that \mathfrak{H}_t^+ is admissible with respect to H_t^- . The natural embedding operator is given by $J_t := J_t^-J_t^+ : \mathfrak{H}_t^+ \rightarrow \mathfrak{H}_t^-$, we obtain:

$$U_t^-(\sigma)J_t f = J_t U_t^+(\sigma)f, \quad f \in \mathfrak{H}_t^+, \quad \sigma \in \mathbb{R}, \quad t \in \mathbb{R},$$

which shows that

$$\text{dom}(H^+(t)) = \{f \in \text{dom}(H^-(t)) : H^-(t)f \in \mathfrak{H}_t^+\}.$$

Moreover, regarding the operator $H(t)$ as an operator acting from \mathfrak{H}_t^+ into \mathfrak{H}_t^- , one finds that $H(t)$ can be extended to a *contraction* $B(t)$ acting from \mathfrak{H}_t^+ into \mathfrak{H}_t^- . Indeed, this follows from the estimate

$$\begin{aligned} \|B(t)f\|_t^- &= \\ \|(H(t) + I)^{-1/2}H(t)f\|_t &\leq \|H(t)^{1/2}f\|_t \leq \|f\|_t^+, \quad f \in \text{dom}(H(t)). \end{aligned} \tag{5.7}$$

Finally, taking into account (5.3)-(5.6) we get the relations:

$$U_t^+(\sigma)^* = U_t^-(-\sigma) \quad \text{and} \quad U_t^-(\sigma)^* = U_t^+(-\sigma), \quad \sigma \in \mathbb{R}, \quad t \in \mathbb{R}.$$

5.2 Auxiliary evolution equation

We consider the Hilbert space

$$X := \mathfrak{H}_{t=0}^- \quad \text{with} \quad \|\cdot\|_X := \|\cdot\|_{t=0}^-$$

and the auxiliary bidirectional evolution equation

$$\frac{\partial}{\partial t}u(t) + iH^-(t)u(t) = 0 \tag{5.8}$$

on \mathbb{R} . To apply results from Section 4 we set $A(t) = iH^-(t)$, $t \in \mathbb{R}$. Obviously, $\{A(t)\}_{t \in \mathbb{R}}$ is a family of group generators in X . Further, we set

$$Y := \mathfrak{H}_{t=0}^+ \quad \text{with} \quad \|\cdot\|_Y := \|\cdot\|_{t=0}^+. \tag{5.9}$$

It turns out that the Hilbert space $Y = \mathfrak{H}_0^+$ is densely and continuous embedded into X and admissible with respect to $\{A(t)\}_{t \in \mathbb{R}}$.

Lemma 5.1 *Let $\{H(t)\}_{t \in \mathbb{R}}$ be a measurable family of non-negative self-adjoint operators defined in a separable Hilbert space \mathfrak{H} such that $\text{dom}(H(t)^{1/2})$ is independent of $t \in \mathbb{R}$. If \mathcal{I} is a bounded open interval and*

$$c_{\mathcal{I}} := \sup_{(t,s) \in \mathcal{I} \times \mathcal{I}} c(t,s) < \infty,$$

then there are constants $M_{\mathcal{I}}$ and $\beta_{\mathcal{I}}$ such that $\{A(t)\}_{t \in \mathcal{I}}$ is a measurable family of group generators belonging to $\mathcal{G}(M_{\mathcal{I}}, \beta_{\mathcal{I}})$.

If the Hilbert space Y is given by (5.9) and there is a constant $\gamma_{\mathcal{I}} > 0$ such that

$$c(t,s) \leq e^{\gamma_{\mathcal{I}}|t-s|}, \quad t, s \in \mathcal{I}, \tag{5.10}$$

holds, then the families $\{A(t)\}_{t \in \mathcal{I}}$ obey the forward and backward Kato conditions, respectively.

Proof. The measurability of the family $\{A(t)\}_{t \in \mathcal{I}}$ follows from the equivalence of weak and strong measurability, see e.g. [1]. Next, we have

$$\begin{aligned} \|e^{\sigma A(t)}x\|_X &= \|e^{i\sigma H^-(t)}x\|_0^- \leq c(0,t)\|e^{i\sigma H^-(t)}x\|_t^- \\ &\leq c(0,t)\|x\|_t^- \leq c(0,t)c(t,0)\|x\|_0^- = c(0,t)c(t,0)\|x\|_X, \end{aligned}$$

$\sigma \in \mathbb{R}$. Hence,

$$\|e^{\sigma A(t)}x\|_X \leq M_{\mathcal{I}}\|x\|_X, \quad x \in X, \quad \sigma \in \mathbb{R}, \quad t \in \mathcal{I},$$

where $M_{\mathcal{I}} := c_{\mathcal{I}}^2$, which yields that $A(t)$ generates a group of the class $\mathcal{G}(M_{\mathcal{I}}, 0)$.

If condition (5.10) is satisfied, then the forward and backward stability of $\{A(t)\}_{t \in \mathcal{I}}$ follows from [43, Theorem 4.3.2].

To prove the measurability of $\{A(t)\}_{t \in \mathcal{I}}$ we note that Y is admissible for a.e. $t \in \mathcal{I}$. Using (5.2) we obtain that the generator $\widehat{A}(t)$ of the induced group (Definition 3.5) belongs to $\mathcal{G}(M_{\mathcal{I}}, 0)$, too. The measurability of the induced family $\{\widehat{A}(t)\}_{t \in \mathcal{I}}$ follows from the equivalence of strong and weak measurability.

The forward and backward stability of $\{\widehat{A}(t)\}_{t \in \mathcal{I}}$ follows again from condition (5.10) and [43, Theorem 4.3.2].

The condition $Y \subseteq \text{dom}(A(t))$ for a.e. $t \in \mathcal{I}$ is obtained from (5.5). The condition $A(\cdot) \upharpoonright Y \in L^\infty(\mathcal{I}, \mathcal{B}(Y, X))$ follows from (5.7). \square

Theorem 5.2 *Let $\{H(t)\}_{t \in \mathbb{R}}$ be a measurable family of non-negative self-adjoint operators defined in a separable Hilbert space \mathfrak{H} such that the domain $\text{dom}(H(t)^{1/2})$ is independent of $t \in \mathbb{R}$. If for any bounded open interval \mathcal{I} the condition (5.10) is satisfied, then the auxiliary bidirectional evolution problem (5.8) is well-posed on \mathbb{R} for $p \in (1, \infty)$ and has a unique solution $\{G^-(t, s)\}_{(t, s) \in \mathbb{R} \times \mathbb{R}}$ obeying the estimate*

$$\|G^-(t, s)x\|_t^- \leq e^{\gamma_{\mathcal{I}}(t-s)} \|x\|_s^-, \quad x \in \mathfrak{H}_s^-, \quad (5.11)$$

for all $(t, s) \in \mathcal{I} \times \mathcal{I}$.

Proof. Since $Y = \mathfrak{H}_0^+$ is a Hilbert space, all conditions (A_1) - (A_3) are satisfied. Using Lemma 5.1 and Theorem (4.14) one gets that the bidirectional evolution equation (5.8) has a unique solution $\{G^-(t, s)\}_{(t, s) \in \mathbb{R} \times \mathbb{R}}$ on \mathbb{R} .

By Corollary 4.7 there is a subsequence $\{n_k\}_{k \in \mathbb{N}}$ such that one has

$$U_{\mathcal{I}}^-(s + \sigma, s)x = s - \lim_{k \rightarrow \infty} e^{-i\frac{\sigma}{n}H^-(s + \frac{n_k-1}{n_k}\sigma)} e^{-i\frac{\sigma}{n}H^-(s + \frac{n_k-2}{n_k}\sigma)} \dots e^{-i\frac{\sigma}{n}H^-(s)}x$$

for each $x \in \mathfrak{H}_s^-$ and a.e. $s \in (a, b - \sigma)$, $0 \leq \sigma \leq b - a$, where $U_{\mathcal{I}}^-(\cdot, \cdot) := G^-(\cdot, \cdot) \upharpoonright \Delta_{\mathcal{I}}$. This yields the estimate

$$\|U_{\mathcal{I}}^-(s + \sigma, t)x\|_{s+\sigma}^- \leq e^{\gamma_{\mathcal{I}}\sigma} \|x\|_s^-, \quad x \in \mathfrak{H}_s^-,$$

for a.e. $s \in (a, b - \sigma)$, $0 \leq \sigma \leq b - a$. Since $U_{\mathcal{I}}^-(\cdot, \cdot)$ is strongly continuous, this holds for any $s \in (a, b - \sigma)$. Setting $t := s + \sigma$ we obtain

$$\|U_{\mathcal{I}}^-(t, s)x\|_t^- \leq e^{\gamma_{\mathcal{I}}|t-s|} \|x\|_s^-, \quad x \in \mathfrak{H}_t^-, \quad (t, s) \in \Delta_{\mathcal{I}}. \quad (5.12)$$

Similarly, using Corollary 4.12 we obtain

$$\|V_{\mathcal{I}}^-(s - \sigma, t)x\|_{s-\sigma}^- \leq e^{\gamma_{\mathcal{I}}\sigma} \|x\|_s^-, \quad x \in \mathfrak{H}_s^-,$$

for $s \in (a + \sigma, b)$, $0 \geq \sigma \geq b - a$, where $V_{\mathcal{I}}^-(\cdot, \cdot) := G^-(\cdot, \cdot) \upharpoonright \nabla_{\mathcal{I}}$. Hence one gets the inequality:

$$\|V_{\mathcal{I}}^-(t, s)x\|_t^- \leq e^{\gamma_{\mathcal{I}}|t-s|} \|x\|_s^-, \quad x \in \mathfrak{H}_t^-, \quad (t, s) \in \nabla_{\mathcal{I}}. \quad (5.13)$$

Using (5.12) and (5.13) we immediately obtain (5.11). \square

5.3 Back to the original problem

Our Theorem 5.2 gives no information about *solvability* of the bidirectional evolution equation (1.4) on \mathbb{R} . This goes back to the fact that in general the evolution equation might be *not* well-posed. In fact, it may happen that the cross-sections of the sets

$$\text{dom}(\widetilde{K}_{\mathcal{I}}) := \text{dom}(D_{\mathcal{I}}) \cap \text{dom}(H_{\mathcal{I}}) = H_a^{1,p}(\mathcal{I}, \mathfrak{H}) \cap \text{dom}(H_{\mathcal{I}})$$

and

$$\text{dom}(\tilde{K}^{\mathcal{I}}) = \text{dom}(D^{\mathcal{I}}) \cap \text{dom}(H_{\mathcal{I}}) = H_b^{1,p}(\mathcal{I}, \mathfrak{H}) \cap \text{dom}(H_{\mathcal{I}}),$$

$p \in (1, \infty)$, are *not dense* in \mathfrak{H} for intervals $\mathcal{I} = (a, b) \subseteq \mathbb{R}$. Recall that $H_{\mathcal{I}}$ is defined as the multiplication operator induced by the family $\{H(t)\}_{t \in \mathcal{I}}$ in $L^p(\mathcal{I}, \mathfrak{H})$.

To avoid this situation we assume in the following that the bidirectional evolution problem (5.1) is well-posed on \mathbb{R} . Naturally, then we face up to the question: whether under this condition the evolution equation (5.1) admits a solution on \mathbb{R} ?

Lemma 5.3 *Let $\{H(t)\}_{t \in \mathbb{R}}$ be a measurable family of non-negative self-adjoint operators defined in the separable Hilbert space \mathfrak{H} such that $\text{dom}(H(t)^{1/2})$ is independent of $t \in \mathbb{R}$. If for any bounded open interval \mathcal{I} the condition (5.10) is satisfied, then there is a unitary bidirectional propagator $\{G(t, s)\}_{(t,s) \in \mathbb{R}^2}$ on \mathfrak{H} , such that*

$$J_0^- G(t, s) = G^-(t, s) J_0^-, \quad (t, s) \in \mathbb{R}^2. \quad (5.14)$$

Moreover, there is a bidirectional propagator $\{G^+(t, s)\}_{(t,s) \in \mathbb{R}^2}$ on \mathfrak{H}_0^+ , such that

$$J_0 G^+(t, s) = G^-(t, s) J_0, \quad (t, s) \in \mathbb{R}^2 \quad (5.15)$$

and

$$J_0^+ G^+(t, s) = G(t, s) J_0^+, \quad (t, s) \in \mathbb{R}^2. \quad (5.16)$$

Proof. Let $J^+ := J_0^+$, $J^- := J_0^-$ and $J := J_0$. We consider the forward case. Let $\mathcal{I} = (a, b)$ be a bounded open interval of \mathbb{R} and let $0 \leq \sigma \leq b - a$. By Corollary 4.7 we get that

$$U^-(\cdot + \sigma, \cdot) J^- x_0 = \lim_{n \rightarrow \infty} \int_s^{L^p(\mathcal{I}_\sigma, X)} e^{-i \frac{\sigma}{n} H^-(\cdot + \frac{n-1}{n} \sigma)} e^{-i \frac{\sigma}{n} H^-(\cdot + \frac{n-2}{n} \sigma)} \dots e^{-i \frac{\sigma}{n} H^-(\cdot)} J^- x_0,$$

$\mathcal{I}_\sigma := (a, b - \sigma)$, for each $x_0 \in \mathfrak{H}$. Since

$$\begin{aligned} e^{-i \frac{\sigma}{n} H^-(s + \frac{n-1}{n} \sigma)} e^{-i \frac{\sigma}{n} H^-(s + \frac{n-2}{n} \sigma)} \dots e^{-i \frac{\sigma}{n} H^-(s)} J^- x_0 = \\ J^- e^{-i \frac{\sigma}{n} H(s + \frac{n-1}{n} \sigma)} e^{-i \frac{\sigma}{n} H(s + \frac{n-2}{n} \sigma)} \dots e^{-i \frac{\sigma}{n} H(s)} x_0 \end{aligned}$$

for a.e. $s \in \mathcal{I}_\sigma$ and since $\{e^{-i \frac{\sigma}{n} H(\cdot + \frac{n-1}{n} \sigma)} e^{-i \frac{\sigma}{n} H(\cdot + \frac{n-2}{n} \sigma)} \dots e^{-i \frac{\sigma}{n} H(\cdot)}\}_{n \in \mathbb{N}}$ is bounded in $L^2(\mathcal{I}_\sigma, \mathfrak{H})$, we obtain that the weak limit

$$U(\cdot + \sigma, \cdot) x_0 := w \int_s^{L^p(\mathcal{I}_\sigma, \mathfrak{H})} \lim_{n \rightarrow \infty} e^{-i \frac{\sigma}{n} H(\cdot + \frac{n-1}{n} \sigma)} e^{-i \frac{\sigma}{n} H(\cdot + \frac{n-2}{n} \sigma)} \dots e^{-i \frac{\sigma}{n} H(\cdot)} x_0$$

exists for each $x_0 \in \mathfrak{H}$ and for each $\sigma \in (0, b - a)$. Hence, we obtain

$$J^- U(s + \sigma, s) x_0 = U^-(s + \sigma, s) J^- x_0$$

for a.e. $s \in \mathcal{I}_\sigma$, $\sigma \in (0, b - a)$ and any $x_0 \in \mathfrak{H}$. We note that

$$\|U(s + \sigma, s) x_0\|_{\mathfrak{H}} \leq \|x_0\|_{\mathfrak{H}}$$

for a.e. $s \in \mathcal{I}_\sigma$ and $\sigma \in (0, b - a)$, $x_0 \in \mathfrak{H}$. Taking into account that the propagator $\{U^-(t, s)\}_{(t,s) \in \Delta_{\mathcal{I}}}$ is strongly continuous, one gets that $\{U(t, s)\}_{(t,s) \in \Delta_{\mathcal{I}}}$ is a weakly continuous family of contractions obeying

$$J^- U(t, s) x_0 = U^-(t, s) J^- x_0 \quad (5.17)$$

for any $(t, s) \in \Delta_{\mathcal{I}}$ and for each $x_0 \in \mathfrak{H}$. Similarly one proves that there is a weakly continuous family of contractions $\{V(t, s)\}_{(t,s) \in \nabla_{\mathcal{I}}}$ such that

$$J^- V(t, s) x_0 = V^-(t, s) J^- x_0 \quad (5.18)$$

holds for $(t, s) \in \nabla_{\mathcal{I}}$ and $x_0 \in \mathfrak{H}$. Setting $G(t, s) := U(t, s)$, $(t, s) \in \Delta_{\mathcal{I}}$, and $G(t, s) := V(t, s)$, $(t, s) \in \nabla_{\mathcal{I}}$, and taking into account that \mathcal{I} is arbitrary, we obtain a weakly continuous family $\{G(t, s)\}_{(t,s) \in \mathbb{R} \times \mathbb{R}}$ of contractions obeying

$$G(t, s) = G(s, t)^{-1}, \quad (t, s) \in \mathbb{R} \times \mathbb{R}.$$

Since for $(t, s) \in \mathbb{R} \times \mathbb{R}$ and any $x_0 \in \mathfrak{H}$ one has

$$\|x_0\|_{\mathfrak{H}} = \|G(s, t)G(t, s)x_0\|_{\mathfrak{H}} \leq \|G(t, s)x_0\|_{\mathfrak{H}} \leq \|x_0\|_{\mathfrak{H}},$$

$\|G(t, s)x_0\|_{\mathfrak{H}} = \|x_0\|_{\mathfrak{H}}$, which shows that $\{G(t, s)\}_{(t,s) \in \mathbb{R} \times \mathbb{R}}$ is a weakly continuous family of unitary operators. However, this immediately yields that $\{G(t, s)\}_{(t,s) \in \mathbb{R} \times \mathbb{R}}$ is in fact a strongly continuous family of unitary operators obeying

$$J^- G(t, s) = G^-(t, s) J^-, \quad (t, s) \in \mathbb{R} \times \mathbb{R}, \quad (5.19)$$

which yields that $\{G(t, s)\}_{(t,s) \in \mathbb{R} \times \mathbb{R}}$ is a unitary propagator.

Now we put

$$V^+(s, t) := U^-(t, s)^*, \quad (t, s) \in \Delta_{\mathbb{R}}, \quad \text{and} \quad U^+(s, t) := V^-(t, s)^*, \quad (t, s) \in \nabla_{\mathbb{R}},$$

as well as

$$G^+(s, t) := G^-(t, s)^*, \quad (t, s) \in \mathbb{R}^2.$$

Then one can easily verify that $\{G^+(t, s)\}_{(t,s) \in \mathcal{I} \times \mathcal{I}}$ is weakly continuous propagator for any bounded interval \mathcal{I} . Taking into account (5.11) and (5.12) we obtain

$$\|V^+(s, t)y\|_s^+ \leq e^{\gamma(t-s)} \|y\|_t^+, \quad y \in \mathfrak{H}_t^+,$$

and

$$\|U^+(t, s)y\|_t^+ \leq e^{\gamma(t-s)} \|y\|_s^+, \quad y \in \mathfrak{H}_s^+,$$

for $s \leq t$. Using the scalar product $(f, g)_s^+ := (\sqrt{H(s) + If}, \sqrt{H(s) + Ig})$, $f, g \in \mathfrak{D}^+$, we get:

$$(\|U^+(t, s)y - y\|_s^+)^2 = (\|U^+(t, s)y\|_s^+)^2 + (\|y\|_s^+)^2 - 2\Re(U^+(t, s)y, y)_s^+.$$

Now, using (5.11) we find

$$\|U^+(t, s)y\|_s^+ \leq e^{\gamma(t-s)} \|U^+(t, s)y\|_t^+ \leq e^{2\gamma(t-s)} \|y\|_s^+,$$

which implies

$$(\|U^+(t, s)y - y\|_s^+)^2 \leq e^{4\gamma(t-s)} (\|y\|_s^+)^2 + (\|y\|_s^+)^2 - 2\Re(U^+(t, s)y, y)_s^+.$$

By the weak continuity of the forward propagator $\{U^+(t, s)\}_{(t,s) \in \Delta_{\mathbb{R}}}$ we obtain $\lim_{t \rightarrow s+0} U^+(t, s) = I$. Hence, $\lim_{t \rightarrow s+0} \|U^+(t, s)y - y\|_s^+ = 0$ for each $y \in \mathfrak{H}_s^+$. Since the norms $\|\cdot\|_t^+$ and $\|\cdot\|_0^+$ are equivalent, we find $\lim_{t \rightarrow s+0} \|U^+(t, s)y - y\|_0^+ = 0$ for each $y \in Y = \mathfrak{H}_0^+$. Similarly we prove $\lim_{t \rightarrow s-0} \|V^+(t, s)y - y\|_0^+ = 0$ for each $y \in Y = \mathfrak{H}_0^+$. Using the representation:

$$G^+(t, s) = G^+(t, 0)G^+(0, s),$$

where

$$G^+(t, 0) = \begin{cases} U^+(t, 0), & t \geq 0, \\ V^+(t, 0), & t \leq 0, \end{cases} \quad \text{and} \quad G^+(0, s) = \begin{cases} V^+(0, s), & s \geq 0, \\ U^+(0, s), & s \leq 0, \end{cases}$$

one proves the strong continuity of the families: $\{G^+(t, 0)\}_{t \in \mathbb{R}}$ and $\{G^+(0, s)\}_{s \in \mathbb{R}}$, which yields the strong continuity of $\{G^+(t, s)\}_{(t,s) \in \mathbb{R}^2}$.

Finally, by $(J^-)^* = J^+$ and $J = J^- J^+$ we find the equation

$$J^+ G^+(s, t) = G(s, t) J^+, \quad (s, t) \in \mathbb{R} \times \mathbb{R},$$

which by virtue of (5.19) proves (5.16). Hence we get that

$$\begin{aligned} JG^+(s, t) &= J^- J^+ G^+(s, t) = \\ J^- G(s, t) J^+ &= G^-(s, t) J^- J^+ = G^-(s, t) J, \quad (s, t) \in \mathbb{R} \times \mathbb{R}, \end{aligned}$$

which proves (5.15). \square

Now it is useful to introduce the following definition.

Definition 5.4 Let $\{G(t, s)\}_{(t,s) \in \mathbb{R} \times \mathbb{R}}$ be a bidirectional propagator in a separable Banach space X and let Y be a separable Banach space, which is densely and continuously embedded into X . The Banach space Y is called admissible with respect to the family $\{G(t, s)\}_{(t,s) \in \mathbb{R} \times \mathbb{R}}$ if there is a bidirectional propagator $\{\widehat{G}(t, s)\}_{(t,s) \in \mathbb{R} \times \mathbb{R}}$ in Y such that

$$G(t, s) J = J \widehat{G}(t, s), \quad (t, s) \in \mathbb{R} \times \mathbb{R}, \quad (5.20)$$

holds where J is the embedding operator of Y into X .

The following theorem generalizes Theorem 8.1 of [24]. Our proof is quite independent from the that in [24].

Theorem 5.5 Let $\{H(t)\}_{t \in \mathbb{R}}$ be a measurable family of non-negative self-adjoint operators defined in the separable Hilbert space \mathfrak{H} such that $\text{dom}(H(t)^{1/2})$ is independent of $t \in \mathbb{R}$. If the bidirectional evolution equation (5.1) is well-posed on \mathbb{R} for some $p \in (1, \infty)$ and the condition (5.10) is satisfied for any bounded open interval, then the bidirectional evolution equation (5.1) admits on \mathbb{R} a unitary solution $\{G(t, s)\}_{(t,s) \in \mathbb{R} \times \mathbb{R}}$ for which the Hilbert space \mathfrak{H}_0^+ is admissible. Moreover, if for any bounded open interval $\mathcal{I} = (a, b)$ the sets

$$H_a^{1,p}(\mathcal{I}, \mathfrak{H}_0^+) \cap \text{dom}(H_{\mathcal{I}}) \quad \text{and} \quad H_b^{1,p}(\mathcal{I}, \mathfrak{H}_0^+) \cap \text{dom}(H_{\mathcal{I}}), \quad p \in (1, \infty), \quad (5.21)$$

are dense in $H_a^{1,p}(\mathcal{I}, \mathfrak{H}_0^+)$ and $H_b^{1,p}(\mathcal{I}, \mathfrak{H}_0^+)$, respectively, then there is only one unitary solution for which the Hilbert space \mathfrak{H}_0^+ is admissible.

Proof. We have to show that the evolution operator $\widetilde{K}_{\mathcal{I}}$,

$$\widetilde{K}_{\mathcal{I}} f = D_{\mathcal{I}} + i H_{\mathcal{I}} f, \quad f \in \text{dom}(\widetilde{K}_{\mathcal{I}}) = \text{dom}(D_{\mathcal{I}}) \cap \text{dom}(H_{\mathcal{I}}),$$

which is associated with the forward evolution equation (5.1), can be extended to a forward generator. Let $\widetilde{K}_{\mathcal{I}}^-$ be evolution operator:

$$\widetilde{K}_{\mathcal{I}}^- g = D_{\mathcal{I}}^- g + i H_{\mathcal{I}}^- g, \quad g \in \text{dom}(\widetilde{K}_{\mathcal{I}}^-) = \text{dom}(D_{\mathcal{I}}^-) \cap \text{dom}(H_{\mathcal{I}}^-),$$

associated with (5.8), where $D_{\mathcal{I}}^-$ is the anti-generator of the right-shift semigroup in $L^p(\mathcal{I}, \mathfrak{H}_0^-)$, and let $H_{\mathcal{I}}^-$ be multiplication operator induced by $\{H^-(t)\}_{t \in \mathcal{I}}$. By \mathcal{J}^- we denote the embedding operator of $L^p(\mathcal{I}, \mathfrak{H})$, $p \in (1, \infty)$, into $L^p(\mathcal{I}, \mathfrak{H}_0^-)$, defined as:

$$(\mathcal{J}^- f)(t) = J^- f(t), \quad f \in L^p(\mathcal{I}, \mathfrak{H})$$

where $J^- := J_0^-$. One can easily verify that $\mathcal{J}^- \text{dom}(\widetilde{K}_{\mathcal{I}}) \subseteq \text{dom}(\widetilde{K}_{\mathcal{I}}^-)$ and

$$\widetilde{K}_{\mathcal{I}}^- \mathcal{J}^- f = \mathcal{J}^- \widetilde{K}_{\mathcal{I}} f, \quad f \in \text{dom}(\widetilde{K}_{\mathcal{I}}).$$

By Theorem 5.2 the forward evolution equation (5.8) is uniquely solvable. This means that the operator $\tilde{K}_{\mathcal{I}}^-$ admits only one extension $K_{\mathcal{I}}^-$, which is a forward generator. In fact, it has been already proven that the closure of $\tilde{K}_{\mathcal{I}}^-$ coincides with $K_{\mathcal{I}}^-$.

By Lemma 5.3 there is a forward generator $\{U_{\mathcal{I}}(t, s)\}_{(t, s) \in \Delta_{\mathcal{I}}}$, $U_{\mathcal{I}}(t, s) := G(t, s) \upharpoonright \Delta_{\mathcal{I}}$ obeying (5.14). By the relation

$$(e^{-\sigma K_{\mathcal{I}}} f)(t) = U_{\mathcal{I}}(t, t - \sigma) \chi_{\mathcal{I}}(t - \sigma) f(t - \sigma), \quad f \in L^p(\mathcal{I}, \mathfrak{H}),$$

one defines a forward generator $K_{\mathcal{I}}$ in $L^p(\mathcal{I}, \mathfrak{H})$. Obviously, we have

$$e^{-\sigma K_{\mathcal{I}}} \mathcal{J}^- f = \mathcal{J}^- e^{-\sigma K_{\mathcal{I}}} f, \quad f \in L^p(\mathcal{I}, \mathfrak{H}).$$

Hence

$$\mathcal{J}^- \text{dom}(K_{\mathcal{I}}) \subseteq \text{dom}(K_{\mathcal{I}}^-)$$

and

$$K_{\mathcal{I}}^- \mathcal{J}^- f = \mathcal{J}^- K_{\mathcal{I}} f, \quad f \in \text{dom}(K_{\mathcal{I}}).$$

Notice that

$$e^{-\sigma K_{\mathcal{I}}^-} g = g - \int_0^\sigma d\tau e^{-\tau K_{\mathcal{I}}^-} K_{\mathcal{I}}^- g, \quad g \in L^p(\mathcal{I}, \mathfrak{H}^-).$$

Then choosing $g = \mathcal{J}^- f$, $f \in \text{dom}(\tilde{K}_{\mathcal{I}}^-)$, we obtain

$$\mathcal{J}^- e^{-\sigma K_{\mathcal{I}}} f = \mathcal{J}^- f - \mathcal{J}^- \int_0^\sigma d\tau e^{-\tau K_{\mathcal{I}}} \tilde{K}_{\mathcal{I}} f$$

which yields

$$e^{-\sigma K_{\mathcal{I}}} f = f - \int_0^\sigma d\tau e^{-\tau K_{\mathcal{I}}} \tilde{K}_{\mathcal{I}} f, \quad f \in \text{dom}(\tilde{K}_{\mathcal{I}}).$$

Therefore, $\tilde{K}_{\mathcal{I}} \subseteq K_{\mathcal{I}}$, which shows that $\{U_{\mathcal{I}}(t, s)\}_{(t, s) \in \Delta_{\mathcal{I}}}$ is a solution of the forward evolution equation (5.1) on \mathcal{I} . The same procedure can be applied to the backward evolution equation (5.1) on \mathcal{I} . Hence the unitary bidirectional propagator $\{G(t, s)\}_{(t, s) \in \mathbb{T} \times \mathbb{R}}$ defined by (5.14) is, in fact, a solution of the bidirectional evolution equation (5.1) on \mathbb{R} .

Assume now that $\{Z(t, s)\}_{(t, s) \in \mathbb{R} \times \mathbb{R}}$ is another unitary solution of the bidirectional evolution equation (5.1) such that Hilbert space \mathfrak{H}_0^+ is admissible with respect to $\{Z(t, s)\}_{(t, s) \in \mathbb{R} \times \mathbb{R}}$. Then from

$$J^+ \widehat{Z}(t, s) = Z(t, s) J^+, \quad (t, s) \in \mathbb{R} \times \mathbb{R},$$

we obtain

$$\widehat{Z}(t, s)^* J^- = J^- Z(t, s)^*, \quad (t, s) \in \mathbb{R} \times \mathbb{R},$$

where it is used that $J^- = (J^+)^*$. We set $Z^-(t, s) := \widehat{Z}(s, t)^*$, $(t, s) \in \mathbb{R} \times \mathbb{R}$. Since $\{Z(t, s)\}_{(t, s) \in \mathbb{R} \times \mathbb{R}}$ is unitary, we have $Z(t, s) = Z(s, t)^*$. By this we find

$$Z^-(t, s) J^- = J^- Z(t, s), \quad (t, s) \in \mathbb{R} \times \mathbb{R}.$$

Since $\{Z(t, s)\}_{(t, s) \in \mathbb{R} \times \mathbb{R}}$ and $\{Z^+(t, s)\}_{(t, s) \in \mathbb{R} \times \mathbb{R}}$ are bidirectional propagators in \mathfrak{H} and \mathfrak{H}_0^+ , respectively, one easily gets that $\{Z^-(t, s)\}_{(t, s) \in \mathbb{R} \times \mathbb{R}}$ is a bidirectional propagator in \mathfrak{H}_0^- . For any bounded interval \mathcal{I} in \mathbb{R} a forward generator $L_{\mathcal{I}}^-$ corresponds to the forward propagator $\{Z_{\mathcal{I}}^-(t, s)\}_{(t, s) \in \Delta_{\mathcal{I}}}$, $Z_{\mathcal{I}}^-(\cdot, \cdot) := Z^-(\cdot, \cdot) \upharpoonright \mathcal{I} \times \mathcal{I}$ by relation:

$$(e^{-\sigma L_{\mathcal{I}}^-} f)(t) := Z_{\mathcal{I}}^-(t, t - \sigma) \chi_{\mathcal{I}}(t - \sigma) f(t - \sigma), \quad t \in \mathcal{I}, \quad f \in L^p(\mathcal{I}, \mathfrak{H}_0^-).$$

It is obvious that

$$e^{-\sigma L_{\mathcal{I}}^-} \mathcal{J}^- = \mathcal{J}^- e^{-\sigma L_{\mathcal{I}}}, \quad \sigma \geq 0,$$

where $L_{\mathcal{I}}$ denotes the forward generator, which corresponds to $\{Z_{\mathcal{I}}(t, s)\}_{(t, s) \in \Delta_{\mathcal{I}}}$, $Z_{\mathcal{I}}(t, \cdot) := Z(\cdot, \cdot) \upharpoonright \Delta_{\mathcal{I}}$. Hence, $\mathcal{J}^- \text{dom}(L_{\mathcal{I}}) \subseteq \text{dom}(L_{\mathcal{I}}^-)$ and

$$L_{\mathcal{I}}^- \mathcal{J}^- f = \mathcal{J}^- L_{\mathcal{I}} f, \quad f \in \text{dom}(K_{\mathcal{I}}).$$

Since $L_{\mathcal{I}}$ is an extension of $\tilde{K}_{\mathcal{I}}$, we obtain

$$L_{\mathcal{I}}^- \mathcal{J}^- f = \mathcal{J}^- \tilde{K}_{\mathcal{I}} f, \quad f \in \text{dom}(\tilde{K}_{\mathcal{I}}),$$

which shows that $L_{\mathcal{I}}^-$ is an extension of $\tilde{L}_{\mathcal{I}}^- := L_{\mathcal{I}}^- \upharpoonright \mathcal{J}^- \text{dom}(\tilde{K}_{\mathcal{I}})$. Since

$$K_{\mathcal{I}}^- \mathcal{J}^- f = \mathcal{J}^- \tilde{K}_{\mathcal{I}} f, \quad f \in \text{dom}(\tilde{K}_{\mathcal{I}}).$$

holds one gets that $K_{\mathcal{I}}^-$ is also an extension of $\tilde{L}_{\mathcal{I}}^-$. Since the intersection $H_a^{1,p}(\mathcal{I}, \mathfrak{H}_0^+) \cap \text{dom}(H_{\mathcal{I}})$, cf. (5.21), is dense in $H_a^{1,p}(\mathcal{I}, \mathfrak{H}_0^+)$, the domain $\text{dom}(\tilde{L}_{\mathcal{I}}^-)$ of the closure $\bar{L}_{\mathcal{I}}^-$ of $\tilde{L}_{\mathcal{I}}^-$ contains $H_a^{1,p}(\mathcal{I}, \mathfrak{H}_0^+)$. By Remark 3.9 the set $H_a^{1,p}(\mathcal{I}, \mathfrak{H}_0^+)$ is a *core* of $K_{\mathcal{I}}^-$, which shows that $K_{\mathcal{I}}^- = \bar{L}_{\mathcal{I}}^-$. Hence $L_{\mathcal{I}}^- = K_{\mathcal{I}}^-$, which yields $Z_{\mathcal{I}}^-(t, s) = \bar{U}_{\mathcal{I}}^-(t, s)$, $(t, s) \in \Delta_{\mathcal{I}}$, for any bounded interval \mathcal{I} of \mathbb{R} . The same can be proven for the backward evolution equation, which ensures that the bidirectional evolution (5.1) admits only *one* solution for which the Hilbert space \mathfrak{H}_0^+ is admissible. \square

6 Examples

6.1 Point interactions with varying coupling constant

We consider a family $\{H(t)\}_{t \in \mathbb{R}}$ of self-adjoint operators associated in the Hilbert space $\mathfrak{H} = L^2(\mathbb{R})$ with the sesquilinear forms

$$\begin{aligned} \mathfrak{h}_t[f, g] := & \int_{\mathbb{R}} \left\{ \frac{1}{2m(x)} f'(x) \overline{g'(x)} \right\} + V(x) f(x) \overline{g(x)} + \sum_{j=1}^N \kappa_j(t) f(x_j) \overline{g(x_j)}, \end{aligned} \quad (6.1)$$

where $f, g \in \text{dom}(\mathfrak{h}_t) := H^{1,2}(\mathbb{R})$, $1 \leq N \leq \infty$. We assume that

$$m(x) > 0, \quad \frac{1}{m} + m \in L^\infty(\mathbb{R}), \quad \text{and} \quad V \in L^\infty(\mathbb{R})$$

$x_j \in \mathbb{R}$, $j = 1, 2, \dots, N$, and that the coupling constants $\kappa_j(\cdot) : \mathbb{R} \rightarrow \mathbb{R}_+$ are measurable functions. The family $\{H(t)\}_{t \in \mathbb{R}}$ is uniformly semibounded from below. Indeed, we have

$$H(t) \geq -\|V\|_{L^\infty(\mathbb{R})}, \quad t \in \mathbb{R}.$$

Therefore, without loss of generality we assume that $V(x) \geq 0$ for a.e. $x \in \mathbb{R}$, which yields that $\{H(t)\}_{t \in \mathbb{R}}$ is a family of non-negative self-adjoint operators. Moreover, one can easily verify that $\{H(t)\}_{t \in \mathbb{R}}$ is a measurable family of self-adjoint operators. For finite N the domain $\text{dom}(H(t))$ admits an explicit description. Indeed, in this case the operators $H(t)$ are given by the sum of operators in the form-sense (6.1):

$$H(t) = -\frac{1}{2} \frac{d}{dx} \frac{1}{m(x)} \frac{d}{dx} \dot{+} V(x) \dot{+} \sum_{j=1}^N \kappa_j(t) \delta(x - x_j)$$

with domain defined by

$$\begin{aligned} \text{dom}(H(t)) := & \left\{ f \in H^{1,2}(\mathbb{R}) : \begin{array}{l} \frac{1}{m} f' \in H^{1,2}(\mathbb{R} \setminus \bigcup_{j=1}^N \{x_j\}), \\ \left(\frac{1}{2m} f' \right) (x_j - 0) - \left(\frac{1}{2m} f' \right) (x_j + 0) = \kappa_j(t) f(x_j), \\ j = 1, 2, \dots, N < \infty \end{array} \right\} \end{aligned} \quad (6.2)$$

for $t \in \mathcal{I}$. In the following we assume (*convergence condition*) that

$$\sup_{t \in \mathcal{I}} \sum_{j=1}^N \kappa_j(t) < \infty, \quad 1 \leq N \leq \infty, \quad (6.3)$$

for each bounded subinterval $\mathcal{I} \subset \mathbb{R}$. Furthermore, we assume (*continuity condition*) that for each bounded subinterval $\mathcal{I} \subset \mathbb{R}$ there is a constant $C_{\mathcal{I}} > 0$ such that

$$\sum_{j=1}^N |\kappa_j(t) - \kappa_j(s)| \leq C_{\mathcal{I}} |t - s|, \quad t, s \in \mathcal{I}. \quad (6.4)$$

Since $\mathfrak{D}^+ := \text{dom}(H(t)^{1/2}) = \text{dom}(\mathfrak{h}_t) = H^{1,2}(\mathbb{R})$ is independent of $t \in \mathbb{R}$, Theorem 5.2 is applicable in this case: the *auxiliary* bidirectional evolution equation (5.8) admits a unique solution, if the estimate (5.10) is satisfied for each bounded subinterval $\mathcal{I} \subset \mathbb{R}$.

To show this it is sufficient to verify that the estimate

$$\|\sqrt{H(t) + If}\| \leq e^{\gamma_{\mathcal{I}}|t-s|} \|\sqrt{H(s) + If}\|, \quad f, g \in \mathfrak{D}^+ \quad (6.5)$$

holds for any $t, s \in \mathcal{I}$. Indeed, one obviously has

$$|f(x_j)|^2 = 2\Re \left\{ \int_{-\infty}^{x_j} f'(x) \overline{f(x)} dx \right\}, \quad f \in H^{1,2}(\mathbb{R}), \quad j \in 1, 2, \dots, N,$$

which yields

$$|f(x_j)|^2 \leq \int_{\mathbb{R}} \{|f'(x)|^2 + |f(x)|^2\} dx, \quad j = 1, 2, \dots, N. \quad (6.6)$$

Hence

$$|f(x_j)|^2 \leq \max\{1, 2\|m\|_{L^\infty}\} \|\sqrt{H(s) + If}\|^2, \quad j = 1, 2, \dots, N. \quad (6.7)$$

Therefore, we have

$$\left| \|\sqrt{H(t) + If}\|^2 - \|\sqrt{H(s) + If}\|^2 \right| \leq \sum_{j=1}^N |\kappa_j(t) - \kappa_j(s)| |f(x_j)|^2.$$

and consequently, by (6.6) we obtain:

$$\begin{aligned} & \left| \|\sqrt{H(t) + If}\|^2 - \|\sqrt{H(s) + If}\|^2 \right| \leq \\ & \max\{1, 2\|m\|_{L^\infty}\} \|\sqrt{H(s) + If}\|^2 \sum_{j=1}^N |\kappa_j(t) - \kappa_j(s)|. \end{aligned}$$

Using (6.4) we get

$$\left| \|\sqrt{H(t) + If}\|^2 - \|\sqrt{H(s) + If}\|^2 \right| \leq 2\gamma_{\mathcal{I}} |t - s| \|\sqrt{H(s) + If}\|^2 \quad (6.8)$$

for $t, s \in \mathcal{I}$, where

$$\gamma_{\mathcal{I}} := \frac{1}{2} C_{\mathcal{I}} \max\{1, 2\|m\|_{L^\infty}\}.$$

From (6.8) it follows that

$$\|\sqrt{H(t) + If}\| \leq \sqrt{1 + 2\gamma_{\mathcal{I}} |t - s|} \|\sqrt{H(s) + If}\|,$$

which yields

$$\|\sqrt{H(t) + If}\| \leq (1 + \gamma_{\mathcal{I}} |t - s|) \|\sqrt{H(s) + If}\|,$$

for $t, s \in \mathcal{I}$. Since $1 + \gamma_{\mathcal{I}} |t - s| \leq e^{\gamma_{\mathcal{I}} |t - s|}$, for any $t, s \in \mathcal{I}$, we obtain (6.5).

Then by Theorem 5.5 the *original* bidirectional evolution equation (5.1) admits a solution for which the Hilbert space $H^{1,2}(\mathbb{R})$ is admissible. It is more complicated to solve the problem whether this solution of the *original* problem is *unique*. To this end one has to verify the additional condition (5.21) of Theorem 5.5. This condition is satisfied if the sets $(I + H_{\mathcal{I}})^{-1}H_a^{1,2}(\mathcal{I}, \mathfrak{H})$ and $(I + H_{\mathcal{I}})^{-1}H_b^{1,2}(\mathcal{I}, \mathfrak{H})$ are dense in $H_a^{1,2}(\mathcal{I}, \mathfrak{H}_0^+)$ and $H_b^{1,2}(\mathcal{I}, \mathfrak{H}_0^+)$, for any bounded interval $\mathcal{I} = (a, b)$, respectively.

To prove this we introduce linear operators $C_j : L^2(\mathbb{R}) \rightarrow \mathbb{C}$ defined by

$$C_j f := ((I + H(0))^{-1/2} f)(x_j), \quad f \in L^2(\mathbb{R}), \quad j = 1, 2, \dots, N.$$

Using the estimate (6.7) we find $|C_j f| \leq C \|f\|_{L^2(\mathbb{R})}$, where C is given by $C := \max\{1, 2\|m\|_{L^\infty}\}$. Setting $B_j := C_j^* C_j$ we obtain the representation

$$(I + H(t))^{-1} = (I + H(0))^{-1/2} R(t) (I + H(0))^{-1/2}, \quad t \in \mathbb{R},$$

where

$$R(t) := \left(I + \sum_{j=1}^N \kappa_j(t) B_j \right)^{-1}, \quad t \in \mathbb{R}.$$

Since the coupling constants are locally Lipschitz continuous, see (6.4), we get that $R(t)x \in H_a^{1,2}(\mathcal{I}, \mathfrak{H})$, $x \in \mathfrak{H}$, for any bounded open interval $\mathcal{I} \subseteq \mathbb{R}$. Hence, $R(t)f(t) \in H_a^{1,2}(\mathcal{I}, \mathfrak{H})$ for $f \in H_a^{1,2}(\mathcal{I}, \mathfrak{H})$ and any bounded open interval $\mathcal{I} \subseteq \mathbb{R}$. Hence we get $(I + H(t))^{-1}f(t) \in H_a^{1,2}(\mathcal{I}, \mathfrak{H}_0^+)$ for $f \in H_a^{1,2}(\mathcal{I}, \mathfrak{H})$ and $\mathcal{I} \subseteq \mathbb{R}$. Now we show that the set of elements $(I + H(t))^{-1}f(t)$, $f \in H_a^{1,2}(\mathcal{I}, \mathfrak{H})$, is *dense* in $H_a^{1,2}(\mathcal{I}, \mathfrak{H}_0^+)$. Note that the standard norm of $H_a^{1,2}(\mathcal{I}, \mathfrak{H}_0^+)$ is equivalent to the norm

$$\|f\|_{H_a^{1,2}(\mathcal{I}, \mathfrak{H}_0^+)} = \left(\int_{\mathcal{I}} \|\sqrt{I + H(0)} f'(t)\|_{\mathfrak{H}}^2 dt \right)^{1/2}.$$

If the elements $(I + H(t))^{-1}f(t)$, $f \in H_a^{1,2}(\mathcal{I}, \mathfrak{H})$, are not dense in $H_a^{1,2}(\mathcal{I}, \mathfrak{H}_0^+)$, then there is an element $g \in H_a^{1,2}(\mathcal{I}, \mathfrak{H}_0^+)$ such that

$$\int_{\mathcal{I}} \left(\sqrt{I + H(0)} \left((I + H(0))^{-1/2} R(t) (I + H(0))^{-1/2} f(t) \right)', \sqrt{I + H(0)} g'(t) \right) dt = 0$$

for any $f \in H_a^{1,2}(\mathcal{I}, \mathfrak{H})$. Hence we obtain

$$\int_{\mathcal{I}} (R'(t)(I + H(0))^{-1/2} f(t) + R(t)(I + H(0))^{-1/2} f'(t), \sqrt{I + H(0)} g'(t)) dt = 0.$$

Setting $h(t) := (I + H(0))^{-1/2} f(t) \in H_a^{1,2}(\mathcal{I}, \mathfrak{H}_0^+)$ and $k(t) := \sqrt{I + H(0)} g'(t) \in L^2(\mathcal{I}, \mathfrak{H})$ we find that

$$\int_{\mathcal{I}} (R'(t)h(t) + R(t)h'(t), k(t)) dt = 0 \tag{6.9}$$

for any $h \in H_a^{1,2}(\mathcal{I}, \mathfrak{H}_0^+)$. Since $H_a^{1,2}(\mathcal{I}, \mathfrak{H}_0^+)$ is dense in $H_a^{1,2}(\mathcal{I}, \mathfrak{H})$ one gets that (6.9) holds for any $h \in H_a^{1,2}(\mathcal{I}, \mathfrak{H})$. From (6.9) we obtain

$$\int_{\mathcal{I}} (h'(t), R(t)k(t)) dt = - \int_{\mathcal{I}} (h(t), R'(t)k(t)) dt$$

for any $h \in H_a^{1,2}(\mathcal{I}, \mathfrak{H})$, which yields $z(t) := R(t)k(t) \in H_b^{1,2}(\mathcal{I}, \mathfrak{H})$ and

$$\frac{d}{dt} R(t)k(t) - R'(t)k(t) = 0 \tag{6.10}$$

for a.e. $t \in \mathcal{I}$. From the representation

$$k(t) = \left(I + \sum_{j=1}^N \kappa_j(t) B_j \right) z(t)$$

and condition (6.4) we obtain that $k(t) \in H_b^{1,2}(\mathcal{I}, \mathfrak{H})$. Taking into account this last observation we get from (6.10) that $R(t)k'(t) = 0$ for a.e. $t \in \mathcal{I}$. Since $\ker(R(t)) = \{0\}$ for $t \in \mathcal{I}$, we find that $k'(t) = 0$, which implies $k(t) = \text{const}$. But since $k(b) = 0$, we get $k(t) = 0$ for $t \in \mathcal{I}$. Hence $g'(t) = 0$ for $t \in \mathcal{I}$, which yields $g(t) = 0$ for $t \in \mathcal{I}$. Consequently, the set $(I + H_{\mathcal{I}})^{-1}H_a^{1,2}(\mathcal{I}, \mathfrak{H})$ is dense in $H_a^{1,2}(\mathcal{I}, \mathfrak{H}_0^+)$ for any bounded open interval $\mathcal{I} = (a, b)$.

Similarly, one proves that the set $(I + H_{\mathcal{I}})^{-1}H_b^{1,2}(\mathcal{I}, \mathfrak{H})$ is dense in $H_b^{1,2}(\mathcal{I}, \mathfrak{H}_0^+)$ for any bounded open interval $\mathcal{I} = (a, b)$.

Taking into account the second part of Theorem 5.5 one finds that there is a *unique* solution of the original problem (5.1) such that \mathfrak{H}_0^+ is admissible.

Therefore, summing up this line of reasoning we obtain the proof of the following theorem:

Theorem 6.1 *Let $0 \leq V \in L^\infty(\mathbb{R})$, $m > 0$ and $1/m + m \in L^\infty(\mathbb{R})$. Further, let $\{x_j\}_{j \in \mathbb{N}}$ be a (infinite) sequence of real numbers which are mutually different and let $\kappa_j(\cdot) : \mathbb{R} \rightarrow \mathbb{R}_+$ be non-negative locally Lipschitz continuous functions. Moreover, let $\{H(t)\}_{t \in \mathbb{R}}$ be a family of non-negative self-adjoint Schrödinger operators associated with the sesquilinear forms (6.1). If the conditions (6.3) and (6.4) are satisfied, then the bidirectional evolution equation (5.1) is well-posed on \mathbb{R} for $p = 2$ and possesses a unique solution $\{G(t, s)\}_{(t,s) \in \mathbb{R} \times \mathbb{R}}$ such that $H^{1,2}(\mathbb{R})$ is admissible.*

A similar problem was treated in three dimensions by [41] for the case of finite point interactions and $m(x) = \text{const}$. In contrast to Theorem 6.1 their results concern the case of coupling constants $\kappa_j(t)$ which are *twice continuously differentiable*, cf. [41, Theorem 1]. In this case the bidirectional evolution equation is verified in the strong sense. Moreover, only the existence of a bidirectional propagator was established under the weaker assumption that the coupling constants $\kappa_j(t)$ are locally L^∞ -function, cf. [41, Theorem 2]. The first results was improved in [5], where the smoothness of the coupling constants was reduced to a certain Hölder continuity. However, it seems to be difficult to extend the technique used [5, 41] to the case of an *infinite* number of point interactions and to a *non-smooth* position dependent effective mass m .

In conclusion we would like to remark that Theorem 6.1 covers rather *bizarre* situations. For instance, let $\{x_j\}_{j \in \mathbb{N}}$ be an enumeration of the rational numbers \mathbb{Q} and let $\{\kappa_j(t)\}$ be a sequence of coupling constants such that conditions (6.3) and (6.4) are satisfied. Moreover, let us assume that for any $t \in \mathbb{R}$ the values $\kappa_j(t)$ are pairwise different. In this case one has $\bigcap_{t \in \mathcal{I}} \text{dom}(H(t)) = \{0\}$ for any bounded open interval $\mathcal{I} \subseteq \mathbb{R}$. Nevertheless, the sets $(I + H_{\mathcal{I}})^{-1}H_a^{1,2}(\mathcal{I}, \mathfrak{H})$ and $(I + H_{\mathcal{I}})^{-1}H_b^{1,2}(\mathcal{I}, \mathfrak{H})$ are dense in $H_a^{1,2}(\mathcal{I}, \mathfrak{H}_0^+)$ and $H_b^{1,2}(\mathcal{I}, \mathfrak{H}_0^+)$, respectively !

6.2 Moving potentials

In this section we consider an example, which is more involved than that we studied above. Here we consider the Hamiltonian of two *moving point particles*:

$$H(t) = -\frac{1}{2} \frac{d^2}{dx^2} + \kappa_1(t) \delta(x - x_1(t)) + \kappa_2(t) \delta(x - x_2(t)), \quad (6.11)$$

which domain is described by

$$\text{dom}(H(t)) := \left\{ f \in H^{1,2}(\mathbb{R}) : \begin{array}{l} f' \in H^{1,2}(\mathbb{R} \setminus \{x_1(t), x_2(t)\}), \\ (f'/2)(x_1(t) - 0) - (f'/2)(x_1(t) + 0) = \kappa_1(t)f(x_1(t)), \\ (f'/2)(x_2(t) - 0) - (f'/2)(x_2(t) + 0) = \kappa_2(t)f(x_2(t)), \end{array} \right\} \quad (6.12)$$

in the Hilbert space $L^2(\mathbb{R})$. In the following we assume that $\kappa_j(\cdot) : \mathbb{R} \rightarrow \mathbb{R}_+$ are *continuous differentiable* functions. Moreover, we suppose that

$$x_1(t) < x_2(t) \tag{6.13}$$

for $t \in \mathbb{R}$. The sesquilinear form associated with $H(t)$ is given by

$$\begin{aligned} \mathfrak{h}_t[f, g] = & \\ & \frac{1}{2} \int_{\mathbb{R}} f'(x) \overline{g'(x)} dx + \kappa_1(t) f(x_1(t)) \overline{g(x_1(t))} + \kappa_2(t) f(x_2(t)) \overline{g(x_2(t))}, \end{aligned}$$

$f, g \in \text{dom}(\mathfrak{h}_t) := H^{1,2}(\mathbb{R})$. Notice that the sesquilinear form \mathfrak{h}_t is non-negative.

To handle this case we start with some formal manipulations. Using the momentum operator P ,

$$Pf = \frac{1}{i} \frac{\partial}{\partial x} f(x), \quad f \in \text{dom}(P) := H^{1,2}(\mathbb{R}),$$

we get the representation

$$\mathfrak{h}_t[f, g] = \frac{1}{2} (Pf, Pg) + \kappa_1(t) f(x_1(t)) \overline{g(x_1(t))} + \kappa_2(t) f(x_2(t)) \overline{g(x_2(t))},$$

$f, g \in H^{1,2}(\mathbb{R})$. The momentum operator generates the right-shift group $S(\tau) := e^{-i\tau P}$, $\tau \in \mathbb{R}$, acting as

$$(S(\tau)f)(x) = f(x - \tau), \quad f \in L^2(\mathbb{R}), \quad \tau \in \mathbb{R}.$$

Obviously, one has that

$$S(\tau)^{-1} H(t) S(\tau) = \frac{1}{2} P^2 + \kappa_1(t) \delta(x - x_1(t) + \tau) + \kappa_2(t) \delta(x - x_2(t) + \tau).$$

In particular, for $y(t) := \frac{1}{2}(x_1(t) + x_2(t))$ we obtain:

$$\begin{aligned} H^S(t) &:= S(y(t))^{-1} H(t) S(y(t)) = \\ & e^{iy(t)P} H(t) e^{-iy(t)P} = \frac{1}{2} P^2 + \kappa_1(t) \delta(x + x(t)) + \kappa_2(t) \delta(x - x(t)), \end{aligned}$$

where the relative coordinate obeys

$$x(t) := \frac{x_2(t) - x_1(t)}{2} > 0, \quad t \in \mathbb{R}.$$

by (6.13). Further, we define the unitary transformations $W(\theta) : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$, $\theta > 0$,

$$W(\theta)f(x) := \sqrt{\theta} f(\theta x), \quad f \in L^2(\mathbb{R}).$$

Let X be multiplication operator $(Xf) := xf(x)$ in $L^2(\mathbb{R})$. Then

$$L = \frac{1}{2} (XP + PX)$$

is a so-called *dilation operator*, which is self-adjoint in $L^2(\mathbb{R})$. The operator iL generates *dilation group* given by

$$(e^{isL}f)(x) = e^{s/2} f(e^s x), \quad f \in L^2(\mathbb{R}), \quad s \in \mathbb{R}.$$

Then we obviously get $W(\theta) = e^{i \ln(\theta)L}$, $\theta > 0$ and

$$\begin{aligned} W(\theta)^{-1} H^S(t) W(\theta) = & \\ & -\frac{\theta^2}{2} \frac{d^2}{dx^2} + \kappa_1(t) \theta \delta(x + \theta x(t)) + \kappa_2(t) \theta \delta(x - \theta x(t)). \end{aligned}$$

If we set $\theta = 1/x(t)$, then

$$\begin{aligned} H^{SW}(t) &:= W(1/x(t))^{-1} H^S(t) W(1/x(t)) = \\ &e^{i \ln(x(t))L} H^S(t) e^{-i \ln(x(t))L} = \frac{1}{2x(t)^2} P^2 + \varkappa_1(t) \delta(x+1) + \varkappa_2(t) \delta(x-1), \end{aligned}$$

where

$$\varkappa_1(t) := \frac{\kappa_1(t)}{x(t)} \quad \text{and} \quad \varkappa_2(t) := \frac{\kappa_2(t)}{x(t)}.$$

Relation between this Hamiltonian and (6.11) has the form:

$$H(t) = e^{-iy(t)P} e^{-i \ln(x(t))L} H^{SW}(t) e^{i \ln(x(t))L} e^{iy(t)P}.$$

Now we introduce in the Hilbert space $L^2(\mathbb{R}, \mathfrak{H})$, $\mathfrak{H} := L^2(\mathbb{R})$, the operator

$$(Df)(t, x) = \left(\frac{1}{i} \frac{\partial}{\partial t} f \right)(t, x), \quad \text{dom}(D) := H^{1,2}(\mathbb{R}, \mathfrak{H}).$$

The multiplication operator $S := M(S(y(t)))$, $y(t) = \frac{1}{2}(x_1(t) + x_2(t))$ (i.e., $(Sf)(t, x) := (S(y(t))f)(t, x) = f(t, x - y(t))$, see (2.1)), defines a unitary operator on $L^2(\mathbb{R}, \mathfrak{H})$, and we have that

$$D^S := S^{-1} D S = D - \dot{y}(t)P.$$

Similarly, the multiplication operator $W := M(W(1/x(t)))$, $x(t) = \frac{1}{2}(x_2(t) - x_1(t))$, induces a unitary operator on $L^2(\mathbb{R}, \mathfrak{H})$. We set

$$D^{SW} := W^{-1} D^S W.$$

Since the multiplication operator $W = M(e^{-i \ln(x(t))L})$, by the commutation relation $LP - PL = iP$ one gets that

$$D^{SW} = D - i \frac{\dot{x}(t)}{x(t)} L - i \frac{\dot{y}(t)}{x(t)} P.$$

Now we set

$$H^{SW} := W^{-1} S^{-1} H S W$$

and

$$\tilde{K}^{SW} := D^{SW} + H^{SW}$$

with domain $\text{dom}(\tilde{K}^{SW}) := \text{dom}(D^{SW}) \cap \text{dom}(H^{SW})$. Then a straightforward computation gives that this operator is equal to

$$\tilde{K}^{SW} := D + L_0$$

with domain $\text{dom}(\tilde{K}^{SW}) = \text{dom}(D) \cap \text{dom}(L_0)$, where

$$\begin{aligned} L_0(t) &:= \frac{1}{2x(t)^2} (P - x(t)(\dot{x}(t)X + \dot{y}(t)))^2 - \frac{1}{2}(\dot{x}(t)X + \dot{y}(t))^2 \\ &+ \varkappa_1(t) \delta(x+1) + \varkappa_2(t) \delta(x-1). \end{aligned}$$

Finally, let us introduce the gauge transformation

$$(\Gamma(t)f)(x) := e^{i \int_0^t ((\dot{x}(s)x + \dot{y}(s))^2 + x^2) ds/2} f(x), \quad f \in L^2(\mathbb{R}),$$

which induces the multiplication operator $\Gamma := M(\Gamma(t))$ on $L^2(\mathcal{I}, \mathfrak{H})$. Then we find

$$\tilde{\mathbf{K}} := \tilde{K}^{SW\Gamma} := \Gamma^{-1} \tilde{K}^{SW} \Gamma = D + L,$$

where operator

$$L(t) := \frac{1}{2x(t)^2} (P + \beta_1(t)X + \beta_0(t))^2 + \frac{1}{2}X^2 + \varkappa_1(t)\delta(x+1) + \varkappa_2(t)\delta(x-1)$$

with

$$\beta_1(t) := \int_0^t (\dot{x}(s)^2 + 1)ds - x(t)\dot{x}(t)$$

and

$$\beta_0(t) := \int_0^t \dot{y}(s)\dot{x}(s)ds - x(t)\dot{y}(t).$$

As above the family $\{L(t)\}_{t \in \mathbb{R}}$, is measurable and defines a densely defined self-adjoint multiplication operator $L := M(L(t))$ on $L^2(\mathcal{I}, \mathfrak{H})$. Then the operators $\tilde{K} := D + H$ and $\tilde{\mathbf{K}} = D + L$ are related by

$$\tilde{K} = S W \Gamma \tilde{\mathbf{K}} \Gamma^{-1} W^{-1} S^{-1}. \quad (6.14)$$

Instead to solve the bidirectional evolution equation (5.1) we consider the modified bidirectional evolution equation

$$\frac{1}{i} \frac{\partial}{\partial t} u(t) + L(t)u(t) = 0. \quad (6.15)$$

Following Section 5 we introduce the family of quadratic forms $\mathfrak{I}_t[\cdot, \cdot]$

$$\begin{aligned} \mathfrak{I}_t[f, g] &:= \frac{1}{2x(t)^2} (Pf + \beta_1(t)Xf + \beta_0(t)f, Pf + \beta_1(t)Xg + \beta_0(t)g) + \\ &\quad \frac{1}{2} (Xf, Xg) + \varkappa_1(t)f(-1)\overline{g(-1)} + \varkappa_2(t)f(1)\overline{g(1)} + (f, g), \end{aligned}$$

$f, g \in \text{dom}(\mathfrak{I}_t) := \text{dom}(P) \cap \text{dom}(X)$ corresponding to operators $L(t)$, and define the norm

$$\|f\|_t^+ := \|\sqrt{L(t) + I}f\| = \sqrt{\mathfrak{I}_t[f, f] + \|f\|^2},$$

$f \in \text{dom}(\sqrt{L(t) + I}) = \text{dom}(\mathfrak{I}_t) = \text{dom}(P) \cap \text{dom}(X)$. It is easy to check that the domain $\text{dom}(\mathfrak{I}_t)$ is independent of $t \in \mathbb{R}$. By \mathfrak{L}_t^+ we denote the Hilbert space, which arises when we endow the domain $\text{dom}(\mathfrak{I}_t)$ with the scalar product $(f, g)_t^+ := \mathfrak{I}_t[f, g] + (f, g)$. Note that the norm $\|\cdot\|_t^+$ is equivalent to the norm $\|f\|_{PX} = \sqrt{\|Pf\|^2 + \|Xf\|^2}$, $f \in \text{dom}(P) \cap \text{dom}(X)$.

Now we proceed as in the previous section. First we find

$$\begin{aligned} \frac{d}{dt} (\|f\|_t^+)^2 &= \frac{\dot{x}(t)}{x(t)^3} \|Pf + \beta_1(t)Xf + \beta_0(t)f\|^2 \\ &\quad + \frac{2}{x(t)^2} \Re(Pf + \beta_1(t)Xf + \beta_0(t)f, \dot{\beta}_1(t)Xf + \dot{\beta}_0(t)f) \\ &\quad + \dot{\varkappa}_1(t)|f(-1)|^2 + \dot{\varkappa}_2(t)|f(1)|^2. \end{aligned}$$

A straightforward computation shows that for any bounded interval \mathcal{I} there is a constant $\gamma_{\mathcal{I}}$ such that

$$\left| \frac{d}{dt} (\|f\|_t^+)^2 \right| \leq 2\gamma_{\mathcal{I}} (\|f\|_t^+)^2$$

for $t \in \mathcal{I}$ which yields

$$-\gamma_{\mathcal{I}} \leq \frac{d}{dt} \ln(\|f\|_t^+) \leq \gamma_{\mathcal{I}}.$$

Hence we obtain the estimate:

$$-\gamma_{\mathcal{I}}(t-s) \leq \ln(\|f\|_t^+) - \ln(\|f\|_s^+) \leq \gamma_{\mathcal{I}}(t-s)$$

for $t, s \in \mathcal{I}$ and $s \leq t$, which yields

$$\|f\|_t^+ \leq e^{\gamma_{\mathcal{I}}(t-s)} \|f\|_s^+, \quad t, s \in \mathcal{I}, \quad s \leq t.$$

The last relation implies (5.10). By virtue of Theorem 5.2 we get that the *auxiliary* bidirectional evolution equation

$$\frac{\partial}{\partial t} u(t) + iL^-(t)u(t) = 0$$

admits a unique solution $\{\Lambda^-(t, s)\}_{(t,s) \in \mathbb{R} \times \mathbb{R}}$ on \mathbb{R} . By Theorem 5.5 the *original* bidirectional evolution equation (6.15) admits a solution for which the Hilbert space \mathfrak{L}_0^+ is admissible. By the same line of reasoning as for non-moving point interactions one can prove that there is unique unitary solution $\{\Lambda(t, s)\}_{(t,s) \in \mathbb{R} \times \mathbb{R}}$ of the bidirectional evolution equation (6.15) for which \mathfrak{L}_0^+ is admissible.

These results allow to prove that the original forward evolution equation (5.8) on \mathbb{R} admits a solution. To this end one has to verify that for any bounded interval \mathcal{I} the extension of the forward generator $\mathbf{K}_{\mathcal{I}}$ of $\tilde{\mathbf{K}}_{\mathcal{I}}$ defines an extension of the forward generator $K_{\mathcal{I}}$ of $\tilde{K}_{\mathcal{I}}$ defined by $K_{\mathcal{I}} := S W \Gamma \mathbf{K}_{\mathcal{I}} \Gamma^{-1} W^{-1} S^{-1}$. However, this is evident since it follows from the representation (6.14). Similarly, one proves that for any bounded interval \mathcal{I} the backward generator extension $\mathbf{K}^{\mathcal{I}}$ of $\tilde{\mathbf{K}}^{\mathcal{I}}$ defines a backward generator extension $K^{\mathcal{I}}$ of $\tilde{K}^{\mathcal{I}}$ by $K^{\mathcal{I}} := S W \Gamma \mathbf{K}^{\mathcal{I}} \Gamma^{-1} W^{-1} S^{-1}$. By these we immediately obtain that the bidirectional propagator $\{G(t, s)\}_{(t,s) \in \mathbb{R} \times \mathbb{R}}$ defined by

$$G(t, s) := e^{-iy(t)P} e^{-i \ln(x(t))L} \Gamma(t) \Lambda(t, s) \Gamma(s)^{-1} e^{i \ln(x(s))L} e^{iy(s)P}, \quad (6.16)$$

for any $(t, s) \in \mathbb{R} \times \mathbb{R}$, is a solution of the bidirectional evolution equation (5.1).

It remains only to identify the subspace which is admissible with respect to $\{G(t, s)\}_{(t,s) \in \mathbb{R} \times \mathbb{R}}$. We recall that \mathfrak{L}_0^+ is the subspace which is admissible with respect to $\{\Lambda(t, s)\}_{(t,s) \in \mathbb{R} \times \mathbb{R}}$. If we set

$$H^{\Gamma}(t) := \Gamma(t)L(t)\Gamma(t)^{-1}, \quad t \in \mathbb{R},$$

then a straightforward computation shows that

$$\begin{aligned} H^{\Gamma}(t) &:= \frac{1}{2x(t)^2} (P - x(t)\dot{x}(t)X - x(t)\dot{y}(t))^2 + \\ &\quad \frac{1}{2} X^2 + \varkappa_1(t)\delta(x+1) + \varkappa_2(t)\delta(x-1). \end{aligned}$$

Further, setting

$$H^{\Gamma W}(t) := e^{-i \ln(x(t))L} \Gamma(t)L(t)\Gamma(t)^{-1} e^{i \ln(x(t))L}$$

we find that

$$H^{\Gamma W}(t) = \frac{1}{2} \left(P - \frac{\dot{x}(t)}{x(t)} X - \dot{y}(t) \right)^2 + \frac{1}{2x(t)^2} X^2 + \kappa_1(t)\delta(x+x(t)) + \kappa_2(t)\delta(x-x(t)).$$

Finally, we introduce the family:

$$H^{\Gamma W S}(t) := e^{-iy(t)P} e^{-i \ln(x(t))L} \Gamma(t)L(t)\Gamma(t)^{-1} e^{i \ln(x(t))L} e^{iy(t)P}$$

which implies

$$\begin{aligned} H^{\Gamma W S}(t) &= \frac{1}{2} \left(P - \frac{\dot{x}(t)}{x(t)} (X - y(t)) \right)^2 + \\ &\quad \frac{1}{2x(t)^2} (X - y(t))^2 + \kappa_1(t)\delta(x-x_1(t)) + \kappa_2(t)\delta(x-x_2(t)). \end{aligned}$$

For a shorthand let $Z(t) := H^{\Gamma W S}(t)$. Then quadratic form associated with $Z(t)$ we denote by $\mathfrak{z}_t[\cdot, \cdot]$. One can easily verify that the domain $\text{dom}(\mathfrak{z}_t)$ is independent of $t \in \mathbb{R}$. The Hilbert space

which is associated with \mathfrak{z}_t is denoted by \mathfrak{Z}_t^+ . A straightforward computation shows that for any $t \in \mathbb{R}$ the Hilbert space \mathfrak{Z}_t^+ can be identified with $\mathfrak{H}_{PX} := \{\text{dom}(P) \cap \text{dom}(X), \|\cdot\|_{PX}\}$. It is obvious, that the Hilbert space \mathfrak{H}_{PX} is admissible for the bidirectional propagator $\{G(t, s)\}_{(t,s) \in \mathbb{R}}$ defined by (6.16). Summing up one gets the following theorem:

Theorem 6.2 *Let $\kappa_j(\cdot) : \mathbb{R} \rightarrow \mathbb{R}_+$ and $x_j(\cdot) : \mathbb{R} \rightarrow \mathbb{R}$ be continuously differentiable functions. Further, let $\{H(t)\}_{t \in \mathbb{R}}$ be the family of non-negative self-adjoint operators given by (6.11) and (6.12). If the condition (6.13) is satisfied for any $t \in \mathbb{R}$, then the bidirectional evolution equation (5.1) is well-posed on \mathbb{R} for $p = 2$ and possesses a unique unitary solution for which the Hilbert space \mathfrak{H}_{PX} is admissible.*

Acknowledgments

This papers results from numerous discussions and mutual visits of the co-authors to Berlin and Marseille. H.N. thanks the CPT (Marseille) for hospitality and financial support. V.A.Z. is thankful to WIAS (Berlin) for invitations and financial support of his several visits.

References

- [1] *Functional analysis and semi-groups*. American Mathematical Society Colloquium Publications, vol. 31. American Mathematical Society, Providence, R. I., 1957. rev. ed.
- [2] Pilar Cembranos and José Mendoza. *Banach spaces of vector-valued functions*, volume 1676 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1997.
- [3] Adrian Constantin. The construction of an evolution system in the hyperbolic case and applications. *Math. Nachr.*, 224:49–73, 2001.
- [4] G. Da Prato and P. Grisvard. Sommes d’opérateurs linéaires et équations différentielles opérationnelles. *J. Math. Pures Appl. (9)*, 54(3):305–387, 1975.
- [5] G. F. Dell’Antonio, R. Figari, and A. Teta. A limit evolution problem for time-dependent point interactions. *J. Funct. Anal.*, 142(1):249–275, 1996.
- [6] G. F. Dell’Antonio, R. Figari, and A. Teta. The Schrödinger equation with moving point interactions in three dimensions. In *Stochastic processes, physics and geometry: new interplays, I (Leipzig, 1999)*, volume 28 of *CMS Conf. Proc.*, pages 99–113. Amer. Math. Soc., Providence, RI, 2000.
- [7] Gianfausto Dell’Antonio. Point interactions. In *Mathematical physics in mathematics and physics (Siena, 2000)*, volume 30 of *Fields Inst. Commun.*, pages 139–150. Amer. Math. Soc., Providence, RI, 2001.
- [8] J. Diestel and J. J. Uhl, Jr. *Vector measures*. American Mathematical Society, Providence, R.I., 1977. Mathematical Surveys, No. 15.
- [9] John D. Dollard and Charles N. Friedman. Asymptotic behavior of solutions of linear ordinary differential equations. *J. Math. Anal. Appl.*, 66(2):394–398, 1978.
- [10] J. R. Dorroh. A simplified proof of a theorem of Kato on linear evolution equations. *J. Math. Soc. Japan*, 27(3):474–478, 1975.
- [11] Joanne Elliott. The equation of evolution in a Banach space. *Trans. Amer. Math. Soc.*, 103:470–483, 1962.

- [12] David E. Evans. Time dependent perturbations and scattering of strongly continuous groups on Banach spaces. *Math. Ann.*, 221(3):275–290, 1976.
- [13] Jerome A. Goldstein. Time dependent hyperbolic equations. *J. Functional Analysis*, 4:31–49, 1969.
- [14] Jerome A. Goldstein. On the absence of necessary conditions for linear evolution operators. *Proc. Amer. Math. Soc.*, 64(1):77–80, 1977.
- [15] Matthew Hackman. The abstract time-dependent Cauchy problem. *Trans. Amer. Math. Soc.*, 133:1–50, 1968.
- [16] Eugen Heyn. Die Differentialgleichung $dT/dt = P(t)T$ für Operatorfunktionen. *Math. Nachr.*, 24:281–330, 1962.
- [17] James S. Howland. Stationary scattering theory for time-dependent Hamiltonians. *Math. Ann.*, 207:315–335, 1974.
- [18] Susumu Ishii. An approach to linear hyperbolic evolution equations by the Yosida approximation method. *Proc. Japan Acad. Ser. A Math. Sci.*, 54(1):17–20, 1978.
- [19] Tosio Kato. Integration of the equation of evolution in a Banach space. *J. Math. Soc. Japan*, 5:208–234, 1953.
- [20] Tosio Kato. On linear differential equations in Banach spaces. *Comm. Pure Appl. Math.*, 9:479–486, 1956.
- [21] Tosio Kato. *Perturbation theory for linear operators*. Die Grundlehren der mathematischen Wissenschaften, Band 132. Springer-Verlag New York, Inc., New York, 1966.
- [22] Tosio Kato. Linear evolution equations of “hyperbolic” type. *J. Fac. Sci. Univ. Tokyo Sect. I*, 17:241–258, 1970.
- [23] Tosio Kato. Linear evolution equations of “hyperbolic” type. II. *J. Math. Soc. Japan*, 25:648–666, 1973.
- [24] J. Kiszyński. Sur les opérateurs de Green des problèmes de Cauchy abstraits. *Studia Math.*, 23:285–328, 1963/1964.
- [25] Kazuo Kobayasi. On a theorem for linear evolution equations of hyperbolic type. *J. Math. Soc. Japan*, 31(4):647–654, 1979.
- [26] Yukio Kōmura. On linear evolution operators in reflexive Banach spaces. *J. Fac. Sci. Univ. Tokyo Sect. I A Math.*, 17:529–542, 1970.
- [27] S. G. Kreĭn. *Lineinye uravneniya v banakhovom prostranstve*. Voronež. Gosudarstv. Univ., Voronezh, 1968.
- [28] Yuri Latushkin and Stephen Montgomery-Smith. Evolutionary semigroups and Lyapunov theorems in Banach spaces. *J. Funct. Anal.*, 127(1):173–197, 1995.
- [29] Sigeru Mizohata. Le problème de Cauchy pour les équations paraboliques. *J. Math. Soc. Japan*, 8:269–299, 1956.
- [30] Sylvie Monniaux and Abdelaziz Rhandi. Semigroup methods to solve non-autonomous evolution equations. *Semigroup Forum*, 60(1):122–134, 2000.
- [31] H. Neidhardt. *Integration of Evolutionsgleichungen mit Hilfe von Evolutionshalbgruppen*. Dissertation, AdW der DDR. Berlin 1979.
- [32] Hagen Neidhardt. On abstract linear evolution equations. I. *Math. Nachr.*, 103:283–298, 1981.

- [33] Hagen Neidhardt. On abstract linear evolution equations. II. Technical report, Prepr., Akad. Wiss. DDR, Inst. Math. P-MATH-07/81, Berlin, 1981.
- [34] Hagen Neidhardt. On linear evolution equations. III: Hyperbolic case. Technical report, Prepr., Akad. Wiss. DDR, Inst. Math. p-MATH-05/82, Berlin, 1982.
- [35] Gregor Nickel. Evolution semigroups and product formulas for nonautonomous Cauchy problems. *Math. Nachr.*, 212:101–116, 2000.
- [36] Gregor Nickel and Roland Schnaubelt. An extension of Kato’s stability condition for nonautonomous Cauchy problems. *Taiwanese J. Math.*, 2(4):483–496, 1998.
- [37] R. S. Phillips. Perturbation theory for semi-groups of linear operators. *Trans. Amer. Math. Soc.*, 74:199–221, 1953.
- [38] Andrea Posilicano. The Schrödinger equation with a moving point interaction in three dimensions. *Proc. Amer. Math. Soc.*, 135(6):1785–1793 (electronic), 2007.
- [39] Frank Rübiger, Abdelaziz Rhandi, and Roland Schnaubelt. Perturbation and an abstract characterization of evolution semigroups. *J. Math. Anal. Appl.*, 198(2):516–533, 1996.
- [40] Frank Rübiger and Roland Schnaubelt. The spectral mapping theorem for evolution semigroups on spaces of vector-valued functions. *Semigroup Forum*, 52(2):225–239, 1996.
- [41] M. R. Sayapova and D. R. Yafaev. Scattering theory for potentials of zero radius which are periodic with respect to time. In *Spectral theory. Wave processes*, volume 10 of *Probl. Mat. Fiz.*, pages 252–266, 301. Leningrad. Univ., Leningrad, 1982.
- [42] M. R. Sayapova and D. R. Yafaev. The evolution operator for time-dependent potentials of zero radius. *Trudy Mat. Inst. Steklov.*, 159:167–174, 1983. Boundary value problems of mathematical physics, 12.
- [43] Hiroki Tanabe. *Equations of evolution*, volume 6 of *Monographs and Studies in Mathematics*. Pitman (Advanced Publishing Program), Boston, Mass., 1979.
- [44] Naoki Tanaka. Generation of linear evolution operators. *Proc. Amer. Math. Soc.*, 128(7):2007–2015, 2000.
- [45] Naoki Tanaka. A characterization of evolution operators. *Studia Math.*, 146(3):285–299, 2001.
- [46] Naoki Tanaka. Nonautonomous abstract Cauchy problems for strongly measurable families. *Math. Nachr.*, 274/275:130–153, 2004.
- [47] H. F. Trotter. On the product of semi-groups of operators. *Proc. Amer. Math. Soc.*, 10:545–551, 1959.
- [48] D. R. Yafaev. Scattering theory for time-dependent zero-range potentials. *Ann. Inst. H. Poincaré Phys. Théor.*, 40(4):343–359, 1984.
- [49] Atsushi Yagi. On a class of linear evolution equations of “hyperbolic” type in reflexive Banach spaces. *Osaka J. Math.*, 16(2):301–315, 1979.
- [50] Atsushi Yagi. Remarks on proof of a theorem of Kato and Kobayasi on linear evolution equations. *Osaka J. Math.*, 17(1):233–243, 1980.
- [51] Kôzaku Yosida. Time dependent evolution equations in a locally convex space. *Math. Ann.*, 162:83–86, 1965/1966.
- [52] Kôzaku Yosida. *Functional analysis*. Second edition. Die Grundlehren der mathematischen Wissenschaften, Band 123. Springer-Verlag New York Inc., New York, 1968.