

Weierstraß-Institut
für Angewandte Analysis und Stochastik
Leibniz-Institut im Forschungsverbund Berlin e. V.

Preprint

ISSN 0946 – 8633

**Passing from bulk to bulk/surface evolution in the Allen–Cahn
equation**

Matthias Liero

submitted: January 9, 2012

Weierstrass Institute
Mohrenstr. 39
10117 Berlin
Germany
E-Mail: matthias.liero@wias-berlin.de

No. 1676
Berlin 2012



2010 *Mathematics Subject Classification.* 35K55,35K20,82C26.

Key words and phrases. Gradient flow, Allen–Cahn equation, dynamic boundary condition, energy balance, Γ -convergence.

The author was supported by the DFG Research Center MATHEON under project D22.

Edited by
Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS)
Leibniz-Institut im Forschungsverbund Berlin e. V.
Mohrenstraße 39
10117 Berlin
Germany

Fax: +49 30 2044975
E-Mail: preprint@wias-berlin.de
World Wide Web: <http://www.wias-berlin.de/>

Abstract

In this paper we formulate a boundary layer approximation for an Allen–Cahn-type equation involving a small parameter ε . Here, ε is related to the thickness of the boundary layer and we are interested in the limit when ε tends to 0 in order to derive nontrivial boundary conditions. The evolution of the system is written as an energy balance formulation of the L^2 -gradient flow with the corresponding Allen–Cahn energy functional. By transforming the boundary layer to a fixed domain we show the convergence of the solutions to a solution of a limit system. This is done by using concepts related to Γ - and Mosco convergence. By considering different scalings in the boundary layer we obtain different boundary conditions.

1 Introduction

In the recent years there has been a growing interest in the coupling of bulk and surface processes. One important example is the theory of spinodal decomposition of binary mixtures where dynamic boundary conditions are used to model the effective short-range interaction between the two mixture components and the wall (i.e., the boundary), see e.g. [Kra95, PuF97] and the references therein. Moreover, we refer to [KE*01, RaZ01, MiZ05, CFP06, FRG*06, CGM08, GGM08, SpW10] for an (incomplete) list of articles related to the mathematical analysis of dynamic boundary conditions for various evolutionary systems including the heat equation, the iso- and non-isothermal Allen–Cahn equation, the Cahn–Hilliard equation and the Caginalp system. In addition, we point out to the book [Tai09] for the connection to Feller semigroups and Markov processes.

In this paper we discuss the question whether such dynamic boundary conditions can be obtained as a limit of a family of bulk systems in the case of the Allen–Cahn equation. More precisely, for a domain Ω with C^2 -boundary Γ we introduce a boundary layer of thickness $\varepsilon > 0$, denoted by Σ_ε , that shrinks to Γ as ε tends to 0 (see Figure 1). In the domains Ω and Σ_ε we consider the following system of (bulk) Allen–Cahn-type equations

$$\begin{aligned}\tau_b \partial_t u_\varepsilon - A_b \Delta u_\varepsilon + W'_b(u_\varepsilon) &= 0 \quad \text{in } \Omega, \\ \tau_\varepsilon \partial_t u_\varepsilon - A_\varepsilon \Delta u_\varepsilon + \frac{1}{\varepsilon} W'_s(u_\varepsilon) &= 0 \quad \text{in } \Sigma_\varepsilon,\end{aligned}$$

subject to natural continuity and transmission conditions (see (2.1)) at the interface Γ . Here, W_b and W_s are given, in general nonconvex, bulk and surface potentials.

In order to derive nontrivial boundary conditions when ε goes to 0 we assume that the relaxation time τ_ε and the diffusion coefficient A_ε depend on ε in the boundary layer Σ_ε . This amounts

to different length and time scales in the bulk and in the boundary layer. We then show that the solutions of this system converge (up to subsequences) to a solution of a limit system which describes the coupling of bulk and surface evolution. The specific form of the derived limit system depends on the scalings of the coefficients τ_ε and A_ε . In particular, we will derive a hierarchy of dynamic and static boundary conditions depending on the scalings.

This approach is quite common in the derivation of lower-dimensional models in static elasticity, see e.g. [Cia00, FJM06]. Furthermore, we refer to [ScT10] for the derivation of models for conductive thin sheets using asymptotic expansion and to [CoR90] for the (non-rigorous) derivation of boundary conditions for the heat equation.

Here, however, we give a rigorous convergence proof which is based on an *energy balance formulation* of the underlying gradient flow structure of the Allen–Cahn equation. More precisely, by defining the Allen–Cahn energy functionals \mathcal{E}_ε the bulk equations can be written as L^2 -gradient flow in form of a *force balance* between the dissipative forces and the potential restoring forces given by the derivative of \mathcal{E}_ε . This force balance formulation is equivalent to a scalar *energy balance* equation written in terms of the energy functionals and quadratic dissipation potentials \mathcal{R}_ε , which in this case are given by the squares of the L^2 -norm (see also [AGS05, Mie11])

$$\mathcal{E}_\varepsilon(u_\varepsilon(t)) + \int_0^t \mathcal{R}_\varepsilon(\dot{u}_\varepsilon) + \mathcal{R}_\varepsilon^*(-D\mathcal{E}_\varepsilon(u_\varepsilon)) \, ds = \mathcal{E}_\varepsilon(u_\varepsilon(0)),$$

where $\mathcal{R}_\varepsilon^*$ denotes the dual dissipation potential, i.e., the Legendre transform of \mathcal{R}_ε . In particular, it is sufficient that only a lower energy estimate holds since the converse estimate follows from the properties of the Legendre transform and the chain rule for $t \mapsto \mathcal{E}_\varepsilon(u_\varepsilon(t))$.

The energy balance formulation opens the door for the application of notions of variational convergence such as Mosco and Γ -convergence [Att84, Dal93, Bra02]. Here we follow the ideas in [SaS04] (see also [BFG06, KMM06, Kur07]) where a method to prove the convergence of gradient flows for Γ -converging energy functionals was presented and applied to derive the limiting dynamics of vortices for the heat flow of the Ginzburg–Landau energy. However, we emphasize that the convergence of the gradient flow cannot follow from the Γ -convergence of the energy functionals only and extra conditions are required for the interplay of the convergence of the energy and the dissipation potentials. These extra conditions amount to the construction of mutual recovery curves for the energy and dissipation potentials.

Additionally, for λ -convex energy functionals the evolution of the system can be equivalently described by an *evolution variational inequality*

$$\mathcal{E}_\varepsilon(u_\varepsilon(t)) + \langle \mathcal{G}_\varepsilon \dot{u}_\varepsilon(t), u_\varepsilon(t) - \tilde{u} \rangle \leq \mathcal{E}_\varepsilon(\tilde{u}) - \Lambda_\varepsilon(u_\varepsilon(t) - \tilde{u}) \quad \forall \tilde{u},$$

where \mathcal{G}_ε denotes the linear and self-adjoint operator associated with \mathcal{R}_ε and Λ_ε corresponds to the λ -convexity of \mathcal{E}_ε .

We show that we can pass to the limit in the energy balance and the evolution variational inequality, respectively, in order to obtain corresponding limit formulations, written in terms of limit functionals \mathcal{E}_0 and \mathcal{R}_0 , which describe the coupling of bulk and surface evolution.

The paper is organized as follows: In Section 2 we introduce the underlying geometry of the boundary layer approximation and present the system of Allen–Cahn-type equations along with

technical details such as growth conditions, etc. The bulk system will then be cast into the gradient flow framework, in particular in the energy balance formulation. Furthermore, we introduce a change of coordinates in order to transform the system to a fixed domain (see Section 2.3). In this change of coordinates we characterize a point in the boundary layer by its projection and distance onto, resp., to Γ . Therefore we can decompose directions in Σ_ε into tangential and normal parts relative to Γ . The normal direction is then rescaled in order to obtain a fixed domain.

In Section 3 we present the main result of the paper, i.e., the limit passage in the energy balance. This is based on the results in [SaS04] which for the convenience of the reader will be reformulated here. Applied to our specific problem the construction of the mutual recovery curves is akin to the construction of the recovery sequences for the energy functionals in the sense of Γ -convergence.

In the final Section 4 the derived limit models will be discussed. In particular, depending on the scaling of the relaxation time and the diffusion coefficient in the boundary layer we obtain the usual Dirichlet- and Neumann boundary conditions as well as dynamic boundary conditions and boundary conditions that are to our knowledge not addressed in the literature so far, e.g., coupling of the bulk equation to an elliptic equation for the trace on Γ (see (4.4))

Notably, we also obtain the system recently considered in [SpW10] where it was studied regarding existence and uniqueness of global solutions, as well as asymptotic behavior and the existence of a global attractor. The system consists of the following bulk equation and dynamic boundary condition:

$$\begin{aligned} \tau_b \partial_t u - A_b \Delta u + W'_b(u) &= 0 && \text{in } \Omega, \\ \tau_s \partial_t u_\Gamma - A_s \Delta_\Gamma u_\Gamma + A_b \nabla u \cdot \nu + W'_s(u_\Gamma) &= 0 && \text{on } \Gamma, \\ u &= u_\Gamma && \text{on } \Gamma, \end{aligned}$$

where Δ_Γ denotes the Laplace-Beltrami operator on Γ .

Finally, we like to remark that the purpose of this paper is twofold: First, we want to identify the relevant scalings in the boundary layer system for deriving nontrivial boundary conditions. It would be interesting to apply these results to the related problem of deriving interface conditions in reaction-diffusion systems. Interface conditions in semiconductor heterostructures and biological systems are of great importance (see [Sch94, Gli11] and [EIS10]). Especially in organic photovoltaics interfaces are the fundamental building block, see [PoA06, Sect. 8].

Second, the paper contributes to the theory of application of Γ -convergence methods to evolutionary problems, especially to gradient flows. We refer to [MRS08], [Mie08] and [AM*11] for the application of the principles of Γ -convergence to rate-independent evolution, Hamiltonian systems and Wasserstein gradient flows, respectively.

2 Setting of the model

2.1 Definitions and notations

We consider an open and bounded domain $\Omega \subset \mathbb{R}^d$, $d \geq 2$, with a C^2 -boundary denoted by $\Gamma := \partial\Omega$. For a sufficiently small parameter $\varepsilon > 0$ we introduce the domain Ω_ε defined by

$$\Omega_\varepsilon := \{x \in \mathbb{R}^d : \text{dist}(x, \Omega) < \varepsilon\},$$

where $\text{dist}(x, \Omega) := \inf_{y \in \Omega} |x - y|$ denotes the distance to Ω . We call the set $\Sigma_\varepsilon := \Omega_\varepsilon \setminus \bar{\Omega}$ the boundary layer (or ε -neighborhood) of Ω . Obviously, we have the convergence $\Omega_\varepsilon \rightarrow \Omega$ for $\varepsilon \rightarrow 0$ with respect to the Hausdorff distance.

Let $T > 0$ be a finite time horizon. In the domain Ω_ε we consider the following system of Allen–Cahn-type equations:

$$\begin{aligned} \tau_b \partial_t u_\varepsilon - A_b \Delta u_\varepsilon + W'_b(u_\varepsilon) &= 0 & \text{in } [0, T] \times \Omega, \\ \tau_\varepsilon \partial_t u_\varepsilon - A_\varepsilon \Delta u_\varepsilon + \frac{1}{\varepsilon} W'_s(u_\varepsilon) &= 0 & \text{in } [0, T] \times \Sigma_\varepsilon, \end{aligned} \quad (\text{AC}_\varepsilon)$$

where $\tau_b, \tau_\varepsilon > 0$ denote the relaxation times, A_b, A_ε the diffusion coefficients, and W'_b, W'_s are the derivatives of potentials $W_b, W_s \in C^1(\mathbb{R})$ in the bulk and in the boundary layer, respectively. The equations above are subjected to the following natural boundary and transmission conditions at the interface Γ

$$\begin{aligned} A_\varepsilon \nabla u_\varepsilon \cdot \nu &= 0 & \text{on } [0, T] \times \partial\Omega_\varepsilon, \\ A_b \nabla u_\varepsilon \cdot \nu &= A_\varepsilon \nabla u_\varepsilon \cdot \nu & \text{on } [0, T] \times \Gamma, \\ [u_\varepsilon] &= 0 & \text{on } [0, T] \times \Gamma, \end{aligned} \quad (2.1)$$

where ν denotes the outer unit normal on Γ and $\partial\Omega_\varepsilon$ and $[\cdot]$ denotes the jump across the interface Γ . Finally, the system is completed by imposing the initial condition $u_\varepsilon(0) = u_\varepsilon^0$ in Ω_ε .

We assume that in the boundary layer Σ_ε the coefficients satisfy the scalings

$$\tau_\varepsilon = \varepsilon^{-\alpha} \tau_s \quad \text{and} \quad A_\varepsilon = \varepsilon^{-\beta} A_s$$

for given $\tau_s, A_s > 0$ and $\alpha, \beta \in \mathbb{R}$. We will discuss explicit choices for the exponents α and β in the following sections.

The nonlinearities W_b and W_s satisfy the following growth conditions:

$$\left. \begin{aligned} W_b(s) &\geq c|s|^{2-\gamma}, & |W'_b(s)| &\leq C(1+|s|^p) \\ W_s(s) &\geq c|s|^{2-\gamma}, & |W'_s(s)| &\leq C(1+|s|^p) \end{aligned} \right\} p \in [1, q[\text{ with } q = \begin{cases} \infty & d = 2, \\ \frac{d+2}{d-2} & d \geq 3. \end{cases} \quad (W_{\text{Grow}})$$

These are the same growth conditions imposed in [SpW10] for the bulk potential W_b , while we have a stronger growth condition for the boundary potential since we are in the full d -dimensional domain Σ_ε in contrast to the $(d-1)$ -dimensional boundary Γ in [SpW10].

A prominent example for the (nonconvex) potentials W_b and W_s is the double well potential $u \mapsto \frac{1}{4}(1-u^2)^2$, which obviously satisfies the above growth conditions for $d = 2, 3$.

We show that solutions of the system above converge in a certain sense to a solution of a limit system which consists of the bulk equation in Ω in (AC_ε) coupled to an equation posed on the boundary Γ . As we will see, the form of the latter equation will heavily depend on the choices for the scaling exponents α and β .

To put the above system in an abstract framework we introduce the function spaces $V_\varepsilon := H^1(\Omega_\varepsilon)$ and $H_\varepsilon := L^2(\Omega_\varepsilon)$. Then, the weak formulation of the system (AC_ε) reads: Find $u_\varepsilon \in H^1(0, T; H_\varepsilon) \cap L^2(0, T; V_\varepsilon)$ with $u_\varepsilon(0) = u_\varepsilon^0$ such that for all $\varphi \in V_\varepsilon$ and almost all $t \in [0, T]$ we have

$$0 = \int_{\Omega_\varepsilon} \left[\mathbb{G}_\varepsilon(x) \partial_t u_\varepsilon \varphi + \mathbb{A}_\varepsilon(x) \nabla u_\varepsilon \cdot \nabla \varphi + \mathbb{W}'_\varepsilon(x, u_\varepsilon) \varphi \right] dx, \quad (\mathbf{w}\text{-}AC_\varepsilon)$$

where we use the notation

$$\mathbb{G}_\varepsilon(x) = \begin{cases} \tau_b & \text{in } \Omega, \\ \tau_\varepsilon & \text{in } \Sigma_\varepsilon, \end{cases} \quad \mathbb{A}_\varepsilon(x) = \begin{cases} A_b & \text{in } \Omega, \\ A_\varepsilon & \text{in } \Sigma_\varepsilon, \end{cases} \quad \mathbb{W}_\varepsilon(x, \cdot) = \begin{cases} W_b(\cdot) & \text{in } \Omega, \\ \frac{1}{\varepsilon} W_s(\cdot) & \text{in } \Sigma_\varepsilon. \end{cases}$$

The existence of solutions of (AC_ε) , resp. $(\mathbf{w}\text{-}AC_\varepsilon)$, follows from standard arguments, see e.g. [Rou05, SpW10].

Theorem 2.1 (Existence of solutions). *For fixed $\varepsilon > 0$ let $u_\varepsilon^0 \in V_\varepsilon$ be given. Moreover, assume that the growth condition (W_{Growth}) holds. Then, there exists a solution $u_\varepsilon \in H^1(0, T; L^2(\Omega_\varepsilon)) \cap L^\infty(0, T; H^1(\Omega_\varepsilon))$ of the system (AC_ε) .*

2.2 Different formulations of gradient flows

It is well known that equation (AC_ε) is the L^2 -gradient flow of the Allen–Cahn functional $\mathcal{E}_\varepsilon : V_\varepsilon \rightarrow \mathbb{R}$ defined by

$$\mathcal{E}_\varepsilon(u) = \int_{\Omega_\varepsilon} \left[\frac{\mathbb{A}_\varepsilon(x)}{2} |\nabla u|^2 + \mathbb{W}_\varepsilon(x, u) \right] dx.$$

More precisely, by defining the symmetric and positive metric tensor $\mathcal{G}_\varepsilon : H_\varepsilon \rightarrow H_\varepsilon^*$ via $\langle \mathcal{G}_\varepsilon \dot{u}, \dot{v} \rangle = \int_{\Omega_\varepsilon} \mathbb{G}_\varepsilon(x) \dot{u} \dot{v} dx$ the equation in $(\mathbf{w}\text{-}AC_\varepsilon)$ can then be written in the form

$$\mathcal{G}_\varepsilon \dot{u}_\varepsilon(t) = -D\mathcal{E}_\varepsilon(u_\varepsilon(t)), \quad (\mathbf{fb}_\varepsilon)$$

with $D\mathcal{E}_\varepsilon(u)$ denoting the Gâteaux derivative of \mathcal{E}_ε which is well-defined due to (W_{Growth}) . Note that we (notationally) distinguish between H_ε and H_ε^* since the former is the space of velocities \dot{u} , while the latter is the space of forces $\xi = D\mathcal{E}_\varepsilon(u)$. Thus, \mathcal{G}_ε maps velocities to forces. The equation above can be seen as a *force balance* formulation of the gradient flow, where $\mathcal{G}_\varepsilon \dot{u}_\varepsilon$ and $D\mathcal{E}_\varepsilon(u_\varepsilon)$ are the dissipative and potential restoring forces, respectively. Defining the inverse operator $\mathcal{K}_\varepsilon = \mathcal{G}_\varepsilon^{-1} : H_\varepsilon^* \rightarrow H_\varepsilon$, mapping forces to velocities, we can write the force balance $(\mathbf{fb}_\varepsilon)$ as *rate equation* in H_ε

$$\dot{u}_\varepsilon(t) = -\mathcal{K}_\varepsilon D\mathcal{E}_\varepsilon(u_\varepsilon(t)) =: -\nabla_{\mathcal{G}_\varepsilon} \mathcal{E}_\varepsilon(u_\varepsilon(t)), \quad (\mathbf{re}_\varepsilon)$$

where $\nabla_{\mathcal{G}_\varepsilon} \mathcal{E}$ denotes the gradient of \mathcal{E}_ε with respect to the metric tensor \mathcal{G}_ε . Note that we have $\langle \xi, \mathcal{K}_\varepsilon \eta \rangle = \int_{\Omega_\varepsilon} \mathbb{G}_\varepsilon(x)^{-1} \xi \eta \, dx$. The operator \mathcal{G}_ε defines the quadratic dissipation potential $\mathcal{R}_\varepsilon(\dot{u}) = \frac{1}{2} \langle \mathcal{G}_\varepsilon \dot{u}, \dot{u} \rangle$ whose Legendre transform is given by \mathcal{K}_ε , i.e., we have $\mathcal{R}_\varepsilon^*(\xi) = \frac{1}{2} \langle \xi, \mathcal{K}_\varepsilon \xi \rangle$, where ξ denotes the “dual variable” (also called chemical potential or thermodynamically conjugated driving force, see [Mie11]). Furthermore, by using the chain rule we have that

$$\begin{aligned} \mathcal{E}_\varepsilon(u_\varepsilon(0)) - \mathcal{E}_\varepsilon(u_\varepsilon(t)) &= \int_0^t \langle \mathcal{G}_\varepsilon \dot{u}_\varepsilon, \dot{u}_\varepsilon \rangle \, ds \\ &= \int_0^t \langle D\mathcal{E}(u_\varepsilon), \mathcal{K}_\varepsilon D\mathcal{E}_\varepsilon(u_\varepsilon) \rangle \, ds \\ &= \int_0^t \left[\mathcal{R}_\varepsilon(\dot{u}_\varepsilon) + \mathcal{R}_\varepsilon^*(-D\mathcal{E}_\varepsilon(u_\varepsilon)) \right] \, ds. \end{aligned}$$

Hence, we see that the force balance (fb $_\varepsilon$) and the rate equation (re $_\varepsilon$) are equivalent to the *energy balance*

$$\mathcal{E}_\varepsilon(u_\varepsilon(t)) + \int_0^t \mathcal{R}_\varepsilon(\dot{u}_\varepsilon) + \mathcal{R}_\varepsilon^*(-D\mathcal{E}_\varepsilon(u_\varepsilon)) \, ds = \mathcal{E}_\varepsilon(u_\varepsilon(0)). \quad (\text{eb}_\varepsilon)$$

This formulation (whose solutions are also called curves of maximal slope see [AGS05, Sect. 1.3]) is indeed equivalent due to the Legendre Fenchel equivalences for convex potentials, i.e.,

$$\mu = D\mathcal{R}_\varepsilon(v) \iff v = D\mathcal{R}_\varepsilon^*(\mu) \iff \mathcal{R}_\varepsilon(v) + \mathcal{R}_\varepsilon^*(\mu) = \langle v, \mu \rangle.$$

We also used the chain rule $\frac{d}{dt} \mathcal{E}_\varepsilon(u) = \langle D\mathcal{E}_\varepsilon(u), \dot{u} \rangle$.

In fact, in (eb $_\varepsilon$) we only need the lower estimate “ \leq ”, the reverse estimate follows from the definition of the Legendre transform. The advantage of (eb $_\varepsilon$) is that it is a scalar equation in \mathbb{R} in contrast to the equations (fb $_\varepsilon$) and (re $_\varepsilon$) in H_ε^* and H_ε , respectively.

Let us remark here that $2\mathcal{R}_\varepsilon(\dot{u})$ and $2\mathcal{R}_\varepsilon^*(-D\mathcal{E}_\varepsilon(u))$ are nothing but the squares of the so called metric derivative of u and the metric slope of \mathcal{E}_ε calculated with respect to the metric induced by \mathcal{G}_ε , see [AGS05].

If the potentials W_b and W_s are λ_b -convex, resp. λ_s -convex, ($s \mapsto W_{b/s}(s) - \frac{\lambda_{b/s}}{2}|s|^2$ is convex) the energy functional satisfies the convexity estimate

$$\mathcal{E}_\varepsilon(\tilde{u}) \geq \mathcal{E}_\varepsilon(u) + \langle D\mathcal{E}_\varepsilon(u), \tilde{u} - u \rangle + \Lambda_\varepsilon(\tilde{u} - u) \quad \forall \tilde{u} \in V_\varepsilon,$$

where $\Lambda_\varepsilon(w) = \int_\Omega \frac{\lambda_b}{2} |w|^2 \, dx + \int_{\Sigma_\varepsilon} \frac{\lambda_s}{2\varepsilon} |w|^2 \, dx$. Note, that λ_b and λ_s do not have to be positive and therefore W_b and W_s are in general nonconvex. The double well potential $u \mapsto \frac{1}{4}(1-u^2)^2$ is λ -convex with $\lambda = -2$. Moreover, every $W \in C^{1,1}(\mathbb{R})$ is λ -convex.

Using the force balance formulation (fb $_\varepsilon$) we arrive at the equivalent formulation as *evolution variational inequality* (see [AGS05, DaS10])

$$\mathcal{E}_\varepsilon(u_\varepsilon(t)) + \langle \mathcal{G}_\varepsilon \dot{u}_\varepsilon(t), u_\varepsilon(t) - \tilde{u} \rangle \leq \mathcal{E}_\varepsilon(\tilde{u}) - \Lambda_\varepsilon(u_\varepsilon(t) - \tilde{u}) \quad \forall \tilde{u} \in V_\varepsilon. \quad (\text{evi}_\varepsilon)$$

Note that this formulation is written only in terms of the functional \mathcal{E}_ε and the operator \mathcal{G}_ε , and is therefore derivative free.

We study the behavior of the solutions u_ε when $\varepsilon \rightarrow 0$. In this case the boundary layer Σ_ε shrinks to Γ and we show that the “limit” of the sequence $u_\varepsilon|_{\Sigma_\varepsilon}$ describes the evolution on Γ .

2.3 Transformation of the problem

In order to provide a notion of convergence of the solutions u_ε we transform the variable domain Ω_ε to a fixed domain.

For this, note that due to the smoothness of the boundary Γ and for sufficiently small ε a point $x \in \Sigma_\varepsilon$ can be characterized in the following way: there exist unique $y \in \Gamma$ and $\vartheta \in]0, \varepsilon[$ such that $x = y + \vartheta\nu(y)$ (see e.g. [Wlo87, Chap. 2]), where ν denotes the unit outer normal on Γ (see Figure 1). Hence, we introduce the change of coordinates in Σ_ε

$$\begin{aligned} X_\varepsilon(y, \theta) &:= y + \varepsilon\theta\nu(y), & (y, \theta) &\in \Gamma \times]0, 1[, \\ Y_\varepsilon(x) &:= (y_\varepsilon(x), \theta_\varepsilon(x)) := (P_\varepsilon(x), d_\varepsilon(x)/\varepsilon), & x &\in \Omega_\varepsilon, \end{aligned}$$

where P_ε and d_ε denote the projection from Σ_ε on Γ and the distance to Γ , respectively.

With this change of coordinates we define $\Sigma := \Gamma \times]0, 1[$ and for a function $u : \Sigma_\varepsilon \rightarrow \mathbb{R}$ we set $U = u \circ X_\varepsilon : \Sigma \rightarrow \mathbb{R}$. Since the boundary Γ is of class C^2 we have that the outer unit normal satisfies $\nu \in C^1(\Gamma; \mathbb{R}^d)$. Therefore, if $u \in H^1(\Sigma_\varepsilon)$ we have $U \in H^1(\Sigma)$. More precisely, it holds

$$\begin{pmatrix} \nabla_\Gamma U \\ \partial_\theta U \end{pmatrix} = \begin{pmatrix} \mathbb{P}(y) - \varepsilon\theta\mathbb{S}(y) \\ \varepsilon\nu(y)^\top \end{pmatrix} \nabla u, \quad \text{and} \quad \nabla u = \begin{pmatrix} \mathbb{Q}_\varepsilon(x) \Big|_{\frac{1}{\varepsilon}\nu(P_\varepsilon(x))} \\ \frac{1}{\varepsilon} \end{pmatrix} \begin{pmatrix} \nabla_\Gamma U \\ \partial_\theta U \end{pmatrix},$$

where $\nabla_\Gamma U \in \mathcal{T}(\Gamma)$ denotes the tangential gradient of U on Γ , $\mathbb{P}(y)$ is the projection onto the tangential space at $y \in \Gamma$, $\mathbb{S} = -\partial\nu/\partial y$ is the so-called shape operator (see e.g. [dCa76]) and \mathbb{Q}_ε is such that $\mathbb{Q}_\varepsilon(\mathbb{P} - \varepsilon\theta\mathbb{S}) = \mathbb{P}$.

The tangential gradient $\nabla_\Gamma U$ on Γ can be characterized in the following way (see [SaV97, dCa76]): For $V : \Gamma \rightarrow \mathbb{R}$ denote by \tilde{V} a smooth extension of V to \mathbb{R}^d , then $\nabla_\Gamma V(y) = \mathbb{P}(y)[\nabla\tilde{V}]$. It is easy to check that this definition is well-defined and independent of the extension \tilde{V} , moreover, we have that $\mathbb{P} = I - \nu \otimes \nu$. Similarly, the divergence on Γ for vector fields V can be defined as

$$\operatorname{div}_\Gamma V = \operatorname{div}\tilde{V} - \nabla(\tilde{V} \cdot \nu)\nu,$$

where \tilde{V} is again a smooth extension of V . In this framework the Laplace–Beltrami operator Δ_Γ has the simple form $\Delta_\Gamma U = \operatorname{div}_\Gamma(\nabla_\Gamma U)$. For a vector field $V \in L^2(\Gamma; \mathcal{T}(\Gamma))$ such that $\operatorname{div}_\Gamma V \in L^2(\Gamma)$ and $U \in H^1(\Gamma)$ we have Green's formula

$$-\int_\Gamma \nabla_\Gamma U \cdot V \, d\Gamma = \int_\Gamma U \operatorname{div}_\Gamma V \, d\Gamma.$$

In contrast to Σ_ε we leave the bulk domain Ω untransformed. Hence, we introduce the spaces for the bulk variable $u : \Omega \rightarrow \mathbb{R}$ and the surface variable $U : \Sigma \rightarrow \mathbb{R}$

$$\mathcal{V} := \{(u, U) \in H^1(\Omega) \times H^1(\Sigma) : u|_\Gamma = U|_{\{\theta=0\}}\}, \quad \mathcal{H} := L^2(\Omega) \times L^2(\Sigma).$$

The measure on Σ is given by $d\mu = d\Gamma \otimes d\lambda^1$, i.e., the product of the surface measure on Γ and the one-dimensional Lebesgue measure on $]0, 1[$. The space $H^1(\Sigma)$ is defined in the usual way, i.e., the closure of $C^1(\Sigma)$ with respect to the norm $\|\cdot\|_{H^1(\Sigma)}$, where

$$\|U\|_{H^1(\Sigma)}^2 = \int_\Sigma [|U|^2 + |\nabla_\Gamma U|^2 + |\partial_\theta U|^2] \, d\mu.$$




Figure 1: Transformation of the boundary layer.

Now, substituting the above transformations in \mathcal{E}_ε we arrive at the transformed energy functional $E_\varepsilon : \mathcal{V} \rightarrow [0, \infty[$, for $\mathbf{u} = (u, U)$ defined by

$$E_\varepsilon(\mathbf{u}) = \int_{\Omega} \left[\frac{A_b}{2} |\nabla u|^2 + W_b(u) \right] dx + \int_{\Sigma} \left[\frac{A_\varepsilon}{2} \left(\nabla_{\Gamma} U \cdot \mathbb{B}_\varepsilon(y, \theta) \nabla_{\Gamma} U + \frac{1}{\varepsilon^2} |\partial_\theta U|^2 \right) + W_\varepsilon(U) \right] \mathbb{J}_\varepsilon(y, \theta) d\mu,$$

where $\mathbb{B}_\varepsilon = \mathbb{Q}_\varepsilon^\top \mathbb{Q}_\varepsilon$ and \mathbb{J}_ε describes the change of volume due to the transformation. Additionally, the transformed dissipation potential $R_\varepsilon : \mathcal{H} \rightarrow [0, \infty[$ reads

$$R_\varepsilon(\dot{\mathbf{u}}) = \int_{\Omega} \frac{\tau_b}{2} |\dot{u}|^2 dx + \int_{\Sigma} \frac{\tau_\varepsilon}{2} |\dot{U}|^2 \mathbb{J}_\varepsilon(y, \theta) d\mu.$$

We denote by $G_\varepsilon : \mathcal{H} \rightarrow \mathcal{H}^*$ the associated operator, i.e., $R_\varepsilon(\dot{\mathbf{u}}) = \frac{1}{2} \langle G_\varepsilon \dot{\mathbf{u}}, \dot{\mathbf{u}} \rangle$. The inverse operator $K_\varepsilon = G_\varepsilon^{-1} : \mathcal{H}^* \rightarrow \mathcal{H}$ gives the dual dissipation potential R_ε^* , more precisely, for a dual variable $\boldsymbol{\xi} = (\xi, \Xi)$ it reads

$$R_\varepsilon^*(\boldsymbol{\xi}) = \int_{\Omega} \frac{\tau_b^{-1}}{2} |\xi|^2 dx + \int_{\Sigma} \frac{\tau_\varepsilon^{-1}}{2 \mathbb{J}_\varepsilon(y, \theta)} |\Xi|^2 d\mu.$$

Note that although we have that $\mathcal{E}_\varepsilon(u) = E_\varepsilon(\mathbf{u})$ and $\mathcal{R}_\varepsilon(\dot{u}) = R_\varepsilon(\dot{\mathbf{u}})$ it holds that $\mathcal{R}_\varepsilon^*(\xi) \neq R_\varepsilon^*(\boldsymbol{\xi})$. This is due to the fact that the Legendre transform R_ε^* is calculated in the space \mathcal{H} whose norm and scalar product is not inherited from H_ε . For the same reasons we have that $D\mathcal{E}_\varepsilon(u) \neq DE_\varepsilon(\mathbf{u})$. However, it holds that $\mathcal{R}_\varepsilon(-D\mathcal{E}_\varepsilon(u)) = R_\varepsilon(-DE_\varepsilon(\mathbf{u}))$. In particular, the energy balance (eb $_\varepsilon$) is equivalent to

$$E_\varepsilon(\mathbf{u}_\varepsilon(t)) + \int_0^t \left[R_\varepsilon(\dot{\mathbf{u}}_\varepsilon) + R_\varepsilon^*(-DE_\varepsilon(\mathbf{u}_\varepsilon)) \right] ds = E_\varepsilon(\mathbf{u}_\varepsilon(0)). \quad (\text{EB}_\varepsilon)$$

Moreover, in the λ -convex case the evolution of the transformed system is equivalently described by the following evolution variational inequality which corresponds to (evi $_\varepsilon$)

$$E_\varepsilon(\mathbf{u}_\varepsilon(t)) + \langle G_\varepsilon \dot{\mathbf{u}}_\varepsilon(t), \mathbf{u}_\varepsilon(t) - \tilde{\mathbf{u}} \rangle \leq E_\varepsilon(\tilde{\mathbf{u}}) - \Lambda_\varepsilon(\mathbf{u}_\varepsilon(t) - \tilde{\mathbf{u}}), \quad (\text{EVI}_\varepsilon)$$

where $\Lambda_\varepsilon(\mathbf{u}) = \int_\Omega \frac{\lambda_b}{2} |u|^2 dx + \int_\Sigma \frac{\lambda_s}{2} |U|^2 \mathbb{J}_\varepsilon d\mu$.

We will use both formulations, (EB_ε) and (EVI_ε) , for the convergence analysis. Note that (EB_ε) contains the derivative of the energy functional E_ε while (EVI_ε) does not. Conversely, (EVI_ε) contains the derivative of the dissipation potential R_ε while (EB_ε) is free of it.

The following lemma is concerned with the convergences of the geometrical quantities \mathbb{B}_ε and \mathbb{J}_ε .

Lemma 2.2. *It holds that $\mathbb{B}_\varepsilon \rightarrow \mathbb{I}$ uniformly in Σ , with \mathbb{I} denoting the identity in the tangent bundle of Γ , and $\mathbb{J}_\varepsilon/\varepsilon \rightarrow 1$ uniformly in Σ .*

Proof. The easiest (although not most elegant) way to see that the convergence is indeed as stated, is to switch to local coordinates and calculate \mathbb{B}_ε and \mathbb{J}_ε explicitly in terms of the covariant and contravariant basis vectors (see [Cia00] for a related problem in the theory of elastic shells). \square

3 Convergence of the system

Our result is formulated abstractly in terms of Mosco convergence of E_ε towards a limit E_0 and of R_ε towards R_0 . For functionals F_n , defined on a Banach space \mathcal{Q} , the definition of Mosco convergence is as follows:

$$F_n \xrightarrow{M} F \Leftrightarrow \begin{cases} \text{(i) Liminf estimate for weakly converging sequences:} \\ \mathbf{q}_n \rightharpoonup \mathbf{q} \implies F(\mathbf{q}) \leq \liminf_{n \rightarrow \infty} F_n(\mathbf{q}), \\ \text{(ii) Existence of strongly converging recovery sequences:} \\ \forall \hat{\mathbf{q}} \in \mathcal{Q} \exists (\hat{\mathbf{q}}_n)_n : \hat{\mathbf{q}}_n \rightarrow \hat{\mathbf{q}} \text{ and } F(\hat{\mathbf{q}}) \geq \limsup_{n \rightarrow \infty} F_n(\hat{\mathbf{q}}_n). \end{cases}$$

Hence, Mosco convergence is nothing but Γ -convergence in the weak *and* in the strong topology.

Since it is essential to choose the right topology for computing the Γ - or Mosco limits, the first step in our convergence proof is to derive a priori estimates for the solutions $(u_\varepsilon, U_\varepsilon)$. This is addressed in the following lemma.

Lemma 3.1 (A priori estimate). *Let $E_\varepsilon(\mathbf{u}_\varepsilon(0)) \leq C < \infty$. Then, there exist constants $C_1, C_2, C_3, C_4 > 0$, independent of ε , such that*

$$\begin{aligned} \|\dot{u}_\varepsilon\|_{L^2([0,T] \times \Omega)}^2 + \varepsilon^{1-\alpha} \|\dot{U}_\varepsilon\|_{L^2([0,T] \times \Sigma)}^2 &\leq C_1, \\ \|D_u E_\varepsilon(\mathbf{u}_\varepsilon)\|_{L^2([0,T] \times \Omega)}^2 + \frac{1}{\varepsilon^{1-\alpha}} \|D_U E_\varepsilon(\mathbf{u}_\varepsilon)\|_{L^2([0,T] \times \Sigma)}^2 &\leq C_2, \\ \|\nabla u_\varepsilon(t)\|_{L^2(\Omega)}^2 + \varepsilon^{1-\beta} \|\nabla_\Gamma U_\varepsilon(t)\|_{L^2(\Sigma)}^2 + \varepsilon^{-(\beta+1)} \|\partial_\theta U_\varepsilon(t)\|_{L^2(\Sigma)}^2 &\leq C_3, \\ \|u_\varepsilon(t)\|_{L^2(\Omega)}^2 + \|U_\varepsilon(t)\|_{L^2(\Sigma)}^2 &\leq C_4, \end{aligned} \tag{3.1}$$

for all $t \in [0, T]$.

Proof. The estimates in (3.1) are a direct consequence of the energy balance (EB $_\varepsilon$). We remind that the relaxation time and the diffusion coefficient are given by $\tau_\varepsilon = \tau_s \varepsilon^{-\alpha}$, $A_\varepsilon = A_s \varepsilon^{-\beta}$. The energy functional satisfies the estimate

$$\begin{aligned} E_\varepsilon(\mathbf{u}_\varepsilon) \geq C(\|\nabla u_\varepsilon\|_{L^2(\Omega)}^2 + \|u_\varepsilon\|_{L^2(\Omega)}^2 + \|U_\varepsilon\|_{L^2(\Omega)}^2 \\ + \varepsilon^{1-\beta} \|\nabla_\Gamma U_\varepsilon\|_{L^2(\Sigma)}^2 + \varepsilon^{-(\beta+1)} \|\partial_\theta U_\varepsilon\|_{L^2(\Sigma)}^2) - c, \end{aligned}$$

where we have used the quadratic growth of the nonlinearities W_b and W_s as well as Lemma 2.2. The dissipation potential satisfies

$$\begin{aligned} R_\varepsilon(\dot{\mathbf{u}}_\varepsilon) &\geq C(\|\dot{u}_\varepsilon\|_{L^2(\Omega)}^2 + \varepsilon^{1-\alpha} \|\dot{U}_\varepsilon\|_{L^2(\Sigma)}^2), \\ R_\varepsilon^*(\boldsymbol{\xi}_\varepsilon) &\geq C(\|D_u E_\varepsilon(\mathbf{u}_\varepsilon)\|_{L^2(\Omega)}^2 + \varepsilon^{\alpha-1} \|D_U E_\varepsilon(\mathbf{u}_\varepsilon)\|_{L^2(\Sigma)}^2). \end{aligned}$$

By assumption the lefthand-side in the energy balance (EB $_\varepsilon$) is bounded, thus we arrive at (3.1). \square

Remark 3.2. *The a priori estimates show that the critical scaling for the relaxation time $\tau_\varepsilon = \varepsilon^{-\alpha} \tau_s$ is $\alpha=1$. For $\alpha < 1$ we expect the time derivatives in Σ to blow up while the thermodynamically conjugated driving forces tend to 0 in the limit. This means that we have a much faster timescale in the boundary layer, such that in the limit the system is always in equilibrium on the boundary. Conversely, $\alpha > 1$ amounts to a slower timescale in the boundary layer with no evolution. In contrast to these degenerate cases $\alpha=1$ results in a nontrivial dynamic boundary condition as in [SpW10].*

In addition, we find the characteristic values $\beta \in \{-1, +1\}$ for the scalings of the diffusion coefficient $A_\varepsilon = \varepsilon^{-\beta} A_s$ in the boundary layer. For $\beta > 1$ all derivatives have to vanish such that U is constant (in every connected component of Σ). However, it is not fixed and may evolve in time, we refer to this as the fast diffusion case. Conversely, for $\beta < 1$ we expect the tangential derivatives to blow up in the boundary layer (no diffusion case). For $\beta=1$ we expect genuine surface diffusion.

The crucial point is that in all of the cases above the derivative with respect to θ has to vanish. Hence, in the limit the surface variable U is given only by its trace on Γ which allows for the reduction to surface evolution, see Section 4 for the final discussion.

Lemma 3.1 shows that we can extract a (not relabeled) subsequence $\mathbf{u}_\varepsilon = (u_\varepsilon, U_\varepsilon)$ such that for the bulk variable u_ε we have the convergence

$$\begin{aligned} u_\varepsilon &\overset{*}{\rightharpoonup} u && \text{in } L^\infty(0, T; H^1(\Omega)), \\ \dot{u}_\varepsilon &\rightharpoonup \dot{u} && \text{in } L^2([0, T] \times \Omega), \\ DE_b(u_\varepsilon) &\rightharpoonup DE_b(u) && \text{in } L^2([0, T] \times \Omega), \end{aligned} \tag{3.2}$$

while additionally for U_ε we have

$$\begin{aligned} U_\varepsilon &\overset{*}{\rightharpoonup} U && \text{in } L^\infty(0, T; L^2(\Omega)), \\ \partial_\theta U_\varepsilon &\rightarrow 0 && \text{in } L^\infty(0, T; L^2(\Sigma)). \end{aligned} \tag{3.3}$$

Depending on the choice for β we find a subsequence such that the tangential gradients of U_ε satisfy

$$\left. \begin{array}{l} \nabla_\Gamma U_\varepsilon \xrightarrow{*} \nabla_\Gamma U \quad \text{for } \beta = 1 \\ \nabla_\Gamma U_\varepsilon \rightarrow 0 \quad \quad \quad \text{for } \beta > 1 \end{array} \right\} \text{ in } L^\infty(0, T; L^2(\Sigma)). \quad (3.4)$$

Furthermore, we can assume that the following convergences take place

$$\left. \begin{array}{l} \dot{U}_\varepsilon \rightharpoonup \dot{U} \quad \text{and} \quad DE_{s,\varepsilon}(U_\varepsilon) \rightharpoonup \Xi \quad \text{for } \alpha = 1 \\ \dot{U}_\varepsilon \rightarrow 0 \quad \text{and} \quad DE_{s,\varepsilon}(U_\varepsilon) \rightarrow 0 \quad \text{for } \alpha < 1 \end{array} \right\} \text{ in } L^2([0, T] \times \Sigma), \quad (3.5)$$

where $E_{s,\varepsilon}$ is such that $E_\varepsilon(u, U) = E_b(u) + E_{s,\varepsilon}(U)$ and $\Xi \in L^2([0, T] \times \Sigma)$ is to be determined.

Obviously, the energy functionals E_ε blow up if the derivative with respect to θ does not vanish (for $\beta > 1$ the same holds for the tangential derivatives). Thus, we expect the limit problems to be defined on the subspace of functions that are constant in normal direction (and tangential direction for $\beta > 1$).

Let us consider the case $\beta \geq 1$ first: We define the reduced spaces $\mathcal{V}_{\text{tang}}$, $\mathcal{V}_{\text{const}}$ and their closures in \mathcal{H} via

$$\begin{aligned} \mathcal{V}_{\text{tang}} &:= \{(u, U) \in \mathcal{V} : \partial_\theta U = 0 \text{ a.e. in } \Sigma\}, & \mathcal{H}_{\text{tang}} &= \overline{\mathcal{V}_{\text{tang}}}^{\mathcal{H}}, \\ \mathcal{V}_{\text{const}} &:= \{(u, U) \in \mathcal{V} : U = \text{const a.e. in } \Sigma\}, & \mathcal{H}_{\text{const}} &= \overline{\mathcal{V}_{\text{const}}}^{\mathcal{H}}. \end{aligned}$$

In the following theorem we prove the Mosco convergence of the energy functionals E_ε for $\beta \geq 1$ in \mathcal{V} .

Theorem 3.3 (Mosco convergence, Part I). *For $\beta=1$ the energy functionals E_ε converge in the sense of Mosco to the limit functional $E_{\text{tang}} : \mathcal{V} \rightarrow \mathbb{R}_\infty$ given by*

$$E_{\text{tang}}(\mathbf{u}) = \begin{cases} E_b(u) + \int_\Sigma \left[\frac{A_s}{2} |\nabla_\Gamma U|^2 + W_s(U) \right] d\mu & \text{if } \mathbf{u} \in \mathcal{V}_{\text{tang}}, \\ +\infty & \text{otherwise.} \end{cases}$$

For $\beta > 1$ the Mosco limit of E_ε , denoted E_{const} , is given by

$$E_{\text{const}}(\mathbf{u}) = \begin{cases} E_b(u) + \int_\Sigma W_s(U) d\mu & \text{if } \mathbf{u} \in \mathcal{V}_{\text{const}}, \\ +\infty & \text{otherwise.} \end{cases}$$

Proof. Here we only consider the case $\beta = 1$. The result for the other case follows analogously.

Liminf estimate for weak convergence. For all sequences $\mathbf{u}_\varepsilon = (u_\varepsilon, U_\varepsilon) \rightharpoonup \mathbf{u} = (u, U)$ in \mathcal{V} we have to show $E_{\text{tang}}(\mathbf{u}) \leq \liminf_{\varepsilon \rightarrow 0} E_\varepsilon(\mathbf{u}_\varepsilon)$. First, let $\mathbf{u} \notin \mathcal{V}_{\text{tang}}$. Since the norm on \mathcal{V} is weakly lower semicontinuous, we find $\liminf_{\varepsilon \rightarrow 0} \|\partial_\theta U_\varepsilon\|_{L^2(\Sigma)} > 0$. Using the coercivity of E_ε we conclude

$$E_\varepsilon(\mathbf{u}_\varepsilon) \geq \frac{C}{\varepsilon^2} \|\partial_\theta U_\varepsilon\|_{L^2(\Sigma)}^2 - c \rightarrow \infty = E_{\text{tang}}(\mathbf{u}).$$

Hence, we can assume that $\mathbf{u} \in \mathcal{V}_{\text{tang}}$ and $\sup_{0 < \varepsilon < \varepsilon_0} E_\varepsilon(\mathbf{u}_\varepsilon) < \infty$, for a sufficiently small $\varepsilon_0 > 0$.

The compact embedding $\mathcal{V} \subset\subset L^q(\Omega) \times L^q(\Sigma)$, where $q \in [1, \infty[$ for $d = 2$ and $q < 2d/(d-2)$ otherwise, yields the strong convergence $(u_\varepsilon, U_\varepsilon) \rightarrow (u, U)$ in $L^q(\Omega) \times L^q(\Sigma)$. Thus, using the growth conditions for W_b and W_s we conclude that

$$\int_{\Omega} W_b(u_\varepsilon) dx \rightarrow \int_{\Omega} W_b(u) dx \quad \text{and} \quad \int_{\Sigma} W_s(U_\varepsilon) d\mu \rightarrow \int_{\Sigma} W_s(U) d\mu.$$

As before, we denote the bulk and surface energy parts of E_ε by E_b and $E_{s,\varepsilon}$, such that $E_\varepsilon(\mathbf{u}_\varepsilon) = E_b(u_\varepsilon) + E_{s,\varepsilon}(U_\varepsilon)$. It holds that

$$E_\varepsilon(\mathbf{u}_\varepsilon) \geq E_b(u_\varepsilon) + \int_{\Sigma} \left[\frac{A_s}{2} \nabla_{\Gamma} U_\varepsilon \cdot \mathbb{B}_\varepsilon(y, \theta) \nabla_{\Gamma} U_\varepsilon + W_s(U_\varepsilon) \right] \frac{\mathbb{J}_\varepsilon(y, \theta)}{\varepsilon} d\mu.$$

Hence, by the uniform convergence of \mathbb{B}_ε and $\mathbb{J}_\varepsilon/\varepsilon$ we obtain the lim inf estimate.

Limsup estimate for strongly converging recovery sequences. The construction of recovery sequences $\widehat{\mathbf{u}}_\varepsilon$ such that $\widehat{\mathbf{u}}_\varepsilon \rightarrow \mathbf{u}$ and $E_\varepsilon(\widehat{\mathbf{u}}_\varepsilon) \rightarrow E_{\text{tang}}(\mathbf{u})$ is straightforward: For $\mathbf{u} \notin \mathcal{V}_{\text{tang}}$ the result is trivial since $E_{\text{tang}}(\mathbf{u}) = \infty$ and we may take $\widehat{\mathbf{u}}_\varepsilon = \mathbf{u}$ and argue as in the first step.

For $\mathbf{u} \in \mathcal{V}_{\text{tang}}$ we can choose the constant sequence $\widehat{\mathbf{u}}_\varepsilon = \mathbf{u}$ since the derivative with respect to θ does not appear in E_ε and we can conclude

$$E_\varepsilon(\mathbf{u}) = E_b(u) + \int_{\Sigma} \left[\frac{A_s}{2} \nabla_{\Gamma} U \cdot \mathbb{B}_\varepsilon(y, \theta) \nabla_{\Gamma} U + W_s(U) \right] \frac{\mathbb{J}_\varepsilon(y, \theta)}{\varepsilon} d\mu \rightarrow E_{\text{tang}}(\mathbf{u}),$$

where we used Lemma 2.2 again. □

The remaining case $\beta \in]-1, 1[$ is more complicated since we lose the uniform coercivity of the energy functionals on \mathcal{V} . Hence, we have to work in the coarser topology of the bigger space \mathcal{W} defined by

$$\mathcal{W} := \{(u, U) \in H^1(\Omega) \times L^2(\Sigma) : \partial_\theta U \in L^2(\Sigma), u|_{\Gamma=U}|_{\{\theta=0\}}\}.$$

Let us point out here that the existence of the derivative with respect to θ in $L^2(\Sigma)$ suffices for the well-definedness of the trace on Γ since for arbitrary $U \in C^\infty(\Sigma)$ it holds that

$$\|U|_{\{\theta=0\}}\|_{L^2(\Gamma)} \leq C(\|U\|_{L^2(\Sigma)} + \|\partial_\theta U\|_{L^2(\Sigma)}).$$

As before we introduce a reduced space of functions which are constant in normal direction

$$\mathcal{W}_{\text{nodiff}} := \{(u, U) \in \mathcal{W} : \partial_\theta U = 0 \text{ a.e. in } \Sigma\}.$$

Since the convergence of the surface variable U_ε is in general only weak in $L^2(\Sigma)$ and the nonlinearity W_s is allowed to be nonconvex we have to replace W_s in the limit by its convex envelope, denoted W_s^{**} in the following (see e.g. [Bra02, Dal93]).

Theorem 3.4 (Mosco convergence, Part II). *Let $-1 < \beta < 1$. The energy functionals E_ε Γ -converge on \mathcal{W} to the limit functional $E_{\text{nodiff}} : \mathcal{W} \rightarrow \mathbb{R}_\infty$ given by*

$$E_{\text{nodiff}}(\mathbf{u}) = \begin{cases} E_b(u) + \int_{\Sigma} W_s^{**}(U) d\mu & \text{if } \mathbf{u} \in \mathcal{W}_{\text{nodiff}}, \\ +\infty & \text{otherwise.} \end{cases}$$

Proof. Liminf estimate for weak convergence. Let $\mathbf{u}_\varepsilon = (u_\varepsilon, U_\varepsilon) \rightharpoonup \mathbf{u} = (u, U)$ in \mathcal{W} . By arguing as in Theorem 3.3 we can assume that $\mathbf{u} \in \mathcal{W}_{\text{nodiff}}$ and $\sup_{0 < \varepsilon < \varepsilon_0} E_\varepsilon(\mathbf{u}_\varepsilon) < \infty$. We have the estimate

$$E_\varepsilon(\mathbf{u}_\varepsilon) \geq E_b(u_\varepsilon) + \int_{\Sigma} W_s^{**}(U_\varepsilon) \frac{\mathbb{J}_\varepsilon(y, \theta)}{\varepsilon} d\mu.$$

Applying $\liminf_{\varepsilon \rightarrow 0}$ to both sides of the estimate and using the uniform convergence of $\mathbb{J}_\varepsilon/\varepsilon$ and the weak lower semicontinuity of $U \mapsto \int_{\Sigma} W_s^{**}(U) d\mu$ on $L^2(\Sigma)$ we conclude that $\liminf_{\varepsilon \rightarrow 0} E_\varepsilon(\mathbf{u}_\varepsilon) \geq E_{\text{nodiff}}(\mathbf{u})$.

Limsup estimate for recovery sequences. Let $\mathbf{u} \in \mathcal{W}_{\text{nodiff}}$ be such that $E_{\text{nodiff}}(\mathbf{u}) < \infty$. By the density of $\mathcal{V}_{\text{tang}}$ in $\mathcal{W}_{\text{nodiff}}$ we can find a sequence $(\widehat{\mathbf{u}}_\varepsilon)_{\varepsilon > 0} \subset \mathcal{V}_{\text{tang}}$ such that $\widehat{\mathbf{u}}_\varepsilon \rightarrow \mathbf{u}$ (strongly) in \mathcal{W} and $\varepsilon^\sigma \|\nabla_{\Gamma} \widehat{U}_\varepsilon\|_{L^2(\Sigma)}^2 \rightarrow 0$, where $\sigma = 1 - \beta \in]0, 2[$. Since $\widehat{\mathbf{u}}_\varepsilon = (\widehat{u}_\varepsilon, \widehat{U}_\varepsilon)$ converges strongly in \mathcal{W} we can extract a (not relabeled) sequence such that $\widehat{U}_\varepsilon(y, \theta) \rightarrow U(y, \theta)$ a.e. in Σ . Using Fatou's lemma we obtain

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0} E_\varepsilon(\widehat{\mathbf{u}}_\varepsilon) &\leq \limsup_{\varepsilon \rightarrow 0} \left\{ E_b(\widehat{u}_\varepsilon) + \int_{\Sigma} \left[C\varepsilon^\sigma |\nabla_{\Gamma} \widehat{U}_\varepsilon|^2 + W_s(U_\varepsilon) \right] \frac{\mathbb{J}_\varepsilon(y, \theta)}{\varepsilon} d\mu \right\} \\ &\leq E_b(u) + \int_{\Sigma} W_s(U) d\mu. \end{aligned}$$

The left-hand side, also known as Γ -limes superior (or upper Γ -limit), is weakly lower semicontinuous on \mathcal{W} (see [Dal93, Bra02]). Hence, by taking the lower semicontinuous envelope on both sides we arrive at $\limsup_{\varepsilon \rightarrow 0} E_\varepsilon(\mathbf{u}_\varepsilon) \leq E_{\text{nodiff}}(\mathbf{u})$. \square

Let us emphasize here that in last case, also for convex W_s , the energy functional E_{nodiff} is in general not Gâteaux differentiable on $\mathcal{W}_{\text{nodiff}}$. Thus, we restrict ourselves to the case of a quadratic potential, such that $W_s(U) = \frac{\omega_s}{2} |U|^2$ with $\omega_s > 0$. In this much simpler case the (strongly converging) recovery sequences are given by $\widehat{\mathbf{u}}_\varepsilon$ in the proof of Theorem 3.4. Hence, E_ε Mosco converges to E_{nodiff} in \mathcal{W} .

The limits for the dissipation potential R_ε and the dual dissipation potentials R_ε^* for the cases $\alpha=1$, $\alpha>1$ and $\alpha<1$ are easily computed. Note that for the last two cases the uniform coercivity of R_ε^* and R_ε on \mathcal{H}^* and \mathcal{H} , respectively, is lost.

For the nondegenerate case $\alpha=1$ we have the convergence

$$R_\varepsilon \xrightarrow{M} R_{\text{dyn}} \quad \text{with} \quad R_{\text{dyn}}(\dot{\mathbf{u}}) = \int_{\Omega} \frac{\tau_b}{2} |\dot{u}|^2 dx + \int_{\Sigma} \frac{\tau_s}{2} |\dot{U}|^2 d\mu$$

while for the other two cases (the slow and the fast evolution cases, see Remark 3.2) it holds

$$\begin{aligned} \alpha > 1 : R_\varepsilon(\dot{\mathbf{u}}) &\rightarrow R_{\text{slow}}(\dot{\mathbf{u}}) \quad \text{with} \quad R_{\text{slow}}(\dot{u}, \dot{U}) = \begin{cases} \int_{\Omega} \frac{\tau_b}{2} |\dot{u}|^2 dx & \text{if } \dot{U} = 0, \\ \infty & \text{else,} \end{cases} \\ \alpha < 1 : R_\varepsilon(\dot{\mathbf{u}}) &\rightarrow R_{\text{fast}}(\dot{\mathbf{u}}) \quad \text{with} \quad R_{\text{fast}}(\dot{u}, \dot{U}) = \int_{\Omega} \frac{\tau_b}{2} |\dot{u}|^2 dx. \end{aligned}$$

The Legendre transforms are easily computed as

$$R_{\text{slow}}^*(\xi, \Xi) = \int_{\Omega} \frac{\tau_b^{-1}}{2} |\xi|^2 dx \quad \text{and} \quad R_{\text{fast}}^*(\xi, \Xi) = \begin{cases} \int_{\Omega} \frac{\tau_b^{-1}}{2} |\xi|^2 dx & \text{if } \Xi = 0, \\ \infty & \text{else.} \end{cases}$$

We see that the limits for R_ε correspond to the observations made in Remark 3.2. For $\alpha > 1$ we obtain the static condition $\dot{U} = 0$, i.e., fixed (boundary-)evolution. While for $\alpha < 1$ the condition $\Xi = 0$ for the thermodynamically conjugated driving force means that the (boundary-)system is in equilibrium.

3.1 Passing to the limit in the energy balance (EB $_\varepsilon$)

In this subsection we focus on the energy balance formulation (EB $_\varepsilon$) and show that the limit $\mathbf{u} = (u, U)$ in (3.2)–(3.5) is a solution of the limit system (E_0, R_0) with $E_0 = E_{\text{tang}}, E_{\text{const}}, E_{\text{nodiff}}$ and $R_0 = R_{\text{slow}}, R_{\text{dyn}}$. In particular, we do not treat the case $R_0 = R_{\text{fast}}$ since in this limit case the chain rule is not available and the obtained limit energy balance is a too weak formulation. However, we show in the following subsection that for λ -convex energies the EVI-formulation can be used instead.

In particular, we show in this subsection that

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0} \left\{ E_\varepsilon(\mathbf{u}_\varepsilon(t)) + \int_0^t [R_\varepsilon(\dot{\mathbf{u}}_\varepsilon) + R_\varepsilon^*(-DE_\varepsilon(\mathbf{u}_\varepsilon))] ds \right\} \\ \geq E_0(\mathbf{u}(t)) + \int_0^t [R_0(\dot{\mathbf{u}}) + R_0^*(-DE_0(\mathbf{u}))] ds. \end{aligned}$$

Here and subsequently we use the notation $\mathcal{V}_0 = \mathcal{V}_{\text{tang}}, \mathcal{V}_{\text{const}}$ and $\mathcal{W}_{\text{nodiff}}$ when we refer to the domains of the corresponding limit energy functionals $E_0 = E_{\text{tang}}$, etc. Note that the situation for E_{tang} and E_{const} is quite different than that for E_{nodiff} due to the worse compactness properties of the underlying space.

Remark 3.5. *In order to pass to the limit we use the pointwise (in time) weak convergence of the solutions in the space \mathcal{V} (resp. \mathcal{W}), i.e., $\mathbf{u}_\varepsilon(t) \rightharpoonup \mathbf{u}(t)$ in \mathcal{V} (resp. \mathcal{W}). Indeed, let $\mathcal{V}_{\text{weak}}$ denote the space \mathcal{V} endowed with the weak topology then the continuous embedding $L^\infty(0, T; \mathcal{V}) \cap H^1(0, T; \mathcal{H}) \subset C([0, T]; \mathcal{V}_{\text{weak}})$ (see e.g. [Rou05, Sect. 8.3]) implies that the weak* convergence $\mathbf{u}_\varepsilon \overset{*}{\rightharpoonup} \mathbf{u}$ in $L^\infty(0, T; \mathcal{V}) \cap H^1(0, T; \mathcal{H})$ implies $\mathbf{u}_\varepsilon(t) \rightharpoonup \mathbf{u}(t)$ in \mathcal{V} (the same holds for \mathcal{V} replaced by \mathcal{W}). This can be seen by means of a simple contradiction argument.*

Following the ideas in [SaS04] we define for a given curve $u_\varepsilon : [0, T] \rightarrow \mathcal{V}$ with $u_\varepsilon(t) \rightharpoonup u(t)$ in \mathcal{V} (resp. in \mathcal{W}) the *energy excess* $D : [0, T] \rightarrow [0, \infty]$ by

$$D_\varepsilon(t) = E_\varepsilon(u_\varepsilon(t)) - E_0(u(t)), \quad D(t) = \limsup_{\varepsilon \rightarrow 0} D_\varepsilon(t) \geq 0.$$

We call u_ε *well-prepared initially* if $D(0) = 0$.

The additional conditions for the convergence of the gradient flow given in [SaS04] can be directly translated in our case to

- 1 (Lower Bound) There exists $f \in L^1(0, T)$ such that for every $t \in [0, T]$

$$\liminf_{\varepsilon \rightarrow 0} \int_0^t R_\varepsilon(\dot{\mathbf{u}}_\varepsilon) ds \geq \int_0^t [R_0(\dot{\mathbf{u}}) - f(s)D(s)] ds. \quad (3.6)$$

2 (*Construction*) There exists a locally bounded function g on $[0, T]$ such that for any $t_0 \in]0, T[$ and any smooth curve $\hat{\mathbf{u}} :]t_0 - \delta, t_0 + \delta[\rightarrow \mathcal{V}_0$ satisfying $\hat{\mathbf{u}}(t_0) = \mathbf{u}(t_0)$ there exists a $\hat{\mathbf{u}}_\varepsilon :]t_0 - \delta, t_0 + \delta[\rightarrow \mathcal{V}$ such that $\hat{\mathbf{u}}_\varepsilon(t_0) = \mathbf{u}_\varepsilon(t_0)$ and

$$\limsup_{\varepsilon \rightarrow 0} R_\varepsilon(\hat{\mathbf{u}}_\varepsilon(t_0)) \leq R_0(\hat{\mathbf{u}}(t_0)) + g(t_0)D(t_0), \quad (3.7a)$$

$$\liminf_{\varepsilon \rightarrow 0} -\frac{d}{dt} E_\varepsilon(\hat{\mathbf{u}}_\varepsilon)|_{t=t_0} \geq -\frac{d}{dt} E_0(\hat{\mathbf{u}})|_{t=t_0} - g(t_0)D(t_0). \quad (3.7b)$$

The energy excess D should be interpreted as a small perturbation. It is shown in [SaS04] that $D \equiv 0$ holds. While the first condition in (3.6) asks for a liminf estimate for the (integrated) dissipation potential R_ε the second condition in (3.7) can be interpreted as a liminf estimate for the dual dissipation potential along the derivative of the energy functionals. Indeed, adding (3.7a) to (3.7b) we arrive at the following

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0} R_\varepsilon^*(-DE_\varepsilon(\mathbf{u}_\varepsilon)) &\geq \liminf_{\varepsilon \rightarrow 0} \left[-\langle DE_\varepsilon(\mathbf{u}_\varepsilon), \dot{\hat{\mathbf{u}}}_\varepsilon \rangle - R_\varepsilon(\dot{\hat{\mathbf{u}}}_\varepsilon) \right] \\ &\geq -\langle DE_0(\mathbf{u}), \dot{\hat{\mathbf{u}}} \rangle - R_0(\dot{\hat{\mathbf{u}}}) - 2gD. \end{aligned}$$

Taking the supremum over all $\dot{\hat{\mathbf{u}}}$ yields the limit dual dissipation potential $R_{\text{dyn}}(-DE_0(\mathbf{u}))$ at the lefthand side.

Let us point out that the limit system considered in [SaS04] is finite dimensional. Therefore, we have to adapt the results for our purpose. In particular, we have to show that the Gâteaux derivative of the limit energy functional is well-defined in \mathcal{H} .

The main result for $E_0 = E_{\text{tang}}, E_{\text{const}}$ and E_{nodiff} and $R_0 = R_{\text{dyn}}$ reads as follows:

Theorem 3.6 (Convergence of the gradient flow, Part I). *Let \mathbf{u}_ε be a family of solutions of the energy balance (EB $_\varepsilon$) converging as in (3.2)–(3.5) to a limit \mathbf{u} . If $D(0) = 0$, i.e., \mathbf{u}_ε is well prepared initially, then $D \equiv 0$ on $[0, T]$ and \mathbf{u} is the solution of the gradient flow for E_0 and R_{dyn} , i.e., it holds that*

$$E_0(\mathbf{u}(t)) + \int_0^t R_{\text{dyn}}(\dot{\mathbf{u}}) + R_{\text{dyn}}^*(-DE_0(\mathbf{u})) ds \leq E_0(\mathbf{u}(0)). \quad (3.8)$$

Proof. The weak convergence $DE_\varepsilon(\mathbf{u}_\varepsilon) \rightharpoonup \boldsymbol{\xi} = (DE_b(u), \Xi)$ in $L^2(0, T; \mathcal{H}^*)$ implies that $DE_0(\mathbf{u}) \in L^2(0, T; \mathcal{H}_0^*)$, where $\mathcal{H}_0 = \overline{\mathcal{V}_0}^{\mathcal{H}}$. Indeed, multiplying with a fixed $\hat{\mathbf{u}} \in L^2(0, T; \mathcal{V}_0 \cap \mathcal{V})$ leads to the convergence

$$\int_0^T \langle DE_\varepsilon(\mathbf{u}_\varepsilon), \hat{\mathbf{u}} \rangle dt \rightarrow \int_0^T \langle DE_0(\mathbf{u}), \hat{\mathbf{u}} \rangle dt = \int_0^T \langle \boldsymbol{\xi}, \hat{\mathbf{u}} \rangle dt.$$

Here we used the continuity properties of the associated Nemytskii operators $u \mapsto W'_b(u)$ and $U \mapsto W'_s(U)$, respectively (see [Rou05]). The density of $\mathcal{V}_0 \cap \mathcal{V}$ in \mathcal{H}_0 yields now $DE_0(\mathbf{u}) \in L^2(0, T; \mathcal{H}_0^*)$.

We see that $\widehat{\mathbf{u}} \in L^2(0, T; \mathcal{V}_0 \cap \mathcal{V})$ satisfies the conditions (3.7a) and (3.7b): We easily check that $\int_0^t R_\varepsilon(\widehat{\mathbf{u}}) ds \rightarrow \int_0^t R_{\text{dyn}}(\widehat{\mathbf{u}}) ds$ holds and conclude that

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0} \int_0^t R_\varepsilon^*(-DE_\varepsilon(\mathbf{u}_\varepsilon)) ds &\geq \liminf_{\varepsilon \rightarrow 0} \int_0^t [-\langle DE_\varepsilon(\mathbf{u}_\varepsilon), \widehat{\mathbf{u}} \rangle - R_\varepsilon(\widehat{\mathbf{u}})] ds \\ &= \int_0^t [-\langle DE_0(\mathbf{u}), \widehat{\mathbf{u}} \rangle - R_{\text{dyn}}(\widehat{\mathbf{u}})] ds. \end{aligned}$$

Taking the supremum over all $\widehat{\mathbf{u}} \in L^2(0, T; \mathcal{H}_0)$ we arrive at the liminf estimate for the dual dissipation along $DE_\varepsilon(\mathbf{u}_\varepsilon)$.

The Mosco convergence of the energy functionals and Remark 3.5 lead together with the liminf estimate for R_ε to the lower energy estimate

$$E_0(\mathbf{u}(t)) + \int_0^t R_{\text{dyn}}(\dot{\mathbf{u}}) + R_{\text{dyn}}^*(-DE_0(\mathbf{u})) ds \leq E_0(\mathbf{u}(0)),$$

which is actually an equality due to the chain rule for $t \mapsto E_0(\mathbf{u}(t))$ and the characterization of the Legendre transform. \square

The derivation of the corresponding energy balance for $R_0 = R_{\text{slow}}$ is remarkably easier.

Theorem 3.7 (Convergence of gradient flow, Part II). *Let \mathbf{u}_ε be a family of solutions of the energy balance (EB $_\varepsilon$) converging as in (3.2)–(3.5) to a limit \mathbf{u} . If $D(0) = 0$ then $D \equiv 0$ on $[0, T]$ and \mathbf{u} is the solution of the gradient flow for E_0 and R_{slow} , i.e., it holds that*

$$E_b(u(t)) + \int_0^t R_b(\dot{u}) + R_b^*(-DE_b(u)) ds = E_b(u(0)),$$

where E_b and R_b denote the bulk part of the limit energy and dissipation potential, such that $E_0(\mathbf{u}) = E_{b,0}(u) + E_{s,0}(U)$ and $R_{\text{slow}}(\dot{\mathbf{u}}) = R_b(\dot{u})$.

Proof. The prove is analogous to the proof of Theorem 3.6 with $\widehat{\mathbf{u}} = 0$. \square

Remark 3.8. *The well preparedness of the initial conditions $\mathbf{u}_\varepsilon(0)$ can be translated into asking that $E_\varepsilon(\mathbf{u}_\varepsilon(0)) \rightarrow E_0(\mathbf{u}(0))$, i.e., the initial energies converge.*

3.2 Passing to the limit in the variational inequality (EVI $_\varepsilon$)

In order to derive limit systems for the case $R_0 = R_{\text{fast}}$ we turn to the evolution variational inequality (EVI $_\varepsilon$) which is an equivalent formulation in case of λ -convex energy functionals. It reads (integrated over time)

$$\int_0^T [E_\varepsilon(\mathbf{u}_\varepsilon) + \langle G_\varepsilon \dot{\mathbf{u}}_\varepsilon, \mathbf{u}_\varepsilon - \tilde{\mathbf{u}} \rangle] dt \leq \int_0^T [E_\varepsilon(\tilde{\mathbf{u}}) - \Lambda_\varepsilon(\mathbf{u}_\varepsilon - \tilde{\mathbf{u}})] dt \quad (3.9)$$

for all $\tilde{\mathbf{u}} \in L^2(0, T; \mathcal{V})$. Note that we consider here the time-integrated version of (EVI $_\varepsilon$). This is due to the fact that we have no estimates for the time derivative of the surface variable U . Hence, we cannot argue with pointwise in time convergence of the solution.

However, working with the integrated inequality bears problems since the Γ -convergence of the time-integrated functionals is in general not trivial. We refer to [Ste08, Sal84] for the following result.

Proposition 3.9. *Let F_ε denote a sequence of weakly lower semicontinuous functionals on a reflexive and separable Banach space \mathcal{X} satisfying the liminf estimate for the weak convergence in \mathcal{X} . Moreover, let $w_\varepsilon \rightharpoonup w$ (weakly- $*$ if $p = \infty$) in $L^p(0, T; \mathcal{X})$. Then, it holds that*

$$\int_0^T F_0(w(t)) dt \leq \liminf_{\varepsilon \rightarrow 0} \int_0^T F_\varepsilon(w_\varepsilon(t)) dt.$$

The main result for the case $R_0 = R_{\text{fast}}$ reads as follows

Theorem 3.10 (Convergence of the gradient flow, Part III). *Let \mathbf{u}_ε be a family of solutions of the evolution variational inequality (3.9) converging as in (3.2)–(3.5) to the limit \mathbf{u} . Then, \mathbf{u} is the solution of the following evolution variational inequality for E_0 and R_{fast}*

$$\int_0^T E_0(\mathbf{u}) dt + \int_0^T \int_\Omega \tau_b \dot{u}(u - \tilde{u}) dx dt \leq \int_0^T [E_0(\tilde{\mathbf{u}}) - \Lambda(\mathbf{u} - \tilde{\mathbf{u}})] dt \quad (3.10)$$

for all $\tilde{\mathbf{u}} \in L^2(0, T; \mathcal{V}_0)$, where $\Lambda(\mathbf{u}) = \int_\Omega \frac{\lambda_b}{2} |u|^2 dx + \int_\Sigma \frac{\lambda_s}{2} |U|^2 d\mu$.

Proof. Let $\tilde{\mathbf{u}} = (\tilde{u}, \tilde{U}) \in L^2(0, T; \mathcal{V}_0 \cap \mathcal{V})$. It is easy to check that $\int_0^T E_\varepsilon(\tilde{\mathbf{u}}) dt \rightarrow \int_0^T E_0(\tilde{\mathbf{u}}) dt$. Moreover, from the estimates in Lemma 3.1 we infer that $\dot{u}_\varepsilon \rightharpoonup \dot{u}$ in $L^2([0, T] \times \Omega)$ and $\varepsilon^{1-\alpha} \dot{U}_\varepsilon \rightarrow 0$ in $L^2([0, T] \times \Sigma)$. Hence, we have that

$$\begin{aligned} \int_0^T \langle G_\varepsilon \dot{u}_\varepsilon, \mathbf{u}_\varepsilon - \tilde{\mathbf{u}} \rangle dt &= \\ &= \int_0^T \int_\Omega \tau_b \dot{u}_\varepsilon (u_\varepsilon - \tilde{u}) dx dt + \int_0^T \int_\Sigma \tau_s \varepsilon^{1-\alpha} \dot{U}_\varepsilon (U_\varepsilon - \tilde{U}) \frac{\mathbb{J}_\varepsilon}{\varepsilon} d\mu dt \\ &\rightarrow \int_0^T \int_\Omega \tau_b \dot{u} (u - \tilde{u}) dx dt. \end{aligned}$$

Thus, applying liminf to (3.9) and using Proposition 3.9 we obtain (3.10). \square

4 Discussion of the limit models

In this final section we show that the limit models obtained in Section 3 can be reduced to a real bulk/surface evolutionary system in $\bar{\Omega}$. The main observation is that for a pair (u, U) in $\mathcal{V}_{\text{tang}}, \mathcal{V}_{\text{const}}$ or $\mathcal{W}_{\text{nodiff}}$ we can characterize U by a function defined only on the boundary $\Gamma = \partial\Omega$. More precisely, these spaces are isomorph to the spaces $V_{\text{tang}}, V_{\text{const}}$ and W_{nodiff} given by

$$\begin{aligned} V_{\text{tang}} &:= \{(u, u_\Gamma) \in H^1(\Omega) \times H^1(\Gamma) : u|_\Gamma = u_\Gamma\}, \\ V_{\text{const}} &:= \{(u, u_\Gamma) \in H^1(\Omega) \times \mathbb{R}^{N_\Gamma} : u|_{\Gamma_i} = u_\Gamma^i, i = 1, \dots, N_\Gamma\}, \\ W_{\text{nodiff}} &:= \{(u, u_\Gamma) \in H^1(\Omega) \times L^2(\Gamma) : u|_\Gamma = u_\Gamma\} \end{aligned}$$

where $N_\Gamma \in \mathbb{N}$ is the number of connected components $\Gamma_i \subset \Gamma$. We denote by H_{tang} , H_{const} and H_{nodiff} the closures of the spaces above with respect to the L^2 -norm, such that

$$H_{\text{tang}} = H_{\text{nodiff}} = L^2(\Omega) \times L^2(\Gamma) \quad \text{and} \quad H_{\text{const}} = L^2(\Omega) \times \mathbb{R}^{N_\Gamma}.$$

With these characterizations the energy functionals E_{tang} and E_{nodiff} can be reduced by integration over the variable $\theta \in]0, 1[$ while for E_{const} we integrate over y as well. The reduced energy functionals, denoted $\mathcal{E}_{\text{tang}}$, $\mathcal{E}_{\text{const}}$ and $\mathcal{E}_{\text{nodiff}}$ are then given by

$$\begin{aligned} \mathcal{E}_{\text{tang}}(u, u_\Gamma) &:= \mathcal{E}_b(u) + \int_\Gamma \left[\frac{A_s}{2} |\nabla_\Gamma u_\Gamma|^2 + W_s(u_\Gamma) \right] d\Gamma, \\ \mathcal{E}_{\text{const}}(u, u_\Gamma) &:= \mathcal{E}_b(u) + |\Gamma| \sum_{i=1}^{N_\Gamma} W_s(u_\Gamma^i), \\ \mathcal{E}_{\text{nodiff}}(u, u_\Gamma) &:= \mathcal{E}_b(u) + \frac{\omega_s}{2} \|u_\Gamma\|_{L^2(\Gamma)}^2, \end{aligned}$$

where in each case $\mathcal{E}_b(u) = \int_\Omega \left[\frac{A_b}{2} |\nabla u|^2 + W_b(u) \right] dx$ denotes the bulk energy.

Starting with the case $\alpha = 1$ we see that the limit energy balance in (3.8) can be written in terms of $\mathcal{E}_0 \in \{\mathcal{E}_{\text{tang}}, \mathcal{E}_{\text{const}}, \mathcal{E}_{\text{nodiff}}\}$ and the dissipation potential \mathcal{R}_{dyn} . Here, in slight abuse of notation, \mathcal{R}_{dyn} is for each of the energy functionals $\mathcal{E}_{\text{tang}}$, $\mathcal{E}_{\text{const}}$ and $\mathcal{E}_{\text{nodiff}}$ defined on the spaces H_{tang} , H_{const} and H_{nodiff} and obtained as before via integration with respect to the variable θ or (y, θ) , respectively. Thus, the reduced energy balance reads

$$\mathcal{E}_0(u(t), u_\Gamma(t)) + \int_0^t \mathcal{R}_{\text{dyn}}(\dot{u}, \dot{u}_\Gamma) + \mathcal{R}_{\text{dyn}}^*(-D\mathcal{E}_0(u, u_\Gamma)) ds = \mathcal{E}_0(u(0), u_\Gamma(0)).$$

To highlight the structure of the limit systems we now write down the corresponding force balance equation written in terms of the reduced energy and dissipation functional. It consists of two equations for the bulk and the surface variable u and $u_\Gamma = u|_\Gamma$, respectively. Using the chain rule and the Fenchel equivalences we obtain

$$\begin{pmatrix} \tau_b \dot{u} + D_u \mathcal{E}_0(u, u_\Gamma) \\ \tau_s \dot{u}_\Gamma + D_{u_\Gamma} \mathcal{E}_0(u, u_\Gamma) \end{pmatrix} = 0.$$

For each of the energy functionals the first equation is formally equivalent to the well-known Allen–Cahn equation in $[0, T] \times \Omega$

$$\tau_b \partial_t u - A_b \Delta u + W_b'(u) = 0. \quad (\text{AC}_{\text{bulk}})$$

This equation is coupled to the boundary evolution of $u|_\Gamma = u_\Gamma$, which for the energy functional $\mathcal{E}_{\text{tang}}$ is described by

$$\tau_s \partial_t u_\Gamma - A_s \Delta_\Gamma u_\Gamma + A_b \nabla u \cdot \nu + W_s'(u_\Gamma) = 0. \quad (4.1)$$

Hence, we obtain the surface Allen–Cahn equation with a contribution given by the conormal derivative of the bulk variable u . The system $(\text{AC}_{\text{bulk}})$ & (4.1) was studied in [SpW10].




Figure 2: Numerical examples for Allen–Cahn equation (AC_{bulk}) in circular domain with dynamic boundary condition (4.1) (left) and homogeneous Neumann boundary condition (right) for double well potential $W_b(u) = W_s(u) = (1 - u^2)^2$ and $A_b = 1000 \cdot A_s$ and $\tau_b = \tau_s$.

For the energy functional $\mathcal{E}_{\text{const}}$ we obtain a simpler boundary condition, which consists of a system of ordinary differential equations for each of the connected components Γ_i of the boundary Γ , namely

$$\tau_s \partial_t u_\Gamma^i + A_b [\nabla u \cdot \nu]_i + W_s'(u_\Gamma^i) = 0, \quad (4.2)$$

where $[g]_i := \frac{1}{|\Gamma_i|} \int_{\Gamma_i} g \, d\Gamma$ denotes the mean value of $g : \Gamma_i \rightarrow \mathbb{R}$ over $\Gamma_i \subset \Gamma$.

Finally, for $\mathcal{E}_0 = \mathcal{E}_{\text{nodiff}}$ the boundary condition reads

$$\tau_s \partial_t u_\Gamma + A_b \nabla u \cdot \nu + \omega_s u_\Gamma = 0. \quad (4.3)$$

This boundary condition can be found as a special case in [Pet04].

In the case $\alpha < 1$ ($R_0 = R_{\text{slow}}$) we obtain the bulk Allen–Cahn equation (AC_{bulk}) and have no evolution on the boundary, i.e. $\dot{u}_\Gamma = 0$. Which means that the boundary values are fixed by the initial conditions. Since we assumed in the convergence analysis that the initial energies converge, the initial values $(u(0), u_\Gamma(0))$ have to lie in V_{tang} , V_{const} and V_{nodiff} , respectively. In particular, in the first case we have $u|_\Gamma = u|_\Gamma(0) \in H^1(\Gamma)$, while in the second case the boundary values are constant (on each connected component) and in the last case we have $u|_\Gamma = u(0)|_\Gamma \in H^{\frac{1}{2}}(\Gamma)$.

At last we discuss the fast evolution case $\alpha > 1$ ($R_0 = R_{\text{fast}}$). Choosing $\tilde{\mathbf{u}} = \mathbf{u} - h\mathbf{w}$, $h > 0$ in the limit evolution variational inequality (3.10) and letting $h \rightarrow 0$ we obtain the system

$$\begin{pmatrix} \tau_b \dot{u} + D_u \mathcal{E}_0(u, u_\Gamma) \\ D_{u_\Gamma} \mathcal{E}_0(u, u_\Gamma) \end{pmatrix} = 0.$$

Hence, for $\mathcal{E}_0 = \mathcal{E}_{\text{tang}}$ the bulk equation (AC_{bulk}) is coupled to the nonlinear elliptic surface equation

$$-A_s \Delta_\Gamma u_\Gamma + A_b \nabla u \cdot \nu + W_s'(u_\Gamma) = 0. \quad (4.4)$$

While for $\mathcal{E}_0 = \mathcal{E}_{\text{const}}$ we have the following nonlinear equation for each connected component of the boundary Γ

$$A_b [\nabla u \cdot \nu]_i + W_s'(u_\Gamma^i) = 0. \quad (4.5)$$

In the last case $\mathcal{E}_0 = \mathcal{E}_{\text{nodiff}}$ we obtain the usual Robin boundary condition

$$A_b \nabla u \cdot \nu + \omega_s u_\Gamma = 0. \quad (4.6)$$

See Figure 2 for a numerical comparison of dynamic boundary condition and classical Neumann boundary condition in case of a circular domain.

Acknowledgment

The author thanks Alexander Mielke and Annegret Glitzky for stimulating and helpful discussions.

References

- [AGS05] L. AMBROSIO, N. GIGLI, and G. SAVARÉ. *Gradient flows in metric spaces and in the space of probability measures*. Lectures in Mathematics, ETH Zürich. Basel: Birkhäuser., 2005.
- [AM*11] S. ARNRICH, A. MIELKE, M. A. PELETIER, G. SAVARÉ, and M. VENERONI. Passing to the limit in a Wasserstein gradient flow: From diffusion to reaction. *WIAS Preprint*, 1593, 2011.
- [Att84] H. ATTOUCH. *Variational convergence for functions and operators*. Applicable Mathematics Series. Pitman (Advanced Publishing Program), Boston, MA, 1984.
- [BFG06] G. BELLETTINI, G. FUSCO, and N. GUGLIELMI. A concept of solution and numerical experiments for forward-backward diffusion equations. *Discrete Contin. Dyn. Syst.*, 16(4), 783–842, 2006.
- [Bra02] A. BRAIDES. Γ -convergence for beginners. Oxford Lecture Series in Mathematics and its Applications 22. Oxford: Oxford University Press. xii, 2002.
- [CFP06] R. CHILL, E. FAŠANGOVÁ, and J. PRÜSS. Convergence to steady states of solutions of the Cahn-Hilliard and Caginalp equations with dynamic boundary conditions. *Math. Nachr.*, 279(13-14), 1448–1462, 2006.
- [CGM08] L. CHERFILS, S. GATTI, and A. MIRANVILLE. Existence of global solutions to the caginalp phase-field system with dynamic boundary conditions and singular potentials. *Journal of Mathematical Analysis and Applications*, 343(1), 557 – 566, 2008.
- [Cia00] P. G. CIARLET. *Mathematical elasticity. Vol. III*, volume 29 of *Studies in Mathematics and its Applications*. North-Holland Publishing Co., Amsterdam, 2000. Theory of shells.
- [CoR90] P. COLLI and J.-F. RODRIGUES. Diffusion through thin layers with high specific heat. *Asymptotic Anal.*, 3(3), 249–263, 1990.
- [Dal93] G. DAL MASO. *An Introduction to Γ -Convergence*. Birkhäuser Boston Inc., Boston, MA, 1993.
- [DaS10] S. DANERI and G. SAVARÉ. Lecture notes on gradient flows and optimal transport. *Seminaires et Congres, SMF*, 2010. to appear.
- [dCa76] M. P. A. DO CARMO. *Differential geometry of curves and surfaces*. Englewood Cliffs, N. J.: Prentice-Hall, 1976.
- [EIS10] C. M. ELLIOTT and B. STINNER. Modeling and computation of two phase geometric biomembranes using surface finite elements. *J. Comput. Phys.*, 229(18), 6585–6612, 2010.
- [FJM06] G. FRIESECKE, R. D. JAMES, and S. MÜLLER. A hierarchy of plate models derived from nonlinear elasticity by Gamma-convergence. *Arch. Ration. Mech. Anal.*, 180(2), 183–236, 2006.
- [FRG*06] A. FAVINI, G. RUIZ GOLDSTEIN, J. A. GOLDSTEIN, and S. ROMANELLI. The heat equation with nonlinear general Wentzell boundary condition. *Adv. Differ. Equ.*, 11(5), 481–510, 2006.

- [GGM08] C. GAL, M. GRASSELLI, and A. MIRANVILLE. Nonisothermal Allen-Cahn equations with coupled dynamic boundary conditions. Colli, P. (ed.) et al., Proceedings of international conference on: Nonlinear phenomena with energy dissipation. Mathematical analysis, modeling and simulation, Chiba, Japan, November 26–30, 2007. Tokyo: Gakkotosha. Gakuto International Series Mathematical Sciences and Applications 29, 117-139 (2008)., 2008.
- [Gli11] A. GLITZKY. An electronic model for solar cells including active interfaces and energy resolved defect densities. *WIAS Preprint*, 1663, 2011.
- [KE*01] R. KENZLER, F. EURICH, P. MAASS, B. RINN, J. SCHROPP, E. BOHL, and W. DIETERICH. Phase separation in confined geometries: Solving the Cahn-Hilliard equation with generic boundary conditions. *Computer Physics Communications*, 133(2-3), 139 – 157, 2001.
- [KMM06] M. KURZKE, C. MELCHER, and R. MOSER. Domain walls and vortices in thin ferromagnetic films. In *Analysis, modeling and simulation of multiscale problems*, pages 249–298. Springer, Berlin, 2006.
- [Kra95] G. KRAUSCH. Surface induced self assembly in thin polymer films. *Materials Science and Engineering: R: Reports*, 14(1-2), v – 94, 1995.
- [Kur07] M. KURZKE. The gradient flow motion of boundary vortices. *Ann. Inst. Henri Poincaré, Anal. Non Linéaire*, 24(1), 91–112, 2007.
- [Mie08] A. MIELKE. Weak-convergence methods for Hamiltonian multiscale problems. *Discrete Contin. Dyn. Syst.*, 20(1), 53–79, 2008.
- [Mie11] A. MIELKE. A gradient structure for reaction-diffusion systems and for energy-drift-diffusion systems. *Nonlinearity*, 24, 1329–1346, 2011.
- [MiZ05] A. MIRANVILLE and S. ZELIK. Exponential attractors for the Cahn-Hilliard equation with dynamic boundary conditions. *Mathematical Methods in the Applied Sciences*, 28(6), 709–735, 2005.
- [MRS08] A. MIELKE, T. ROUBÍČEK, and U. STEFANELLI. Γ -limits and relaxations for rate-independent evolutionary problems. *Calc. Var. Partial Differ. Equ.*, 31(3), 387–416, 2008.
- [Pet04] J. PETERSSON. A note on quenching for parabolic equations with dynamic boundary conditions. *Nonlinear Anal.*, 58(3-4), 417–423, 2004.
- [PoA06] J. POORTMANS and V. ARKHIPOV. *Thin film solar cells: fabrication, characterization and applications*. Wiley series in materials for electronic and optoelectronic applications. Wiley, 2006.
- [PuF97] S. PURI and H. L. FRISCH. Surface-directed spinodal decomposition: modelling and numerical simulations. *Journal of Physics: Condensed Matter*, 9(10), 2109, 1997.
- [RaZ01] R. RACKE and S. ZHENG. The Cahn-Hilliard equation with dynamic boundary conditions. *Advances Di. Equations*, 8, 8–83, 2001.
- [Rou05] T. ROUBÍČEK. *Nonlinear partial differential equations with applications*. ISNM. International Series of Numerical Mathematics 153. Basel: Birkhäuser, 2005.

- [Sal84] A. SALVADORI. On the M-convergence for integral functionals on L^p_X . *Atti Sem. Mat. Fis. Univ. Modena*, 33, 137–154, 1984.
- [SaS04] E. SANDIER and S. SERFATY. Gamma-convergence of gradient flows with applications to Ginzburg-Landau. *Commun. Pure Appl. Math.*, 57(12), 1627–1672, 2004.
- [SaV97] G. SAVARÉ and A. VISINTÍN. Variational convergence of nonlinear diffusion equations: Applications to concentrated capacity problems with change of phase. *Atti Accad. Naz. Lincei, Cl. Sci. Fis. Mat. Nat., IX. Ser., Rend. Lincei, Mat. Appl.*, 8(1), 49–89, 1997.
- [Sch94] D. SCHROEDER. *Modelling of interface carrier transport for device simulation*. Computational microelectronics. Springer-Verlag, 1994.
- [ScT10] K. SCHMIDT and S. TORDEUX. Asymptotic modelling of conductive thin sheets. *Z. Angew. Math. Phys.*, 61(4), 603–626, 2010.
- [SpW10] J. SPREKELS and H. WU. A note on parabolic equation with nonlinear dynamical boundary condition. *Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods*, 72(6), 3028–3048, 2010.
- [Ste08] U. STEFANELLI. The Brezis – Ekeland principle for doubly nonlinear equations. *SIAM J. Control Optim.*, 47(3), 1615–1642, 2008.
- [Tai09] K. TAIRA. *Boundary value problems and Markov processes. 2nd ed.* Lecture Notes in Mathematics 1499. Berlin: Springer. xii, 186 p., 2009.
- [Wlo87] J. WLOKA. *Partial differential equations. Transl. from the German by C. B. and M. J. Thomas*. Cambridge etc.: Cambridge University Press. XI, 1987.