

Weierstraß-Institut
für Angewandte Analysis und Stochastik
Leibniz-Institut im Forschungsverbund Berlin e. V.

Preprint

ISSN 2198-5855

**Optimal boundary control of a viscous
Cahn–Hilliard system with dynamic boundary condition
and double obstacle potentials**

Pierluigi Colli¹, M. Hassan Farshbaf-Shaker², Gianni Gilardi¹,
Jürgen Sprekels^{2,3}

submitted: September 1, 2014

¹ Dipartimento di Matematica “F. Casorati”
Università di Pavia
via Ferrata, 1
27100 Pavia, Italy
E-Mail: pierluigi.colli@unipv.it,
E-Mail: gianni.gilardi@unipv.it

² Weierstrass Institute
Mohrenstr. 39
10117 Berlin
Germany
E-Mail: hassan.farshbaf-shaker@wias-berlin.de
E-Mail: juergen.sprekels@wias-berlin.de

³ Department of Mathematics
Humboldt-Universität zu Berlin
Unter den Linden 6
10099 Berlin, Germany

No. 2006
Berlin 2014



2010 *Mathematics Subject Classification.* 74M15, 49K20, 35K61.

Key words and phrases. Optimal control, parabolic obstacle problems, MPECs, dynamic boundary conditions, optimality conditions.

This paper was initiated during a visit of JS to the Università di Pavia. The kind hospitality and stimulating atmosphere of the Università di Pavia are gratefully acknowledged. Some financial support comes from the MIUR-PRIN Grant 2010A2TFX2 “Calculus of Variations”.

Edited by
Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS)
Leibniz-Institut im Forschungsverbund Berlin e. V.
Mohrenstraße 39
10117 Berlin
Germany

Fax: +49 30 20372-303
E-Mail: preprint@wias-berlin.de
World Wide Web: <http://www.wias-berlin.de/>

Abstract

In this paper, we investigate optimal boundary control problems for Cahn–Hilliard variational inequalities with a dynamic boundary condition involving double obstacle potentials and the Laplace–Beltrami operator. The cost functional is of standard tracking type, and box constraints for the controls are prescribed. We prove existence of optimal controls and derive first-order necessary conditions of optimality. The general strategy, which follows the lines of the recent approach by Colli, Farshbaf-Shaker, Sprekels (see Appl. Math. Optim., 2014) to the (simpler) Allen–Cahn case, is the following: we use the results that were recently established by Colli, Gilardi, Sprekels in the preprint arXiv:1407.3916 [math.AP] for the case of (differentiable) logarithmic potentials and perform a so-called “deep quench limit”. Using compactness and monotonicity arguments, it is shown that this strategy leads to the desired first-order necessary optimality conditions for the case of (non-differentiable) double obstacle potentials.

1 Introduction

Let $\Omega \subset \mathbb{R}^N$, $2 \leq N \leq 3$, denote some open, connected and bounded domain with smooth boundary Γ and outward unit normal field \mathbf{n} , and let $T > 0$ be a fixed final time. Putting $Q := \Omega \times (0, T)$, $\Sigma := \Gamma \times (0, T)$, we introduce the function spaces

$$\begin{aligned} H &:= L^2(\Omega), \quad V := H^1(\Omega), \quad H_\Gamma := L^2(\Gamma), \quad V_\Gamma := H^1(\Gamma), \\ \mathcal{H} &:= H \times H_\Gamma, \quad \mathcal{V} := \{(y, y_\Gamma) : y \in V \times V_\Gamma : v|_\Gamma = v_\Gamma\}, \end{aligned} \quad (1.1)$$

which are Hilbert spaces when endowed with the topologies induced by their respective natural inner products, denoted by $(\cdot, \cdot)_E$ for $E \in \{H, H_\Gamma, V, V_\Gamma, \mathcal{H}, \mathcal{V}\}$. In the following, we denote the norm in the generic Banach space E by $\|\cdot\|_E$, with the one exception that for convenience the norm of the space H^N will also be denoted by $\|\cdot\|_H$. Moreover, let E^* indicate the dual space of E and let $\langle \cdot, \cdot \rangle_E$ always stand for the duality pairing between elements of E^* and elements of E . It is understood that H is embedded in V^* in the usual way, namely, such that $\langle u, v \rangle_V = (u, v)_H$ for all $u \in H$ and $v \in V$; we then obtain the Hilbert triplet $V \subset H \subset V^*$ with dense and compact embeddings. In the same way, we construct the Hilbert triplets $V_\Gamma \subset H_\Gamma \subset V_\Gamma^*$ and $\mathcal{V} \subset \mathcal{H} \subset \mathcal{V}^*$, with dense and compact embeddings.

Throughout this paper, we generally assume:

(A1) There are given constants $\beta_i \geq 0$, $1 \leq i \leq 5$, which do not all vanish, as well as functions

$$z_Q \in L^2(Q), \quad z_\Sigma \in L^2(\Sigma), \quad z_\Omega \in L^2(\Omega), \quad z_\Gamma \in L^2(\Gamma), \quad \text{and} \\ \tilde{u}_{1\Gamma}, \tilde{u}_{2\Gamma} \in L^\infty(\Sigma) \quad \text{with} \quad \tilde{u}_{1\Gamma} \leq \tilde{u}_{2\Gamma} \quad \text{a. e. on } \Sigma.$$

We then introduce the tracking type cost functional

$$\mathcal{J}((y, y_\Gamma), u_\Gamma) := \frac{\beta_1}{2} \|y - z_Q\|_{L^2(Q)}^2 + \frac{\beta_2}{2} \|y_\Gamma - z_\Sigma\|_{L^2(\Sigma)}^2 \\ + \frac{\beta_3}{2} \|y(T) - z_\Omega\|_{L^2(\Omega)}^2 + \frac{\beta_4}{2} \|y_\Gamma(T) - z_\Gamma\|_{L^2(\Gamma)}^2 + \frac{\beta_5}{2} \|u_\Gamma\|_{L^2(\Sigma)}^2, \quad (1.2)$$

which is meaningful for, e. g., $(y, y_\Gamma) \in \mathcal{V}$ and $u_\Gamma \in H_\Gamma$, and, for $\tau > 0$, the viscous Cahn–Hilliard system with dynamic boundary conditions

$$\partial_t y - \Delta w = 0 \quad \text{in } Q, \quad (1.3)$$

$$w = \tau \partial_t y - \Delta y + \xi + f'_2(y) \quad \text{in } Q, \quad (1.4)$$

$$y|_\Gamma = y_\Gamma, \quad \partial_{\mathbf{n}} y + \partial_t y_\Gamma - \Delta_\Gamma y_\Gamma + \xi_\Gamma + g'_2(y_\Gamma) = u_\Gamma, \quad \partial_{\mathbf{n}} w = 0, \quad \text{on } \Sigma, \quad (1.5)$$

$$\xi \in \partial I_{[-1,1]}(y) \quad \text{a. e. in } Q, \quad \xi_\Gamma \in \partial I_{[-1,1]}(y_\Gamma) \quad \text{a. e. on } \Sigma, \quad (1.6)$$

$$y(\cdot, 0) = y_0 \quad \text{a. e. in } \Omega, \quad y_\Gamma(\cdot, 0) = y_{0\Gamma} \quad \text{a. e. on } \Gamma. \quad (1.7)$$

Moreover, let $M_0 > 0$ denote some given constant, and let

$$\mathcal{U}_{\text{ad}} := \{u_\Gamma \in H^1(0, T; H_\Gamma) \cap L^\infty(\Sigma) : \|\partial_t u_\Gamma\|_{L^2(\Sigma)} \leq M_0, \\ \tilde{u}_{1\Gamma} \leq u_\Gamma \leq \tilde{u}_{2\Gamma} \quad \text{a. e. in } \Sigma \}, \quad (1.8)$$

be the set of admissible controls which is assumed nonempty throughout this paper. Our overall boundary control problem reads as follows:

$$(\mathcal{P}_0) \quad \text{Minimize } \mathcal{J}((y, y_\Gamma), u_\Gamma) \quad \text{subject to the state constraints (1.3)–(1.7)} \\ \text{and to the control constraint } u_\Gamma \in \mathcal{U}_{\text{ad}}.$$

In (1.7), y_0 and $y_{0\Gamma}$ are given initial data with $y_{0|_\Gamma} = y_{0\Gamma}$, where the trace $y|_\Gamma$ (if it exists) of a function y on Γ will throughout be denoted by y_Γ without further comment. Moreover, in the following $\partial_{\mathbf{n}}$, ∇_Γ and Δ_Γ will always stay for the outward normal derivative, the tangential gradient, and the Laplace–Beltrami operator, respectively, on Γ ; in addition, f_2 , g_2 are given smooth nonlinearities, while u_Γ is a boundary control. Since we will confine ourselves to the viscous case $\tau > 0$, we will henceforth assume without loss of generality that $\tau = 1$.

The system (1.3)–(1.7) is an initial-boundary value problem with nonlinear dynamic boundary condition for a Cahn–Hilliard differential inclusion, which (cf. Proposition 2.2 below) under appropriate conditions on the data admits for every $u_\Gamma \in \mathcal{U}_{\text{ad}}$ a solution quintuple $(y, y_\Gamma, w, \xi, \xi_\Gamma)$,

where the solution components $(y, y_\Gamma, \xi_\Gamma)$ are uniquely determined. Hence, the *control-to-state operator* $\mathcal{S}_0 : u_\Gamma \mapsto \mathcal{S}_0(u_\Gamma) := (y, y_\Gamma)$ is well defined on \mathcal{U}_{ad} , and the control problem (\mathcal{P}_0) is equivalent to minimizing the reduced cost functional

$$\mathcal{J}_{\text{red}}(u_\Gamma) := \mathcal{J}(\mathcal{S}_0(u_\Gamma), u_\Gamma) \quad (1.9)$$

over \mathcal{U}_{ad} .

In the physical interpretation, the unknown y usually stands for the (conserved) order parameter of an isothermal phase transition, typically a rescaled fraction of one of the involved phases. In such a situation, it is physically meaningful to require y to attain values in the interval $[-1, 1]$ on both Ω and Γ . A standard technique to meet this requirement is to use the indicator function of the interval $[-1, 1]$,

$$I_{[-1,1]}(y) = \begin{cases} 0 & \text{if } y \in [-1, 1] \\ +\infty & \text{otherwise} \end{cases},$$

so that the non-diffusive parts of the local specific bulk and surface free energies, $F_{\text{bulk}} := I_{[-1,1]} + f_2$ and $F_{\text{surface}} := I_{[-1,1]} + g_2$, are of *double obstacle type*, and the subdifferential $\partial I_{[-1,1]}$, defined by

$$\eta \in \partial I_{[-1,1]}(v) \quad \text{if and only if} \quad \eta \begin{cases} \leq 0 & \text{if } v = -1 \\ = 0 & \text{if } -1 < v < 1 \\ \geq 0 & \text{if } v = 1 \end{cases},$$

is employed in place of the usual derivative. Concerning the selections ξ, ξ_Γ in (1.6), one has to keep in mind that ξ may be not regular enough as to single out its trace on the boundary Γ , and if the trace $\xi|_\Gamma$ exists, it may differ from ξ_Γ , in general.

The optimization problem (\mathcal{P}_0) belongs to the problem class of so-called MPECs (Mathematical Programs with Equilibrium Constraints). It is a well-known fact that the differential inclusion conditions encoded in (1.3)–(1.6), which occur as constraints in (\mathcal{P}_0) , violate all of the known classical nonlinear programming constraint qualifications. Hence, the existence of Lagrange multipliers cannot be inferred from standard theory, and the derivation of first-order necessary condition becomes very difficult.

While numerous papers deal with the well-posedness and asymptotic behavior of Cahn–Hilliard system (cf., e. g., the references given in [13, 14, 7]), there are comparatively few investigations of associated optimal control problems. Usually, these papers treat the non-viscous case $\tau = 0$ and are restricted to differentiable free energies and to the case of distributed controls, with the no-flux condition $(\partial_n y)|_\Gamma = 0$ assumed in place of the more difficult dynamic boundary condition (1.5). In this connection, we refer to [21] and [15], where the latter paper also deals with the case of double obstacle potentials.

Quite recently, also convective Cahn–Hilliard systems have been investigated from the viewpoint of optimal control. In this connection, we refer to [22] and [23], where the latter paper deals with the two-dimensional case. The three-dimensional case with a nonlocal free energy was studied in [19]. There also exist contributions dealing with the more general and difficult

Cahn–Hilliard/Navier–Stokes systems, cf. [17] and [16]. Finally, we mention the papers [5] and [6], in which control problems for a generalized Cahn–Hilliard system introduced in [18] were investigated.

The only existing contribution to the optimal control of viscous or non-viscous Cahn–Hilliard systems with dynamic boundary conditions of the form (1.5) seems to be the recent paper [8] in which three of the present authors investigated the case of differentiable bulk and surface free energies that may have singular derivatives. A typical case to which the analysis in [8] applies is given by the logarithmic form

$$\begin{aligned} F_{\log}(y) &= h(y) + f_2(y), \quad \text{where} \\ h(y) &= \widehat{c}((1+y) \ln(1+y) + (1-y) \ln(1-y)), \quad -1 < y < 1, \end{aligned} \quad (1.10)$$

with some fixed constant $\widehat{c} > 0$. Note that in this case the inclusions (1.6) have to be replaced by the equations $\xi = h'(y)$ and $\xi_\Gamma = h'(y_\Gamma)$, respectively.

In this paper, we aim to employ the results established in [8] to treat the non-differentiable double obstacle case when ξ, ξ_Γ satisfy the inclusions (1.6). Our approach is guided by the strategy used by three of the present authors in their recent paper [9] for a corresponding optimal control problem for the simpler Allen–Cahn equation: in [9], necessary optimality conditions for the double obstacle case could be established by performing a so-called “deep quench limit” in a family of optimal control problems with differentiable nonlinearities of a form that had been previously treated in [10] and for which the corresponding systems had been analyzed in [4].

The general idea is briefly explained as follows: we replace the inclusions (1.6) by

$$\xi = \varphi(\alpha) h'(y), \quad \xi_\Gamma = \psi(\alpha) h'(y), \quad (1.11)$$

where h is defined in (1.10), and where φ, ψ are continuous and positive functions on $(0, 1]$ that satisfy

$$\lim_{\alpha \searrow 0} \varphi(\alpha) = \lim_{\alpha \searrow 0} \psi(\alpha) = 0, \quad \varphi(\alpha) \leq C_{\varphi\psi} \psi(\alpha) \quad \forall \alpha > 0, \quad \text{with some } C_{\varphi\psi} > 0. \quad (1.12)$$

We remark that we could simply choose $\varphi(\alpha) = \psi(\alpha) = \alpha^p$ for some $p > 0$; however, there might be situations (e. g., in the numerical approximation) in which it is advantageous to let φ and ψ have a different behavior as $\alpha \searrow 0$.

Now observe that $h'(y) = \ln\left(\frac{1+y}{1-y}\right)$ and $h''(y) = \frac{2}{1-y^2} > 0$ for $y \in (-1, 1)$. Hence, in particular, we have

$$\begin{aligned} \lim_{\alpha \searrow 0} \varphi(\alpha) h'(y) &= 0 \quad \text{for } -1 < y < 1, \\ \lim_{\alpha \searrow 0} \left(\varphi(\alpha) \lim_{y \searrow -1} h'(y) \right) &= -\infty, \quad \lim_{\alpha \searrow 0} \left(\varphi(\alpha) \lim_{y \nearrow +1} h'(y) \right) = +\infty. \end{aligned} \quad (1.13)$$

Since similar relations hold if φ is replaced by ψ , we may regard the graphs of the functions $\varphi(\alpha) h'$ and $\psi(\alpha) h'$ as approximations to the graph of the subdifferential $\partial I_{[-1,1]}$.

Now, for any $\alpha > 0$ the optimal control problem (later to be denoted by (\mathcal{P}_α)), which results if in (\mathcal{P}_0) the relation (1.6) is replaced by (1.11), is of the type for which in [8] the existence of optimal controls $u_\Gamma^\alpha \in \mathcal{U}_{\text{ad}}$ as well as first-order necessary optimality conditions have been derived. Proving a priori estimates (uniform in $\alpha > 0$), and employing compactness and monotonicity arguments, we will be able to show the following existence and approximation result: whenever $\{u_\Gamma^{\alpha_n}\} \subset \mathcal{U}_{\text{ad}}$ is a sequence of optimal controls for (\mathcal{P}_{α_n}) , where $\alpha_n \searrow 0$ as $n \rightarrow \infty$, then there exist a subsequence of $\{\alpha_n\}$, which is again indexed by n , and an optimal control $\bar{u}_\Gamma \in \mathcal{U}_{\text{ad}}$ of (\mathcal{P}_0) such that

$$u_\Gamma^{\alpha_n} \rightarrow \bar{u}_\Gamma \quad \text{weakly-star in } \mathcal{X} \text{ as } n \rightarrow \infty, \quad (1.14)$$

where here and in the following

$$\mathcal{X} := H^1(0, T; H_\Gamma) \cap L^\infty(\Sigma) \quad (1.15)$$

will always denote the control space. In other words, optimal controls for (\mathcal{P}_α) are for small $\alpha > 0$ likely to be ‘close’ to optimal controls for (\mathcal{P}_0) . It is natural to ask if the reverse holds, i. e., whether every optimal control for (\mathcal{P}_0) can be approximated by a sequence $\{u_\Gamma^{\alpha_n}\}$ of optimal controls for (\mathcal{P}_{α_n}) , for some sequence $\alpha_n \searrow 0$.

Unfortunately, we will not be able to prove such a ‘global’ result that applies to all optimal controls for (\mathcal{P}_0) . However, a ‘local’ result can be established. To this end, let $\bar{u}_\Gamma \in \mathcal{U}_{\text{ad}}$ be any optimal control for (\mathcal{P}_0) . We introduce the ‘adapted’ cost functional

$$\tilde{\mathcal{J}}((y, y_\Gamma), u_\Gamma) := \mathcal{J}((y, y_\Gamma), u_\Gamma) + \frac{1}{2} \|u_\Gamma - \bar{u}_\Gamma\|_{L^2(\Sigma)}^2 \quad (1.16)$$

and consider for every $\alpha \in (0, 1]$ the *adapted control problem* of minimizing $\tilde{\mathcal{J}}$ subject to $u_\Gamma \in \mathcal{U}_{\text{ad}}$ and to the constraint that (y, y_Γ) solves the approximating system (1.3)–(1.5), (1.7), (1.11). It will then turn out that the following is true:

(i) There are some sequence $\alpha_n \searrow 0$ and minimizers $\bar{u}_\Gamma^{\alpha_n} \in \mathcal{U}_{\text{ad}}$ of the adapted control problem associated with α_n , $n \in \mathbb{N}$, such that

$$\bar{u}_\Gamma^{\alpha_n} \rightarrow \bar{u}_\Gamma \quad \text{strongly in } L^2(\Sigma) \text{ as } n \rightarrow \infty. \quad (1.17)$$

(ii) It is possible to pass to the limit as $\alpha \searrow 0$ in the first-order necessary optimality conditions corresponding to the adapted control problems associated with $\alpha \in (0, 1]$ in order to derive first-order necessary optimality conditions for problem (\mathcal{P}_0) .

The paper is organized as follows: in Section 2, we give a precise statement of the problem under investigation, and we derive some results concerning the state system (1.3)–(1.7) and its α -approximation which is obtained if in (\mathcal{P}_0) the relations (1.6) are replaced by the relations (1.11). In Section 3, we then prove the existence of optimal controls and the approximation result formulated above in (i). The final Section 4 is devoted to the derivation of the first-order necessary optimality conditions, where the strategy outlined in (ii) is employed.

During the course of this analysis, we will make repeated use of the elementary Young's inequality

$$ab \leq \gamma|a|^2 + \frac{1}{4\gamma}|b|^2 \quad \forall a, b \in \mathbb{R} \quad \forall \gamma > 0,$$

and we will use the following notation: for functions $v \in V^*$ and $w \in L^1(0, T; V^*)$ we define their generalized mean values as

$$v^\Omega := \frac{1}{|\Omega|} \langle v, 1 \rangle_V, \quad \text{and} \quad w^\Omega(t) := (w(t))^\Omega \quad \text{for a.e. } t \in (0, T). \quad (1.18)$$

Clearly, (1.18) gives the usual mean values when elements of H or of $L^1(0, T; H)$, respectively, are involved. We also recall Poincaré's inequality

$$\|v\|_V \leq C_P (\|\nabla v\|_H + |v^\Omega|) \quad \forall v \in V, \quad (1.19)$$

with a constant $C_P > 0$ that only depends on Ω .

2 General assumptions and state equations

In this section, we formulate the general assumptions of the paper, and we state some preparatory results for the state system (1.3)–(1.7) and its α -approximations.

We make the following general assumptions:

(A2) $f_2, g_2 \in C^3([-1, 1])$.

(A3) $y_0 \in H^2(\Omega)$, $y_{0\Gamma} := y_0|_\Gamma \in H^2(\Gamma)$, $\partial_n y_0|_\Gamma = 0$, and we have

$$-1 < y_0(x) < 1 \quad \forall x \in \bar{\Omega}. \quad (2.1)$$

(A4) There exist $\xi_0 \in H$ and $\xi_{\Gamma,0} \in H_\Gamma$ such that

$$\xi_0 \in I_{[-1,1]}(y_0) \quad \text{a.e. in } \Omega, \quad \xi_{\Gamma,0} \in I_{[-1,1]}(y_{0\Gamma}) \quad \text{a.e. on } \Gamma. \quad (2.2)$$

Now observe that the set \mathcal{U}_{ad} is a bounded subset of \mathcal{X} . Hence, there exists a bounded open ball in \mathcal{X} that contains \mathcal{U}_{ad} . For later use it is convenient to fix such a ball once and for all, noting that any other such ball could be used instead. In this sense, the following assumption is rather a denotation:

(A5) \mathcal{U} is a nonempty open and bounded subset of \mathcal{X} containing \mathcal{U}_{ad} , and the constant $R > 0$ satisfies

$$\|u_\Gamma\|_{H^1(0,T;H_\Gamma)} + \|u_\Gamma\|_{L^\infty(\Sigma)} \leq R \quad \forall u_\Gamma \in \mathcal{U}. \quad (2.3)$$

Next, we introduce our notion of solution to the problem (1.3)–(1.7) in the abstract setting introduced above.

Definition 2.1: A quintuple $(y, y_\Gamma, w, \xi, \xi_\Gamma)$ such that

$$y \in H^1(0, T; H) \cap L^\infty(0, T; V) \cap L^2(0, T; H^2(\Omega)), \quad (2.4)$$

$$y_\Gamma \in H^1(0, T; H_\Gamma) \cap L^\infty(0, T; V_\Gamma) \cap L^2(0, T; H^2(\Gamma)), \quad (2.5)$$

$$y \in [-1, 1] \text{ a. e. in } Q, \quad y_\Gamma \in [-1, 1] \text{ a. e. on } \Sigma, \quad (2.6)$$

$$\xi \in L^2(0, T; H) \text{ and } \xi \in \partial I_{[-1,1]}(y) \text{ a. e. in } Q, \quad (2.7)$$

$$\xi_\Gamma \in L^2(0, T; H_\Gamma) \text{ and } \xi_\Gamma \in \partial I_{[-1,1]}(y_\Gamma) \text{ a. e. on } \Sigma, \quad (2.8)$$

$$w \in L^2(0, T; V), \quad (2.9)$$

as well as $y_\Gamma = y|_\Gamma$, $y(0) = y_0$, $y_\Gamma(0) = y_{0\Gamma}$, is called a solution to (1.3)–(1.7) if and only if it satisfies for almost every $t \in (0, T)$ the variational equations

$$\int_\Omega \partial_t y(t) v \, dx + \int_\Omega \nabla w(t) \cdot \nabla v \, dx = 0 \quad \text{for every } v \in V, \quad (2.10)$$

$$\begin{aligned} \int_\Omega w(t) v \, dx &= \int_\Omega \partial_t y(t) v \, dx + \int_\Omega \nabla y(t) \cdot \nabla v \, dx + \int_\Omega (\xi(t) + f'_2(y(t))) v \, dx \\ &+ \int_\Gamma \partial_t y_\Gamma(t) v_\Gamma \, d\Gamma + \int_\Gamma \nabla_\Gamma y_\Gamma(t) \cdot \nabla_\Gamma v_\Gamma \, d\Gamma + \int_\Gamma (\xi_\Gamma(t) + g'_2(y_\Gamma(t)) - u_\Gamma(t)) v_\Gamma \, d\Gamma \end{aligned} \quad (2.11)$$

for every $(v, v_\Gamma) \in \mathcal{V}$.

It is worth noting that (recall the notation (1.18))

$$\begin{aligned} (\partial_t y(t))^\Omega &= 0 \quad \text{for a. e. } t \in (0, T), \text{ and } y(t)^\Omega = m_0 \text{ for every } t \in [0, T], \\ \text{where } m_0 &= (y_0)^\Omega \text{ is the mean value of } y_0, \end{aligned} \quad (2.12)$$

as usual for the Cahn–Hilliard equation. Notice that **(A3)** implies $-1 < m_0 < 1$ so that $h'(m_0)$ is finite.

The following existence and uniqueness result follows from [7, Theorems 2.2 and 2.4].

Proposition 2.2: Assume that **(A2)–(A4)** are fulfilled. Then there exists for any $u_\Gamma \in \mathcal{X}$ a quintuple $(y, y_\Gamma, w, \xi, \xi_\Gamma)$ solving problem (1.3)–(1.7) in the sense of Definition 2.1. For any such solution, we have the additional regularity properties

$$\begin{aligned} y &\in W^{1,\infty}(0, T; H) \cap H^1(0, T; V) \cap L^\infty(0, T; H^2(\Omega)), \\ y_\Gamma &\in W^{1,\infty}(0, T; H_\Gamma) \cap H^1(0, T; V_\Gamma) \cap L^\infty(0, T; H^2(\Gamma)), \\ \xi_\Gamma &\in L^\infty(0, T; H_\Gamma). \end{aligned}$$

Moreover, any two solution quintuples have the same components y, y_Γ, ξ_Γ (while the components w, ξ may not be uniquely determined).

As in the Introduction, we denote the control-to-state operator, which assigns to every $u_\Gamma \in \mathcal{X}$ the (uniquely determined) first two components (y, y_Γ) of the associated solution quintuple, by \mathcal{S}_0 .

We now turn our attention to the approximating state equations. As announced in the Introduction, we choose a special approximation of (1.3)–(1.7); namely, for $\alpha \in (0, 1]$ we consider the system

$$\partial_t y^\alpha - \Delta w^\alpha = 0 \quad \text{a. e. in } Q, \quad (2.13)$$

$$w^\alpha = \partial_t y^\alpha - \Delta y^\alpha + \varphi(\alpha) h'(y^\alpha) + f'_2(y^\alpha) \quad \text{a. e. in } Q, \quad (2.14)$$

$$\begin{aligned} y_\Gamma^\alpha = y_\Gamma^\alpha, \quad \partial_n y^\alpha + \partial_t y_\Gamma^\alpha - \Delta_\Gamma y_\Gamma^\alpha + \psi(\alpha) h'(y_\Gamma^\alpha) + g'_2(y_\Gamma^\alpha) &= u_\Gamma, \\ \partial_n w^\alpha &= 0 \quad \text{a. e. on } \Sigma, \end{aligned} \quad (2.15)$$

$$y^\alpha(\cdot, 0) = y_0 \quad \text{a. e. in } \Omega, \quad y_\Gamma^\alpha(\cdot, 0) = y_{0_\Gamma} \quad \text{a. e. on } \Gamma, \quad (2.16)$$

where h is defined in (1.10) and φ, ψ are positive and continuous functions on $(0, 1]$ that satisfy (1.12). Observe that as in (2.10), (2.11) the notion of a solution to (2.13)–(2.16) has to be understood in the sense that for almost every $t \in (0, T)$ the following variational equations are satisfied:

$$\int_\Omega \partial_t y^\alpha(t) v \, dx + \int_\Omega \nabla w^\alpha(t) \cdot \nabla v \, dx = 0 \quad \text{for every } v \in V, \quad (2.17)$$

$$\begin{aligned} \int_\Omega w^\alpha(t) v \, dx &= \int_\Omega \partial_t y^\alpha(t) v \, dx + \int_\Omega \nabla y^\alpha(t) \cdot \nabla v \, dx + \int_\Omega (\varphi(\alpha) h'(y^\alpha(t)) + f'_2(y^\alpha(t))) v \, dx \\ &+ \int_\Gamma \partial_t y_\Gamma^\alpha(t) v_\Gamma \, d\Gamma + \int_\Gamma \nabla_\Gamma y_\Gamma^\alpha(t) \cdot \nabla_\Gamma v_\Gamma \, d\Gamma + \int_\Gamma (\psi(\alpha) h'(y_\Gamma^\alpha(t)) + g'_2(y_\Gamma^\alpha(t)) - u_\Gamma(t)) v_\Gamma \, d\Gamma \end{aligned}$$

for every $(v, v_\Gamma) \in \mathcal{V}$. (2.18)

Since the functions $f^\alpha(y) := \varphi(\alpha) h(y) + f_2(y)$ and $f_\Gamma^\alpha(y) := \psi(\alpha) h(y) + g_2(y)$ fulfill on $(-1, 1)$ the conditions (2.3)–(2.7) in [8], we can infer from [8, Thm. 2.1] that the system (2.13)–(2.16) admits for every $u_\Gamma \in \mathcal{U}$ a unique solution triple $(y^\alpha, y_\Gamma^\alpha, w^\alpha)$ having the following properties:

$$y^\alpha \in W^{1,\infty}(0, T; H) \cap H^1(0, T; V) \cap L^\infty(0, T; H^2(\Omega)), \quad (2.19)$$

$$y_\Gamma^\alpha \in W^{1,\infty}(0, T; H_\Gamma) \cap H^1(0, T; V_\Gamma) \cap L^\infty(0, T; H^2(\Gamma)), \quad (2.20)$$

$$w^\alpha \in L^\infty(0, T; H^2(\Omega)), \quad (2.21)$$

$$r_-^\alpha \leq y^\alpha \leq r_+^\alpha \quad \text{a. e. in } Q, \quad r_-^\alpha \leq y_\Gamma^\alpha \leq r_+^\alpha \quad \text{a. e. on } \Sigma, \quad (2.22)$$

with suitable constants $r_-^\alpha, r_+^\alpha \in (-1, 1)$ that only depend on $\Omega, T, y_0, y_{0_\Gamma}, f_2, g_2, \alpha$, and the constant $R > 0$ introduced in **(A5)**. In particular, the control-to-state mapping for the system

(2.13)–(2.16), $\mathcal{S}_\alpha : u_\Gamma \mapsto \mathcal{S}_\alpha(u_\Gamma) := (y^\alpha, y_\Gamma^\alpha)$, for $u_\Gamma \in \mathcal{X}$, is well defined. Observe that the separation property (2.22) cannot be expected to hold uniformly in $\alpha \in (0, 1]$, in general; indeed, it cannot be excluded that there exists some sequence $\{\alpha_n\} \subset (0, 1]$ with $\alpha_n \searrow 0$ such that $r_-^{\alpha_n} \searrow -1$ and/or $r_+^{\alpha_n} \nearrow +1$ as $n \rightarrow \infty$.

We now aim to derive some a priori estimates for $(y^\alpha, y_\Gamma^\alpha)$ which are independent of α . Prior to this, we recall a functional analytic framework which is customary in the context of Cahn–Hilliard systems. We define

$$\text{dom } \mathcal{N} := \{v_* \in V^* : v_*^\Omega = 0\} \quad \text{and} \quad \mathcal{N} : \text{dom } \mathcal{N} \rightarrow \{v \in V : v^\Omega = 0\} \quad (2.23)$$

by setting for $v_* \in \text{dom } \mathcal{N}$

$$\mathcal{N}v_* \in V, \quad (\mathcal{N}v_*)^\Omega = 0, \quad \text{and} \quad \int_\Omega \nabla \mathcal{N}v_* \cdot \nabla z \, dx = \langle v_*, z \rangle_V \quad \forall z \in V, \quad (2.24)$$

that is, $\mathcal{N}v_*$ is the (unique) solution to the generalized Neumann problem $-\Delta v = v_*$ in Ω , $\partial_n v = 0$ on Γ , that satisfies $v^\Omega = 0$. Since Ω is a bounded connected domain with smooth boundary, it turns out that (2.24) yields a well-defined isomorphism that also fulfills, for all $s \geq 0$,

$$\begin{aligned} \mathcal{N}v_* \in H^{s+2}(\Omega) \quad \text{and} \quad \|\mathcal{N}v_*\|_{H^{s+2}(\Omega)} \leq C_s \|v_*\|_{H^s(\Omega)} \\ \text{for all } v_* \in H^s(\Omega) \cap \text{dom } \mathcal{N}, \end{aligned} \quad (2.25)$$

where the constant $C_s > 0$ depends only on Ω and s . Moreover, if we define the mapping $\|\cdot\|_* : V^* \rightarrow [0, +\infty)$ through the formula

$$\|v_*\|_*^2 := \|\nabla \mathcal{N}(v_* - v_*^\Omega)\|_H^2 + |v_*^\Omega|^2 \quad \forall v_* \in V^*, \quad (2.26)$$

then it is straightforward to prove that $\|\cdot\|_*$ defines a norm on V^* which turns out to be equivalent to the usual norm of V^* . We thus have, with a constant $C_* > 0$ that depends only on Ω ,

$$|\langle v_*, v \rangle_V| \leq C_* \|v_*\|_* \|v\|_V \quad \forall v_* \in V^*, \quad \forall v \in V. \quad (2.27)$$

Moreover, it follows from (2.24) and (2.26) that

$$\langle v_*, \mathcal{N}v_* \rangle_V = \|v_*\|_*^2 \quad \forall v_* \in \text{dom } \mathcal{N}, \quad (2.28)$$

and we have

$$\langle u_*, \mathcal{N}v_* \rangle_V = \langle v_*, \mathcal{N}u_* \rangle_V = \int_\Omega (\nabla \mathcal{N}v_*) \cdot (\nabla \mathcal{N}u_*) \, dx \quad \forall u_*, v_* \in \text{dom } \mathcal{N}, \quad (2.29)$$

whence also

$$2 \langle \partial_t v_*(t), \mathcal{N}v_*(t) \rangle_V = \frac{d}{dt} \int_\Omega |\nabla \mathcal{N}v_*(t)|^2 \, dx = \frac{d}{dt} \|v_*(t)\|_*^2 \quad \text{for all } t \in (0, T), \quad (2.30)$$

for any $v_* \in H^1(0, T; V^*)$ satisfying $v_*^\Omega(t) = 0$ for a. e. $t \in (0, T)$.

The next step is to prove a priori estimates uniformly in $\alpha \in (0, 1]$ for the solution $(y^\alpha, y_\Gamma^\alpha)$ of (2.13)–(2.16). We have the following result.

Proposition 2.3: *Suppose that (A2)–(A5) are satisfied. Then there is some constant $K_1^* > 0$, which only depends on $\Omega, T, y_0, y_{0\Gamma}, f_2, g_2$, and R , such that we have: whenever $(y^\alpha, y_\Gamma^\alpha) = \mathcal{S}_\alpha(u_\Gamma)$ for some $u_\Gamma \in \mathcal{U}$ and some $\alpha \in (0, 1]$, then it holds*

$$\|y^\alpha\|_{H^1(0,T;H) \cap L^\infty(0,T;V) \cap L^2(0,T;H^2(\Omega))} + \|y_\Gamma^\alpha\|_{H^1(0,T;H_\Gamma) \cap L^\infty(0,T;V_\Gamma) \cap L^2(0,T;H^2(\Gamma))} \leq K_1^*. \quad (2.31)$$

PROOF: Suppose that $u_\Gamma \in \mathcal{U}$ and $\alpha \in (0, 1]$ are arbitrarily chosen, and let $(y^\alpha, y_\Gamma^\alpha) = \mathcal{S}_\alpha(u_\Gamma)$. The result will be established in a series of a priori estimates. To this end, we will in the following denote by $C_i, i \in \mathbb{N}$, positive constants which may depend on the quantities mentioned in the statement, but not on $\alpha \in (0, 1]$. We remark that the subsequent estimates follow the same pattern as the a priori estimates in the proof of [7, Thm. 2.3], but since not all of these estimates are standard, we detail them here for the reader's convenience.

First a priori estimate: First, note that (cf. (2.12)) $y^\alpha(t)^\Omega = m_0$ for all $t \in [0, T]$, so that $(y^\alpha(t) - m_0) \in \text{dom } \mathcal{N}$. We thus may choose in (2.17) $v = \mathcal{N}(y^\alpha(t) - m_0)$, and in (2.18) $v = -(y^\alpha(t) - m_0)$. Adding the resulting equalities, then inserting two additional terms on both sides for convenience, and integrating over $[0, t]$, where $t \in [0, T]$ is arbitrary, we arrive at the identity

$$\begin{aligned} & \frac{1}{2} (\|y^\alpha(t) - m_0\|_*^2 + \|y^\alpha(t) - m_0\|_H^2 + \|y_\Gamma^\alpha(t) - m_0\|_{H_\Gamma}^2) + \int_0^t \int_\Omega |\nabla y^\alpha|^2 dx ds \\ & + \int_0^t \int_\Gamma |\nabla y_\Gamma^\alpha|^2 d\Gamma ds + \int_0^t \int_\Omega \varphi(\alpha)(h'(y^\alpha) - h'(m_0))(y^\alpha - m_0) dx ds \\ & + \int_0^t \int_\Gamma \psi(\alpha)(h'(y_\Gamma^\alpha) - h'(m_0))(y_\Gamma^\alpha - m_0) d\Gamma ds \\ & = \frac{1}{2} (\|y_0 - m_0\|_*^2 + \|y_0 - m_0\|_H^2 + \|y_{0\Gamma} - m_0\|_{H_\Gamma}^2) \\ & - \psi(\alpha)h'(m_0) \int_0^t \int_\Gamma (y_\Gamma^\alpha - m_0) d\Gamma ds \\ & - \int_0^t \int_\Omega f_2'(y^\alpha)(y^\alpha - m_0) dx ds + \int_0^t \int_\Gamma (u_\Gamma - g_2'(y_\Gamma^\alpha))(y_\Gamma^\alpha - m_0) d\Gamma ds. \quad (2.32) \end{aligned}$$

By the monotonicity of h' , all of the terms on the left-hand side of (2.32) are nonnegative, while the first term on the right-hand side is obviously bounded. Since also, in view of (A2) and (2.6),

$$\max_{0 \leq i \leq 3} \left(\|f_2^{(i)}(y^\alpha)\|_{L^\infty(Q)} + \|g_2^{(i)}(y_\Gamma^\alpha)\|_{L^\infty(\Sigma)} \right) \leq C_1 \quad \forall \alpha \in (0, 1], \quad (2.33)$$

it follows from Young's inequality and Gronwall's lemma that

$$\|y^\alpha\|_{L^\infty(0,T;H) \cap L^2(0,T;V)} + \|y_\Gamma^\alpha\|_{L^\infty(0,T;H_\Gamma) \cap L^2(0,T;V_\Gamma)} \leq C_2 \quad \forall \alpha \in (0, 1]. \quad (2.34)$$

Second a priori estimate: Recalling (2.12), we may insert $v = \mathcal{N}(\partial_t y^\alpha(t))$ in (2.17) and $v = -\partial_t y^\alpha(t)$ in (2.18). Adding the resulting equations, integrating over $[0, t]$, and using (2.24) and (2.26), we obtain the identity

$$\begin{aligned}
& \int_0^t \|\partial_t y^\alpha(s)\|_*^2 ds + \int_0^t \int_\Omega |\partial_t y^\alpha|^2 dx ds + \int_0^t \int_\Gamma |\partial_t y_\Gamma^\alpha|^2 d\Gamma ds \\
& + \frac{1}{2} (\|\nabla y^\alpha(t)\|_H^2 + \|\nabla_\Gamma y_\Gamma^\alpha(t)\|_{H_\Gamma}^2) + \int_\Omega \varphi(\alpha) h(y^\alpha(t)) dx + \int_\Gamma \psi(\alpha) h(y_\Gamma^\alpha(t)) d\Gamma \\
& = \frac{1}{2} (\|\nabla y_0\|_H^2 + \|\nabla_\Gamma y_{0\Gamma}\|_{H_\Gamma}^2) + \int_\Omega \varphi(\alpha) h(y_0) dx + \int_\Gamma \psi(\alpha) h(y_{0\Gamma}) d\Gamma \\
& - \int_0^t \int_\Omega f_2'(y^\alpha) \partial_t y^\alpha dx ds + \int_0^t \int_\Gamma (u_\Gamma - g_2'(y_\Gamma^\alpha)) \partial_t y_\Gamma^\alpha d\Gamma ds. \tag{2.35}
\end{aligned}$$

Obviously, the last two terms on the left-hand side are bounded from below and the four terms containing the initial data on the right-hand side of (2.35) are bounded. Thus, invoking (2.33) and Young's inequality, we can easily conclude from (2.35) the estimate

$$\|y^\alpha\|_{H^1(0,T;H) \cap L^\infty(0,T;V)} + \|y_\Gamma^\alpha\|_{H^1(0,T;H_\Gamma) \cap L^\infty(0,T;V_\Gamma)} \leq C_3 \quad \forall \alpha \in (0, 1]. \tag{2.36}$$

Third a priori estimate: Next, we insert $v = w^\alpha(t) - (w^\alpha(t))^\Omega$ in (2.17) and apply Young's inequality, (2.27), and Poincaré's inequality (1.19) to find the estimate

$$\begin{aligned}
& \int_\Omega |\nabla w^\alpha(t)|^2 dx = \int_\Omega |\nabla (w^\alpha(t) - (w^\alpha(t))^\Omega)|^2 dx \leq |\langle \partial_t y^\alpha(t), w^\alpha(t) - (w^\alpha(t))^\Omega \rangle_V| \\
& \leq C_* \|\partial_t y^\alpha(t)\|_* \|w^\alpha(t) - (w^\alpha(t))^\Omega\|_V \leq \frac{1}{2} \int_\Omega |\nabla w^\alpha(t)|^2 dx + C_4 \|\partial_t y^\alpha(t)\|_*^2. \tag{2.37}
\end{aligned}$$

Now recall that the embedding $H \subset V^*$ is continuous. Hence, we can infer from estimate (2.36) that

$$\|\nabla w^\alpha\|_{L^2(0,T;H)} \leq C_5 \quad \forall \alpha \in (0, 1]. \tag{2.38}$$

Next, we aim to establish a bound for the mean value of w^α in $L^2(0, T)$. To this end, we insert $v \equiv 1$ in (2.18). It follows:

$$\begin{aligned}
& \int_\Omega w^\alpha(t) dx = \int_\Omega \partial_t y^\alpha(t) dx + \int_\Gamma \partial_t y_\Gamma^\alpha(t) d\Gamma + \int_\Omega f_2'(y^\alpha(t)) dx \\
& + \int_\Gamma (g_2'(y_\Gamma^\alpha(t)) - u_\Gamma(t)) d\Gamma + \int_\Omega \varphi(\alpha) h'(y^\alpha(t)) dx + \int_\Gamma \psi(\alpha) h'(y_\Gamma^\alpha(t)) d\Gamma. \tag{2.39}
\end{aligned}$$

By virtue of (2.33) and (2.36), the first four integrals on the right-hand side of (2.39) define functions that are bounded in $L^2(0, T)$, uniformly in $\alpha \in (0, 1]$. In order to handle the two remaining terms on the right-hand side, we insert $v = \mathcal{N}(y^\alpha(t) - m_0)$ in (2.17) and $v = -(y^\alpha(t) - m_0)$ in (2.18) and add the resulting equations to obtain

$$\begin{aligned}
& \int_\Omega |\nabla y^\alpha(t)|^2 dx + \int_\Gamma |\nabla_\Gamma y_\Gamma^\alpha(t)|^2 d\Gamma + \int_\Omega \varphi(\alpha) h'(y^\alpha(t))(y^\alpha(t) - m_0) dx \\
& + \int_\Gamma \psi(\alpha) h'(y_\Gamma^\alpha(t))(y_\Gamma^\alpha(t) - m_0) d\Gamma = G^\alpha(t), \tag{2.40}
\end{aligned}$$

where

$$G^\alpha(t) := - \int_{\Omega} \partial_t y^\alpha(t) \mathcal{N}(y^\alpha(t) - m_0) dx - \int_{\Omega} (\partial_t y^\alpha(t) + f_2'(y^\alpha(t)))(y^\alpha(t) - m_0) dx - \int_{\Gamma} (\partial_t y_\Gamma^\alpha(t) + g_2'(y_\Gamma^\alpha(t)) - u_\Gamma(t))(y^\alpha(t) - m_0) d\Gamma. \quad (2.41)$$

Now, we may employ (2.26)–(2.27) and (2.33)–(2.34) to see that

$$|G^\alpha(t)| \leq C_6 (1 + \|\partial_t y^\alpha(t)\|_* \|y^\alpha(t) - m_0\|_* + \|\partial_t y^\alpha(t)\|_H + \|\partial_t y_\Gamma^\alpha(t)\|_{H_\Gamma}), \quad (2.42)$$

for a. e. $t \in (0, T)$, and it follows from (2.36) that G^α is bounded in $L^2(0, T)$, uniformly in $\alpha \in (0, 1]$.

At this point, we claim that there are $\widehat{\delta} > 0$ and $\widehat{C} > 0$ such that, for all $r \in (-1, 1)$,

$$h'(r)(r - m_0) \geq \widehat{\delta} |h'(r)| - \widehat{C}. \quad (2.43)$$

Indeed, since $-1 < m_0 < 1$, we may employ exactly the same argument as that used in [13, p. 908] to prove a corresponding estimate. From (2.43) it immediately follows that there is some $C_7 > 0$ such that for all $\alpha \in (0, 1]$ we have

$$\begin{aligned} \varphi(\alpha) h'(r)(r - m_0) &\geq \widehat{\delta} |\varphi(\alpha) h'(r)| - C_7 \quad \text{and} \\ \psi(\alpha) h'(r)(r - m_0) &\geq \widehat{\delta} |\psi(\alpha) h'(r)| - C_7 \quad \text{for all } r \in (-1, 1). \end{aligned} \quad (2.44)$$

Consequently, we deduce that

$$\begin{aligned} &\int_{\Omega} \varphi(\alpha) h'(y^\alpha(t))(y^\alpha(t) - m_0) dx + \int_{\Gamma} \psi(\alpha) h'(y_\Gamma^\alpha(t))(y_\Gamma^\alpha(t) - m_0) d\Gamma \\ &\geq \widehat{\delta} \int_{\Omega} |\varphi(\alpha) h'(y^\alpha(t))| dx + \widehat{\delta} \int_{\Gamma} |\psi(\alpha) h'(y_\Gamma^\alpha(t))| d\Gamma - C_8, \end{aligned} \quad (2.45)$$

and we can infer from (2.39) that

$$\|(w^\alpha)^\Omega\|_{L^2(0, T)} \leq C_9 \quad \forall \alpha \in (0, 1], \quad (2.46)$$

whence, recalling (2.38) and Poincaré's inequality,

$$\|w^\alpha\|_{L^2(0, T; V)} \leq C_{10} \quad \forall \alpha \in (0, 1]. \quad (2.47)$$

Fourth a priori estimate: Next, observe that in view of (2.19), (2.20) and (2.22) we have $(v, v_\Gamma) \in \mathcal{V}$ for $v = \varphi(\alpha) h'(y^\alpha)$. Hence, we may insert $v = \varphi(\alpha) h'(y^\alpha)$ in (2.18) to obtain

$$\begin{aligned} &\int_0^t \int_{\Omega} \varphi(\alpha) h''(y^\alpha) |\nabla y^\alpha|^2 dx ds + \int_0^t \int_{\Gamma} \varphi(\alpha) h''(y_\Gamma^\alpha) |\nabla_\Gamma y_\Gamma^\alpha|^2 d\Gamma ds \\ &\quad + \int_0^t \int_{\Omega} |\varphi(\alpha) h'(y^\alpha)|^2 dx ds + \int_0^t \int_{\Gamma} \varphi(\alpha) \psi(\alpha) |h'(y_\Gamma^\alpha)|^2 d\Gamma ds \\ &= \int_0^t \int_{\Omega} \varphi(\alpha) h'(y^\alpha) (w^\alpha - f_2'(y^\alpha) - \partial_t y^\alpha) dx ds \\ &\quad + \int_0^t \int_{\Gamma} \varphi(\alpha) h'(y_\Gamma^\alpha) (u_\Gamma - g_2'(y_\Gamma^\alpha) - \partial_t y_\Gamma^\alpha) d\Gamma ds. \end{aligned} \quad (2.48)$$

Now notice that $h'' > 0$ in $(-1, 1)$, which implies that the two integrals in which h'' occurs in the integrands, are both nonnegative. Moreover, (1.12) implies that

$$\int_0^t \int_{\Gamma} \varphi(\alpha) \psi(\alpha) |h'(y_{\Gamma}^{\alpha})|^2 d\Gamma ds \geq \frac{1}{C_{\varphi\psi}} \int_0^t \int_{\Gamma} (\varphi(\alpha))^2 |h'(y_{\Gamma}^{\alpha})|^2 d\Gamma ds.$$

Therefore the boundary integral

$$\int_0^t \int_{\Gamma} \varphi(\alpha) h'(y_{\Gamma}^{\alpha}) (u_{\Gamma} - g_2'(y_{\Gamma}^{\alpha}) - \partial_t y_{\Gamma}^{\alpha}) d\Gamma ds$$

can be handled using Young's inequality. Now applying (2.33), (2.36), (2.47) and Young's inequality, we find that

$$\|\varphi(\alpha) h'(y^{\alpha})\|_{L^2(0,T;H)} \leq C_{11} \quad \forall \alpha \in (0, 1]. \quad (2.49)$$

Fifth a priori estimate: Now observe that the variational equality (2.18) implies that y^{α} solves (2.14) at least in the sense of distributions. Since all other terms have been proved to be bounded in $L^2(0, T; H)$, we must have

$$\|\Delta y^{\alpha}\|_{L^2(0,T;H)} \leq C_{12} \quad \forall \alpha \in (0, 1]. \quad (2.50)$$

Next, we use [3, Thm. 3.2, p. 1.79] to conclude that

$$\int_0^T \|y^{\alpha}(t)\|_{H^{3/2}(\Omega)}^2 dt \leq C_{13} \int_0^T (\|\Delta y^{\alpha}(t)\|_H^2 + \|y_{\Gamma}^{\alpha}(t)\|_{V_{\Gamma}}^2) dt,$$

whence it follows that

$$\|y^{\alpha}\|_{L^2(0,T;H^{3/2}(\Omega))} \leq C_{14} \quad \forall \alpha \in (0, 1]. \quad (2.51)$$

Hence, by the trace theorem [3, Thm. 2.27, p. 1.64], we have

$$\|\partial_{\mathbf{n}} y^{\alpha}\|_{L^2(0,T;H_{\Gamma})} \leq C_{15} \quad \forall \alpha \in (0, 1]. \quad (2.52)$$

From the above estimates it follows that all the terms occurring in the integration by parts formula for the Laplace operator are functions, and we deduce that the variational equation (2.18) also implies that the second identity in (2.15) holds at least in a generalized sense, in principle. Therefore, the preceding estimates yield that, by letting $G_{\Gamma}^{\alpha} := u_{\Gamma} - \partial_{\mathbf{n}} y^{\alpha} - \partial_t y_{\Gamma}^{\alpha} - g_2'(y_{\Gamma}^{\alpha})$, we can write

$$-\Delta_{\Gamma} y_{\Gamma}^{\alpha} + \psi(\alpha) h'(y_{\Gamma}^{\alpha}) = G_{\Gamma}^{\alpha} \text{ on } \Sigma, \text{ where } \|G_{\Gamma}^{\alpha}\|_{L^2(\Sigma)} \leq C_{16} \quad \forall \alpha \in (0, 1]. \quad (2.53)$$

Testing the above equation by $\psi(\alpha) h'(y_{\Gamma}^{\alpha})$, we obtain

$$\begin{aligned} & \int_0^t \int_{\Gamma} \psi(\alpha) h''(y_{\Gamma}^{\alpha}) |\nabla_{\Gamma} y_{\Gamma}^{\alpha}|^2 d\Gamma ds + \int_0^t \int_{\Gamma} |\psi(\alpha) h'(y_{\Gamma}^{\alpha})|^2 d\Gamma ds \\ &= \int_0^t \int_{\Gamma} \psi(\alpha) h'(y_{\Gamma}^{\alpha}) G_{\Gamma}^{\alpha} d\Gamma ds, \end{aligned} \quad (2.54)$$

and a simple application of Young's inequality shows that

$$\|\psi(\alpha)h'(y_\Gamma^\alpha)\|_{L^2(0,T;H_\Gamma)} \leq C_{17} \quad \forall \alpha \in (0, 1], \quad (2.55)$$

whence also

$$\|\Delta_\Gamma y_\Gamma^\alpha\|_{L^2(0,T;H_\Gamma)} \leq C_{18} \quad \forall \alpha \in (0, 1]. \quad (2.56)$$

The boundary version of the elliptic regularity theory then yields

$$\|y_\Gamma^\alpha\|_{L^2(0,T;H^2(\Gamma))} \leq C_{19} \quad \forall \alpha \in (0, 1], \quad (2.57)$$

and consequently it follows from standard elliptic estimates that

$$\|y^\alpha\|_{L^2(0,T;H^2(\Omega))} \leq C_{20} \quad \forall \alpha \in (0, 1]. \quad (2.58)$$

With this, the assertion is completely proved. ■

3 Existence and approximation of optimal controls

Our first aim in this section is to prove the following existence result:

Theorem 3.1: *Suppose that the assumptions (A1)–(A5) are satisfied. Then the optimal control problem (\mathcal{P}_0) admits a solution.*

Before proving Theorem 3.1, we introduce the solution space

$$\mathcal{Y} := \left\{ (y, y_\Gamma) \in \mathcal{Y} : y \in H^1(0, T; H) \cap L^\infty(0, T; V) \cap L^2(0, T; H^2(\Omega)), \right. \\ \left. y_\Gamma = y|_\Gamma, \quad y_\Gamma \in H^1(0, T; H_\Gamma) \cap L^\infty(0, T; V_\Gamma) \cap L^2(0, T; H^2(\Gamma)) \right\}, \quad (3.1)$$

and a family of auxiliary optimal control problems (\mathcal{P}_α) , which is parametrized by $\alpha \in (0, 1]$. In what follows, we will always assume that h is given by (1.10) and that φ and ψ are functions that are positive and continuous on $(0, 1]$ and satisfy the conditions (1.12). For $\alpha \in (0, 1]$, let us denote by \mathcal{S}_α the operator mapping $u_\Gamma \in \mathcal{U}_{\text{ad}}$ into the unique solution $(y^\alpha, y_\Gamma^\alpha) \in \mathcal{Y}$ to the variational problem (2.16)–(2.18). We define:

$$(\mathcal{P}_\alpha) \quad \text{Minimize } \mathcal{J}((y, y_\Gamma), u_\Gamma) \quad \text{over } \mathcal{Y} \times \mathcal{U}_{\text{ad}} \quad \text{subject to the condition that} \\ (2.16)–(2.18) \text{ are satisfied.}$$

The following result is a consequence of [8, Thm. 2.2].

Lemma 3.2: *Suppose that the assumptions (A1)–(A5) and (1.10), (1.12) are fulfilled, and let $\alpha \in (0, 1]$ be given. Then the optimal control problem (\mathcal{P}_α) admits a solution.*

PROOF OF THEOREM 3.1: Let $\{\alpha_n\} \subset (0, 1]$ be any sequence such that $\alpha_n \searrow 0$ as $n \rightarrow \infty$. By virtue of Lemma 3.2, for any $n \in \mathbb{N}$ we may pick an optimal pair for the optimal control problem (\mathcal{P}_{α_n}) ,

$$((y^{\alpha_n}, y_\Gamma^{\alpha_n}), u_\Gamma^{\alpha_n}) \in \mathcal{Y} \times \mathcal{U}_{\text{ad}}$$

where $(y^{\alpha_n}, y_\Gamma^{\alpha_n}, w^{\alpha_n})$ is the unique solution to (2.16)–(2.18), written for $\alpha = \alpha_n$, which satisfies (2.19)–(2.22). In particular, $(y^{\alpha_n}, y_\Gamma^{\alpha_n}) = \mathcal{S}_{\alpha_n}(u_\Gamma^{\alpha_n})$ for all $n \in \mathbb{N}$. Moreover, Proposition 2.3 implies that (2.31) holds for any $\alpha_n, n \in \mathbb{N}$. From this and from (2.47) we may without loss of generality assume that there are $u_\Gamma \in \mathcal{U}_{\text{ad}}, w$, and (y, y_Γ) such that

$$u_\Gamma^{\alpha_n} \rightharpoonup u_\Gamma \quad \text{weakly-star in } \mathcal{X}, \quad (3.2)$$

$$w^{\alpha_n} \rightharpoonup w \quad \text{weakly in } L^2(0, T; V), \quad (3.3)$$

$$y^{\alpha_n} \rightharpoonup y \quad \text{weakly-star in } H^1(0, T; H) \cap L^\infty(0, T; V) \cap L^2(0, T; H^2(\Omega)), \quad (3.4)$$

$$y_\Gamma^{\alpha_n} \rightharpoonup y_\Gamma \quad \text{weakly-star in } H^1(0, T; H_\Gamma) \cap L^\infty(0, T; V_\Gamma) \cap L^2(0, T; H^2(\Gamma)). \quad (3.5)$$

By the continuity of the embedding $H^1(0, T; H) \cap L^2(0, T; H^2(\Omega)) \subset C^0([0, T]; V)$, we have in fact $y \in C^0([0, T]; V)$, and, by the same token, $y_\Gamma \in C^0([0, T]; V_\Gamma)$. Owing to the Aubin-Lions lemma (see [20, Sect. 8, Cor. 4]), we also have

$$y^{\alpha_n} \rightarrow y \quad \text{strongly in } C^0([0, T]; H) \cap L^2(0, T; V), \quad (3.6)$$

$$y_\Gamma^{\alpha_n} \rightarrow y_\Gamma \quad \text{strongly in } C^0([0, T]; H_\Gamma) \cap L^2(0, T; V_\Gamma). \quad (3.7)$$

In particular, it holds $y(\cdot, 0) = y_0$, as well as $y_\Gamma(\cdot, 0) = y_{0\Gamma}$. In addition, the Lipschitz continuity of f'_2 and g'_2 on $[-1, 1]$ yields that

$$f'_2(y^{\alpha_n}) \rightarrow f'_2(y) \quad \text{strongly in } C^0([0, T]; H), \quad (3.8)$$

$$g'_2(y_\Gamma^{\alpha_n}) \rightarrow g'_2(y_\Gamma) \quad \text{strongly in } C^0([0, T]; H_\Gamma). \quad (3.9)$$

Moreover, (2.49) and (2.55) show that without loss of generality we may also assume that

$$\varphi(\alpha_n) h'(y^{\alpha_n}) \rightharpoonup \xi \quad \text{weakly in } L^2(0, T; H), \quad (3.10)$$

$$\psi(\alpha_n) h'(y_\Gamma^{\alpha_n}) \rightharpoonup \xi_\Gamma \quad \text{weakly in } L^2(0, T; H_\Gamma), \quad (3.11)$$

for some weak limits ξ and ξ_Γ .

Combining the above convergences, we may pass to the limit as $n \rightarrow \infty$ in (2.17) and (2.18) (written for α_n) to find that the quintuple $(y, y_\Gamma, w, \xi, \xi_\Gamma)$ is a solution to (2.10)–(2.11), and obviously the properties (2.4)–(2.6) and (2.9) are satisfied. In order to show that the quintuple $(y, y_\Gamma, w, \xi, \xi_\Gamma)$ is a solution to problem (1.3)–(1.7) in the sense of Definition 2.1, it remains to show that $\xi \in \partial I_{[-1, 1]}(y)$ a. e. in Q and $\xi_\Gamma \in \partial I_{[-1, 1]}(y_\Gamma)$ a. e. in Σ . Once this will be shown, we can conclude that $(y, y_\Gamma) = \mathcal{S}_0(u_\Gamma)$, i. e., that the pair $((y, y_\Gamma, w, \xi, \xi_\Gamma), u_\Gamma)$ is admissible for (\mathcal{P}_0) .

Now, recalling (1.10) and owing to the convexity of h , we have, for every $n \in \mathbb{N}$,

$$\begin{aligned} & \int_0^T \int_\Omega \varphi(\alpha_n) h(y^{\alpha_n}) \, dx \, dt + \int_0^T \int_\Omega \varphi(\alpha_n) h'(y^{\alpha_n}) (z - y^{\alpha_n}) \, dx \, dt \\ & \leq \int_0^T \int_\Omega \varphi(\alpha_n) h(z) \, dx \, dt \quad \text{for all } z \in \mathcal{K} = \{v \in L^2(Q) : |v| \leq 1 \text{ a.e. in } Q\}. \end{aligned} \quad (3.12)$$

Thanks to (1.12), the integral on the right-hand side and the first integral on the left-hand side of (3.12) tend to zero as $n \rightarrow \infty$, since h is a bounded function. Hence, invoking (3.6) and (3.10), the passage to the limit as $n \rightarrow \infty$ yields

$$\int_0^T \int_{\Omega} \xi (y - z) \, dx \, dt \geq 0 \quad \forall z \in \mathcal{K}. \quad (3.13)$$

Inequality (3.13) entails that ξ is an element of the subdifferential of the extension \mathcal{I} of $I_{[-1,1]}$ to $L^2(Q)$, which means that $\xi \in \partial \mathcal{I}(y)$ or, equivalently (cf. [2, Ex. 2.3.3., p. 25]), $\xi \in \partial I_{[-1,1]}(y)$ a. e. in Q . Similarly we prove that $\xi_{\Gamma} \in \partial I_{[-1,1]}(y_{\Gamma})$ a. e. in Σ .

It remains to show that $((y, y_{\Gamma}, w, \xi, \xi_{\Gamma}), u_{\Gamma})$ is in fact optimal for (\mathcal{P}_0) . To this end, let $v_{\Gamma} \in \mathcal{U}_{\text{ad}}$ be arbitrary. In view of the convergence properties (3.2) and (3.4)–(3.7), and using the weak sequential lower semicontinuity properties of the cost functional, we have

$$\begin{aligned} \mathcal{J}((y, y_{\Gamma}), u_{\Gamma}) &= \mathcal{J}(\mathcal{S}_0(u_{\Gamma}), u_{\Gamma}) \leq \liminf_{n \rightarrow \infty} \mathcal{J}(\mathcal{S}_{\alpha_n}(u_{\Gamma}^{\alpha_n}), u_{\Gamma}^{\alpha_n}) \\ &\leq \liminf_{n \rightarrow \infty} \mathcal{J}(\mathcal{S}_{\alpha_n}(v_{\Gamma}), v_{\Gamma}) = \lim_{n \rightarrow \infty} \mathcal{J}(\mathcal{S}_{\alpha_n}(v_{\Gamma}), v_{\Gamma}) = \mathcal{J}(\mathcal{S}_0(v_{\Gamma}), v_{\Gamma}), \end{aligned} \quad (3.14)$$

where for the last equality the continuity of the cost functional with respect to the first variable was used. With this, the assertion is completely proved. \blacksquare

Corollary 3.3: *Let the general assumptions (A1)–(A5) and (1.10), (1.12) be satisfied, and let sequences $\{\alpha_n\} \subset (0, 1]$ and $\{u_{\Gamma}^{\alpha_n}\} \subset \mathcal{U}$ be given such that, as $n \rightarrow \infty$, $\alpha_n \searrow 0$ and $u_{\Gamma}^{\alpha_n} \rightarrow u_{\Gamma}$ weakly-star in \mathcal{X} . Then we have*

$$\mathcal{S}_{\alpha_n}(u_{\Gamma}^{\alpha_n}) \rightarrow \mathcal{S}_0(u_{\Gamma}) \quad \text{weakly-star in } \mathcal{Y}, \quad (3.15)$$

$$\lim_{n \rightarrow \infty} \mathcal{J}(\mathcal{S}_{\alpha_n}(v_{\Gamma}), v_{\Gamma}) = \mathcal{J}(\mathcal{S}_0(v_{\Gamma}), v_{\Gamma}) \quad \forall v_{\Gamma} \in \mathcal{U}. \quad (3.16)$$

PROOF: By the same arguments as in the first part of the proof of Theorem 3.1, we can conclude that (3.15) holds at least for some subsequence. But the limit is given by the first two components of a solution quintuple in the sense of Definition 2.1 to the state system (1.3)–(1.7), which, according to Proposition 2.2, are uniquely determined. Hence, the limit is the same for all convergent subsequences and (3.15) is true for the entire sequence. Now, let $v_{\Gamma} \in \mathcal{U}$ be arbitrary. Then (see (3.6)–(3.7)) $\mathcal{S}_{\alpha_n}(v_{\Gamma})$ converges strongly to $\mathcal{S}_0(v_{\Gamma})$ in $(C^0([0, T]; H) \cap L^2(0, T; V)) \times (C^0([0, T]; H_{\Gamma}) \cap L^2(0, T; V_{\Gamma}))$, so that (3.16) follows from the continuity properties of the cost functional with respect to its first argument. \blacksquare

Theorem 3.1 does not yield any information on whether every solution to the optimal control problem (\mathcal{P}_0) can be approximated by a sequence of solutions to the problems (\mathcal{P}_{α}) . As already announced in the Introduction, we are not able to prove such a general ‘global’ result. Instead, we can only give a ‘local’ answer for every individual optimizer of (\mathcal{P}_0) . For this purpose, we employ a trick due to Barbu [1]. To this end, let $\bar{u}_{\Gamma} \in \mathcal{U}_{\text{ad}}$ be an arbitrary optimal

control for (\mathcal{P}_0) , and let $(\bar{y}, \bar{y}_\Gamma, \bar{w}, \bar{\xi}, \bar{\xi}_\Gamma)$ be an associated solution quintuple to the state system (1.3)–(1.7) in the sense of Definition 2.1. In particular, $(\bar{y}, \bar{y}_\Gamma) = \mathcal{S}_0(\bar{u}_\Gamma)$. We associate with this optimal control the *adapted cost functional*

$$\tilde{\mathcal{J}}((y, y_\Gamma), u_\Gamma) := \mathcal{J}((y, y_\Gamma), u_\Gamma) + \frac{1}{2} \|u_\Gamma - \bar{u}_\Gamma\|_{L^2(\Sigma)}^2 \quad (3.17)$$

and a corresponding *adapted optimal control problem*

$$(\tilde{\mathcal{P}}_\alpha) \quad \text{Minimize } \tilde{\mathcal{J}}((y, y_\Gamma), u_\Gamma) \text{ over } \mathcal{Y} \times \mathcal{U}_{\text{ad}} \text{ subject to the condition that (2.13)–(2.16) be satisfied.}$$

With a standard direct argument that needs no repetition here, we can show the following result.

Lemma 3.4: *Suppose that the assumptions (A1)–(A5) and (1.10), (1.12) are satisfied, and let $\alpha \in (0, 1]$. Then the optimal control problem $(\tilde{\mathcal{P}}_\alpha)$ admits a solution.*

We are now in the position to give a partial answer to the question raised above. We have the following result.

Theorem 3.5: *Let the general assumptions (A1)–(A5) and (1.10), (1.12) be fulfilled, and suppose that $\bar{u}_\Gamma \in \mathcal{U}_{\text{ad}}$ is an arbitrary optimal control of (\mathcal{P}_0) with associated state quintuple $(\bar{y}, \bar{y}_\Gamma, \bar{w}, \bar{\xi}, \bar{\xi}_\Gamma)$. Then for every sequence $\{\alpha_n\} \subset (0, 1]$ such that $\alpha_n \searrow 0$ as $n \rightarrow \infty$ and for any $n \in \mathbb{N}$ there exists some optimal control $\bar{u}_\Gamma^{\alpha_n} \in \mathcal{U}_{\text{ad}}$ of the adapted problem $(\tilde{\mathcal{P}}_{\alpha_n})$ with associated state triple $(\bar{y}^{\alpha_n}, \bar{y}_\Gamma^{\alpha_n}, \bar{w}^{\alpha_n})$ such that, as $n \rightarrow \infty$,*

$$\bar{u}_\Gamma^{\alpha_n} \rightarrow \bar{u}_\Gamma \text{ strongly in } H_\Gamma, \quad (3.18)$$

$$\bar{y}^{\alpha_n} \rightarrow \bar{y} \text{ weakly-star in } H^1(0, T; H) \cap L^\infty(0, T; V) \cap L^2(0, T; H^2(\Omega)), \quad (3.19)$$

$$\bar{y}_\Gamma^{\alpha_n} \rightarrow \bar{y}_\Gamma \text{ weakly-star in } H^1(0, T; H_\Gamma) \cap L^\infty(0, T; V_\Gamma) \cap L^2(0, T; H^2(\Gamma)), \quad (3.20)$$

$$\tilde{\mathcal{J}}((\bar{y}^{\alpha_n}, \bar{y}_\Gamma^{\alpha_n}), \bar{u}_\Gamma^{\alpha_n}) \rightarrow \mathcal{J}((\bar{y}, \bar{y}_\Gamma), \bar{u}_\Gamma). \quad (3.21)$$

PROOF: Let $\alpha_n \searrow 0$ as $n \rightarrow \infty$. For any $n \in \mathbb{N}$, we pick an optimal control $\bar{u}_\Gamma^{\alpha_n} \in \mathcal{U}_{\text{ad}}$ for the adapted problem $(\tilde{\mathcal{P}}_{\alpha_n})$ and denote by $(\bar{y}^{\alpha_n}, \bar{y}_\Gamma^{\alpha_n}, \bar{w}^{\alpha_n})$ the associated solution triple of problem (2.13)–(2.16); in particular, we have $(\bar{y}^{\alpha_n}, \bar{y}_\Gamma^{\alpha_n}) = \mathcal{S}_{\alpha_n}(\bar{u}_\Gamma^{\alpha_n})$, and (2.19)–(2.22) are satisfied. By the boundedness of \mathcal{U}_{ad} , we have for some subsequence of $\{\alpha_n\}$, which is again indexed by n , that it holds

$$\bar{u}_\Gamma^{\alpha_n} \rightarrow u_\Gamma \text{ weakly-star in } \mathcal{X} \text{ as } n \rightarrow \infty, \quad (3.22)$$

with some $u_\Gamma \in \mathcal{U}_{\text{ad}}$. Owing to Corollary 3.3, we have

$$(\bar{y}^{\alpha_n}, \bar{y}_\Gamma^{\alpha_n}) = \mathcal{S}_{\alpha_n}(\bar{u}_\Gamma^{\alpha_n}) \rightarrow \mathcal{S}_0(u_\Gamma) =: (y, y_\Gamma) \text{ weakly-star in } \mathcal{Y}. \quad (3.23)$$

In particular, y, y_Γ are the first two components of a quintuple $(y, y_\Gamma, w, \xi, \xi_\Gamma)$ solving the state system associated with u_Γ , which implies that $((y, y_\Gamma, w, \xi, \xi_\Gamma), u_\Gamma)$ is admissible for (\mathcal{P}_0) .

We now aim to prove that $u_\Gamma = \bar{u}_\Gamma$. Once this will be shown, the uniqueness result of Proposition 2.2 yields that also $(y, y_\Gamma) = (\bar{y}, \bar{y}_\Gamma)$, which shows that (3.19) and (3.20) hold at least for the subsequence; but since the limit is the same for any subsequence, we have (3.19), (3.20) for the entire sequence $\{\alpha_n\}$. By the same token, also (3.22) will hold for the entire sequence.

Indeed, we have, owing to the weak sequential lower semicontinuity of $\tilde{\mathcal{J}}$, and in view of the optimality property of $((\bar{y}, \bar{y}_\Gamma), \bar{u}_\Gamma)$ for problem (\mathcal{P}_0) ,

$$\begin{aligned} \liminf_{n \rightarrow \infty} \tilde{\mathcal{J}}((\bar{y}^{\alpha_n}, \bar{y}_\Gamma^{\alpha_n}), \bar{u}_\Gamma^{\alpha_n}) &\geq \mathcal{J}((y, y_\Gamma), u_\Gamma) + \frac{1}{2} \|u_\Gamma - \bar{u}_\Gamma\|_{L^2(\Sigma)}^2 \\ &\geq \mathcal{J}((\bar{y}, \bar{y}_\Gamma), \bar{u}_\Gamma) + \frac{1}{2} \|u_\Gamma - \bar{u}_\Gamma\|_{L^2(\Sigma)}^2. \end{aligned} \quad (3.24)$$

On the other hand, the optimality property of $((\bar{y}^{\alpha_n}, \bar{y}_\Gamma^{\alpha_n}), \bar{u}_\Gamma^{\alpha_n})$ for problem $(\tilde{\mathcal{P}}_{\alpha_n})$ yields that for any $n \in \mathbb{N}$ we have

$$\tilde{\mathcal{J}}((\bar{y}^{\alpha_n}, \bar{y}_\Gamma^{\alpha_n}), \bar{u}_\Gamma^{\alpha_n}) = \tilde{\mathcal{J}}(\mathcal{S}_{\alpha_n}(\bar{u}_\Gamma^{\alpha_n}), \bar{u}_\Gamma^{\alpha_n}) \leq \tilde{\mathcal{J}}(\mathcal{S}_{\alpha_n}(\bar{u}_\Gamma), \bar{u}_\Gamma), \quad (3.25)$$

whence, taking the limes superior as $n \rightarrow \infty$ on both sides and invoking (3.16) in Corollary 3.3,

$$\begin{aligned} \limsup_{n \rightarrow \infty} \tilde{\mathcal{J}}((\bar{y}^{\alpha_n}, \bar{y}_\Gamma^{\alpha_n}), \bar{u}_\Gamma^{\alpha_n}) &\leq \tilde{\mathcal{J}}(\mathcal{S}_0(\bar{u}_\Gamma), \bar{u}_\Gamma) = \tilde{\mathcal{J}}((\bar{y}, \bar{y}_\Gamma), \bar{u}_\Gamma) \\ &= \mathcal{J}((\bar{y}, \bar{y}_\Gamma), \bar{u}_\Gamma). \end{aligned} \quad (3.26)$$

Combining (3.24) with (3.26), we have thus shown that $\frac{1}{2} \|u_\Gamma - \bar{u}_\Gamma\|_{L^2(\Sigma)}^2 = 0$, so that $u_\Gamma = \bar{u}_\Gamma$ and thus also $(y, y_\Gamma) = (\bar{y}, \bar{y}_\Gamma)$. Moreover, (3.24) and (3.26) also imply that

$$\begin{aligned} \mathcal{J}((\bar{y}, \bar{y}_\Gamma), \bar{u}_\Gamma) &= \tilde{\mathcal{J}}((\bar{y}, \bar{y}_\Gamma), \bar{u}_\Gamma) = \liminf_{n \rightarrow \infty} \tilde{\mathcal{J}}((\bar{y}^{\alpha_n}, \bar{y}_\Gamma^{\alpha_n}), \bar{u}_\Gamma^{\alpha_n}) \\ &= \limsup_{n \rightarrow \infty} \tilde{\mathcal{J}}((\bar{y}^{\alpha_n}, \bar{y}_\Gamma^{\alpha_n}), \bar{u}_\Gamma^{\alpha_n}) = \lim_{n \rightarrow \infty} \tilde{\mathcal{J}}((\bar{y}^{\alpha_n}, \bar{y}_\Gamma^{\alpha_n}), \bar{u}_\Gamma^{\alpha_n}), \end{aligned} \quad (3.27)$$

which proves (3.21) and, at the same time, also (3.18). The assertion is thus completely checked. ■

4 The optimality system

In this section our aim is to establish first-order necessary optimality conditions for the optimal control problem (\mathcal{P}_0) . This will be achieved by passage to the limit as $\alpha \searrow 0$ in the (recently in [8]) derived first-order necessary optimality conditions for the adapted optimal control problems (\mathcal{P}_α) . It will turn out that in the limit certain generalized first-order necessary conditions of optimality result. To fix things once and for all, we will throughout the entire section assume that h is given by (1.10) and that (1.12) and the general assumptions **(A1)**–**(A5)** are satisfied; we

also assume that a fixed optimal control $\bar{u}_\Gamma \in \mathcal{U}_{\text{ad}}$ for (\mathcal{P}_0) , along with a solution quintuple $(\bar{y}, \bar{y}_\Gamma, \bar{w}, \bar{\xi}, \bar{\xi}_\Gamma)$ of the associated state system (1.3)–(1.7), is given. In addition, we make the following compatibility assumption:

(A6) It holds $\beta_3 = \beta_4 = 0$.

We remark that in [8, Remark 5.6] it has been pointed out that this assumption is dispensable at the expense of less regularity of the adjoint state variables; in order to keep the technicalities at a reasonable level, we here confine ourselves to the case $\beta_3 = \beta_4 = 0$.

4.1 The optimality conditions for $(\tilde{\mathcal{P}}_\alpha)$

We begin our analysis by formulating the adjoint state system for the adapted control problem $(\tilde{\mathcal{P}}_\alpha)$. To this end, let us assume that $\bar{u}_\Gamma^\alpha \in \mathcal{U}_{\text{ad}}$ is an arbitrary optimal control for $(\tilde{\mathcal{P}}_\alpha)$ and that $(\bar{y}^\alpha, \bar{y}_\Gamma^\alpha, \bar{w}^\alpha)$ is the solution triple to the associated state system (2.13)–(2.16). In particular, $(\bar{y}^\alpha, \bar{y}_\Gamma^\alpha) = \mathcal{S}_\alpha(\bar{u}_\Gamma^\alpha)$, and the solution has the regularity properties (2.19)–(2.22). It then follows (see [8, Eqs. (5.7)–(5.9)]) that the corresponding adjoint state variables $q^\alpha, q_\Gamma^\alpha, p^\alpha$ solve the following backward-in-time variational problem:

$$\int_{\Omega} q^\alpha(t) v \, dx = \int_{\Omega} \nabla p^\alpha(t) \cdot \nabla v \, dx \quad \text{for all } v \in V \text{ and } t \in (0, T), \quad (4.1)$$

$$\begin{aligned} & - \int_{\Omega} \partial_t (q^\alpha(t) + p^\alpha(t)) v \, dx + \int_{\Omega} \nabla q^\alpha(t) \cdot \nabla v \, dx + \int_{\Gamma} \nabla_{\Gamma} q_\Gamma^\alpha(t) \cdot \nabla_{\Gamma} v_{\Gamma} \, d\Gamma \\ & - \int_{\Gamma} \partial_t q_\Gamma^\alpha v_{\Gamma} \, d\Gamma + \int_{\Omega} (\varphi(\alpha) h''(\bar{y}^\alpha(t)) + f_2''(\bar{y}^\alpha(t))) q^\alpha(t) v \, dx \\ & + \int_{\Gamma} (\psi(\alpha) h''(\bar{y}_\Gamma^\alpha(t)) + g_2''(\bar{y}_\Gamma^\alpha(t))) q_\Gamma^\alpha(t) v_{\Gamma} \, d\Gamma \\ & = \int_{\Omega} \beta_1 (\bar{y}^\alpha(t) - z_Q(t)) v \, dx + \int_{\Gamma} \beta_2 (\bar{y}_\Gamma^\alpha(t) - z_\Sigma(t)) v_{\Gamma} \, d\Gamma \\ & \text{for every } (v, v_{\Gamma}) \in \mathcal{V} \text{ and a. a. } t \in (0, T), \end{aligned} \quad (4.2)$$

$$\int_{\Omega} (q^\alpha(T) + p^\alpha(T)) v \, dx + \int_{\Gamma} q_\Gamma^\alpha(T) v_{\Gamma} \, d\Gamma = 0 \quad \text{for every } (v, v_{\Gamma}) \in \mathcal{V}. \quad (4.3)$$

In [8, Thm. 2.4] it has been shown that the system (4.1)–(4.2) has for every $\alpha \in (0, 1]$ a unique solution triple $(q^\alpha, q_\Gamma^\alpha, p^\alpha)$ such that

$$(q^\alpha, q_\Gamma^\alpha) \in \mathcal{V}, \quad p^\alpha \in H^1(0, T; H^2(\Omega)) \cap L^2(0, T; H^4(\Omega)), \quad (4.4)$$

and we may regard $(q^\alpha, q_\Gamma^\alpha, p^\alpha)$ as a solution to the linear PDE system

$$-\Delta p^\alpha = q^\alpha \quad \text{in } Q, \quad \partial_{\mathbf{n}} p^\alpha = 0 \quad \text{on } \Sigma, \quad (4.5)$$

$$-\partial_t(q^\alpha + p^\alpha) - \Delta q^\alpha + (\varphi(\alpha)h''(\bar{y}^\alpha) + f_2''(\bar{y}^\alpha))q^\alpha = \beta_1(\bar{y}^\alpha - z_Q) \quad \text{in } Q, \quad (4.6)$$

$$-\partial_t q_\Gamma^\alpha + \partial_{\mathbf{n}} q^\alpha - \Delta_\Gamma q_\Gamma^\alpha + (\psi(\alpha)h''(\bar{y}_\Gamma^\alpha) + g_2''(\bar{y}_\Gamma^\alpha))q_\Gamma^\alpha = \beta_2(\bar{y}_\Gamma^\alpha - z_\Sigma) \\ \text{and } q_\Gamma^\alpha = q_\Gamma^\alpha \quad \text{on } \Sigma, \quad (4.7)$$

$$q^\alpha(T) + p^\alpha(T) = 0 \quad \text{in } \Omega, \quad q_\Gamma^\alpha(T) = 0 \quad \text{on } \Gamma. \quad (4.8)$$

Moreover, as we are now dealing with $(\tilde{\mathcal{P}}_\alpha)$ instead of (\mathcal{P}_α) , the variational inequality given by [8, Thm. 2.5] has to be modified as follows:

$$\int_0^T \int_\Gamma (q_\Gamma^\alpha + \beta_5 \bar{u}_\Gamma^\alpha + (\bar{u}_\Gamma^\alpha - \bar{u}_\Gamma)) (v_\Gamma - \bar{u}_\Gamma) \, d\Gamma \, dt \geq 0 \quad \forall v_\Gamma \in \mathcal{U}_{\text{ad}}. \quad (4.9)$$

In order to pave the road for the limit process as $\alpha \searrow 0$ in the optimality conditions for $(\tilde{\mathcal{P}}_\alpha)$, we employ an idea that was developed in [8]. Namely, it is possible to show that the system (4.1)–(4.3) is equivalent to a decoupled problem that can be solved by first finding q^α and then reconstructing p^α . We briefly motivate this approach. First, standard embedding results yield that $q^\alpha \in C^0([0, T]; V)$, and it immediately follows from inserting $v \equiv 1$ in (4.1) that $(q^\alpha(t))^\Omega = 0$ for all $t \in [0, T]$. Hence $q^\alpha(t) \in \text{dom } \mathcal{N}$, and, with the mean value function $(p^\alpha)^\Omega \in C^0([0, T])$, the function $(p^\alpha - (p^\alpha)^\Omega)(t)$ satisfies for every $t \in [0, T]$ the identity (2.24) with $v_* = q^\alpha(t)$. In other words, we have

$$(p^\alpha - (p^\alpha)^\Omega)(t) = \mathcal{N}(q^\alpha(t)) \quad \forall t \in [0, T]. \quad (4.10)$$

On the other hand, $(p^\alpha(t))^\Omega$ is for any fixed $t \in [0, T]$ a constant function and thus orthogonal in H to the subspace of functions having zero mean value. Consequently, p^α is completely eliminated from (4.2) if we confine ourselves to the use of test functions having zero mean value. Similar remarks apply for the final condition on $q^\alpha + p^\alpha$ appearing in (4.3). In this way, we may try to first construct $(q^\alpha, q_\Gamma^\alpha)$ and then recover p^α from (4.10), where the calculation of $(p^\alpha(t))^\Omega$ is an easy task, since simple integration of (4.6) over $\Omega \times [t, T]$, using (4.8) and the fact that $q^\alpha(t)$ has zero mean value, immediately yields that

$$(p^\alpha(t))^\Omega = \frac{1}{|\Omega|} \int_t^T \int_\Omega (-\Delta q^\alpha + (\varphi(\alpha)h''(\bar{y}^\alpha) + f_2''(\bar{y}^\alpha)) q^\alpha - \beta_1(\bar{y}^\alpha - z_Q)) \, dx \, ds.$$

We now make this approach precise. Since our test functions will have zero mean value, we introduce the linear spaces

$$\mathcal{H}_\Omega := \{(v, v_\Gamma) \in \mathcal{H} : v^\Omega = 0\}, \quad \mathcal{V}_\Omega := \mathcal{H}_\Omega \cap \mathcal{V}, \quad (4.11)$$

and we define on \mathcal{H}_Ω and \mathcal{V}_Ω the inner products

$$((u, u_\Gamma), (v, v_\Gamma))_{\mathcal{H}_\Omega} := ((u, u_\Gamma), (v, v_\Gamma))_{\mathcal{H}} = \int_\Omega u v \, dx + \int_\Gamma u_\Gamma v_\Gamma \, d\Gamma, \quad (4.12)$$

$$((u, u_\Gamma), (v, v_\Gamma))_{\mathcal{V}_\Omega} := \int_\Omega \nabla u \cdot \nabla v \, dx + \int_\Gamma \nabla_\Gamma u_\Gamma \cdot \nabla_\Gamma v_\Gamma \, d\Gamma, \quad (4.13)$$

where $(u, u_\Gamma), (v, v_\Gamma)$ are generic elements of \mathcal{H}_Ω (resp., \mathcal{V}_Ω). Note that it follows from Poincaré's inequality (1.19) that (4.13) actually defines an inner product in \mathcal{V}_Ω whose associated norm is equivalent to the standard one.

Next, we infer from [8, Lemma 5.1 and Cor. 5.3] that

$$V_\Gamma = \{v_\Gamma : (v, v_\Gamma) \in \mathcal{V}_\Omega\}, \quad \text{and } \mathcal{V}_\Omega \text{ is dense in } \mathcal{H}_\Omega. \quad (4.14)$$

Therefore, we can construct the Hilbert triple $\mathcal{V}_\Omega \subset \mathcal{H}_\Omega \subset \mathcal{V}_\Omega^*$ with dense and compact embeddings, that is, we identify \mathcal{H}_Ω with a subspace of \mathcal{V}_Ω^* in such a way that

$$\langle (u, u_\Gamma), (v, v_\Gamma) \rangle_{\mathcal{V}_\Omega} = ((u, u_\Gamma), (v, v_\Gamma))_{\mathcal{H}_\Omega} \quad \forall (u, u_\Gamma) \in \mathcal{H}_\Omega, \quad \forall (v, v_\Gamma) \in \mathcal{V}_\Omega. \quad (4.15)$$

Observe that, because of the zero mean value condition, the first components v of the elements $(v, v_\Gamma) \in \mathcal{V}_\Omega$ cannot span the whole space $C_0^\infty(\Omega)$; consequently, variational equalities with test functions in \mathcal{V}_Ω cannot immediately be interpreted as equations in the sense of distributions. We obviously have the following result:

Lemma 4.1: *Let the general assumptions (A1)–(A6) and (1.10), (1.12) be satisfied. Then the pair $(q, q_\Gamma) = (q^\alpha, q_\Gamma^\alpha)$ is a solution to the variational system*

$$\begin{aligned} & - \int_\Omega \partial_t (\mathcal{N}(q(t)) + q(t)) v \, dx + \int_\Omega \nabla q(t) \cdot \nabla v \, dx + \int_\Gamma \nabla_\Gamma q_\Gamma(t) \cdot \nabla_\Gamma v_\Gamma \, d\Gamma \\ & - \int_\Gamma \partial_t q_\Gamma v_\Gamma \, d\Gamma + \int_\Omega (\varphi(\alpha) h''(\bar{y}^\alpha(t)) + f_2''(\bar{y}^\alpha(t))) q(t) v \, dx \\ & + \int_\Gamma (\psi(\alpha) h''(\bar{y}_\Gamma^\alpha(t)) + g_2''(\bar{y}_\Gamma^\alpha(t))) q_\Gamma(t) v_\Gamma \, d\Gamma \\ & = \int_\Omega \beta_1(\bar{y}^\alpha(t) - z_Q(t)) v \, dx + \int_\Gamma \beta_2(\bar{y}_\Gamma^\alpha(t) - z_\Sigma(t)) v_\Gamma \, d\Gamma \\ & \text{for every } (v, v_\Gamma) \in \mathcal{V}_\Omega \quad \text{and for a. a. } t \in (0, T), \end{aligned} \quad (4.16)$$

$$\int_\Omega (\mathcal{N}(q) + q)(T) v \, dx + \int_\Gamma q_\Gamma(T) v_\Gamma \, d\Gamma = 0 \quad \text{for every } (v, v_\Gamma) \in \mathcal{V}_\Omega. \quad (4.17)$$

Notice that we may insert $(v, v_\Gamma) = (q^\alpha(T), q_\Gamma^\alpha(T)) \in \mathcal{V}_\Omega$ in the end point condition (4.17), which, in view of (2.28), yields that

$$\|q^\alpha(T)\|_*^2 + \|q^\alpha(T)\|_H^2 + \|q_\Gamma^\alpha(T)\|_{H_\Gamma}^2 = 0;$$

we thus may replace (4.17) by the simpler condition

$$q^\alpha(T) = 0 \quad \text{a. e. in } \Omega, \quad \mathcal{N}(q^\alpha(T)) = 0 \quad \text{a. e. in } \Omega, \quad q_\Gamma^\alpha(T) = 0 \quad \text{a. e. on } \Gamma, \quad (4.18)$$

where the second equation simply follows from the fact that $q^\alpha(T)$ belongs to the domain of the operator \mathcal{N} .

Remark 4.2: In [8, Theorems 2.5 and 5.4] it has been shown that there is only one solution to problem (4.16)–(4.17) (namely, $(q^\alpha, q_\Gamma^\alpha)$) that has zero mean value and belongs to \mathcal{Y} .

We now prove an a priori estimate which will be fundamental for the derivation of the optimality conditions for (\mathcal{P}_0) . To this end, we introduce some further function spaces. At first, we put

$$\mathcal{W} := H^1(0, T; \mathcal{V}_\Omega^*) \cap L^2(0, T; \mathcal{V}_\Omega). \quad (4.19)$$

Then we define

$$\mathcal{W}_0 := \{(\eta, \eta_\Gamma) \in \mathcal{W} : (\eta(0), \eta_\Gamma(0)) = (0, 0)\}. \quad (4.20)$$

Observe that both these spaces are Banach spaces when equipped with the natural norm of \mathcal{W} . Moreover, \mathcal{W} is continuously embedded in $C^0([0, T]; \mathcal{H}_\Omega)$, so that the initial condition encoded in (4.20) is meaningful. We also point out that \mathcal{W}_0 is dense in $L^2(0, T; \mathcal{V}_\Omega)$ for it contains the dense subspace $H_0^1(0, T; \mathcal{V}_\Omega)$. Therefore, the dual space $L^2(0, T; \mathcal{V}_\Omega^*) = (L^2(0, T; \mathcal{V}_\Omega))^*$ can be identified with a subspace of the dual space \mathcal{W}_0^* in the usual way, i.e., in order that

$$\begin{aligned} \langle\langle (z, z_\Gamma), (\eta, \eta_\Gamma) \rangle\rangle &= \int_0^T \langle (z(t), z_\Gamma(t)), (\eta(t), \eta_\Gamma(t)) \rangle_{\mathcal{V}_\Omega} dt \\ &\text{for all } (z, z_\Gamma) \in L^2(0, T; \mathcal{V}_\Omega^*) \text{ and } (\eta, \eta_\Gamma) \in \mathcal{W}_0, \end{aligned} \quad (4.21)$$

where $\langle\langle \cdot, \cdot \rangle\rangle$ denotes the duality pairing between \mathcal{W}_0^* and \mathcal{W}_0 . Next, we put

$$\mathcal{Z} := L^\infty(0, T; \mathcal{H}) \cap L^2(0, T; \mathcal{V}), \quad (4.22)$$

which is a Banach space when equipped with its natural norm.

Proposition 4.3: *Let the general assumptions (A1)–(A6) and (1.10), (1.12) be satisfied and let*

$$(\lambda^\alpha, \lambda_\Gamma^\alpha) := (\varphi(\alpha) h''(\bar{y}^\alpha) q^\alpha, \psi(\alpha) h''(\bar{y}_\Gamma^\alpha) q_\Gamma^\alpha) \quad \forall \alpha \in (0, 1]. \quad (4.23)$$

Then there exists a constant $K_2^ > 0$, which only depends on the data of the system and on R , such that for all $\alpha \in (0, 1]$ it holds*

$$\begin{aligned} &\|(q^\alpha, q_\Gamma^\alpha)\|_{\mathcal{Z}} + \|(\lambda^\alpha, \lambda_\Gamma^\alpha)\|_{\mathcal{W}_0^*} + \|\mathcal{N}(q^\alpha)\|_{L^\infty(0, T; H^2(\Omega)) \cap L^2(0, T; H^3(\Omega))} \\ &+ \left\| (\mathcal{N}(q^\alpha))|_\Gamma \right\|_{L^\infty(0, T; H^{3/2}(\Gamma)) \cap L^2(0, T; H^{5/2}(\Gamma))} \\ &+ \|(\partial_t(\mathcal{N}(q^\alpha) + q^\alpha), \partial_t((\mathcal{N}(q^\alpha))|_\Gamma + q_\Gamma^\alpha))\|_{\mathcal{W}_0^*} \leq K_2^*. \end{aligned} \quad (4.24)$$

PROOF: In the following, $C_i, i \in \mathbb{N}$, denote positive constants which are independent of $\alpha \in (0, 1]$. To show the boundedness of the adjoint variables, we insert $(v, v_\Gamma) = (q^\alpha(t), q_\Gamma^\alpha(t)) \in \mathcal{V}_\Omega$ in (4.16), written for $(q, q_\Gamma) = (q^\alpha, q_\Gamma^\alpha)$, and integrate over $[s, T]$ where $s \in [0, T]$. First,

note that

$$\begin{aligned}
& - \int_s^T \int_{\Omega} \partial_t (\mathcal{N}(q^\alpha) + q^\alpha) q^\alpha \, dx \, dt = \int_{\Omega} \left(\mathcal{N}(q^\alpha(s)) q^\alpha(s) + \frac{1}{2} |q^\alpha(s)|^2 \right) dx \\
& + \int_s^T \int_{\Omega} \mathcal{N}(q^\alpha) \partial_t q^\alpha \, dx \, dt = \frac{1}{2} (\|q^\alpha(s)\|_H^2 + \|q^\alpha(s)\|_*^2)
\end{aligned} \tag{4.25}$$

since $\partial_t \mathcal{N}(q^\alpha) \in L^2(0, T; H^2(\Omega))$ by (4.4) and (4.10), and the integration by parts with respect to time can be done in view of (2.28), (2.30), and (4.18). We thus obtain the equation

$$\begin{aligned}
& \frac{1}{2} (\|q^\alpha(s)\|_H^2 + \|q^\alpha(s)\|_*^2 + \|q_\Gamma^\alpha(s)\|_{H_\Gamma}^2) + \int_s^T \int_{\Omega} |\nabla q^\alpha|^2 \, dx \, dt \\
& + \int_s^T \int_{\Gamma} |\nabla_\Gamma q_\Gamma^\alpha|^2 \, d\Gamma \, dt + \int_s^T \int_{\Omega} \lambda^\alpha q^\alpha \, dx \, dt + \int_s^T \int_{\Omega} \lambda_\Gamma^\alpha q_\Gamma^\alpha \, d\Gamma \, dt \\
& = - \int_s^T \int_{\Omega} f_2''(\bar{y}^\alpha) |q^\alpha|^2 \, dx \, dt + \int_s^T \int_{\Gamma} g_2''(\bar{y}_\Gamma^\alpha) |q_\Gamma^\alpha|^2 \, d\Gamma \, dt \\
& + \int_s^T \int_{\Omega} \beta_1 (\bar{y}^\alpha - z_Q) q^\alpha \, dx \, dt + \int_s^T \int_{\Gamma} \beta_2 (\bar{y}_\Gamma^\alpha - z_\Sigma) q_\Gamma^\alpha \, d\Gamma \, dt.
\end{aligned} \tag{4.26}$$

By (4.23) and the positivity of h'' , the last two integrals in the second line of the left-hand side of (4.26) are nonnegative, while, owing to (2.33) and **(A1)**, the right-hand side of (4.26) can obviously be bounded by an expression of the form

$$C_1 \left(1 + \int_s^T \int_{\Omega} |q^\alpha|^2 \, dx \, dt + \int_s^T \int_{\Gamma} |q_\Gamma^\alpha|^2 \, d\Gamma \, dt \right).$$

Hence, invoking Gronwall's inequality, we find the estimate

$$\|(q^\alpha, q_\Gamma^\alpha)\|_{L^\infty(0, T; \mathcal{H}) \cap L^2(0, T; \mathcal{V})} \leq C_2 \quad \forall \alpha \in (0, 1]. \tag{4.27}$$

Moreover, first using (2.25) and then the trace theorem, we find that

$$\|\mathcal{N}(q^\alpha)\|_{L^\infty(0, T; H^2(\Omega)) \cap L^2(0, T; H^3(\Omega))} \leq C_3 \quad \forall \alpha \in (0, 1], \tag{4.28}$$

$$\|(\mathcal{N}(q^\alpha))|_\Gamma\|_{L^\infty(0, T; H^{3/2}(\Gamma)) \cap L^2(0, T; H^{5/2}(\Gamma))} \leq C_4 \quad \forall \alpha \in (0, 1]. \tag{4.29}$$

Next, we derive the bound for the time derivatives. To this end, let $(\eta, \eta_\Gamma) \in \mathcal{W}_0$ be arbitrary. Using (4.18), the initial condition for (η, η_Γ) , and the estimates (4.27)–(4.29), we obtain from

integration by parts and (4.21) that

$$\begin{aligned}
& \langle\langle -(\partial_t(\mathcal{N}(q^\alpha) + q^\alpha), \partial_t((\mathcal{N}(q^\alpha))_{|\Gamma} + q_\Gamma^\alpha)), (\eta, \eta_\Gamma) \rangle\rangle \\
&= -\int_0^T \int_\Omega \partial_t(\mathcal{N}(q^\alpha) + q^\alpha) \eta \, dx \, dt - \int_0^T \int_\Gamma \partial_t((\mathcal{N}(q^\alpha))_{|\Gamma} + q_\Gamma^\alpha) \eta_\Gamma \, d\Gamma \, dt \\
&= \int_0^T \langle \partial_t \eta(t), \mathcal{N}(q^\alpha(t)) + q^\alpha(t) \rangle_V \, dt + \int_0^T \langle \partial_t \eta_\Gamma(t), (\mathcal{N}(q^\alpha(t)) + q^\alpha(t)) \rangle_{V_\Gamma} \, dt \\
&\leq \int_0^T \|\partial_t \eta(t)\|_{V^*} \|\mathcal{N}(q^\alpha(t)) + q^\alpha(t)\|_V \, dt \\
&\quad + \int_0^T \|\partial_t \eta_\Gamma(t)\|_{V_\Gamma^*} \|(\mathcal{N}(q^\alpha(t))_{|\Gamma} + q_\Gamma^\alpha(t))\|_{V_\Gamma} \, dt \\
&\leq C_5 \|(\eta, \eta_\Gamma)\|_{\mathcal{W}_0^*}, \quad \text{for all } \alpha \in (0, 1]. \tag{4.30}
\end{aligned}$$

We thus have shown that

$$\|(\partial_t(\mathcal{N}(q^\alpha) + q^\alpha), \partial_t((\mathcal{N}(q^\alpha))_{|\Gamma} + q_\Gamma^\alpha))\|_{\mathcal{W}_0^*} \leq C_5 \quad \forall \alpha \in (0, 1]. \tag{4.31}$$

Finally, by recalling (4.23) and the estimates (4.27)–(4.29), (4.31), a comparison in (4.16) yields that

$$\|(\lambda^\alpha, \lambda_\Gamma^\alpha)\|_{\mathcal{W}_0^*} \leq C_6 \quad \forall \alpha \in (0, 1] \tag{4.32}$$

as well, and the assertion is proved. \blacksquare

4.2 The optimality conditions for (\mathcal{P}_0)

We now establish first-order necessary optimality conditions for (\mathcal{P}_0) by performing a limit as $\alpha \searrow 0$ in the approximating problems. To this end, recall that a fixed optimal control $\bar{u}_\Gamma \in \mathcal{U}_{\text{ad}}$ for (\mathcal{P}_0) , along with a solution quintuple $(\bar{y}, \bar{y}_\Gamma, \bar{w}, \bar{\xi}, \bar{\xi}_\Gamma)$ of the associated state system (1.3)–(1.7) is given.

We draw some consequences from the previously established results. First recall that by Theorem 3.5 for any sequence $\{\alpha_n\} \subset (0, 1]$ with $\alpha_n \searrow 0$ as $n \rightarrow \infty$, and for any $n \in \mathbb{N}$ we can find an optimal control $\bar{u}_\Gamma^{\alpha_n} \in \mathcal{U}_{\text{ad}}$ for $(\tilde{\mathcal{P}}_{\alpha_n})$ and an associated state triple $(\bar{y}^{\alpha_n}, \bar{y}_\Gamma^{\alpha_n}, \bar{w}^{\alpha_n})$ such that the convergences (3.18)–(3.21) hold. As in the proof of Theorem 3.1, we may without loss of generality assume that

$$f_2''(\bar{y}^{\alpha_n}) \rightarrow f_2''(\bar{y}) \quad \text{strongly in } C^0([0, T]; H), \tag{4.33}$$

$$g_2''(\bar{y}_\Gamma^{\alpha_n}) \rightarrow g_2''(\bar{y}_\Gamma) \quad \text{strongly in } C^0([0, T]; H_\Gamma). \tag{4.34}$$

Also, by virtue of Lemma 4.1 and Proposition 4.3, we may without loss of generality assume that there exist the corresponding adjoint state variables $(q^{\alpha_n}, q_{\Gamma}^{\alpha_n}) \in \mathcal{Y}$ that satisfy

$$(q^{\alpha_n}, q_{\Gamma}^{\alpha_n}) \rightarrow (q, q_{\Gamma}) \quad \text{weakly-star in } \mathcal{Z}, \quad (4.35)$$

$$\mathcal{N}(q^{\alpha_n}) \rightarrow \mathcal{N}(q) \quad \text{weakly-star in } L^{\infty}(0, T; H^2(\Omega)) \cap L^2(0, T; H^3(\Omega)), \quad (4.36)$$

$$(\mathcal{N}(q^{\alpha_n}))|_{\Gamma} \rightarrow (\mathcal{N}(q))|_{\Gamma} \quad \text{weakly-star in } L^{\infty}(0, T; H^{3/2}(\Gamma)) \cap L^2(0, T; H^{5/2}(\Gamma)), \quad (4.37)$$

$$(\lambda^{\alpha_n}, \lambda_{\Gamma}^{\alpha_n}) \rightarrow (\lambda, \lambda_{\Gamma}) \quad \text{weakly in } \mathcal{W}_0(0, T)^*, \quad (4.38)$$

for suitable limits (q, q_{Γ}) and $(\lambda, \lambda_{\Gamma})$. Therefore, passing to the limit as $n \rightarrow \infty$ in the variational inequality (4.9), written for α_n , $n \in \mathbb{N}$, and recalling (3.18), we obtain that (q, q_{Γ}) satisfies

$$\int_0^T \int_{\Gamma} (q_{\Gamma} + \beta_5 \bar{u}_{\Gamma}) (v_{\Gamma} - \bar{u}_{\Gamma}) \, d\Gamma \, dt \geq 0 \quad \forall v_{\Gamma} \in \mathcal{U}_{\text{ad}}. \quad (4.39)$$

Next, we will show that in the limit as $n \rightarrow \infty$ a limiting adjoint system for (\mathcal{P}_0) is satisfied. To this end, we insert an arbitrary $(\eta, \eta_{\Gamma}) \in \mathcal{W}_0(0, T)$ in (4.16), written for α_n , $n \in \mathbb{N}$, and integrate the resulting equation over $[0, T]$. Integrating by parts with respect to t , and invoking (4.18) and the zero initial conditions for (η, η_{Γ}) , we arrive at the identity

$$\begin{aligned} & \int_0^T \int_{\Omega} \lambda^{\alpha_n} \eta \, dx \, dt + \int_0^T \int_{\Gamma} \lambda_{\Gamma}^{\alpha_n} \eta_{\Gamma} \, d\Gamma \, dt + \int_0^T \langle \partial_t \eta(t), \mathcal{N}(q^{\alpha_n}(t)) + q^{\alpha_n}(t) \rangle_V \, dt \\ & + \int_0^T \langle \partial_t \eta_{\Gamma}(t), q_{\Gamma}^{\alpha_n}(t) \rangle_{V_{\Gamma}} \, dt + \int_0^T \int_{\Omega} \nabla q^{\alpha_n} \cdot \nabla \eta \, dx \, dt + \int_0^T \int_{\Gamma} \nabla_{\Gamma} q_{\Gamma}^{\alpha_n} \cdot \nabla_{\Gamma} \eta_{\Gamma} \, d\Gamma \, dt \\ & + \int_0^T \int_{\Omega} f_2''(\bar{y}^{\alpha_n}) q^{\alpha_n} \eta \, dx \, dt + \int_0^T \int_{\Gamma} g_2''(\bar{y}_{\Gamma}^{\alpha_n}) q_{\Gamma}^{\alpha_n} \eta_{\Gamma} \, d\Gamma \, dt \\ & = \beta_1 \int_0^T \int_{\Omega} (\bar{y}^{\alpha_n} - z_Q) \eta \, dx \, dt + \beta_2 \int_0^T \int_{\Gamma} (\bar{y}_{\Gamma}^{\alpha_n} - z_{\Sigma}) \eta_{\Gamma} \, d\Gamma \, dt. \end{aligned} \quad (4.40)$$

Now, by virtue of the convergences (3.19), (3.20), and (4.33)–(4.38), we may pass to the limit as $n \rightarrow \infty$ in (4.40) to obtain, for all $(\eta, \eta_{\Gamma}) \in \mathcal{W}_0(0, T)$,

$$\begin{aligned} & \langle (\lambda, \lambda_{\Gamma}), (\eta, \eta_{\Gamma}) \rangle + \int_0^T \langle \partial_t \eta(t), \mathcal{N}(q(t)) + q(t) \rangle_V \, dt + \int_0^T \langle \partial_t \eta_{\Gamma}(t), q_{\Gamma}(t) \rangle_{V_{\Gamma}} \, dt \\ & + \int_0^T \int_{\Omega} \nabla q \cdot \nabla \eta \, dx \, dt + \int_0^T \int_{\Gamma} \nabla_{\Gamma} q_{\Gamma} \cdot \nabla_{\Gamma} \eta_{\Gamma} \, d\Gamma \, dt \\ & + \int_0^T \int_{\Omega} f_2''(\bar{y}) q \eta \, dx \, dt + \int_0^T \int_{\Gamma} g_2''(\bar{y}_{\Gamma}) q_{\Gamma} \eta_{\Gamma} \, d\Gamma \, dt \\ & = \beta_1 \int_0^T \int_{\Omega} (\bar{y} - z_Q) \eta \, dx \, dt + \beta_2 \int_0^T \int_{\Gamma} (\bar{y}_{\Gamma} - z_{\Sigma}) \eta_{\Gamma} \, d\Gamma \, dt. \end{aligned} \quad (4.41)$$

Next, we show that the limit pair $((\lambda, \lambda_\Gamma), (q, q_\Gamma))$ satisfies some sort of a complementarity slackness condition. To this end, observe that for all $n \in \mathbb{N}$ we obviously have

$$\int_0^T \int_\Omega \lambda^{\alpha_n} q^{\alpha_n} dx dt = \int_0^T \int_\Omega \varphi(\alpha_n) h''(\bar{y}^{\alpha_n}) |q^{\alpha_n}|^2 dx dt \geq 0.$$

An analogous inequality holds for the corresponding boundary terms. We thus have

$$\liminf_{n \rightarrow \infty} \int_0^T \int_\Omega \lambda^{\alpha_n} q^{\alpha_n} dx dt \geq 0, \quad \liminf_{n \rightarrow \infty} \int_0^T \int_\Gamma \lambda_\Gamma^{\alpha_n} q_\Gamma^{\alpha_n} d\Gamma dt \geq 0. \quad (4.42)$$

Finally, we derive a relation which gives some indication that the limit $(\lambda, \lambda_\Gamma)$ should somehow be concentrated on the set where $|\bar{y}| = 1$ and $|\bar{y}_\Gamma| = 1$ (which, however, we cannot prove rigorously). To this end, we test the pair $(\lambda^{\alpha_n}, \lambda_\Gamma^{\alpha_n})$ by the function $((1 - (\bar{y}^{\alpha_n})^2) \phi, (1 - (\bar{y}_\Gamma^{\alpha_n})^2) \phi_\Gamma)$ that belongs to \mathcal{V}_Ω since (ϕ, ϕ_Γ) is any smooth test function satisfying

$$(\phi(0), \phi_\Gamma(0)) = (0, 0), \quad \int_\Omega (1 - (\bar{y}^{\alpha_n}(t))^2) \phi(t) dx = 0 \quad \forall t \in [0, T]. \quad (4.43)$$

As $h''(r) = 2/(1 - r^2)$ for every $r \in (-1, 1)$, we obtain

$$\begin{aligned} & \lim_{n \rightarrow \infty} \left(\int_0^T \int_\Omega \lambda^{\alpha_n} (1 - (\bar{y}^{\alpha_n})^2) \phi dx dt, \int_0^T \int_\Gamma \lambda_\Gamma^{\alpha_n} (1 - (\bar{y}_\Gamma^{\alpha_n})^2) \phi_\Gamma d\Gamma dt \right) \\ &= \lim_{n \rightarrow \infty} \left(2 \int_0^T \int_\Omega \varphi(\alpha_n) q^{\alpha_n} \phi dx dt, 2 \int_0^T \int_\Gamma \psi(\alpha_n) q_\Gamma^{\alpha_n} \phi_\Gamma d\Gamma dt \right) = (0, 0). \end{aligned} \quad (4.44)$$

We now collect the results established above, especially in Theorem 3.5. We have the following statement.

Theorem 4.4: *Let the assumptions (A1)–(A6) be satisfied, let h be given by (1.10), and let φ, ψ be positive and continuous functions on $(0, 1]$ fulfilling (1.12). Moreover, let $\bar{u}_\Gamma \in \mathcal{U}_{\text{ad}}$ be an optimal control for (\mathcal{P}_0) with associated solution quintuple $(\bar{y}, \bar{y}_\Gamma, \bar{w}, \bar{\xi}, \bar{\xi}_\Gamma)$ to the corresponding state system (1.3)–(1.7) in the sense of Definition 2.1. Then the following assertions hold true:*

(i) *For every sequence $\{\alpha_n\} \subset (0, 1]$, with $\alpha_n \searrow 0$ as $n \rightarrow \infty$, and for any $n \in \mathbb{N}$, there exists a solution $\bar{u}_\Gamma^{\alpha_n} \in \mathcal{U}_{\text{ad}}$ to the adapted control problem $(\tilde{\mathcal{P}}_{\alpha_n})$ such that, with the associated solution triple $(\bar{y}^{\alpha_n}, \bar{y}_\Gamma^{\alpha_n}, \bar{w}^{\alpha_n})$ of the corresponding state system (2.13)–(2.16), the convergences (3.18)–(3.21) hold as $n \rightarrow \infty$.*

(ii) *Whenever sequences $\{\alpha_n\} \subset (0, 1]$ and $\{(\bar{y}^{\alpha_n}, \bar{y}_\Gamma^{\alpha_n}, \bar{u}_\Gamma^{\alpha_n})\}$ having the properties described in (i) are given, then the following holds true: to any subsequence $\{n_k\}_{k \in \mathbb{N}}$ of \mathbb{N} there are a subsequence $\{n_{k_\ell}\}_{\ell \in \mathbb{N}}$ and some $((\lambda, \lambda_\Gamma), (q, q_\Gamma)) \in \mathcal{W}_0(0, T)^* \times \mathcal{Z}$ such that*

- the relations (4.35)–(4.38), (4.42), and (4.44) hold (where the sequences are indexed by n_{k_ℓ} and the limits are taken as $\ell \rightarrow \infty$), and
- the variational inequality (4.39) and the adjoint equation (4.41) are satisfied.

Remark 4.5: Unfortunately, we are not able to show that the limit pair (q, q_Γ) solving the adjoint problem associated with the optimal triple $(\bar{y}, \bar{y}_\Gamma, \bar{u}_\Gamma)$ is uniquely determined. Therefore, it may well happen that the limiting pairs differ for different subsequences. However, it follows from the variational inequality (4.39) that for any such limit pair (q, q_Γ) it holds, with the orthogonal projection $\mathbb{P}_{\mathcal{U}_{\text{ad}}}$ onto \mathcal{U}_{ad} with respect to the standard inner product in H_Γ , that for $\beta_5 > 0$ we have

$$\bar{u}_\Gamma = \mathbb{P}_{\mathcal{U}_{\text{ad}}}(-\beta_5^{-1}q_\Gamma) . \quad (4.45)$$

Standard arguments then yield that if the function $\bar{u}_\Gamma \in L^2(\Sigma)$ defined by

$$\bar{u}_\Gamma(x, t) = \begin{cases} \tilde{u}_{2_\Gamma}(x, t) & \text{if } -\beta_5^{-1}q_\Gamma(x, t) > \tilde{u}_{2_\Gamma}(x, t) \\ \tilde{u}_{1_\Gamma}(x, t) & \text{if } -\beta_5^{-1}q_\Gamma(x, t) < \tilde{u}_{1_\Gamma}(x, t) \\ -\beta_5^{-1}q_\Gamma(x, t) & \text{otherwise} \end{cases} \quad \text{for a. a. } (x, t) \in \Sigma , \quad (4.46)$$

belongs to \mathcal{U}_{ad} (i.e., its time derivative actually exists and satisfies the bound prescribed in (1.8)), then $\bar{u}_\Gamma = \bar{u}_\Gamma$ and \bar{u}_Γ turns out to be a pointwise projection.

References

- [1] V. BARBU, Necessary conditions for nonconvex distributed control problems governed by elliptic variational inequalities, *J. Math. Anal. Appl.* **80** (1981) 566–597.
- [2] H. BREZIS, “Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert”, North-Holland Math. Stud. **5**, North-Holland, Amsterdam, 1973.
- [3] F. BREZZI, G. GILARDI, Chapters 1-3 in “Finite Element Handbook”, H. Kardestuncer and D. H. Norrie (eds.), McGraw-Hill Book Co., New York, 1987.
- [4] L. CALATRONI, P. COLLI, Global solution to the Allen–Cahn equation with singular potentials and dynamic boundary conditions, *Nonlinear Anal.* **79** (2013) 12–27.
- [5] P. COLLI, G. GILARDI, P. PODIO-GUIDUGLI, J. SPREKELS, Distributed optimal control of a nonstandard system of phase field equations, *Cont. Mech. Thermodyn.* **24** (2012) 437–459.
- [6] P. COLLI, G. GILARDI, J. SPREKELS, Analysis and boundary control of a nonstandard system of phase field equations, *Milan J. Math.* **80** (2012) 119–149.
- [7] P. COLLI, G. GILARDI, J. SPREKELS, On the Cahn–Hilliard equation with dynamic boundary conditions and a dominating boundary potential, *J. Math. Anal. Appl.* **419** (2014) 972–994.

- [8] P. COLLI, G. GILARDI, J. SPREKELS, A boundary control problem for the viscous Cahn–Hilliard equation with dynamic boundary conditions, preprint arXiv:1407.3916 [math.AP] (2014), pp. 1–27, submitted.
- [9] P. COLLI, M. H. FARSHBAF-SHAKER, J. SPREKELS, A deep quench approach to the optimal control of an Allen–Cahn equation with dynamic boundary conditions and double obstacles, *Appl. Math. Optim.*, published online May 2014, DOI 10.1007/s00245-014-9250-8 (see also WIAS Preprint No. 1838 (2013), pp. 1–23).
- [10] P. COLLI, J. SPREKELS, Optimal control of an Allen–Cahn equation with singular potentials and dynamic boundary condition, preprint arXiv:1212.2359 [math.AP] (2012), pp. 1–24, submitted.
- [11] M. H. FARSHBAF-SHAKER, A penalty approach to optimal control of Allen–Cahn variational inequalities: MPEC-view, *Numer. Funct. Anal. Optim.* **33** (2012) 1321–1349.
- [12] M. H. FARSHBAF-SHAKER, A relaxation approach to vector-valued Allen–Cahn MPEC problems, Universität Regensburg, Preprintreihe der Fakultät für Mathematik, No. 27 (2011), pp. 1–27.
- [13] G. GILARDI, A. MIRANVILLE, G. SCHIMPERNA, On the Cahn–Hilliard equation with irregular potentials and dynamic boundary conditions, *Commun. Pure Appl. Anal.* **8** (2009) 881–912.
- [14] G. GILARDI, A. MIRANVILLE, G. SCHIMPERNA, Long-time behavior of the Cahn–Hilliard equation with irregular potentials and dynamic boundary conditions, *Chin. Ann. Math. Ser. B* **31** (2010) 679–712.
- [15] M. HINTERMÜLLER, D. WEGNER, Distributed optimal control of the Cahn–Hilliard system including the case of a double-obstacle homogeneous free energy density, *SIAM J. Control Optim.* **50** (2012) 388–418.
- [16] M. HINTERMÜLLER, D. WEGNER, Optimal control of a semi-discrete Cahn–Hilliard–Navier–Stokes system, *SIAM J. Control Optim.* **52** (2014) 747–772.
- [17] M. HINZE, C. KAHLE, A nonlinear predictive concept for control of two-phase flows governed by the Cahn–Hilliard Navier–Stokes system, *System Modeling and Optimization*, 25th IFIP TC 7 Conference 2011, IFIP AICT 391 (2012) 348–357.
- [18] P. PODIO-GUIDUGLI, Models of phase segregation and diffusion of atomic species on a lattice, *Ric. Mat.* **55** (2006) 105–118.
- [19] E. ROCCA, J. SPREKELS, Optimal distributed control of a nonlocal convective Cahn–Hilliard equation by the velocity in 3D, WIAS Preprint No. 1942 (2014), pp. 1–24, submitted.
- [20] J. SIMON, Compact sets in the space $L^p(0, T; B)$, *Ann. Mat. Pura Appl. (4)* **146** (1987) 65–96.
- [21] Q.-F. WANG, S.-I. NAKAGIRI, Optimal control of distributed parameter system given by Cahn–Hilliard equation, *Nonlinear Funct. Anal. Appl.* **19** (2014) 19–33.

- [22] X. P. ZHAO, C. C. LIU, Optimal control of the convective Cahn–Hilliard equation, *Appl. Anal.* **92** (2013) 1028–1045.
- [23] X. P. ZHAO, C. C. LIU, Optimal control for the convective Cahn–Hilliard equation in 2D case, *Appl. Math. Optim.* **70** (2014) 61–82.