

Weierstraß-Institut
für Angewandte Analysis und Stochastik
Leibniz-Institut im Forschungsverbund Berlin e. V.

Preprint

ISSN 2198-5855

Exit time risk-sensitive control for systems of cooperative agents

Paul Dupuis¹, Vaios Laschos², Kavita Ramanan¹

submitted: June 14, 2017 (revision: February 12, 2018)

¹ Division of Applied Mathematics
Brown University
182 George Street
Providence, RI 02912
USA
E-Mail: Paul_Dupuis@Brown.edu
Kavita_Ramanan@Brown.edu

² Weierstrass Institute
Mohrenstr. 39
10117 Berlin
Germany
E-Mail: vaios.laschos@wias-berlin.de
(WIAS member until December 31, 2017)

No. 2407
Berlin 2018



2010 *Mathematics Subject Classification.* 93E20.

Key words and phrases. Risk-sensitive stochastic control, mean-field interaction, min-max theorems.

Edited by
Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS)
Leibniz-Institut im Forschungsverbund Berlin e. V.
Mohrenstraße 39
10117 Berlin
Germany

Fax: +49 30 20372-303
E-Mail: preprint@wias-berlin.de
World Wide Web: <http://www.wias-berlin.de/>

Exit time risk-sensitive control for systems of cooperative agents

Paul Dupuis, Vaios Laschos, Kavita Ramanan

Abstract

We study sequences, parametrized by the number of agents, of exit time stochastic control problems with risk-sensitive costs structures generate by unbounded costs. We identify a fully characterizing assumption, under which, each of them corresponds to a risk-neutral stochastic control problem with additive cost, and also to a risk-neutral stochastic control problem on the simplex, where the specific information about the state of each agent can be discarded. We finally prove that, under some additional assumptions, the sequence of value functions converges to the value function of a deterministic control problem, which can be used for the design of nearly optimal controls for the original problem, when the number of agents is sufficiently large.

1 Introduction

In this paper, we study many agent exit time stochastic control problems with risk-sensitive cost. The reader with background in physics or chemistry can consider particles in place of agents. Each agent occupies a state that takes values in a finite set \mathcal{X} , and by controlling the transition rates between states for each agent, we try to keep the system away from a “ruin” set \mathcal{K} , for as long as possible and with the least cost. We prove, under suitable assumptions, that for every finite number n of agents the control problem is equivalent to one with an ordinary (additive) cost. Moreover, when $\mathcal{K} \subset \mathcal{X}^n$ can be identified with a subset of the simplex of probability measures $\mathcal{P}(\mathcal{X})$ (in the sense that for every permutation $\sigma \in \mathbb{S}_n$ we have $\sigma\mathcal{K} = \mathcal{K}$), then we can replace the original problem by one on $\mathcal{P}^n(\mathcal{X}) = \mathcal{P}(\mathcal{X}) \cap \frac{1}{n}\mathbb{Z}^d$, getting in this way a control problem whose state is the empirical measure on the states of the individual agents. We also study the behavior as $n \rightarrow \infty$ of the sequence of suitable renormalized value functions, and prove uniform convergence to the value function of a deterministic control problem.

We first describe the model without control, which we call the “base” or “nominal” model. Let $\mathcal{X} = \{e_1, \dots, e_d\}$, where e_i is the i th unit vector in \mathbb{R}^d . Let also $\gamma = \{\gamma_{xy}\}_{(x,y) \in \mathcal{X} \times \mathcal{X}}$ denote the rates of an ergodic Markov process on \mathcal{X} . This process has the generator

$$\mathcal{L}_\gamma[f](x) = \sum_{y \in \mathcal{X}} \gamma_{xy} [f(y) - f(x)]. \quad (1)$$

For $n \in \mathbb{N}$, consider n agents that independently and randomly occupy different states x_i^n among the elements of $\mathcal{X} = \{e_1, \dots, e_d\}$, and let $\mathbf{x}^n = (x_1^n, \dots, x_n^n)$. This process takes values in \mathcal{X}^n and has the generator

$$\mathcal{L}_\gamma^n[f](\mathbf{x}^n) = \sum_{i=1}^n \sum_{y \in \mathcal{Z}_{x_i^n}} \gamma_{x_i^n y} \left[f(\mathbf{x}^n + \mathbf{v}_{i,x_i^n}^n y) - f(\mathbf{x}^n) \right], \quad (2)$$

Here $\mathcal{Z} \doteq \{(x, y) \in \mathcal{X} \times \mathcal{X} : \gamma_{xy} > 0\}$, $\mathcal{Z}_x \doteq \{y \in \mathcal{X} : (x, y) \in \mathcal{Z}\}$ is the set of allowed transitions from x , and $\mathbf{v}_{i,x_i^n}^n = (0, \dots, 0, \mathbf{v}_{xy}, 0, \dots, 0)$ is a $d \times n$ matrix with all columns equal to zero apart from the i th column, which is identically equal to the vector $\mathbf{v}_{xy} \doteq y - x$. Since the process is ergodic, \mathcal{Z} generates the hyperplane

$$\mathcal{H} \doteq \left\{ \sum_{(x,y) \in \mathcal{Z}} a_{xy} \mathbf{v}_{xy} : a_{xy} > 0, (x, y) \in \mathcal{Z} \right\}, \quad (3)$$

which coincides with the hyperplane through the origin that is parallel to $\mathcal{P}(\mathcal{X})$.

We claim that the set \mathcal{H} does not change if the a_{xy} are allowed to be arbitrary real numbers. By ergodicity, for any two states $(x, y) \in \mathcal{Z}$ there is a sequence of states $x = x_1, \dots, x_j = y$ with $y = x_2$ and with the property that $(x_i, x_{i+1}) \in \mathcal{Z}$ for $i = 1, \dots, j-1$, and hence $\sum_{i=1}^{j-1} \mathbf{v}_{x_i x_{i+1}} = 0$. Repeating this for every possible $(x, y) \in \mathcal{Z}$, there are strictly positive integers b_{xy} such that $\sum_{(x,y) \in \mathcal{Z}} b_{xy} \mathbf{v}_{xy} = 0$, which implies the claim.

Next we introduce the empirical measure process. This process is obtained by projecting from \mathcal{X}^n onto $\mathcal{P}^n(\mathcal{X}) = \mathcal{P}(\mathcal{X}) \cap \frac{1}{n}\mathbb{Z}^d \subset \mathcal{P}(\mathcal{X})$, and has the generator

$$\mathcal{N}_\gamma^n[f](\mathbf{m}) = n \sum_{(x,y) \in \mathcal{Z}} \gamma_{xy} m_x \left[f\left(\mathbf{m} + \frac{1}{n}\mathbf{v}_{xy}\right) - f(\mathbf{m}) \right]. \quad (4)$$

One can interpret the model introduced above as a collection of independent agents with each evolving according to the transition rates γ . This is the “preferred” or “nominal” dynamic, and is what would occur if no “outside influence” or other form of control acts on the agents. If a controller should wish to change this behavior, then it must pay a cost to do so. We would like to model the situation where limited information on the system state, and in particular information relating only to the empirical measure of the states of all agents, is used to produce a desired behavior of the group of agents, which again will be characterized in terms of their empirical measure (which is used to characterize how the collective “loads” the system).

Thus we consider for each $n \in \mathbb{N}$ “reward” functions $R^n : \mathcal{P}(\mathcal{X}) \rightarrow [0, \infty)$, where we recall

$$\mathcal{P}(\mathcal{X}) \doteq \left\{ \mathbf{m} \in \mathbb{R}^{\mathcal{X}} : m_x \geq 0 \text{ for all } x \in \mathcal{X} \text{ and } \sum_{x \in \mathcal{X}} m_x = 1 \right\} \quad (5)$$

is the simplex of probability measures on \mathcal{X} . It is assumed that the R^n are continuous and that they converge uniformly to some R^∞ . We also have a sequence of unbounded “cost” functions $\mathbf{C}^n = \{C_{xy}^n : [0, \infty) \rightarrow [0, \infty]\}_{(x,y) \in \mathcal{Z}}$ that converge on $(0, \infty)$ to some \mathbf{C}^∞ in a sense that will be defined in the sequel. In the controlled setting, the jump rates of each agent can be perturbed from γ to \mathbf{u} . Let χ^n denote the controlled state occupied by the collection of agents, and for $\mathbf{x}^n = \{x_i^n\}_{i \leq n} \in \mathcal{X}^n$ define

$$L(\mathbf{x}^n) \doteq \sum_{i=1}^n \delta_{x_i^n}. \quad (6)$$

If the problem is of interest over the interval $[0, T]$ then there is a collective risk-sensitive cost (paid by the coordinating controller) equal to

$$\mathbb{E}_{\mathbf{x}^n} \left[\exp \left(\int_0^T \left(\sum_{i=1}^n \sum_{y \in \mathcal{Z}_{\chi_i^n(t)}} \gamma_{\chi_i^n(t)y} C_{\chi_i^n(t)y}^n \left(\frac{u_{\chi_i^n(t)y}(t,i)}{\gamma_{\chi_i^n(t)y}} \right) - nR^n(L(\chi^n(t))) \right) dt \right) \right]. \quad (7)$$

Here the control process \mathbf{u} takes values in a space that will be defined later, and for a collection of $n|\mathcal{Z}|$ independent Poisson random measures (PRM) $\{N_{i,xy}^1\}_{1 \leq i \leq n, (x,y) \in \mathcal{Z}}$ with intensity measure equal to Lebesgue measure, the controlled dynamics are given by

$$\chi_i^n(t) = x_i^n + \sum_{(x,y) \in \mathcal{Z}} \mathbf{v}_{xy} \int_{(0,t]} \int_{[0,\infty)} 1_{[0,1_x(\chi_i^n(s-))u_{xy}(s,i)]}(r) N_{i,xy}^1(dsdr). \quad (8)$$

Thus χ_i^n changes from state x to y with rate u_{xy} . The formulation of the dynamics in terms of a stochastic differential equation will be convenient in the analysis to follow.

In this paper we present three results. The first is that, under additional assumptions on the cost \mathbf{C}^n , for each n the risk-sensitive control problem is equivalent to an ordinary control problem the cost function $\mathbf{F}^n = \{F_{xy}^n\}_{(x,y) \in \mathcal{Z}}$, where F_{xy}^n is defined by

$$F_{xy}^n(q) \doteq \sup_{u \in (0,\infty)} G_{xy}^n(u, q) \quad \text{and} \quad G_{xy}^n(u, q) \doteq \left[u \ell \left(\frac{q}{u} \right) - \gamma_{xy} C_{xy}^n \left(\frac{u}{\gamma_{xy}} \right) \right], \quad (9)$$

$$\ell(q) \doteq q \log q - q + 1 \text{ for } q \geq 0. \quad (10)$$

Under the additional conditions we do not end up with a *stochastic game*, as one might expect, but rather a control problem with additive cost. Control problems are often substantially simpler than games, and in particular are often more tractable from a computational perspective.

The last contribution, again under additional assumptions on the sequence of costs $\{\mathbf{C}^n\}_n$, is that the sequence of value functions, suitably renormalized, converges to the value function of a deterministic control problem. This is also helpful in the construction of controls. We work with sequences of cost functions instead of a fixed one, because we would like to note that the limit deterministic problem exists regardless of perturbations that may appear on the cost function when the number of agents change, as long as we have some convergence to a limit cost function.

Example 1 *As an example consider the issue of modeling the users of a resource such as energy. Here the agents would be households or similar entities. The state of an agent indicates their use of the common resource, and this usage evolves in a Markovian fashion. In exchange for a cost paid by the central controller to the individual agents, the agents agree to modify their behavior based on the current loading of the system. Thus an energy consumer would agree to give up control on if or when certain activities take place, thus altering the evolution of their own state, but will be compensated for doing so. The goal of the central controller, and the motivation for paying this cost, is to manage the group behavior so as to keep the system, as characterized by the empirical measure, in a desired operating region for as long as possible and with minimal cost. In this context, the use of risk sensitive cost is motivated in part by the resulting properties of robustness with respect to model error.*

1.1 Literature

For ordinary discrete-time and continuous-time stochastic control problems (also known as Markov decision processes in discrete time [1, 17, 24, 28, 14]), one controls a random processes to optimize an expected cost. The most common optimality criteria for continuous time escape (or ruin) stochastic control problems are of the form

$$J_T(x_0, \pi) = \mathbb{E}_{x_0, \pi} \left[\int_0^T C(X_t, u_t) dt + P(X_T) \right], \quad (11)$$

where C is some cost function that depends on the state $x \in X$ and the control/action $u \in U$, and π is a policy or strategy. For risk-sensitive stochastic control problems one deals with optimality criteria of the form

$$J_T(x_0, \pi) = g^{-1} \left(\mathbb{E}_{x_0, \pi} \left[g \left(\int_0^T C(X_t, U_t) dt + P(X_T) \right) \right] \right), \quad (12)$$

where g is a monotone convex/concave function. One motivation behind the use of risk-sensitive cost structures is that, depending on the type of monotonicity, variation from the average is penalized (risk-averting behavior) or ignored (risk-seeking behavior). One of the most studied cases is the entropic risk measure corresponding to $g_\theta(x) = e^{\theta x}$ (see [2, 5, 6, 7, 8, 13, 16, 18, 20, 21] for discrete time and [9, 11, 15] for continuous time). The function $g_\theta(x) = e^{\theta x}$ is special because it satisfies the property

$$\frac{1}{\theta} \log (\mathbb{E} [\exp (\theta X)]) = \tilde{X} + \frac{1}{\theta} \log \left(\mathbb{E} \left[e^{\theta(X-\tilde{X})} \right] \right)$$

where X is a random variable and \tilde{X} its expectation. This property implies that the weight that is given to deviations from the expectation depends only on the difference from the expectation and not the expectation itself. It can be proved that the exponential is the only function that satisfies such a property (see [27]). Furthermore, exponential integrals have a variational characterization involving entropy, which turns out to be convenient from mathematical point of view, and which also allows for an explicit analysis of the robust and model insensitivity properties of the resulting controls [10, 23]. In our problem θ is integrated in the choice of cost C , and this explains partially our choice to work with sequences of cost functions instead of a fixed cost function, since it shows that it is allowed to tune the value of θ , along with the number of agents, as long as we do not violate the main assumptions and the sequence of θ converges. In recent years, following the seminal work of [25], new risk-sensitive criteria have been studied that allow a mixture of risk-seeking and risk avoiding behavior in different regimes.

2 Notation and definitions

For a locally compact Polish space \mathcal{S} , the space of positive Borel measures on \mathcal{S} is denoted by $\mathcal{M}(\mathcal{S})$. With the subscripts f, c we denote, respectively, the space of finite measures, and the space of measures that are finite on every compact subset. Letting $C_c(\mathcal{S})$ denote the space of continuous functions with compact support, we equip $\mathcal{M}_c(\mathcal{S})$, with the weakest topology such that for every $f \in C_c(\mathcal{S})$, the function $\nu \rightarrow \int_{\mathcal{S}} f d\nu$, $\nu \in \mathcal{M}_c(\mathcal{S})$, is continuous. $\mathcal{B}(\mathcal{S})$ is the Borel σ -algebra on \mathcal{S} and $\mathcal{P}(\mathcal{S})$ the set of probability measures on $(\mathcal{S}, \mathcal{B}(\mathcal{S}))$. Finally, for a second Polish space \mathcal{S}' , we let

$$\mathcal{F}(\mathcal{S}; \mathcal{S}') = \{f : \mathcal{S} \rightarrow \mathcal{S}' : f \text{ measurable}\} \quad (13)$$

denote the space of measurable functions from \mathcal{S} to \mathcal{S}' .

For the finite set \mathcal{X} , let

$$\mathcal{P}_*(\mathcal{X}) = \left\{ \mathbf{m} \in \mathbb{R}^{\mathcal{X}} : m_x > 0 \text{ for all } x \in \mathcal{X} \text{ and } \sum_{x \in \mathcal{X}} m_x = 1 \right\}, \quad (14)$$

and

$$\mathcal{P}_a(\mathcal{X}) = \left\{ \mathbf{m} \in \mathbb{R}^{\mathcal{X}} : m_x \geq a \text{ for all } x \in \mathcal{X} \text{ and } \sum_{x \in \mathcal{X}} m_x = 1 \right\}. \quad (15)$$

For a set $K \subset \mathcal{P}(\mathcal{X})$, the closure \bar{K} , the complement K^c and the interior K° , will be considered with respect to the restriction of the Euclidean topology on the set $\mathcal{P}(\mathcal{X})$. $\mathcal{D}([0, \infty); \mathcal{S})$ denotes the space of cadlag functions on \mathcal{S} , equipped with the Skorohod topology (see [4, Section 16]), i.e., the Skorohod space. This space is separable and complete [4, Theorem 16.3], and a set is relatively compact in $\mathcal{D}([0, \infty); \mathcal{S})$, if and only if for every $M < \infty$, its natural projection on $\mathcal{D}([0, M]; \mathcal{S})$, is relatively compact [4, Theorem 16.4].

For $\bar{\mathcal{M}} = \mathcal{M}_c([0, \infty)^2)$, let \mathbb{P} be the probability measure on $(\bar{\mathcal{M}}, \mathcal{B}(\bar{\mathcal{M}}))$, under which the canonical map $N(\omega) = \omega$ is a Poisson measure with intensity measure equal to Lebesgue measure on $[0, \infty)^2$. Let $\mathcal{G}_t = \sigma\{N((0, s] \times A) : 0 \leq s \leq t, A \in \mathcal{B}([0, \infty))\}$, and let \mathcal{F}_t be the completion of \mathcal{G}_t under \mathbb{P} . Let \mathcal{P} be the corresponding predictable σ -field in $[0, \infty) \times \bar{\mathcal{M}}$. Similarly, for natural numbers k, k' we similarly define a measure $\mathbb{P}^{k, k'}$ on $(\bar{\mathcal{M}}^{k, k'}, \mathcal{B}(\bar{\mathcal{M}}^{k, k'}))$ under which the maps $N_i(\omega) = \omega_i$, $1 \leq i \leq k'$, are independent Poisson measures with intensity measure equal to k times the Lebesgue measure on $[0, \infty)^2$. $\{\mathcal{G}_t^{k, k'}\}$, $\{\mathcal{F}_t^{k, k'}\}$, and $\mathcal{P}^{k, k'}$ are defined analogously. Let \mathcal{A} be the class of $\mathcal{P} \setminus \mathcal{B}([0, \infty))$ measurable maps $\phi : [0, \infty) \times \bar{\mathcal{M}} \rightarrow [0, \infty)$, and \mathcal{A}_b the subset of these functions that are uniformly bounded from below away from zero and above by a positive constant. Similarly we define $\mathcal{A}^{k, k'}$ to be the set of $\mathcal{P}^{k, k'} \setminus \mathcal{B}([0, \infty)^{k'})$ measurable maps $\phi : [0, \infty) \times \bar{\mathcal{M}}^{k'} \rightarrow [0, \infty)^{k'}$, and $\mathcal{A}_b^{k, k'}$ the subset of these functions that all entries are uniformly bounded from below and above by positive constants.

2.1 The many particle control problem

For a subset \mathcal{K} of \mathcal{X}^n , we define a risk-sensitive cost $\mathcal{J}_{\mathcal{K}}^n : \mathcal{X}^n \times \mathcal{A}_b^{1, n|\mathcal{Z}|} \rightarrow [0, \infty]$ that corresponds to cost/reward up to the first time of hitting of \mathcal{K} by

$$\mathcal{J}_{\mathcal{K}}^n(\mathbf{x}^n, \mathbf{u}) \doteq \mathbb{E}_{\mathbf{x}^n} \left[\exp \left(\int_0^{T_{\mathcal{K}}} \left(\sum_{i=1}^n \sum_{y \in \mathcal{Z}_{\mathcal{X}_i^n}(t)} \gamma_{\mathcal{X}_i^n}(t)y C_{\mathcal{X}_i^n}(t)y \left(\frac{u_{\mathcal{X}_i^n}(t)y(t, i)}{\gamma_{\mathcal{X}_i^n}(t)y} \right) - nR^n(L(\mathcal{X}^n(t))) \right) dt \right) \right], \quad (16)$$

where $\mathbb{E}_{\mathbf{x}^n}$ denotes expected value given $\mathcal{X}^n(0) = \mathbf{x}^n$, the dynamics are given in (8), and

$$T_{\mathcal{K}} \doteq \inf \{t \in [0, \infty) : \mathcal{X}^n(t) \in \mathcal{K}\}. \quad (17)$$

We define the value function $\mathcal{W}_{\mathcal{K}}^n : \mathcal{X}^n \rightarrow [0, \infty]$ by

$$\mathcal{W}_{\mathcal{K}}^n(\mathbf{x}^n) \doteq \inf_{\mathbf{u} \in \mathcal{A}_b^{1, n|\mathcal{Z}|}} \mathcal{J}_{\mathcal{K}}^n(\mathbf{x}^n, \mathbf{u}). \quad (18)$$

Similarly, for a set $\mathcal{K} \subset \mathcal{X}^n$ we define the ordinary cost $\mathcal{J}_{\mathcal{K}}^n : \mathcal{X}^n \times \mathcal{A}_b^{1, n|\mathcal{Z}|} \rightarrow [0, \infty]$ and corresponding value function $\mathcal{V}_{\mathcal{K}}^n : \mathcal{X}^n \rightarrow [0, \infty]$ by

$$\mathcal{J}_{\mathcal{K}}^n(\mathbf{x}^n, \mathbf{q}) \doteq \mathbb{E}_{\mathbf{x}^n} \left[\int_0^{T_{\mathcal{K}}} \left(\frac{1}{n} \sum_{i=1}^n \sum_{y \in \mathcal{Z}_{\mathcal{X}_i^n}(t)} F_{\mathcal{X}_i^n}^n(t)y(q_{\mathcal{X}_i^n}(t)y(t, i)) + R^n(L(\mathcal{X}^n(t))) \right) dt \right], \quad (19)$$

where F^n is as in (9), and

$$\mathcal{V}_{\mathcal{K}}^n(\mathbf{x}^n) \doteq \inf_{\mathbf{q} \in \mathcal{A}_b^{1, n|\mathcal{Z}|}} \mathcal{J}_{\mathcal{K}}^n(\mathbf{x}^n, \mathbf{q}), \quad (20)$$

where the dynamics are given by (8) with \mathbf{u} replaced by \mathbf{q} , and the stopping time by (17). We remark that the reason for two different notations for controls is to aid the reader, by associating one with the risk sensitive problem and one with the

regular control problem. Moreover, there are occasions that both variables appear at the same time, as in the definition of \mathbf{F}^n or that of the Hamiltonian. Specific conditions on the cost functions will be given in Section 3.1, and properties of \mathbf{F}^n will be proved in Lemma 4. Note that for the many agent systems there are $n|\mathcal{Z}|$ PRMs, each with intensity 1.

2.2 The mean-field control problems

Suppose that \mathcal{K} can be identified with a subset of the simplex of probability measures $\mathcal{P}(\mathcal{X})$, in the sense that for every permutation $\sigma \in \mathbb{S}_n$ we have $\sigma\mathcal{K} = \mathcal{K}$. Then we can replace a control problem on \mathcal{X}^n by one on $\mathcal{P}(\mathcal{X})$. In this case $\mathcal{W}_{\mathcal{K}}^n$ and $\mathcal{V}_{\mathcal{K}}^n$ can be considered as functions on $\mathcal{P}^n(\mathcal{X})$, in the sense that we can find $W_K^n, V_K^n : \mathcal{P}^n(\mathcal{X}) \rightarrow [0, \infty]$, such that $\mathcal{W}_{\mathcal{K}}^n(\mathbf{x}^n) = W_K^n(L(\mathbf{x}^n))$ and $\mathcal{V}_{\mathcal{K}}^n(\mathbf{x}^n) = V_K^n(L(\mathbf{x}^n))$. To see this, pick a starting point $\mathbf{x}^n \in \mathcal{X}^n$ and some permutation σ . Then for any admissible control \mathbf{u} , the total cost generated starting at \mathbf{x}^n is the same as starting from \mathbf{x}_{σ}^n and picking \mathbf{u}_{σ} as the control. Therefore, for every $\mathbf{x}^n \in \mathcal{X}^n$, $\sigma \in \mathbb{S}_n$, we have $\mathcal{V}_{\mathcal{K}}(\mathbf{x}^n) = \mathcal{V}_{\mathcal{K}}(\mathbf{x}_{\sigma}^n)$.

Define $h^n : \mathcal{D}([0, \infty); \mathcal{P}^n(\mathcal{X})) \times \mathcal{A}_b^{n, |\mathcal{Z}|} \times \mathcal{P}^n(\mathcal{X}) \times \bar{\mathcal{M}}^{n, |\mathcal{Z}|} \rightarrow \mathcal{D}([0, \infty); \mathbb{R}^d)$ by

$$h^n \left(\boldsymbol{\mu}, \mathbf{u}, \mathbf{m}, \frac{1}{n} \mathbf{N}^n \right) (t) \doteq \mathbf{m} + \sum_{(x,y) \in \mathcal{Z}} \mathbf{v}_{xy} \int_{(0,t]} \int_{[0,\infty)} 1_{[0, \mu_x(-s)u_{xy}(s)]}(r) \frac{1}{n} N_{xy}^n(dsdr).$$

Since $\mathbf{u} \in \mathcal{A}_b^{n, |\mathcal{Z}|}$ implies the rates $u_{xy}(s)$ are uniformly bounded, one can explicitly construct a unique $\mathcal{D}([0, \infty); \mathcal{P}^n(\mathcal{X}))$ -valued process that satisfies

$$\boldsymbol{\mu} = h^n \left(\boldsymbol{\mu}, \mathbf{u}, \mathbf{m}, \frac{1}{n} \mathbf{N}^n \right). \quad (21)$$

[12]. Here $\boldsymbol{\mu}$ is the controlled process, \mathbf{u} is the control, \mathbf{m} is an initial condition, and \mathbf{N}^n/n is scaled noise.

Now with $T_K \doteq \inf \{t \in [0, \infty] : \boldsymbol{\mu}(t) \in K\}$, the functions $I_K^n, J_V^n : \mathcal{P}^n(\mathcal{X}) \times \mathcal{A}_b^{n, |\mathcal{Z}|} \rightarrow [0, \infty]$ and $W_K^n, V_K^n : \mathcal{P}^n(\mathcal{X}) \rightarrow [0, \infty]$ are given by

$$W_K^n(\mathbf{m}) \doteq \inf_{\mathbf{u} \in \mathcal{A}_b^{n, |\mathcal{Z}|}} I_K^n(\mathbf{m}, \mathbf{u}) \quad (22)$$

$$I_K^n(\mathbf{m}, \mathbf{u}) \doteq \mathbb{E}_{\mathbf{m}} \left[e^{n \int_0^{T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(t) \gamma_{xy} C_{xy}^n \left(\frac{u_{xy}(t)}{\gamma_{xy}} \right) - R^n(\boldsymbol{\mu}(t)) \right) dt} : \boldsymbol{\mu} = h^n \left(\boldsymbol{\mu}, \mathbf{u}, \mathbf{m}, \frac{1}{n} \mathbf{N}^n \right) \right], \quad (23)$$

and

$$V_K^n(\mathbf{m}) \doteq \inf_{\mathbf{q} \in \mathcal{A}_b^{n, |\mathcal{Z}|}} J_K^n(\mathbf{m}, \mathbf{q}) \quad (24)$$

$$J_K^n(\mathbf{m}, \mathbf{q}) \doteq \mathbb{E}_{\mathbf{m}} \left[\int_0^{T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(t) F_{xy}^n(q_{xy}(t)) + R^n(\boldsymbol{\mu}(t)) \right) dt : \boldsymbol{\mu} = h^n \left(\boldsymbol{\mu}, \mathbf{q}, \mathbf{m}, \frac{1}{n} \mathbf{N}^n \right) \right]. \quad (25)$$

For these control problems, there are $|\mathcal{Z}|$ PRMs, each with intensity n .

3 The equivalence of the control problems

In this section we prove that after a natural renormalization, the value function $\mathcal{W}_{\mathcal{K}}^n$ defined in (18) is linked to $\mathcal{V}_{\mathcal{K}}^n$ defined in (20) which, as noted before, it is the value function of an ordinary stochastic control problem with a new cost function. Specifically, we show that $-\log(\mathcal{W}_{\mathcal{K}}^n)/n$ equals $\mathcal{V}_{\mathcal{K}}^n$, and that the many particle and the mean field control problem are equivalent:

$$-\frac{1}{n} \log(\mathcal{W}_{\mathcal{K}}^n(L(\mathbf{x}^n))) = V_K^n(L(\mathbf{x}^n)) = \mathcal{V}_{\mathcal{K}}^n(\mathbf{x}^n) = -\frac{1}{n} \log(\mathcal{W}_{\mathcal{K}}^n(\mathbf{x}^n)). \quad (26)$$

3.1 The cost functions

One of the aims of this paper is to identify cost structures that make sense for the problem formulation and for which the risk-sensitive problem is equivalent to a control problem (rather than a game). The only place where restrictions are needed are in the costs C^n paid by the centralized controller to the agents for deviating from the nominal rates γ . To see what conditions will be needed, we first discuss briefly the strategy to be used for the proof of (26). The proof will use a related Bellman equation. Let $H^n : \mathcal{P}(\mathcal{X}) \times \mathbb{R}^{|\mathcal{Z}|} \rightarrow \mathbb{R}$ be given by

$$H^n(\mathbf{m}, \boldsymbol{\xi}) \doteq \inf_{\mathbf{q} \in (0, \infty)^{|\mathcal{Z}|}} \left\{ \sum_{(x,y) \in \mathcal{Z}} m_x (q_{xy} \xi_{xy} + F_{xy}^n(q_{xy})) \right\}, \quad (27)$$

where

$$F_{xy}^n(q) \doteq \sup_{u \in (0, \infty)} G_{xy}^n(u, q) \quad \text{and} \quad G_{xy}^n(u, q) \doteq \left[ul \left(\frac{q}{u} \right) - \gamma_{xy} C_{xy}^n \left(\frac{u}{\gamma_{xy}} \right) \right]. \quad (28)$$

We will show that the equation

$$H^n(\mathbf{m}, \Delta^n V(\mathbf{m})) + R^n(\mathbf{m}) = 0 \quad \text{in } \mathcal{P}^n(\mathcal{X}) \setminus K, \quad (29)$$

and boundary condition $V(\mathbf{m}) = 0$ for $\mathbf{m} \in K$ has V_K^n as the unique solution, where by $\Delta^n V(\mathbf{m})$ we denote the $|\mathcal{Z}|$ -dimensional vector $n(V(\mathbf{m} + \frac{\mathbf{v}_{xy}}{n}) - V(\mathbf{m}))$, and by $\Delta_{xy}^n V(\mathbf{m})$ the component $n(V(\mathbf{m} + \frac{\mathbf{v}_{xy}}{n}) - V(\mathbf{m}))_{xy}$, $(x, y) \in \mathcal{Z}$.

We are also going to prove that W_K^n is the unique solution to

$$\sup_{\mathbf{u} \in (0, \infty)^{|\mathcal{Z}|}} \left\{ \sum_{(x,y) \in \mathcal{Z}} m_x \left(u_{xy} \left(\frac{W(\mathbf{m}) - W(\mathbf{m} + \frac{\mathbf{v}_{xy}}{n})}{W(\mathbf{m})} \right) - \gamma_{xy} C_{xy}^n \left(\frac{u_{xy}}{\gamma_{xy}} \right) \right) \right\} = -R^n(\mathbf{m}) \quad (30)$$

for $\mathbf{m} \in \mathcal{P}^n(\mathcal{X}) \setminus K$ and $W(\mathbf{m}) = 1$ for $\mathbf{m} \in K$. In the proof of $-\frac{1}{n} \log(W_K^n) = V_K^n$, we will use the following lemma.

Lemma 1 *If $\tilde{V} : \mathcal{P}^n(\mathcal{X}) \rightarrow [0, \infty)$ is a solution to (29) and $\tilde{V}(\mathbf{m}) = 0$ for $\mathbf{m} \in K$, then $\tilde{W} = e^{-n\tilde{V}} : \mathcal{P}^n(\mathcal{X}) \rightarrow (0, \infty)$ is a solution of (30) and $W(\mathbf{m}) = 1$ for $\mathbf{m} \in K$.*

For Lemma 1 to be true, we will need the following to be valid.

$$\begin{aligned} H^n(\mathbf{m}, \boldsymbol{\xi}) &\doteq \inf_{\mathbf{q} \in [0, \infty)^{|\mathcal{Z}|}} \left\{ \sum_{(x,y) \in \mathcal{Z}} m_x (q_{xy} \xi_{xy} + F_{xy}^n(q_{xy})) \right\} \\ &= \inf_{\mathbf{q} \in [0, \infty)^{|\mathcal{Z}|}} \sup_{\mathbf{u} \in (0, \infty)^{|\mathcal{Z}|}} \left\{ \sum_{(x,y) \in \mathcal{Z}} m_x (q_{xy} \xi_{xy} + G_{xy}^n(u_{xy}, q_{xy})) \right\} \\ &= \sup_{\mathbf{u} \in (0, \infty)^{|\mathcal{Z}|}} \inf_{\mathbf{q} \in [0, \infty)^{|\mathcal{Z}|}} \left\{ \sum_{(x,y) \in \mathcal{Z}} m_x (q_{xy} \xi_{xy} + G_{xy}^n(u_{xy}, q_{xy})) \right\} \\ &= \sup_{\mathbf{u} \in (0, \infty)^{|\mathcal{Z}|}} \left\{ \sum_{(x,y) \in \mathcal{Z}} m_x \left(u_{xy} (1 - e^{-\xi_{xy}}) - \gamma_{xy} C_{xy}^n \left(\frac{u_{xy}}{\gamma_{xy}} \right) \right) \right\} \\ &= \sum_{(x,y) \in \mathcal{Z}} m_x \gamma_{xy} (C_{xy}^n)^* (1 - e^{-\xi_{xy}}), \end{aligned} \quad (31)$$

where $(C_{xy}^n)^* : (-\infty, 1) \rightarrow \mathbb{R}$ is given by $(C_{xy}^n)^*(z) = \sup_{u > 0} [zu - C_{xy}^n(u)]$. The crucial step is the third equality, which requires that the Isaac's condition hold, i.e., the supremum and infimum can be exchanged. For this, we will apply Sion's Theorem (Corollary 3.3 in [26]), which states that if continuous $F(u, q)$ is quasi-concave for every u in some convex set U and quasi-convex for every q in some convex set Q , and if one of the two sets is compact, then we can exchange the supremum with the infimum. We would like to apply Sion's Theorem to

$$L_{xy}^n(u, q) = q\xi + ul \left(\frac{q}{u} \right) - \gamma_{xy} C_{xy}^n \left(\frac{u}{\gamma_{xy}} \right).$$

Since ℓ is convex, L_{xy}^n is convex with respect to q . It is easy to see that $L_{xy}^n(u, q)$ is not concave with respect to u . However it is possible for L_{xy}^n to be quasi-concave, with respect to u , for every $q \geq 0$. For $L_{xy}^n(\cdot, q)$ to be quasi-concave for each q means that it changes monotonicity at most one time. By differentiating with respect to u we get

$$\partial_u L_{xy}^n(u, q) = -\frac{q}{u} + 1 - (C_{xy}^n)' \left(\frac{u}{\gamma_{xy}} \right),$$

and so what is needed is that for every $q \geq 0$ there is u_q^n such that

$$-\frac{q}{u} + 1 - (C_{xy}^n)' \left(\frac{u}{\gamma_{xy}} \right) \geq 0 \text{ for } u \leq u_q^n \quad \text{and} \quad -\frac{q}{u} + 1 - (C_{xy}^n)' \left(\frac{u}{\gamma_{xy}} \right) \leq 0 \text{ for } u \geq u_q^n.$$

This is equivalent to the existence of u_q^n such that

$$u(C_{xy}^n)'(u) - u \leq -q \text{ for } u \leq u_q^n \quad \text{and} \quad u(C_{xy}^n)'(u) - u \geq -q \text{ for } u \geq u_q^n.$$

We set $Q(u) = u(C_{xy}^n)'(u) - u$. Now, the above can be reformulated as, every level set $\{u \in (0, \infty) : Q(u) = \hat{q}\}$ of Q , for negative \hat{q} , is crossed once and after it has been "exited", the graph of Q stays above it. The latter, implies monotonicity of Q until level-set zero is reached (i.e. when $(C_{xy}^n)'(u) = 1$). After level-set zero is reached, Q can take any value as long as $u(C_{xy}^n)'(u) - u \geq 0$, since we have no restriction for $\hat{q} > 0$. From the above, we conclude that

$$u(C_{xy}^n)'(u) - u \text{ is increasing until } (C_{xy}^n)'(u) \geq 1, \quad (32)$$

and then $(C_{xy}^n)'(u) \geq 1$ remains bigger than one. By taking another derivative, we have

$$(C_{xy}^n)'(u) + u(C_{xy}^n)''(u) - 1 \geq 0, \quad (33)$$

from which we further conclude, that while $(C_{xy}^n)'(u) < 1$, the function C_{xy}^n is also convex. In fact, as it is apparent from the last line in (31), the values of C_{xy}^n , after $(C_{xy}^n)'$ gets bigger than one, are irrelevant, and therefore (C_{xy}^n) does not have to be convex on the whole $[0, \infty)$. However for simplicity we will assume that (C_{xy}^n) is convex, and that it satisfies (32) everywhere where it is finite. In Theorem 24, it is argued that under some extra smoothness assumption on C_{xy}^n , (32) is the weakest condition such that in the definition of the Hamiltonian (31) one can exchange the supremum with the infimum, and therefore the application of Sion's theorem is optimal (there is no better result that we could have used). Now we provide the main assumption for C_{xy}^n .

Assumption 2 For each $n \in \mathbb{N}$, $R^n : \mathcal{P}(\mathcal{X}) \rightarrow [0, \infty)$ is a continuous function. Moreover, for every $(x, y) \in \mathcal{Z}$, $C_{xy}^n : [0, \infty) \rightarrow [0, \infty]$ is a convex function that satisfies the following:

- 1 $u(C_{xy}^n)'(u) - u$ is increasing on the maximal open interval where C_{xy}^n is finite.
- 2 $C_{xy}^n(1) = 0$.

Assumption 2.2 is not necessary, but it simplifies the analysis and it is appropriate for the situation being modeled to have zero cost when there is no change from the nominal rates. Under Assumption 2, we will prove in Lemma 23 that the Isaac's condition is satisfied, and therefore the equality (31) is true.

Lemma 3 Under Assumption 2, the cost functions C_{xy}^n satisfy the following on $(u_{1,xy}^n, u_{2,xy}^n)$:

- 1 for every $(x, y) \in \mathcal{Z}$ we have $(C_{xy}^n)'(u) \geq 1 - \frac{1}{u}$ for $u > 1$, and therefore $\liminf_{u \rightarrow \infty} (C_{xy}^n)'(u) \geq 1$,
- 2 for every $(x, y) \in \mathcal{Z}$ and $u \in (0, \infty)$ we have $C_{xy}^n(u) \geq -\log u + u - 1$.

Proof. It follows from the monotonicity that $u(C_{xy}^n)'(u) - u \geq -1$ for $u > 1$, which gives the first statement. The second follows by comparing $(C_{xy}^n)(u)$ with $\int_1^u [1 - \frac{1}{s}] ds$ and using $C_{xy}^n(1) = 0$. ■

Example 2 The family of functions $C_{xy}^n(u) = \frac{1}{pu^p} + \frac{u^q}{q} - \frac{p+q}{pq}$, where $p \geq 1$ and $q \geq 1$, satisfy Assumption 2. Taking the derivative of $C_{xy}^n(u)$ gives $-\frac{1}{u^{p+1}} + u^{q-1}$, and multiplying with u and subtracting u yields $-\frac{1}{u^p} + u^q - u$. Taking the derivative again gives $\frac{p}{u^{p+1}} + qu^{q-1} - 1$, which is always bigger than zero, since $\frac{p}{u^{p+1}}$ and qu^{q-1} are everywhere positive and bigger than one on the intervals $[0, 1]$ and $[1, \infty)$, respectively.

Before proceeding with the proof, we state some properties of F_{xy}^n .

Lemma 4 For every $n \in \mathbb{N}$ and for every $(x, y) \in \mathcal{Z}$, let F_{xy}^n be as in (9), where $\{C_{xy}^n\}$ satisfy Assumption 2. Then the following hold.

$$1. F_{xy}^n(q) \geq \gamma_{xy} \ell\left(\frac{q}{\gamma_{xy}}\right) \geq 0, \quad 2. F_{xy}^n(\gamma_{xy}) = 0, \quad 3. F_{xy}^n \text{ is convex on } [0, \infty).$$

3.2 Equivalence of the stochastic problems

Theorem 5 Let $n \in \mathbb{N}$, $\mathcal{K} \subset \mathcal{X}^n$, (resp. $K \subset \mathcal{P}^n(\mathcal{X})$), and C, R be as in Assumption 2. Then

$$V_K^n(\mathbf{m}) = -\frac{1}{n} \log(W_K^n(\mathbf{m})) \quad (34)$$

and

$$\mathcal{V}_{\mathcal{K}}^n(\mathbf{x}^n) = -\frac{1}{n} \log(W_{\mathcal{K}}^n(\mathbf{x}^n)). \quad (35)$$

If in addition $\mathcal{K} \subset \mathcal{X}^n$ is invariant under permutations, and therefore can be identified with a subset of $\mathcal{P}^n(\mathcal{X})$, then

$$-\frac{1}{n} \log(W_K^n(L(\mathbf{x}^n))) = V_K^n(L(\mathbf{x}^n)) = \mathcal{V}_{\mathcal{K}}^n(\mathbf{x}^n) = -\frac{1}{n} \log(W_{\mathcal{K}}^n(\mathbf{x}^n)). \quad (36)$$

The proof of this result appears later in this section. Also, we will only prove the first equality and note that the third follows in a similar manner.

Lemma 6 Let $n \in \mathbb{N}$, $\emptyset \neq K \subset \mathcal{P}^n(\mathcal{X})$, and C, R be as in Assumption 2. Then, the equation (29) has at least one solution.

Proof. For the proof we use the equivalent discrete time stochastic control problem. We consider the following set of controls

$$A_a(\mathbf{m}) = \left\{ \mathbf{q} \in [0, \infty)^{|\mathcal{Z}|} : \frac{1}{a} \geq \sum_{(x,y) \in \mathcal{Z}} m_x q_{xy}(\mathbf{m}) \geq a \right\} \quad \text{and} \quad A_+(\mathbf{m}) = \cup_{a>0} A_a(\mathbf{m}). \quad (37)$$

For such a control the probability of moving from state \mathbf{m} to state $\mathbf{m} + \frac{1}{n} \mathbf{v}_{\bar{x}, \bar{y}}$ will be given by

$$\frac{m_{\bar{x}} q_{\bar{x}\bar{y}}(\mathbf{m})}{\sum_{(x,y) \in \mathcal{Z}} m_x q_{xy}(\mathbf{m})},$$

and the (conditional) expected cost till the time of transition is given by

$$\frac{\sum_{(x,y) \in \mathcal{Z}} m_x F_{xy}^n(q_{xy}(\mathbf{m})) + R(\mathbf{m})}{n \sum_{(x,y) \in \mathcal{Z}} m_x q_{xy}(\mathbf{m})}.$$

Also, with some abuse of notation, we define the set of feedback controls

$$A_a = \{\mathbf{q} \in [0, \infty)^{|\mathcal{P}^n(\mathcal{X}) \times \mathcal{Z}|} : \mathbf{q}(\mathbf{m}) \in A_a(\mathbf{m})\} \quad \text{and} \quad A_+ = \cup_{a>0} A_a. \quad (38)$$

Given controlled transition probabilities as above, let $\boldsymbol{\mu}(i)$ be the corresponding controlled discrete time process. We define the value function $\bar{V}_K^n(\mathbf{m}) : \mathcal{P}(\mathbb{R}^d) \rightarrow [0, \infty)$ by

$$\bar{V}_K^n(\mathbf{m}) \doteq \inf_{\mathbf{q} \in A_+} \mathbb{E}_{\mathbf{m}} \left[\sum_{i=1}^{T_K} \frac{\sum_{(x,y) \in \mathcal{Z}} \mu_x(i) F_{xy}^n(q_{xy}(\boldsymbol{\mu}(i))) + R(\boldsymbol{\mu}(i))}{n \sum_{(x,y) \in \mathcal{Z}} \mu_x(i) q_{xy}(\boldsymbol{\mu}(i))} \right], \quad (39)$$

where $\mathbb{E}_{\mathbf{m}}$ denotes expected value given $\boldsymbol{\mu}(0) = \mathbf{m}$ and $T_K \doteq \inf\{i \in \mathbb{N} : \boldsymbol{\mu}(i) \in K\}$.

To see that $\bar{V}_K^n(\mathbf{m})$ is finite, we just have to use the original rates and note that the total cost is proportional to the expected exit time, which is finite by classical results on Markov chains. Since $F_{xy}, R \geq 0$, and F_{xy} is convex with $\gamma_{xy} \ell\left(\frac{\cdot}{\gamma_{xy}}\right)$ as a lower bound (see Lemma 4), one can see that we can find a constant $a_0 > 0$ such that only controls in A_{a_0} (or any $a < a_0$) should be considered. More specifically to see that a term in the sum appearing on the RHS of (39) gets big when $\sum_{(x,y) \in \mathcal{Z}} \mu_x(i) q_{xy}(\boldsymbol{\mu}(i))$ gets small we bound the denominator by $|\mathcal{Z}|$ times the biggest term and the nominator by the same term and then we use the fact that $F_{xy}(0) \geq \gamma_{xy}$. For the other bound we use the superlinearity of F_{xy} . Now by [3, Proposition 1.1 in Chapter 3], we have that this value function satisfies

$$\bar{V}_K^n(\mathbf{m}) = \inf_{\mathbf{q} \in A_{a_0}(\mathbf{m})} \left\{ \frac{\sum_{(x,y) \in \mathcal{Z}} m_x F_{xy}^n(q_{xy}) + R(\mathbf{m})}{n \sum_{(x,y) \in \mathcal{Z}} m_x q_{xy}} + \sum_{(\bar{x}, \bar{y}) \in \mathcal{Z}} \frac{m_{\bar{x}} q_{\bar{x}\bar{y}}}{\sum_{(x,y) \in \mathcal{Z}} m_x q_{xy}} \bar{V}_K^n\left(\mathbf{m} + \frac{1}{n} \mathbf{v}_{\bar{x}\bar{y}}\right) \right\}.$$

It then follows that $\bar{V}_K^n(\mathbf{m})$ satisfies the last display if and only if [with $\Delta_{xy}^n \bar{V}_K^n(\mathbf{m}) \doteq n(\bar{V}_K^n(\mathbf{m} + \frac{\mathbf{v}_{xy}}{n}) - \bar{V}_K^n(\mathbf{m}))$]

$$\inf_{\mathbf{q} \in A_{a_0}(\mathbf{m})} \left\{ \sum_{(x,y) \in \mathcal{Z}} m_x (q_{xy} \Delta_{xy}^n \bar{V}_K^n(\mathbf{m}) + F_{xy}^n(q_{xy})) \right\} + R(\mathbf{m}) = 0.$$

Since a_0 can be chosen arbitrary small and the left side on the previous display is continuous with respect to \mathbf{q} , we get

$$\inf_{\mathbf{q} \in [0, \infty)^{|\mathcal{Z}|}} \left\{ \sum_{(x,y) \in \mathcal{Z}} m_x (q_{xy} \Delta_{xy}^n \bar{V}_K^n(\mathbf{m}) + F_{xy}^n(q_{xy})) \right\} + R(\mathbf{m}) = 0.$$

Then using the definition (27) this is the same as

$$H^n(\mathbf{m}, \Delta^n \bar{V}_K^n(\mathbf{m})) + R(\mathbf{m}) = 0,$$

and we also have the boundary condition $\bar{V}_K^n(\mathbf{m}) = 0$ for all $\mathbf{m} \in K$. ■

Proof of Lemma 1. Let \tilde{V} be a solution to (29). We then have $H^n(\mathbf{m}, \Delta^n \tilde{V}(\mathbf{m})) + R^n(\mathbf{m}) = 0$, or by using the second from the bottom line in (31),

$$\sup_{\mathbf{u} \in (0, \infty)^{|\mathcal{Z}|}} \left\{ \sum_{(x,y) \in \mathcal{Z}} m_x \left(u_{xy} \left(1 - e^{-n(\tilde{V}(\mathbf{m} + \frac{\mathbf{v}_{xy}}{n}) - \tilde{V}(\mathbf{m}))} \right) - \gamma_{xy} C_{xy}^n \left(\frac{u_{xy}}{\gamma_{xy}} \right) \right) \right\} + R^n(\mathbf{m}) = 0.$$

By making the substitution we have

$$\sup_{\mathbf{u} \in (0, \infty)^{|\mathcal{Z}|}} \left\{ \sum_{(x,y) \in \mathcal{Z}} m_x \left(u_{xy} \left(1 - \frac{\tilde{W}(\mathbf{m} + \frac{\mathbf{v}_{xy}}{n})}{\tilde{W}(\mathbf{m})} \right) - \gamma_{xy} C_{xy}^n \left(\frac{u_{xy}}{\gamma_{xy}} \right) \right) \right\} + R^n(\mathbf{m}) = 0,$$

which is the same as (30). ■

Lemma 7 Let $f : \mathcal{P}^n(\mathcal{X}) \rightarrow \mathbb{R}$, $\mathbf{m} \in \mathcal{P}^n(\mathcal{X})$, and $\mathbf{q} \in \mathcal{A}_b^{n, |\mathcal{Z}|}$ be given, and let $\boldsymbol{\mu}$ solve (21). Then

$$\begin{aligned} f(\boldsymbol{\mu}(t)) - f(\mathbf{m}) - \int_0^t \sum_{(x,y) \in \mathcal{Z}} \mu_x(s) q_{xy}(s) \Delta_{xy}^n f(\boldsymbol{\mu}(s)) ds, \\ f(\boldsymbol{\mu}(t \wedge T_K)) - f(\mathbf{m}) - \int_0^{t \wedge T_K} \sum_{(x,y) \in \mathcal{Z}} \mu_x(s) q_{xy}(s) \Delta_{xy}^n f(\boldsymbol{\mu}(s)) ds, \\ f(\boldsymbol{\mu}(t \wedge T_K)) - f(\boldsymbol{\mu}(t' \wedge T_K)) - \int_{t' \wedge T_K}^{t \wedge T_K} \sum_{(x,y) \in \mathcal{Z}} \mu_x(s) q_{xy}(s) \Delta_{xy}^n f(\boldsymbol{\mu}(s)) ds, \end{aligned}$$

are martingales with respect to the filtration \mathcal{F}_t .

Proof. By the construction of $\boldsymbol{\mu}$ (also see Ito's formula [19, Chapter 2, Theorem 5.1]), we have

$$f(\boldsymbol{\mu}(t)) - f(\mathbf{m}) - \frac{1}{n} \int_{(0,t]} \sum_{(x,y) \in \mathcal{Z}} \int_{[0,\infty)} \mathbf{1}_{[0,\mu_x(s-)q_{xy}(s)]}(r) \Delta_{xy}^n f(\boldsymbol{\mu}(s-)) N_{xy}^n(dsdr) = 0.$$

Indeed, the right hand side simply records each jump in $f(\boldsymbol{\mu}(s))$ for $0 < s \leq t$. Since the intensity measure of $N_{xy}^n(dsdr)$ is n times Lebesgue measure, by [19, Chapter 2, Theorem 3.1], for each $(x, y) \in \mathcal{Z}$

$$\int_0^t \mu_x(s-)q_{xy}(s)\Delta_{xy}^n f(\boldsymbol{\mu}(s-))ds - \frac{1}{n} \int_{(0,t]} \int_{[0,\infty)} \mathbf{1}_{[0,\mu_x(s-)q_{xy}(s)]}(r) \Delta_{xy}^n f(\boldsymbol{\mu}(s-)) N_{xy}^n(dsdr),$$

is a martingale. By combining the last two displays and using that $s-$ in the ordinary integral can be replaced by s due to left continuity,

$$f(\boldsymbol{\mu}(t)) - f(\mathbf{m}) - \int_0^t \sum_{(x,y) \in \mathcal{Z}} \mu_x(s)q_{xy}(s)\Delta_{xy}^n f(\boldsymbol{\mu}(s))ds$$

is a martingale. The second and third formulas then follow from standard properties of martingales. ■

Lemma 8 Let $g : \mathcal{P}^n(\mathcal{X}) \rightarrow (0, \infty)$, $\mathbf{m} \in \mathcal{P}^n(\mathcal{X})$, and $\mathbf{u} \in \mathcal{A}_b^{n,|\mathcal{Z}|}$ be given, and let $\boldsymbol{\mu}$ solve (21). Then

$$\begin{aligned} & \frac{g(\boldsymbol{\mu}(t))}{g(\mathbf{m})} \exp \left\{ - \int_0^t \sum_{(x,y) \in \mathcal{Z}} \mu_x(s)u_{xy}(s) \frac{\Delta_{xy}^n g(\boldsymbol{\mu}(s))}{g(\boldsymbol{\mu}(s))} ds \right\}, \\ & \frac{g(\boldsymbol{\mu}(t \wedge T_K))}{g(\mathbf{m})} \exp \left\{ - \int_0^{t \wedge T_K} \sum_{(x,y) \in \mathcal{Z}} \mu_x(s)u_{xy}(s) \frac{\Delta_{xy}^n g(\boldsymbol{\mu}(s))}{g(\boldsymbol{\mu}(s))} ds \right\}, \\ & \frac{g(\boldsymbol{\mu}(t \wedge T_K))}{g(\boldsymbol{\mu}(t' \wedge T_K))} \exp \left\{ - \int_{t' \wedge T_K}^{t \wedge T_K} \sum_{(x,y) \in \mathcal{Z}} \mu_x(s)u_{xy}(s) \frac{\Delta_{xy}^n g(\boldsymbol{\mu}(s))}{g(\boldsymbol{\mu}(s))} ds \right\}, \end{aligned} \quad (40)$$

are martingales with respect to the filtration \mathcal{F}_t .

Proof. The proof is a direct application of the corollary in [22, Page 66]. ■

Lemma 9 Let $\mathbf{m} \in \mathcal{P}^n(\mathcal{X})$ and $\mathbf{u} \in \mathcal{A}_b^{n,|\mathcal{Z}|}$. There exists a constant $c > 0$, that depends only on the bounds on \mathbf{u} , the dimension d , the constant $R_{\max}^n = \max\{R^n(\mathbf{m}) : \mathbf{m} \in \mathcal{P}^n(\mathcal{X})\}$, and the number n of agents, such that for every $t \geq t' \geq 0$,

$$\mathbb{E}_{\mathbf{m}} \left[e^{-nR_{\max}^n(t \wedge T_K - t' \wedge T_K)} \Big| \mathcal{F}_{t'} \right] \geq c.$$

Furthermore it is true that

$$T_K < \infty \text{ a.s.}, \text{ and } \mathbb{E}_{\mathbf{m}} \left[e^{-nR_{\max}^n(T_K - t' \wedge T_K)} \Big| \mathcal{F}_{t'} \right] \geq c.$$

Proof. We claim there exists g such that for all s

$$\sum_{(x,y) \in \mathcal{Z}} \mu_x(s)u_{xy}(s) \frac{\Delta_{xy}^n g(\boldsymbol{\mu}(s))}{g(\boldsymbol{\mu}(s))} \geq nR_{\max}^n. \quad (41)$$

To show the existence of such a g we use the following procedure. Since the one particle process with generator given in (1) is ergodic, we have that the process on \mathcal{X}^n , with generator given in (2), as well as the one on $\mathcal{P}^n(\mathcal{X})$, with generator given in (4), are also ergodic. We split $\mathcal{P}^n(\mathcal{X})$ into sets $\{K_i\}_{0 \leq i \leq i_{\max}}$, where $K_0 = K$, and K_{i+1} is generated inductively as the set of all points in $\mathcal{P}^n(\mathcal{X})$ that do not belong to K_i but such that the process with generator (4) can reach K_i in one jump. Since the original process has d states, it is easy to see that $i_{\max} \leq d^n$. Since $\mathbf{u} \in \mathcal{A}_b^{n,|\mathcal{Z}|}$, there exist constants $0 < c_1 \leq c_2 < \infty$ such that $c_1 \leq u_{xy}(t) \leq c_2$ for all $t \geq 0$ a.s. Let g be defined by

$$g(\mathbf{m}) \doteq \left(\frac{nR_{\max}^n + nd^2c_2 + c_1}{c_1} \right)^{i_{\max} - i}, \text{ for } \mathbf{m} \in K_i.$$

Let $\boldsymbol{\mu}(\cdot)$ be the process with control \mathbf{u} . For $0 \leq s \leq t$ suppose that $\boldsymbol{\mu}(s) \in K_i$ for some $i \geq 1$. Then there exists at least one $(\tilde{x}, \tilde{y}) \in \mathcal{Z}$ such that $\boldsymbol{\mu}(s) + \frac{v_{\tilde{x}\tilde{y}}}{n} \in K_{i-1}$. Therefore

$$\begin{aligned} \sum_{(x,y) \in \mathcal{Z}} \mu_x(s) u_{xy}(s) \frac{\Delta_{xy}^n g(\boldsymbol{\mu}(s))}{g(\boldsymbol{\mu}(s))} &= \mu_{\tilde{x}}(s) u_{\tilde{x}\tilde{y}}(s) \frac{\Delta_{\tilde{x}\tilde{y}}^n g(\boldsymbol{\mu}(s))}{g(\boldsymbol{\mu}(s))} + n \sum_{(x,y) \in \mathcal{Z}, (x,y) \neq (\tilde{x}, \tilde{y})} \frac{g(\boldsymbol{\mu}(s) + \frac{v_{xy}}{n})}{g(\boldsymbol{\mu}(s))} \mu_x(s) u_{xy}(s) \\ &- n \sum_{(x,y) \in \mathcal{Z}, (x,y) \neq (\tilde{x}, \tilde{y})} \frac{g(\boldsymbol{\mu}(s))}{g(\boldsymbol{\mu}(s))} \mu_x(s) u_{xy}(s) \geq \mu_{\tilde{x}}(s) u_{\tilde{x}\tilde{y}}(s) \frac{\Delta_{\tilde{x}\tilde{y}}^n g(\boldsymbol{\mu}(s))}{g(\boldsymbol{\mu}(s))} - n \sum_{(x,y) \in \mathcal{Z}} \mu_x(s) u_{xy}(s) \\ &\geq c_1 \left(\frac{nR_{\max}^n + nd^2c_2 + c_1}{c_1} - 1 \right) - nc_2d^2 \geq nR_{\max}^n, \end{aligned}$$

where in the next to last inequality we used the fact that $\mu_{\tilde{x}}(s) \geq \frac{1}{n}$ (because otherwise there is no particle at \tilde{x} to move), and that $\Delta_{xy}^n V(\mathbf{m}) = n(V(\mathbf{m} + \frac{v_{xy}}{n}) - V(\mathbf{m}))$.

Using the last martingale in Lemma 8, we have

$$\mathbb{E}_{\mathbf{m}} \left[\frac{g(\boldsymbol{\mu}(t \wedge T_K))}{g(\boldsymbol{\mu}(t' \wedge T_K))} \exp \left\{ - \int_{t' \wedge T_K}^{t \wedge T_K} \sum_{(x,y) \in \mathcal{Z}} \mu_x(s) u_{xy}(s) \frac{\Delta_{xy}^n g(\boldsymbol{\mu}(s))}{g(\boldsymbol{\mu}(s))} ds \right\} \middle| \mathcal{F}_{t'} \right] = 1,$$

from which we get

$$\mathbb{E}_{\mathbf{m}} \left[\exp \left\{ - \int_{t' \wedge T_K}^{t \wedge T_K} \sum_{(x,y) \in \mathcal{Z}} \mu_x(s) u_{xy}(s) \frac{\Delta_{xy}^n g(\boldsymbol{\mu}(s))}{g(\boldsymbol{\mu}(s))} ds \right\} \middle| \mathcal{F}_{t'} \right] \geq c \doteq \frac{\min_{\mathcal{P}^n(x)} g}{\max_{\mathcal{P}^n(x)} g}.$$

By applying equation (41)

$$\mathbb{E}_{\mathbf{m}} \left[e^{-nR_{\max}^n(t \wedge T_K - t' \wedge T_K)} \middle| \mathcal{F}_{t'} \right] \geq c.$$

Now choose now $\tau > 0$ such that $e^{-nR_{\max}^n\tau} \leq c/2$. We claim that

$$(T_K \leq t' + \tau) \Leftrightarrow (T_K \wedge (t' + 2\tau) - t' \wedge T_K) \leq \tau.$$

Indeed if $t' \geq T_K$, then both parts are trivially true. Let assume that $t' \leq T_K$, and $T_K \leq t' + \tau$. Then $T_K \wedge (t' + 2\tau) = T_K$, and $t' \wedge T_K = t'$, and therefore $(T_K \wedge (t' + 2\tau) - t' \wedge T_K) = T_K - t' \leq \tau$. If on the other hand $t' \leq T_K$ and $(T_K \wedge (t' + 2\tau) - t' \wedge T_K) \leq \tau$, we get $(T_K \wedge (t' + 2\tau)) \leq \tau + t'$, which gives that $T_K \leq (t' + 2\tau)$, and therefore $T_K = (T_K \wedge (t' + 2\tau)) \leq t' + \tau$. Using the claim just proved gives

$$\begin{aligned} \mathbb{P}_{\mathbf{m}}(T_K \leq t' + \tau | \mathcal{F}_{t'}) &= \mathbb{P}_{\mathbf{m}}(T_K \wedge (t' + 2\tau) - t' \wedge T_K \leq \tau | \mathcal{F}_{t'}) \\ &= \mathbb{P}_{\mathbf{m}} \left(e^{-nR_{\max}^n(T_K \wedge (t' + 2\tau) - t' \wedge T_K)} \geq e^{-nR_{\max}^n\tau} \middle| \mathcal{F}_{t'} \right). \end{aligned}$$

Let $E_1 \doteq \{e^{-nR_{\max}^n(T_K \wedge (t' + 2\tau) - t' \wedge T_K)} \geq e^{-nR_{\max}^n\tau}\}$ and $E_2 \doteq E_1^c$. Then since $T_K \wedge (t' + 2\tau) - t' \wedge T_K \geq 0$

$$\begin{aligned} \mathbb{E}_{\mathbf{m}} \left[e^{-nR_{\max}^n(T_K \wedge (t' + 2\tau) - t' \wedge T_K)} \middle| \mathcal{F}_{t'} \right] &= \mathbb{E}_{\mathbf{m}} \left[\mathbf{1}_{E_1} e^{-nR_{\max}^n(T_K \wedge (t' + 2\tau) - t' \wedge T_K)} \middle| \mathcal{F}_{t'} \right] \\ &+ \mathbb{E}_{\mathbf{m}} \left[\mathbf{1}_{E_2} e^{-nR_{\max}^n(T_K \wedge (t' + 2\tau) - t' \wedge T_K)} \middle| \mathcal{F}_{t'} \right] \leq \mathbb{E}_{\mathbf{m}} \left[\mathbf{1}_{E_1} \middle| \mathcal{F}_{t'} \right] + e^{-nR_{\max}^n\tau}. \end{aligned}$$

From this, the first part of the lemma and $e^{-nR_{\max}^n\tau} \leq c/2$, we get

$$\begin{aligned} \mathbb{P}_{\mathbf{m}} \left(e^{-nR_{\max}^n(T_K \wedge (t' + 2\tau) - t' \wedge T_K)} \geq e^{-nR_{\max}^n\tau} \middle| \mathcal{F}_{t'} \right) \\ \geq \mathbb{E}_{\mathbf{m}} \left[e^{-nR_{\max}^n(T_K \wedge (t' + 2\tau) - t' \wedge T_K)} \middle| \mathcal{F}_{t'} \right] - e^{-nR_{\max}^n\tau} \geq c - \frac{c}{2} = \frac{c}{2}. \end{aligned}$$

Now we have

$$\begin{aligned} \mathbb{P}_{\mathbf{m}}(T_K = \infty) &= \lim_{k \rightarrow \infty} \mathbb{P}_{\mathbf{m}}(T_K > k\tau) = \mathbb{P}_{\mathbf{m}}(T_K > 0) \lim_{k \rightarrow \infty} \prod_{k'=0}^k (1 - \mathbb{P}_{\mathbf{m}}(T_K \leq (k'+1)\tau | T_K > k'\tau)) \\ &\leq \lim_{k \rightarrow \infty} \left(1 - \frac{c}{2}\right)^k = 0, \end{aligned}$$

where in the second inequality we iteratively used the formula for conditional probability. The remaining inequality is just an application of the monotone convergence theorem. ■

Lemma 10 Given $\mathbf{m} \in \mathcal{P}^n(\mathcal{X})$, $\epsilon > 0$ and $\mathbf{u} \in \mathcal{A}_b^{n,|\mathcal{Z}|}$ with

$$\mathbb{E}_{\mathbf{m}} \left[e^{n \int_0^{T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(t) C_{xy}^n \left(\frac{u_{xy}(t)}{\gamma_{xy}} \right) - R^n(\boldsymbol{\mu}(t)) \right) dt} \right] < \infty,$$

there exists $\tilde{\mathbf{u}} \in \mathcal{A}_b^{n,|\mathcal{Z}|}$ and $\tau < \infty$, such that

$$\sum_{(x,y) \in \mathcal{Z}} \tilde{\mu}_x(t) \gamma_{xy} C_{xy}^n \left(\frac{\tilde{u}_{xy}(t)}{\gamma_{xy}} \right) - R^n(\tilde{\boldsymbol{\mu}}(t)) \leq 0$$

for every $t > \tau$, and

$$I_K^n(\mathbf{m}, \tilde{\mathbf{u}}) \leq I_K^n(\mathbf{m}, \mathbf{u}) + \epsilon.$$

Proof. Let such $\mathbf{m} \in \mathcal{P}^n(\mathcal{X})$, $\epsilon > 0$, and $\mathbf{u} \in \mathcal{A}_b^{n,|\mathcal{Z}|}$ be given, and let $c > 0$ from Lemma 9 be such that

$$\mathbb{E}_{\mathbf{m}} \left[e^{n R_{\max}^n(T_K - t' \wedge T_K)} \Big| \mathcal{F}_{t'} \right] \geq c \quad (43)$$

for $t' \in [0, \infty)$. Since by Lemma 9 T_K is finite a.s., we can find $\tau < \infty$ such that

$$\mathbb{E}_{\mathbf{m}} \left[I_{\{T_K \geq \tau\}} e^{n \int_0^{T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(t) \gamma_{xy} C_{xy}^n \left(\frac{u_{xy}(t)}{\gamma_{xy}} \right) - R^n(\boldsymbol{\mu}(t)) \right) dt} \right] \leq \epsilon c.$$

Now set $\tilde{\mathbf{u}}(t) = \mathbf{u}(t)$ for $t \leq \tau$, and $\tilde{\mathbf{u}}(t) = \boldsymbol{\gamma}$ so that $C_{xy}^n(\tilde{u}_{xy}(t)/\gamma_{xy}) = 0$ for $t \geq \tau$. Let $\tilde{\boldsymbol{\mu}}$ and \tilde{T}_K be the corresponding controlled process and stopping time. Then the first claim of the lemma follows. The remaining claim follows from the following display, where the first inequality uses again that $C_{xy}^n(1) = 0$, the following equality uses that $(\tilde{\mathbf{u}}, \tilde{\boldsymbol{\mu}}, \tilde{T}_K)$ had the same distribution as the original versions up till time τ , and the second inequality uses (43):

$$\begin{aligned} I_K^n(\mathbf{m}, \tilde{\mathbf{u}}) &= \mathbb{E}_{\mathbf{m}} \left[e^{n \int_0^{T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \tilde{\mu}_x(t) \gamma_{xy} C_{xy}^n \left(\frac{\tilde{u}_{xy}(t)}{\gamma_{xy}} \right) - R^n(\tilde{\boldsymbol{\mu}}(t)) \right) dt} \right] \\ &\leq \mathbb{E}_{\mathbf{m}} \left[I_{\{T_K \leq \tau\}} e^{n \int_0^{T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \tilde{\mu}_x(t) \gamma_{xy} C_{xy}^n \left(\frac{\tilde{u}_{xy}(t)}{\gamma_{xy}} \right) - R^n(\tilde{\boldsymbol{\mu}}(t)) \right) dt} \right] \\ &\quad + \mathbb{E}_{\mathbf{m}} \left[I_{\{T_K \geq \tau\}} e^{n \int_0^{T_K \wedge \tau} \left(\sum_{(x,y) \in \mathcal{Z}} \tilde{\mu}_x(t) \gamma_{xy} C_{xy}^n \left(\frac{\tilde{u}_{xy}(t)}{\gamma_{xy}} \right) - R^n(\tilde{\boldsymbol{\mu}}(t)) \right) dt} \right] \\ &= \mathbb{E}_{\mathbf{m}} \left[I_{\{T_K \leq \tau\}} e^{n \int_0^{T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(t) \gamma_{xy} C_{xy}^n \left(\frac{u_{xy}(t)}{\gamma_{xy}} \right) - R^n(\boldsymbol{\mu}(t)) \right) dt} \right] \\ &\quad + \mathbb{E}_{\mathbf{m}} \left[I_{\{T_K \geq \tau\}} e^{n \int_0^{T_K \wedge \tau} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(t) \gamma_{xy} C_{xy}^n \left(\frac{u_{xy}(t)}{\gamma_{xy}} \right) - R^n(\boldsymbol{\mu}(t)) \right) dt} \right] \\ &\quad \times \frac{\mathbb{E}_{\mathbf{m}} \left[e^{n \int_{T_K \wedge \tau}^{T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(t) \gamma_{xy} C_{xy}^n \left(\frac{u_{xy}(t)}{\gamma_{xy}} \right) - R^n(\boldsymbol{\mu}(t)) \right) dt} \Big| \mathcal{F}_{\tau} \right]}{\mathbb{E}_{\mathbf{m}} \left[e^{n \int_{T_K \wedge \tau}^{T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(t) \gamma_{xy} C_{xy}^n \left(\frac{u_{xy}(t)}{\gamma_{xy}} \right) - R^n(\boldsymbol{\mu}(t)) \right) dt} \Big| \mathcal{F}_{\tau} \right]} \Big] \\ &\leq \mathbb{E}_{\mathbf{m}} \left[e^{n \int_0^{T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(t) C_{xy}^n \left(\frac{u_{xy}(t)}{\gamma_{xy}} \right) - R^n(\boldsymbol{\mu}(t)) \right) dt} \right] \\ &\quad + \frac{1}{c} \mathbb{E}_{\mathbf{m}} \left[I_{\{T_K \geq \tau\}} e^{n \int_0^{T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(t) \gamma_{xy} C_{xy}^n \left(\frac{u_{xy}(t)}{\gamma_{xy}} \right) - R^n(\boldsymbol{\mu}(t)) \right) dt} \right] \\ &\leq I_K^n(\mathbf{m}, \mathbf{u}) + \epsilon. \end{aligned}$$

■

Proof of Theorem 5. We are first going to prove that \tilde{V}_K^n is the unique solution to (29). Let \tilde{V} be any solution to (29), and let $\mathbf{m} \in \mathcal{P}(\mathcal{X})$. Let also $\mathbf{q} \in \mathcal{A}_b^{n,|\mathcal{Z}|}$ be given and let $\boldsymbol{\mu}$ solve (21). By Lemma 7,

$$\tilde{V}(\boldsymbol{\mu}(t \wedge T_K)) - \tilde{V}(\mathbf{m}) - \int_0^{t \wedge T_K} \sum_{(x,y) \in \mathcal{Z}} \mu_x(s) q_{xy}(s) \Delta^n \tilde{V}(\boldsymbol{\mu}(s)) ds$$

is a martingale. Taking expectation gives

$$\mathbb{E}_{\mathbf{m}} \left[\tilde{V}(\boldsymbol{\mu}(t \wedge T_K)) \right] - \mathbb{E}_{\mathbf{m}} \left[\int_0^{t \wedge T_K} \sum_{(x,y) \in \mathcal{Z}} \mu_x(s) q_{xy}(s) \Delta^n \tilde{V}(\boldsymbol{\mu}(s)) ds \right] = \tilde{V}(\mathbf{m}),$$

and since \tilde{V} is a solution to (29) and by (27),

$$\mathbb{E}_{\mathbf{m}} \left[\tilde{V}(\boldsymbol{\mu}(t \wedge T_K)) \right] + \mathbb{E}_{\mathbf{m}} \left[\int_0^{t \wedge T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(s) F_{xy}^n(q_{xy}(s)) + R^n(\boldsymbol{\mu}(s)) \right) ds \right] \geq \tilde{V}(\mathbf{m}).$$

By Lemma 9, $T_K < \infty$ almost surely. Letting $t \rightarrow \infty$, Lemma 4 and the monotone convergence theorem imply

$$J_K^n(\mathbf{m}, \mathbf{q}) = \mathbb{E}_{\mathbf{m}} \left[\int_0^{T_K} \sum_{(x,y) \in \mathcal{Z}} \mu_x(t) F_{xy}^n(q_{xy}(s)) + R^n(\boldsymbol{\mu}(s)) ds \right] \geq \tilde{V}(\mathbf{m}).$$

Since $\mathbf{q} \in \mathcal{A}_b^{n,|\mathcal{Z}|}$ was arbitrary we get $V_K^n(\mathbf{m}) \geq \tilde{V}(\mathbf{m})$.

Since $\mathbf{q} \in \mathcal{A}_b^{n,|\mathcal{Z}|}$ was arbitrary we get $V_K^n(\mathbf{m}) \geq \tilde{V}(\mathbf{m})$. For each $\mathbf{m} \in \mathcal{P}^n(\mathcal{X})$ let $\bar{\mathbf{q}}(\mathbf{m})$ satisfy

$$\sum_{(x,y) \in \mathcal{Z}} \left(\bar{q}_{xy}(\mathbf{m}) n \left(\tilde{V} \left(\mathbf{m} + \frac{1}{n} v_{xy} \right) - \tilde{V}(\mathbf{m}) \right) + m_x F_{xy}(\bar{q}_{xy}(\mathbf{m})) \right) + R(\mathbf{m}) \leq \epsilon \sum_{(x,y) \in \mathcal{Z}} m_x F_{xy}(\bar{q}_{xy}(\mathbf{m})). \quad (44)$$

To see that such a $\bar{\mathbf{q}}(\mathbf{m})$ exists and it is actually bounded away from zero, we take a minimizing sequence of $\bar{\mathbf{q}}_n(\mathbf{m})$, where each $q_{xy,n}$ can be considered positive by continuity of both sides of (44) with respect to $\bar{\mathbf{q}}_n(\mathbf{m})$. With no loss of generality we can assume that the sequence is converging. If all elements converge to the original rates we just take those and the inequality is satisfied trivially. If on the other hand it converges to different values the right hand will be always bounded away from zero while the left hand will converge to zero, therefore for sufficiently large value of n , we will recover the desirable control. We can construct a solution to (21) with \mathbf{u} replaced by the feedback control $\bar{\mathbf{q}}(\boldsymbol{\mu})$, and then obtain $\hat{\mathbf{q}} \in \mathcal{A}_b^{n,|\mathcal{Z}|}$ by setting $\hat{\mathbf{q}}(t) = \bar{\mathbf{q}}(\boldsymbol{\mu}(t))$. Then

$$\mathbb{E}_{\mathbf{m}} \left[\tilde{V}(\boldsymbol{\mu}(t \wedge T_K)) \right] - \mathbb{E}_{\mathbf{m}} \left[\int_0^{t \wedge T_K} \sum_{(x,y) \in \mathcal{Z}} \mu_x(s) \bar{q}_{xy}(\boldsymbol{\mu}(s)) \Delta^n \tilde{V}(\boldsymbol{\mu}(s)) ds \right] = \tilde{V}(\mathbf{m}),$$

and therefore by (44)

$$\mathbb{E}_{\mathbf{m}} \left[\tilde{V}(\boldsymbol{\mu}(t \wedge T_K)) \right] + \mathbb{E}_{\mathbf{m}} \left[\int_0^{t \wedge T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(t) F_{xy}^n(\bar{q}_{xy}(\boldsymbol{\mu}(s))) + R^n(\boldsymbol{\mu}(s)) - \epsilon R_{\min}^n \right) ds \right] \leq \tilde{V}(\mathbf{m}).$$

Again using Lemma 9 and the monotone convergence theorem gives

$$(1 - \epsilon) \mathbb{E}_{\mathbf{m}} \left[\int_0^{T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(t) F_{xy}^n(\bar{q}_{xy}(\boldsymbol{\mu}(s))) + R^n(\boldsymbol{\mu}(s)) \right) ds \right] \leq \tilde{V}(\mathbf{m}),$$

and therefore $V_K^n(\mathbf{m}) \leq J_K^n(\mathbf{m}, \hat{\mathbf{q}}) \leq \frac{1}{1-\epsilon} \tilde{V}(\mathbf{m})$. Since ϵ is arbitrary we get $V_K^n(\mathbf{m}) = \tilde{V}(\mathbf{m})$, which implies the uniqueness of \tilde{V} .

We now proceed with the proof that W_K^n is the unique solution to

$$\sup_{\mathbf{u} \in (0, \infty)^{|\mathcal{Z}|}} \left\{ \sum_{(x,y) \in \mathcal{Z}} \mu_x \left(u_{xy} \left(\frac{W(\boldsymbol{\mu}) - W(\boldsymbol{\mu} + \frac{v_{xy}}{n})}{W(\boldsymbol{\mu})} \right) - \gamma_{xy} C_{xy}^n \left(\frac{u_{xy}}{\gamma_{xy}} \right) \right) \right\} = -R^n(\boldsymbol{\mu}). \quad (45)$$

Since V_K^n is a solution to (29), by Lemma 1 we get that $\frac{1}{n} \log(V_K^n)$ is a solution to (45), and thus uniqueness will imply $\frac{1}{n} \log(V_K^n) = W_K^n$.

Let \tilde{W} be any solution to (45), $m \in \mathcal{P}^n(\mathcal{X})$, and $\mathbf{u} \in \mathcal{A}_b^{n,|\mathcal{Z}|}$, and let $\boldsymbol{\mu}$ solve (21). Further assume that there exists $\tau < \infty$ such that for $t > \tau$

$$\sum_{(x,y) \in \mathcal{Z}} \mu_x(t) \gamma_{xy} C_{xy}^n \left(\frac{u_{xy}(t)}{\gamma_{xy}} \right) - R^n(\boldsymbol{\mu}(t)) \leq 0. \quad (46)$$

To show $J_K^n(\mathbf{m}, \mathbf{u}) \geq \tilde{W}(\mathbf{m})$ we can assume that $J_K^n(\mathbf{m}, \mathbf{u}) < \infty$, since otherwise there is nothing to prove. By Lemma 8

$$\frac{\tilde{W}(\boldsymbol{\mu}(t \wedge T_K))}{\tilde{W}(\mathbf{m})} \exp \left\{ - \int_0^{t \wedge T_K} \sum_{(x,y) \in \mathcal{Z}} \mu_x(s) u_{xy}(s) \frac{\Delta^n \tilde{W}(\boldsymbol{\mu}(s))}{\tilde{W}(\boldsymbol{\mu}(s))} ds \right\}$$

is a martingale. Taking expectations gives

$$\mathbb{E}_m \left[\tilde{W}(\boldsymbol{\mu}(t \wedge T_K)) \exp \left\{ - \int_0^{t \wedge T_K} \sum_{(x,y) \in \mathcal{Z}} \mu_x(s) u_{xy}(s) \frac{\Delta^n \tilde{W}(\boldsymbol{\mu}(s))}{\tilde{W}(\boldsymbol{\mu}(s))} ds \right\} \right] = \tilde{W}(\mathbf{m}),$$

and by (29) and the definition of Δ^n

$$\mathbb{E}_m \left[\tilde{W}(\boldsymbol{\mu}(t \wedge T_K)) \exp \left\{ n \int_0^{t \wedge T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(s) \gamma_{xy} C_{xy}^n \left(\frac{u_{xy}(s)}{\gamma_{xy}} \right) - R^n(\boldsymbol{\mu}(s)) \right) ds \right\} \right] \geq \tilde{W}(\mathbf{m}).$$

We claim that

$$\mathbb{E}_m \left[\tilde{W}(\boldsymbol{\mu}(t \wedge T_K)) \exp \left\{ n \int_0^{\tau \wedge T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(s) \gamma_{xy} C_{xy}^n \left(\frac{u_{xy}(s)}{\gamma_{xy}} \right) - R^n(\boldsymbol{\mu}(s)) \right) ds \right\} \right] < \infty. \quad (47)$$

Since \tilde{W} is uniformly bounded this term can be ignored. One can then bound what remains in (47) by using

$$\infty > J_K^n(\mathbf{m}, \mathbf{u}) = \mathbb{E}_m \left[\exp \left\{ n \int_0^{T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(s) \gamma_{xy} C_{xy}^n \left(\frac{u_{xy}(s)}{\gamma_{xy}} \right) - R^n(\boldsymbol{\mu}(s)) \right) ds \right\} \right],$$

breaking the integral over $[0, T_K]$ into contributions over $[0, \tau \wedge T_K]$ and $[\tau \wedge T_K, T_K]$, and then conditioning on \mathcal{F}_τ and using the lower bound on the term corresponding to $[\tau \wedge T_K, T_K]$ provided by Lemma 9 (as in the proof of Lemma 10). Since (by Lemma 9) T_K is finite almost surely, and (46) holds for $t \geq \tau$, by dominated convergence theorem and (47) it follows that

$$J_K^n(\mathbf{m}, \mathbf{u}) = \mathbb{E} \left[\exp \left\{ n \int_0^{T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(s) \gamma_{xy} C_{xy}^n \left(\frac{u_{xy}(s)}{\gamma_{xy}} \right) - R^n(\boldsymbol{\mu}(s)) \right) ds \right\} \right] \geq \tilde{W}(\mathbf{m}).$$

By minimizing over all \mathbf{u} that satisfy (46) and applying Lemma 10, we get $W_K^n(\mathbf{m}) \geq \tilde{W}(\mathbf{m})$.

Next let $\epsilon \in (0, 1/2)$. For $\mathbf{m} \in \mathcal{P}^n(\mathcal{X})$, $t \geq 0$ we choose $\bar{u}(\mathbf{m}, t)$ such that

$$\sum_{(x,y) \in \mathcal{Z}} m_x \left(\bar{u}_{xy}(\mathbf{m}, t) \left(\frac{\tilde{W}(\mathbf{m}) - \tilde{W}(\mathbf{m} + \frac{\mathbf{v}_{xy}}{n})}{\tilde{W}(\mathbf{m})} \right) - \gamma_{xy} C_{xy}^n \left(\frac{\bar{u}_{xy}(\mathbf{m}, t)}{\gamma_{xy}} \right) \right) \geq -R^n(\mathbf{m}) - \frac{\epsilon}{t^2 + 1}. \quad (48)$$

As before we can solve (21) and then generate a corresponding element \mathbf{u} of $\mathcal{A}_b^{n,|\mathcal{Z}|}$ by composing $\bar{u}_{xy}(\mathbf{m}, t)$ with the solution. It is easy to see that \mathbf{u} is an element of $\mathcal{A}_b^{n,|\mathcal{Z}|}$, since very big or very small values of $\bar{u}_{xy}(\mathbf{m}, t)$ will make the left hand of (48) tend to $-\infty$. Arguing as before, for fixed $t < \infty$

$$\mathbb{E}_m \left[\tilde{W}(\boldsymbol{\mu}(t \wedge T_K)) \exp \left\{ n \int_0^{T_K \wedge t} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(s) \gamma_{xy} C_{xy}^n \left(\frac{\bar{u}_{xy}(\boldsymbol{\mu}(s), s)}{\gamma_{xy}} \right) - R^n(\boldsymbol{\mu}(s)) - \frac{\epsilon}{s^2 + 1} \right) ds \right\} \right] \leq \tilde{W}(\mathbf{m}).$$

By sending $t \rightarrow \infty$ and using the boundary condition, Fatou's lemma gives

$$\mathbb{E}_m \left[\exp \left(\int_0^\infty -\frac{\epsilon}{s^2 + 1} ds \right) \exp \left\{ n \int_0^{T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(s) \gamma_{xy} C_{xy}^n \left(\frac{\bar{u}_{xy}(\boldsymbol{\mu}(s), s)}{\gamma_{xy}} \right) - R^n(\boldsymbol{\mu}(s)) \right) ds \right\} \right] \leq \tilde{W}(\mathbf{m}),$$

from which we get $W_K^n(\mathbf{m}) \leq \tilde{W}(\mathbf{m}) \exp[\epsilon \int_0^\infty 1/(s^2 + 1) ds]$. Sending ϵ to zero shows $W_K^n(\mathbf{m}) \leq \tilde{W}(\mathbf{m})$.

The proof that $\mathcal{V}_K^n(\mathbf{x}^n) = -\frac{1}{n} \log(W_K^n(\mathbf{x}^n))$ is similar and thus omitted. It remains only to prove $V_K^n(L(\mathbf{x}^n)) = \mathcal{V}_K^n(\mathbf{x}^n)$. We have established that V_K^n is the only function that satisfies

$$\inf_{\mathbf{q} \in (0, \infty)^{n|Z|}} \left\{ \sum_{(x,y) \in Z} m_x (q_{xy} \Delta_{xy}^n V_K^n(\mathbf{m}) + F_{xy}^n(q_{xy})) \right\} = -R^n(\mathbf{m}), \quad (49)$$

and that \mathcal{V}_K^n is the only function that satisfies

$$\inf_{\mathbf{q} \in (0, \infty)^{n|Z|}} \left\{ \sum_{i=1}^n \sum_{y \in Z_{x_i^n}} (q_{x_i^n y} \Delta_{i, x_i^n y}^n \mathcal{V}_K^n(\mathbf{x}^n) + F_{x_i^n y}^n(q_{x_i^n y})) \right\} = -nR^n(L(\mathbf{x}^n)). \quad (50)$$

Since $K \subset \mathcal{X}^n$ is invariant under permutations, and therefore can be identified with a subset of $\mathcal{P}^n(\mathcal{X})$, we have that there exists a function $\bar{V} : \mathcal{P}^n(\mathcal{X}) \rightarrow [0, \infty)$ such that $\bar{V}(L(\mathbf{x}^n)) = \mathcal{V}_K^n(\mathbf{x}^n)$, and therefore (50) becomes

$$\inf_{\mathbf{q} \in (0, \infty)^{n|Z|}} \left\{ \sum_{i=1}^n \sum_{y \in Z_{x_i^n}} (q_{x_i^n y} \Delta_{i, x_i^n y}^n \bar{V}(L(\mathbf{x}^n)) + F_{x_i^n y}^n(q_{x_i^n y})) \right\} = -nR^n(L(\mathbf{x}^n)).$$

For $\epsilon > 0$, let $\bar{\mathbf{q}} \in (0, \infty)^{n|Z|}$ satisfy

$$\sum_{i=1}^n \sum_{y \in Z_{x_i^n}} [\bar{q}_{x_i^n y} \Delta_{i, x_i^n y}^n \bar{V}(L(\mathbf{x}^n)) + F_{x_i^n y}^n(\bar{q}_{x_i^n y})] \leq -nR^n(L(\mathbf{x}^n)) + \epsilon.$$

Now pick $\tilde{\mathbf{q}} \in (0, \infty)^{|Z|}$ by requiring $nL_x(\mathbf{x}^n)\tilde{q}_{xy} = \sum_{i=1}^n I_{x_i^n=x} \bar{q}_{x_i^n y}$, so that

$$\sum_{(x,y) \in Z} nL_x(\mathbf{x}^n)\tilde{q}_{xy} \Delta_{xy}^n \bar{V}(L(\mathbf{x}^n)) + \sum_{i=1}^n \sum_{y \in Z_{x_i^n}} F_{x_i^n y}^n(\bar{q}_{x_i^n y}) \leq -nR^n(L(\mathbf{x}^n)) + \epsilon.$$

By using convexity of F_{xy}^n (see Lemma 4) we get

$$\sum_{(x,y) \in Z} L_x(\mathbf{x}^n) [\tilde{q}_{xy} \Delta_{xy}^n \bar{V}(L(\mathbf{x}^n)) + F_{xy}^n(\tilde{q}_{xy})] \leq -R^n(L(\mathbf{x}^n)) + \epsilon/n,$$

and sending $\epsilon \downarrow 0$ gives

$$\inf_{\mathbf{q} \in (0, \infty)^{|Z|}} \left\{ \sum_{(x,y) \in Z} L_x(\mathbf{x}^n) [q_{xy} \Delta_{xy}^n \bar{V}(L(\mathbf{x}^n)) + F_{xy}^n(q_{xy})] \right\} \leq -R^n(L(\mathbf{x}^n)).$$

The other direction is trivial, and follows if in (50) one uses rates that are the same for all agents in the same position. ■

4 Discussion regarding convergence

Before we introduce the deterministic control problem, we define the set of admissible controls and controlled trajectories.

Definition 11 We define the space of paths and controls by

$$\mathcal{C} \doteq \{(\boldsymbol{\mu}, \mathbf{q}) \in \mathcal{D}([0, \infty); \mathcal{P}(\mathcal{X})) \times \mathcal{F}([0, \infty); [0, \infty)^{\otimes Z}) : \mu_x q_{xy} \text{ is locally integrable } \forall (x, y) \in Z\}, \quad (51)$$

where $\mathcal{F}([0, \infty); [0, \infty)^{\otimes Z})$ was defined in (13). We define $\Lambda : \mathcal{C} \times \mathcal{P}(\mathcal{X}) \rightarrow \mathcal{D}([0, \infty); \mathcal{H})$ by

$$\Lambda(\boldsymbol{\mu}, \mathbf{q}, \mathbf{m})(t) \doteq \mathbf{m} + \sum_{(x,y) \in Z} \mathbf{v}_{xy} \int_{[0,t)} \mu_x(s) q_{xy}(s) ds. \quad (52)$$

Also we define the set of all deterministic pairs that correspond to a solution of the equation $\mu = \Lambda(\mu, q, \mathbf{m})$, i.e.,

$$\mathcal{T}_{\mathbf{m}} \doteq \{(\mu, q) \in \mathcal{C} : \mu = \Lambda(\mu, q, \mathbf{m}), \mu(0) = \mathbf{m}\}$$

Finally we introduce the set of controls that generate controlled trajectories

$$\mathcal{U}_{\mathbf{m}} \doteq \{q \in \mathcal{F}([0; \infty); [0, \infty)^{\otimes \mathcal{Z}}) : \exists \mu \in \mathcal{D}([0, \infty); \mathcal{P}(\mathcal{X})) \text{ such that } (\mu, q) \in \mathcal{T}_{\mathbf{m}}\}. \quad (53)$$

Then the deterministic control problems are given by

$$V_K(\mathbf{m}) \doteq \inf_{(\mu, q) \in \mathcal{T}_{\mathbf{m}}} J_K(\mathbf{m}, \mu, q) \quad (54)$$

$$J_K(\mathbf{m}, \mu, q) \doteq \left\{ \int_0^{T_K} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(t) \hat{F}_{xy}^\infty(q_{xy}(t)) + R^\infty(\mu(t)) \right) dt \right\}, \quad (55)$$

$$F^\infty(q) \doteq \inf \left\{ \liminf_{n \rightarrow \infty} F^n(q_n) : \{q^n\} \text{ sequence in } [0, \infty) \text{ with } q_n \rightarrow q \right\}, \quad (56)$$

with

$$\hat{F}^\infty(q) \doteq \sup \{F(q) : F \text{ convex and } F \leq F^\infty\} \quad (57)$$

and $T_K \doteq \inf_{t \in [0, \infty]} \{\mu(t) \in K\}$.

In this section we consider sets $K \subset \mathcal{P}(\mathcal{X})$ that satisfy the following assumption.

Assumption 12 $K = \overline{K^\circ} \neq \emptyset$.

For such sets we show that the sequence of values functions V_K^n converges uniformly to the function V_K . To simplify the notation we will drop the index that corresponds to the set from the stopping time. We split the study of the convergence in two parts. In the first part, without making any extra assumptions on the cost functions and in great generality, we prove that for any sequence $\{\mathbf{m}^n\}$, with $\mathbf{m}^n \in \mathcal{P}^n(\mathcal{X})$ converging in $\mathbf{m} \in \mathcal{P}(\mathcal{X})$,

$$\liminf_{n \rightarrow \infty} V_K^n(\mathbf{m}^n) \geq V_K(\mathbf{m}).$$

The other direction of the inequality, i.e., $\limsup_{n \rightarrow \infty} V_K^n(\mathbf{m}^n) \leq V_K(\mathbf{m})$, is not as straightforward and its analysis can be quite involved. In order to avoid technical issues relating to controllability we will add some assumptions.

Before we present the extra assumptions on $\{C^n\}$ we discuss an almost trivial choice for the cost function that does not depend on n and that will motivate these extra assumptions. As stated in Lemma 3, for every $(x, y) \in \mathcal{Z}$ we have $C_{xy}^n(u) \geq -\log u + u - u$. Actually the function $C_{xy}^n(u) = -\log u + u - 1$ satisfies Assumption 2 and therefore is an eligible cost function.

Setting $C_{xy}^n(u) \equiv \bar{C}(u) = -\log u + u - 1$, we get

$$\begin{aligned} G_{xy}^n(u, q) &= u \ell\left(\frac{q}{u}\right) - \gamma_{xy} C_{xy}^n\left(\frac{u}{\gamma_{xy}}\right) = q \log \frac{q}{u} - q + u + \gamma_{xy} \log \frac{u}{\gamma_{xy}} - u + \gamma_{xy} \\ &= q \log q + (\gamma_{xy} - q) \log u - q + \gamma_{xy}. \end{aligned} \quad (58)$$

Examining (58) and referring to the definition of F_{xy}^n in (9), we observe that if $q_{xy} > \gamma_{xy}$ then the ‘‘maximizing player’’ (the one that picks u), can produce an arbitrarily large cost by making u_{xy} as **small** as needed. If $q_{xy} < \gamma_{xy}$, this player can produce an arbitrarily large cost by making u_{xy} as **big** as needed. Hence the minimizing player must keep $q_{xy} = \gamma_{xy}$, and the value function $V(\mathbf{m})$ is infinite unless the solution of the equation $\dot{\nu}(t) = \nu(t)\gamma$ passes through K for the specific choice of initial data \mathbf{m} . To resolve this difficulty we could start by imposing the following assumption on the cost.

$$\lim_{u \rightarrow 0} \sup_{n \in \mathbb{N}} u(C_{xy}^n)'(u) = -\infty, \quad \lim_{u \rightarrow \infty} \inf_{n \in \mathbb{N}} \{u(C_{xy}^n)'(u) - u\} \geq 0.$$

This Assumption makes \hat{F}^∞ finite on $(0, \infty)$ and allows for some controllability. Specifically, if the first point is true and if $\mathbf{m}, \tilde{\mathbf{m}} \in \mathcal{P}_a(\mathcal{X})$ for some $a > 0$, then one can observe (see the proof of Lemma 4) that the total cost $V_{\{\tilde{\mathbf{m}}\}}(\mathbf{m})$ for moving from point \mathbf{m} to $\tilde{\mathbf{m}}$ is uniformly bounded by $c_a \|\mathbf{m} - \tilde{\mathbf{m}}\|$, where $c_a > 0$ is an appropriate constant, where the minimizing player picks $\tilde{q}_{xy}(t)$ to be uniformly bounded from above, but big enough to reach the desired point. In particular, the maximizing player cannot impose an arbitrarily large cost by taking u_{xy} small. In an analogous fashion, the second point implies that the minimizer can choose controls so that the total cost $V_{\{\tilde{\mathbf{m}}\}}(\mathbf{m})$ for moving from point \mathbf{m} to $\tilde{\mathbf{m}}$ is uniformly bounded by $c'_a \|\mathbf{m} - \tilde{\mathbf{m}}\|$ by picking $\tilde{q}_{xy}(t)$ bounded from below but small enough.

However, if $\tilde{\mathbf{m}}$ is in the natural boundary of the simplex $\mathcal{P}(\mathcal{X})$ an additional complication arises, because to reach the natural boundary it must be true that for at least one $(x, y) \in \mathcal{Z}$ the quantity $\tilde{q}_{xy}(t)$ will scale like $1/\tilde{\mu}_x(t)$. In that case, the first point is not enough for a finite cost, since sending $\tilde{q}_{xy}(t)$ to infinity in order to reach the natural boundary may result in an infinite total cost. Taking all these issues into account we end up with the following assumption.

Assumption 13 For every $n \in \mathbb{N}$, let C^n, R^n be as in Assumption 2. Assume that there exist convex functions $C^\infty : (0, \infty)^{\otimes \mathcal{Z}} \rightarrow [0, \infty]$, $R^\infty : \mathcal{P}(\mathcal{X}) \rightarrow [0, \infty)$, such that $\{R^n\}_n$ converges uniformly to R^∞ , and for all $(x, y) \in \mathcal{Z}$, the following are valid.

1 For every $(x, y) \in \mathcal{Z}$, C_{xy}^n and C_{xy}^∞ have a common maximal interval $(u_{1,xy}, u_{2,xy}) \subset (0, \infty)$ where they are finite, which contain 1 and for which C_{xy}^n converges locally uniformly to C_{xy}^∞ .

2 There exists $p > 0$ such that

$$\limsup_{u \rightarrow 0} \sup_{n \in \mathbb{N}} u^{p+1} (C_{xy}^n)'(u) = -\infty. \quad (59)$$

3

$$\liminf_{u \rightarrow \infty} \inf_{n \in \mathbb{N}} \{u (C_{xy}^n)'(u) - u\} \geq 0. \quad (60)$$

Now we state the second main theorem of the paper.

Theorem 14 Let $\{C^n\}_n, \{R^n\}_n, C^\infty$, and R^∞ satisfy Assumptions 2 and 13. Let also K be a closed subset of $\mathcal{P}(\mathcal{X})$ that satisfies Assumption 12. Finally assume that in every compact subset of K^c , R^∞ , is bounded from below by a positive constant. Then the sequence of functions V_K^n defined in (24) converges uniformly to V_K defined in (54).

Before proceeding with the proof, we state some properties of F_{xy}^n .

Lemma 15 For every $n \in \mathbb{N}$ and for every $(x, y) \in \mathcal{Z}$, let F_{xy}^n be as in (9), where $\{C_{xy}^n\}$ satisfy Assumption 13. Then the following hold.

1 There exists a constant $M \in (0, \infty)$ and a decreasing function $\bar{M} : (0, \infty) \rightarrow (0, \infty)$, such that for every $\epsilon > 0$ and every $q \geq \epsilon$,

$$F_{xy}^n(q) \leq q \log \frac{q}{\min \{ \gamma_{xy} (\gamma_{xy}/q)^{1/p}, M \}} + \bar{M}(\epsilon).$$

2 F_{xy}^n is continuous on the interval $(0, \infty)$, and continuous as an extended function on $[0, \infty)$.

3 F_{xy}^n converges locally uniformly, on the set $(0, \infty)$, to the function

$$\bar{F}_{xy}(q) = \sup_{u \in (0, \infty)} \left\{ u \ell \left(\frac{q}{u} \right) - \gamma_{xy} C_{xy}^\infty \left(\frac{u}{\gamma_{xy}} \right) \right\}.$$

Furthermore, we have $\bar{F}_{xy} = F_{xy}^\infty$, where F^∞ was defined in (56). Finally F_{xy}^∞ is convex on the whole domain $[0, \infty)$ and therefore $\bar{F}_{xy} = F_{xy}^\infty = \hat{F}_{xy}^\infty$.

The proof of the Lemma 15 can be found in Appendix A. It is worth mentioning that it is possible that $F_{xy}^n(0) = \infty$. In the sequel we will make use of the following remark, which states a property proved in [12, Proposition 4.14]

Remark 16 *There exists $D \geq 1$ and $b_1 > 0, b_2 < \infty$ such that for every $\mathbf{m} \in \mathcal{P}(X)$, if $\nu(\mathbf{m}, t)$ is the solution of $\dot{\nu}(t) = \nu(t)\gamma$ with initial point $\nu(0) = \mathbf{m}$, then*

$$1. \forall x \in \mathcal{X}, \nu_x(\mathbf{m}, t) \geq b_1 t^D \quad \text{and} \quad 2. \|\nu(\mathbf{m}, t) - \mathbf{m}\| \leq b_2 t.$$

Before proceeding with the proof of Theorem 14, we prove that the function $V(\mathbf{m})$ is continuous. We will actually prove something stronger. Recall that γ denotes the original unperturbed jump rates and the definitions of $\mathcal{P}_*(\mathcal{X})$ and $\mathcal{P}_a(\mathcal{X})$ in (14), (15).

Theorem 17 *There is a constant \bar{c} that depends only the dimension d and the unperturbed rates γ , such that for every $\mathbf{m} \in \mathcal{P}_*(\mathcal{X})$, $\tilde{\mathbf{m}} \in \mathcal{P}(\mathcal{X})$ there exists a control $\mathbf{q} \in \mathcal{U}_m$, that generates a unique μ with $(\mu, \mathbf{q}) \in \mathcal{T}_m$, satisfying*

- 1 μ is a constant speed parametrization of the straight line that connects \mathbf{m} and $\tilde{\mathbf{m}}$,
- 2 the exit time $T_{\{\tilde{\mathbf{m}}\}}$ is equal to $\|\mathbf{m} - \tilde{\mathbf{m}}\|$,
- 3 $\gamma_{xy} \leq q_{xy}(t)$ and $\mu_x(t)q_{xy}(t) \leq \bar{c}$.

Furthermore, if $\mathbf{m}, \tilde{\mathbf{m}} \in \mathcal{P}_a(\mathcal{X})$ then

$$\gamma_{xy} \leq q_{xy}(t) \leq \frac{\bar{c}}{a}, \quad (61)$$

and we can find a constant $c_a < \infty$ such that the total cost for applying the control is bounded above by $c_a \|\mathbf{m} - \tilde{\mathbf{m}}\|$. Finally, for every $\epsilon > 0$ there exists $\delta > 0$, such that $\|\tilde{\mathbf{m}} - \tilde{\mathbf{m}}\| \leq \delta$ implies $V_{\{\tilde{\mathbf{m}}\}}(\tilde{\mathbf{m}}), V_{\{\tilde{\mathbf{m}}\}}(\tilde{\mathbf{m}}) \leq \epsilon$, and therefore as a function of two variables V is continuous on $\mathcal{P}(\mathcal{X}) \times \mathcal{P}(\mathcal{X})$.

Proof. Recall the definitions above (3), and let $\mathbf{m} \in \mathcal{P}_*(\mathcal{X})$, $\tilde{\mathbf{m}} \in \mathcal{P}(\mathcal{X})$. We can find a positive constant \bar{c} that depend only the dimension d and on the rates γ , and also rates \mathbf{q} such that

$$1. q_{xy} \geq \gamma_{xy}, \quad 2. \sum_{(x,y) \in \mathcal{Z}} m_x q_{xy} \mathbf{v}_{xy} = \frac{\tilde{\mathbf{m}} - \mathbf{m}}{\|\tilde{\mathbf{m}} - \mathbf{m}\|}, \quad 3. \max\{m_x q_{xy}, (x, y) \in \mathcal{Z}\} \leq \bar{c}.$$

Indeed, since $\{a_{xy} \mathbf{v}_{xy} : a_{xy} \geq 0, (x, y) \in \mathcal{Z}\} = \mathcal{H}$, we can find a constant $c < \infty$ such that for every point $\mathbf{m} \in \mathcal{P}_*(\mathcal{X})$, there exist vectors $q_{xy} m_x \mathbf{v}_{xy}$ with $q_{xy} m_x \leq c$, and $\sum_{(x,y) \in \mathcal{Z}} m_x q_{xy} \mathbf{v}_{xy} = \frac{\tilde{\mathbf{m}} - \mathbf{m}}{\|\tilde{\mathbf{m}} - \mathbf{m}\|}$. Now if for some $(x_1, y_1) \in \mathcal{Z}$ we do not have $q_{x_1 y_1} \geq \gamma_{x_1 y_1}$, then by ergodicity we can pick $x_1, x_2 = y_1, x_3, \dots, x_j$, with $j \leq d$, such that $\sum_{i=1}^{j-1} \mathbf{v}_{x_i x_{i+1}} = 0$. If we pick the new $q_{x_i x_{i+1}}$ equal to $\max_{xy} \{\gamma_{xy}\} / m_{x_i}$ plus the original $q_{x_i x_{i+1}}$, then property 2 is still satisfied, but we now also have $q_{x_1 y_1} \geq \gamma_{x_1 y_1}$. We have to repeat the procedure at most $|\mathcal{Z}|$ times to enforce property 1, and can then set $\bar{c} \doteq \max\{m_x q_{xy}, (x, y) \in \mathcal{Z}\}$.

Let

$$\tilde{\mu}(t) = [(\tilde{\mathbf{m}} - \mathbf{m})t / \|\tilde{\mathbf{m}} - \mathbf{m}\| + \mathbf{m}], \quad (62)$$

and define $\tilde{\mathbf{q}} \in \mathcal{U}_m$ by

$$\tilde{\mu}_x(t) \tilde{q}_{xy}(t) = m_x q_{xy} \leq \bar{c}. \quad (63)$$

Then automatically

$$\sum_{(x,y) \in \mathcal{Z}} \mathbf{v}_{xy} \int_{[0,t)} \tilde{\mu}_x(s) \tilde{q}_{xy}(s) ds = t \frac{\tilde{\mathbf{m}} - \mathbf{m}}{\|\tilde{\mathbf{m}} - \mathbf{m}\|} = \tilde{\mu}(t) - \mathbf{m},$$

and thus $(\tilde{\mu}, \tilde{\mathbf{q}}) \in \mathcal{T}_m$. This will lead to hitting $\{\tilde{\mathbf{m}}\}$ in time $T_{\{\tilde{\mathbf{m}}\}} = \|\mathbf{m} - \tilde{\mathbf{m}}\|$. Using properties stated in Lemma 15

we get

$$\begin{aligned}
& \inf_{(\boldsymbol{\mu}, \mathbf{q}) \in \mathcal{T}_{\tilde{\mathbf{m}}}} J_{\{\tilde{\mathbf{m}}\}}(\mathbf{m}, \boldsymbol{\mu}, \mathbf{q}) \leq J_{\{\tilde{\mathbf{m}}\}}(\mathbf{m}, \tilde{\boldsymbol{\mu}}, \tilde{\mathbf{q}}) \leq \sum_{(x,y) \in \mathcal{Z}} \int_0^{T_{\{\tilde{\mathbf{m}}\}}} \tilde{\mu}_x(t) \hat{F}_{xy}^\infty(\tilde{q}_{xy}(t)) + R_{\max} T_{\{\tilde{\mathbf{m}}\}} \\
& \stackrel{(63)}{\leq} \sum_{(x,y) \in \mathcal{Z}} \int_0^{T_{\{\tilde{\mathbf{m}}\}}} \left(\tilde{\mu}_x(t) \tilde{q}_{xy}(t) \log \frac{\tilde{q}_{xy}(t)}{\min\{\gamma_{xy} (\gamma_{xy}/\tilde{q}_{xy}(t))^{1/p}, M\}} + \max_{(x,y) \in \mathcal{Z}} \bar{M}(\gamma_{xy}) \right) dt + R_{\max} T_{\{\tilde{\mathbf{m}}\}} \\
& \leq \sum_{(x,y) \in \mathcal{Z}} \int_0^{T_{\{\tilde{\mathbf{m}}\}}} |\tilde{\mu}_x(t) \tilde{q}_{xy}(t) \log \tilde{q}_{xy}(t)| dt + \sum_{(x,y) \in \mathcal{Z}} \int_0^{T_{\{\tilde{\mathbf{m}}\}}} |\tilde{\mu}_x(t) \tilde{q}_{xy}(t) \log (\gamma_{xy}/\tilde{q}_{xy}(t))^{1/p}| dt + \\
& + \sum_{(x,y) \in \mathcal{Z}} \int_0^{T_{\{\tilde{\mathbf{m}}\}}} |\tilde{\mu}_x(t) \tilde{q}_{xy}(t) \log \gamma_{xy}| dt + \sum_{(x,y) \in \mathcal{Z}} \int_0^{T_{\{\tilde{\mathbf{m}}\}}} |\tilde{\mu}_x(t) \tilde{q}_{xy}(t) \log M| dt + c' T_{\{\tilde{\mathbf{m}}\}} \tag{64} \\
& \stackrel{(63)}{\leq} \bar{c} \sum_{(x,y) \in \mathcal{Z}} \int_0^{T_{\{\tilde{\mathbf{m}}\}}} |\log \tilde{q}_{xy}(t)| dt + \bar{c} \sum_{(x,y) \in \mathcal{Z}} \int_0^{T_{\{\tilde{\mathbf{m}}\}}} |\log (\gamma_{xy}/\tilde{q}_{xy}(t))^{1/p}| dt + c'' T_{\{\tilde{\mathbf{m}}\}} \\
& \stackrel{(63)}{\leq} \bar{c} \sum_{(x,y) \in \mathcal{Z}} \int_0^{T_{\{\tilde{\mathbf{m}}\}}} \left| \log \frac{m_x q_{xy}}{\tilde{\mu}_x(t)} \right| dt + \bar{c} \sum_{(x,y) \in \mathcal{Z}} \int_0^{T_{\{\tilde{\mathbf{m}}\}}} \left| \log (\tilde{\mu}_x(t) \gamma_{xy}/m_x q_{xy})^{1/p} \right| dt + c'' T_{\{\tilde{\mathbf{m}}\}} \\
& \stackrel{(63)}{\leq} \bar{c} \sum_{(x,y) \in \mathcal{Z}} \int_0^{T_{\{\tilde{\mathbf{m}}\}}} |\log \tilde{\mu}_x(t)| dt + \bar{c} \sum_{(x,y) \in \mathcal{Z}} \int_0^{T_{\{\tilde{\mathbf{m}}\}}} \frac{1}{p} |\log \tilde{\mu}_x(t)| dt + c''' T_{\{\tilde{\mathbf{m}}\}},
\end{aligned}$$

where the constants c' , c'' , c''' depend only on γ , \bar{c} and R_{\max} .

Now if $\mathbf{m}, \tilde{\mathbf{m}} \in \mathcal{P}_a(\mathcal{X})$, then all elements are bounded by a constant c_a (that depends on γ , \bar{c} , R_{\max} , and a) times $T_{\{\tilde{\mathbf{m}}\}} = \|\tilde{\mathbf{m}} - \mathbf{m}\|$, and therefore the first part of the theorem follows.

Let $1 > \delta > 0$, and $\bar{\mathbf{m}}, \tilde{\mathbf{m}} \in \mathcal{P}(\mathcal{X})$, with $\|\bar{\mathbf{m}} - \tilde{\mathbf{m}}\| < \delta$. We take $\mathbf{m} = \nu(\bar{\mathbf{m}}, \delta)$, where $\nu(\bar{\mathbf{m}}, t)$ is the solution of $\dot{\nu}(t) = \nu(t)\gamma$, with initial data $\nu(0) = \bar{\mathbf{m}}$. Now by appropriate use of the inequality $\tilde{\mu}_x(t) \geq \min\{m_x, m_x(T_{\{\tilde{\mathbf{m}}\}} - t)\}$, that we get from (62), and using the last display, we get

$$V_{\{\tilde{\mathbf{m}}\}}(\mathbf{m}) \leq c'''' \left(\sum_{(x,y) \in \mathcal{Z}} \int_0^{T_{\{\tilde{\mathbf{m}}\}}} (|\log m_x| + |\log(T_{\{\tilde{\mathbf{m}}\}} - t)|) dt + T_{\{\tilde{\mathbf{m}}\}} \right).$$

By a simple change of variable and Remark 16, we have

$$V_{\{\tilde{\mathbf{m}}\}}(\mathbf{m}) \leq c'''' \left(\sum_{(x,y) \in \mathcal{Z}} \int_0^{b_2 \delta} (|\log b_1 \delta^D| + |\log t|) dt + b_2 \delta \right). \tag{65}$$

Therefore

$$V_{\{\tilde{\mathbf{m}}\}}(\bar{\mathbf{m}}) \leq V_{\{\mathbf{m}\}}(\bar{\mathbf{m}}) + V_{\{\tilde{\mathbf{m}}\}}(\mathbf{m}) \leq \delta R_{\max} + c'''' \left(\sum_{(x,y) \in \mathcal{Z}} \int_0^{b_2 \delta} (|\log b_1 \delta^D| + |\log t|) dt + b_2 \delta \right),$$

and the right hand side can be made as small as desired by making δ small enough. The estimate for $V_{\{\tilde{\mathbf{m}}\}}(\tilde{\mathbf{m}})$ is proved in a symmetric way. This proves the last statement of the theorem. ■

5 Lower bound

For the proof of Theorem 14, we first prove the lower bound: for every sequence $\mathbf{m}^n \in \mathcal{P}^n(\mathcal{X})$ and $\mathbf{m} \in \mathcal{P}(\mathcal{X})$, with $\mathbf{m}^n \rightarrow \mathbf{m}$, we have

$$\liminf_{n \rightarrow \infty} V_K^n(\mathbf{m}^n) \geq V_K(\mathbf{m}).$$

Without loss of generality we can assume that the liminf is actually a limit, otherwise we can just work with a subsequence. If the limit is ∞ then the conclusion is trivial, therefore we can assume that there is $c \in \mathbb{R}$ such that

$$\sup_{n \in \mathbb{N}} V_K^n(\mathbf{m}^n) \leq c. \quad (66)$$

Let $\epsilon \in (0, 1)$. Recalling (24), let $\mathbf{q}^n \in \mathcal{A}_b^{n, |\mathcal{Z}|}$ be such that

$$\mathbb{E}_{\mathbf{m}^n} \left[\int_0^{T^n} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x^n(t) F_{xy}^n(q_{xy}^n(t)) + R^n(\mu^n(t)) \right) dt \right] < V_K^n(\mathbf{m}^n) + \epsilon, \quad (67)$$

where $\mu^n = h^n(\mu^n, \mathbf{q}^n, \mathbf{m}^n, N^n/n)$ and $T^n \doteq \inf \{t \in [0, \infty) : \mu^n(t) \in K\}$. For $\delta > 0$ such that

$$\|\tilde{\mathbf{m}} - \tilde{\mathbf{m}}\| \leq \delta \Rightarrow V_{\tilde{\mathbf{m}}}(\tilde{\mathbf{m}}) \leq \epsilon, \quad (68)$$

we define

$$K_\delta \doteq \{\mathbf{m} : d(\mathbf{m}, K) \leq \delta\} \quad \text{and} \quad T^{n,\delta} \doteq \inf \{t \in [0, \infty) : \mu^n(t) \in K_\delta\}. \quad (69)$$

The existence of such a δ is given by Theorem 17. Now for μ^n, \mathbf{q}^n as in (67) and $T^{n,\delta}$ as above, we define the sequences $\mu^{n,\delta}(t) = \mu^n(t \wedge T^{n,\delta})$,

$$\mathbf{q}^{n,\delta}(t) = \begin{cases} \mathbf{q}^n(t) & t \leq T^{n,\delta} \\ \gamma & T > T^{n,\delta} \end{cases}. \quad (70)$$

We note that for $t > T^\delta$, $\mathbf{q}^{n,\delta}(t)$ does not actually generate μ^n , but we define it this way to simplify some arguments later on. We will show that

$$\begin{aligned} \liminf_{n \rightarrow \infty} \mathbb{E}_{\mathbf{m}^n} \left[\int_0^{T^n} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x^n(t) F_{xy}^n(q_{xy}^n(t)) + R^n(\mu^n(t)) \right) dt \right] &\geq \\ \liminf_{n \rightarrow \infty} \mathbb{E}_{\mathbf{m}^n} \left[\int_0^{T^{n,\delta}} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x^{n,\delta}(t) F_{xy}^n(q_{xy}^{n,\delta}(t)) + R^n(\mu^{n,\delta}(t)) \right) dt \right] &\geq V_{K_\delta}(\mathbf{m}), \end{aligned} \quad (71)$$

and then by an application of Theorem 17 and (67) deduce $\lim_{n \rightarrow \infty} V_K^n(\mathbf{m}^n) + 2\epsilon \geq V_K(\mathbf{m})$. Since ϵ is arbitrary the lower bound will follow. The first inequality in (71) is true since $F_{xy}^n \geq 0, R^n \geq 0$ and $T^{n,\delta} \leq T^n$. Therefore only the second inequality needs to be proved.

Before proceeding we introduce some auxiliary random measures. For $(x, y) \in \mathcal{Z}, q_{xy} \in \mathcal{F}([0, \infty); [0, \infty))$, and $t \in [0, \infty)$, define

$$\eta_{xy}(dr; t) \doteq \delta_{q_{xy}(t)}(dr) \mu_x(t).$$

For each $t \in [0, \infty), (x, y) \in \mathcal{Z}$ we have that $\eta_{xy}(\cdot; t)$ is a subprobability measure on $[0, \infty)$. Also we consider the measures $\theta_{xy}(drdt) = \eta_{xy}(dr; t) dt$ on $[0, \infty) \times [0, \infty)$ as equipped with the topology that generalizes the weak convergence of probability measures to general measures that have at most mass T on $[0, \infty) \times [0, T]$. This can be defined in terms of a distance (a generalization of the Prohorov metric) d_T , and the metric on measures on $[0, \infty) \times [0, \infty)$ is

$$\sum_{T \in \mathbb{N}} 2^{-T} [d_T(\mu|_T, \nu|_T) \vee 1], \quad (72)$$

where $\mu|_T$ denotes the restriction to $[0, T]$ in the last variable.

Let $\theta^{n,\delta} = \{\theta^{n,\delta}\}_{(x,y) \in \mathcal{Z}}$ be the random measures that correspond to $\mu^{n,\delta}, \mathbf{q}^{n,\delta}$, according to the construction above. We observe that

$$\mu^{n,\delta}(t) = \mathbf{m}^n + \sum_{(x,y) \in \mathcal{Z}} \mathbf{v}_{xy} \int_0^{t \wedge T^{n,\delta}} \int_0^\infty r \theta_{xy}^{n,\delta}(drds) + \text{a martingale},$$

where the martingale will converge to zero as $n \rightarrow \infty$, and that for every $(x, y) \in \mathcal{Z}$,

$$\mathbb{E}_{\mathbf{m}^n} \left[\int_0^{T^{n,\delta}} F_{xy}^n(q_{xy}^{n,\delta}(t)) \mu_x^{n,\delta}(t) dt \right] = \mathbb{E}_{\mathbf{m}^n} \left[\int_0^{T^{n,\delta}} \int_0^\infty F_{xy}^n(r) \theta_{xy}^{n,\delta}(drdt) \right]. \quad (73)$$

We will split the proof of (71) in three parts. First we prove that $(\mu^{n,\delta}, \theta^{n,\delta}, T^{n,\delta})$ is tight. Then we show that for every limit point $(\mu^\delta, \theta^\delta, T^\delta)$, θ_{xy}^δ has the decomposition $\theta_{xy}^\delta(dr dt) = \eta_{xy}^\delta(dr; t)dt$, with $\sum_{y \in \mathcal{X}} \eta_{xy}^\delta([0, \infty); t) = \mu_x^\delta(t)$, and for q^δ defined by $\mu_x^\delta(t)q_{xy}^\delta(t) = \int_0^\infty r \eta_{xy}^\delta(dr; t)$, that

$$\mu^\delta(t) = \mathbf{m} + \sum_{(x,y) \in \mathcal{Z}} \mathbf{v}_{xy} \int_0^{t \wedge T^\delta} \int_0^\infty r \theta_{xy}^\delta(dr ds) = \mathbf{m} + \sum_{(x,y) \in \mathcal{Z}} \mathbf{v}_{xy} \int_0^{t \wedge T^\delta} \mu_x^\delta(s) q_{xy}^\delta(s) ds.$$

Finally, by an application of Fatou's Lemma, for such a q^δ , we get

$$\begin{aligned} \liminf_{n \rightarrow \infty} \mathbb{E}_{\mathbf{m}^n} \left[\int_0^{T^{n,\delta}} \int_0^\infty F_{xy}^n(r) \theta_{xy}^{n,\delta}(dr dt) \right] &\geq \mathbb{E}_{\mathbf{m}} \left[\int_0^{T^\delta} \int_0^\infty F_{xy}^\infty(r) \theta_{xy}^\delta(dr dt) \right] \\ &\geq \mathbb{E}_{\mathbf{m}} \left[\int_0^{T^\delta} \int_0^\infty \hat{F}_{xy}^\infty(r) \theta_{xy}^\delta(dr dt) \right] \geq \mathbb{E}_{\mathbf{m}} \left[\int_0^{T^\delta} \int_0^\infty \hat{F}_{xy}^\infty(r) \eta_{xy}^\delta(dr; t) dt \right] \\ &\geq \mathbb{E}_{\mathbf{m}} \left[\int_0^{T^\delta} \hat{F}_{xy}^\infty \left(\int_0^\infty r \frac{\eta_{xy}^\delta(dr; t)}{\eta_{xy}^\delta([0, \infty); t)} \right) \eta_{xy}^\delta([0, \infty); t) dt \right] = \mathbb{E}_{\mathbf{m}} \left[\int_0^{T^\delta} \hat{F}_{xy}^\infty(q_{xy}^\delta(t)) \mu_x^\delta(t) dt \right], \end{aligned}$$

where in the second inequality, we used the fact that $\hat{F}_{xy}^\infty \leq F_{xy}^\infty$, and for the fourth, we applied Jensen's inequality. Together with $\mu^{n,\delta} \rightarrow \mu^\delta$, $\hat{F}_{xy}^\infty \geq 0$, $R^\infty \geq 0$ and another application of Fatou's Lemma, this gives (71).

5.1 Tightness of $(\mu^{n,\delta}, \theta^{n,\delta}, T^{n,\delta})$

First, we prove that $(\mu^{n,\delta}(\cdot), T^{n,\delta})$, which takes values in $D([0, \infty); \mathcal{P}(\mathcal{X})) \times [0, \infty) \subset D([0, \infty); \mathbb{R}^d) \times [0, \infty)$, is tight. For that, we introduce some auxiliary random variables $\tilde{\mu}^{n,\delta}$ in $D([0, \infty); \mathbb{R}^d)$, to compare with $\mu^{n,\delta}$, given by

$$\tilde{\mu}^{n,\delta}(t) = \mathbf{m}^n + \sum_{(x,y) \in \mathcal{Z}} \mathbf{v}_{xy} \int_0^{t \wedge T^{n,\delta}} \mu_x^n(s) q_{xy}^n(s) ds. \quad (74)$$

Since $\gamma_{xy} \ell(\cdot / \gamma_{xy}) \leq F_{xy}^n(\cdot)$, recalling (66), (67) and that for sufficiently large n R^n is bounded away from zero in $K_\delta = \{\mathbf{m} : d(\mathbf{m}, K) \geq \delta\}$ by a constant R_{\min}^δ , we get

$$\mathbb{E}_{\mathbf{m}^n} \left[\int_0^{T^{n,\delta}} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x^n(t) \gamma_{xy} \ell \left(\frac{q_{xy}^n(t)}{\gamma_{xy}} \right) \right) dt + R_{\min}^\delta T^{n,\delta} \right] \leq c + 1, \quad (75)$$

which shows tightness of $\{T^{n,\delta}\}$. By setting $\gamma_{\max} = \max\{\gamma_{xy} : (x, y) \in \mathcal{Z}\}$, we get

$$\mathbb{E}_{\mathbf{m}^n} \left[\int_0^{T^{n,\delta}} \left(\sum_{(x,y) \in \mathcal{Z}} \gamma_{\max} \frac{\mu_x^n(t) \gamma_{xy}}{\gamma_{\max}} \ell \left(\frac{q_{xy}^n(t)}{\gamma_{xy}} \right) \right) dt + R_{\min}^\delta T^{n,\delta} \right] \leq c + 1.$$

Using the fact that ℓ is convex and $\ell(1) = 0$, by Jensen's inequality $a\ell(b) \geq \ell(ab + 1 - a)$ for $a \in [0, 1]$ and $b \geq 0$. By setting $a = \frac{\mu_x^n(t) \gamma_{xy}}{\gamma_{\max}}$, the inequality above gives

$$\mathbb{E}_{\mathbf{m}^n} \left[\int_0^{T^{n,\delta}} \left(\sum_{(x,y) \in \mathcal{Z}} \gamma_{\max} \ell \left(\frac{\mu_x^n(t)}{\gamma_{\max}} q_{xy}^n(t) + 1 - \frac{(\mu_x^n(t) \gamma_{xy})}{\gamma_{\max}} \right) \right) dt + R_{\min}^\delta T^{n,\delta} \right] \leq c + 1.$$

By applying Jensen's inequality once more

$$\mathbb{E}_{\mathbf{m}^n} \left[\int_0^{T^{n,\delta}} |\mathcal{Z}| \gamma_{\max} \ell \left(\frac{1}{|\mathcal{Z}| \gamma_{\max}} \sum_{(x,y) \in \mathcal{Z}} \mu_x^n(t) q_{xy}^n(t) + \sum_{(x,y) \in \mathcal{Z}} \left[1 - \frac{(\mu_x^n(t) \gamma_{xy})}{|\mathcal{Z}| \gamma_{\max}} \right] \right) dt + R_{\min}^\delta T^{n,\delta} \right] \leq c + 1. \quad (76)$$

Now by multiplying with $\frac{1}{|\mathcal{Z}| \gamma_{\max}}$, using (74) and the fact that $q \leq q'$ implies $\ell(q) \leq \ell(q') + 1$, we get

$$\mathbb{E}_{\mathbf{m}^n} \left[\int_0^{T^{n,\delta}} \ell \left(\frac{|\dot{\tilde{\boldsymbol{\mu}}}^{n,\delta}(t)|}{|\mathcal{Z}|\gamma_{\max}} \right) dt + \left(\frac{1}{|\mathcal{Z}|\gamma_{\max}} R_{\min}^\delta - 1 \right) T^{n,\delta} \right] \leq \frac{c+1}{|\mathcal{Z}|\gamma_{\max}}. \quad (77)$$

Finally, by using that for every $\bar{c} > 0$ there exists $c_1 > 0, c_2 < \infty$ such that $\ell(\bar{c}q) \geq c_1 \ell(q) - c_2$, we get

$$\mathbb{E}_{\mathbf{m}^n} \left[\int_0^{T^{n,\delta}} c_1 \ell \left(\frac{|\dot{\tilde{\boldsymbol{\mu}}}^{n,\delta}(t)|}{|\mathcal{Z}|\gamma_{\max}} \right) dt + \left(\frac{1}{|\mathcal{Z}|\gamma_{\max}} R_{\min}^\delta - 1 - c_2 \right) T^{n,\delta} \right] \leq \frac{c+1}{|\mathcal{Z}|\gamma_{\max}},$$

which implies

$$\mathbb{E}_{\mathbf{m}^n} \left[\int_0^{T^{n,\delta}} \ell \left(\frac{|\dot{\tilde{\boldsymbol{\mu}}}^{n,\delta}(t)|}{|\mathcal{Z}|\gamma_{\max} c_1} \right) dt + \frac{1}{|\mathcal{Z}|\gamma_{\max} c_1} R_{\min}^\delta T^{n,\delta} \right] \leq \frac{c+1}{|\mathcal{Z}|\gamma_{\max} c_1} + \frac{(c_2+1)}{c_1} \mathbb{E}_{\mathbf{m}^n} [T^{n,\delta}] \leq c', \quad (78)$$

where

$$c' = \frac{c+1}{|\mathcal{Z}|\gamma_{\max} c_1} + \frac{(c+1)(c_2+1)}{c_1}. \quad (79)$$

It will follow from the following lemma that $\tilde{\boldsymbol{\mu}}^{n,\delta}$ is a tight sequence in $D([0, \infty); \mathbb{R}^d)$. Let \mathcal{S} be the elements $(\boldsymbol{\mu}, T)$ of $C([0, \infty); \mathcal{P}(\mathcal{X})) \times [0, \infty)$ that satisfy $\boldsymbol{\mu}(t) = \boldsymbol{\mu}(T)$ for $t \geq T$.

Lemma 18 *For every positive number a , the function*

$$H(\boldsymbol{\mu}, T) = \begin{cases} \int_0^T \ell(|\dot{\boldsymbol{\mu}}(t)|) dt + aT, & \boldsymbol{\mu} \in AC([0, \infty); \mathbb{R}^d), T \in [0, \infty) \\ \infty, & \text{otherwise,} \end{cases} \quad (80)$$

is a tightness function on \mathcal{S} , where $AC([0, \infty); \mathbb{R}^d)$ is the set of all absolutely continuous functions from $[0, \infty)$ to \mathbb{R}^d .

The proof of this lemma is in Appendix 8 (Is better if we use a reference of some sort). Using the bound (78), it follows from Lemma 18 that $\{\tilde{\boldsymbol{\mu}}^{n,\delta}\}$ is tight in $D([0, \infty); \mathbb{R}^d)$. Now we have that

$$|\boldsymbol{\mu}^{n,\delta}(t) - \tilde{\boldsymbol{\mu}}^{n,\delta}(t)| \leq \sum_{(x,y) \in \mathcal{Z}} \left| \int_0^{t \wedge T^{n,\delta}} \mu_x^n(s) q_{xy}^n(s) ds - \int_0^{t \wedge T^{n,\delta}} \int_0^\infty \mathbf{1}_{[0, \mu_x^n(s) q_{xy}^n(s)]}(r) \frac{1}{n} N_{xy}^n(ds dr) \right|,$$

where the summands on the right side, denoted from now on by $Q_{xy,t}^{n,\delta}$, are all martingales with quadratic variation $\mathbb{Q}_{xy,t}^{n,\delta}$ that is bounded above by

$$\begin{aligned} & \frac{1}{n^2} \mathbb{E}_{\mathbf{m}^n} \left[\int_0^{t \wedge T^{n,\delta}} \int_0^\infty \mathbf{1}_{[0, \mu_x^n(s) q_{xy}^n(s)]}(r) N_{xy}^n(ds dr) \right] = \frac{1}{n} \mathbb{E}_{\mathbf{m}^n} \left[\int_0^{t \wedge T^{n,\delta}} \mu_x^n(s) q_{xy}^n(s) ds \right] \\ & \leq \frac{1}{n} \mathbb{E}_{\mathbf{m}^n} \left[\int_0^{t \wedge T^{n,\delta}} (\ell(\mu_x^n(s) q_{xy}^n(s)) + e) ds \right] \stackrel{(79)}{\leq} \frac{c' + e \mathbb{E}_{\mathbf{m}^n} [T^{n,\delta} \wedge t]}{n} \leq \frac{c' + e \mathbb{E}_{\mathbf{m}^n} [T^{n,\delta}]}{n} \stackrel{(75)}{\leq} \frac{\left(\frac{(c+1)e}{R_{\min}^\delta} + c' \right)}{n}, \end{aligned}$$

where in the first inequality of the last line, the estimate $ab \leq e^a + \ell(b)$, with $a = 1, b = \mu_x^n(s) q_{xy}^n(s)$ was used. By using the Burkholder-Gundy-Davis inequality, for every $T \in (0, \infty)$

$$\mathbb{E}_{\mathbf{m}^n} \left[\sup_{t \in [0, T]} |Q_{xy,t}^{n,\delta}| \right] \leq c_{BGD} \mathbb{E}_{\mathbf{m}^n} [\mathbb{Q}_{xy}^{n,\delta}]_T^{1/2} \leq c_{BGD} \sqrt{\frac{\left(\frac{(c+1)e}{R_{\min}^\delta} + c' \right)}{n}}, \quad (81)$$

from which we get that $\mathbb{E}_{\mathbf{m}^n} [\sup_{t \in [0, T]} |Q_{xy,t}^{n,\delta}|]$ converges to zero as $n \rightarrow \infty$. Recalling that we already proved $\{\tilde{\boldsymbol{\mu}}^{n,\delta}\}$ is tight in $D([0, \infty); \mathbb{R}^d)$, it follows from $\mathbb{E}_{\mathbf{m}^n} [d(\boldsymbol{\mu}^{n,\delta}, \tilde{\boldsymbol{\mu}}^{n,\delta})] \rightarrow 0$ that $\{(\boldsymbol{\mu}^{n,\delta}, T^{n,\delta})\}$ is tight as well.

To show that the variable $\boldsymbol{\theta}^{n,\delta}$ is tight, we combine (73) and (66), (67) to get

$$\mathbb{E}_{\mathbf{m}^n} \left[\sum_{(x,y) \in \mathcal{Z}} \int_0^{T^{n,\delta}} \int_0^\infty F_{xy}^n(r) \theta_{xy}^{n,\delta}(drdt) + \int_0^{T^{n,\delta}} R^n(\boldsymbol{\mu}^{n,\delta}(t)) \right] < c + 1.$$

Since, by part 1 of Lemma 4, we have $\gamma_{xy} \ell(\cdot/\gamma_{xy}) \leq F_{xy}^n(\cdot)$, and $q^{n,\delta} = \gamma$ for $t > T^{n,\delta}$, we get

$$\mathbb{E}_{\mathbf{m}^n} \left[\sum_{(x,y) \in \mathcal{Z}} \int_0^\infty \int_0^\infty \gamma_{xy} \ell\left(\frac{r}{\gamma_{xy}}\right) \theta_{xy}^{n,\delta}(drdt) \right] = \mathbb{E}_{\mathbf{m}^n} \left[\sum_{(x,y) \in \mathcal{Z}} \int_0^{T^{n,\delta}} \int_0^\infty \gamma_{xy} \ell\left(\frac{r}{\gamma_{xy}}\right) \theta_{xy}^{n,\delta}(drdt) \right] < c + 1.$$

Now by using the fact that

$$\tilde{H}(\theta) = \int_0^\infty \int_0^T \gamma_{xy} \ell\left(\frac{r}{\gamma_{xy}}\right) \theta(drdt),$$

is a tightness function on the space of measures on $[0, \infty) \times [0, T]$ with mass no greater than T , we conclude that for every $(x, y) \in \mathcal{Z}$, $\theta_{xy}^{n,\delta}$ is tight with the topology introduced in (72).

5.2 Distributional limits and the lower bound

From the previous two subsections we have that $(\boldsymbol{\mu}^{n,\delta}, \tilde{\boldsymbol{\mu}}^{n,\delta}, \boldsymbol{\theta}^{n,\delta}, T^{n,\delta})$, is tight. For proving the lower bound, we can assume without loss that the sequence has a distributional limit $(\boldsymbol{\mu}^\delta, \tilde{\boldsymbol{\mu}}^\delta, \boldsymbol{\theta}^\delta, T^\delta)$. By using the Skorohod representation theorem we can also assume the sequence of variables is on the same probability space $(\Omega, \mathcal{F}, \mathbb{P})$, and that $(\boldsymbol{\mu}^\delta, \tilde{\boldsymbol{\mu}}^\delta, \boldsymbol{\theta}^\delta, T^\delta)$ is an a.s. pointwise limit.

Consider any $\omega \in \Omega$ for which there is convergence. Since by the definition of $\theta^{n,\delta}$

$$\theta_{xy}^{n,\delta}([0, \infty) \times A) = \int_{A \cap [0, T^{n,\delta}]} \mu_x^{n,\delta}(t) dt, \quad \forall A \in \mathcal{B}(\mathbb{R}),$$

for every continuity set A of $\theta_{xy}^\delta([0, \infty) \times \cdot)$ we have

$$\begin{aligned} \left| \theta_{xy}^\delta([0, \infty) \times A) - \int_{A \cap [0, T^\delta]} \mu_x^\delta(t) dt \right| &\leq \left| \theta_{xy}^\delta([0, \infty) \times A) - \theta_{xy}^{n,\delta}([0, \infty) \times A) \right| + \left| \int_{A \cap [0, T^{n,\delta}]} \mu_x^{n,\delta}(t) dt - \int_{A \cap [0, T^\delta]} \mu_x^\delta(t) dt \right| \\ &\leq \left| \theta_{xy}^\delta([0, \infty) \times A) - \theta_{xy}^{n,\delta}([0, \infty) \times A) \right| + \left| \int_{A \cap [0, T^\delta]} \mu_x^{n,\delta}(t) dt - \int_{A \cap [0, T^\delta]} \mu_x^\delta(t) dt \right| + \int_{A \cap [\min\{T^{n,\delta}, T^\delta\}, \max\{T^\delta, T^{n,\delta}\}]} \mu_x^{n,\delta}(t) dt \\ &\leq \left| \theta_{xy}^\delta([0, \infty) \times A) - \theta_{xy}^{n,\delta}([0, \infty) \times A) \right| + d(\mu_x^{n,\delta}, \mu_x^\delta) + |T^\delta - T^{n,\delta}| \rightarrow 0. \end{aligned}$$

Therefore for every continuity set A of $\theta_{xy}^\delta([0, \infty) \times \cdot)$

$$\theta_{xy}^\delta([0, \infty) \times A) = \int_{A \cap [0, T^\delta]} \mu_x^\delta(t) dt,$$

from which we conclude that for all $(x, y) \in \mathcal{Z}$, θ_{xy}^δ has the decomposition $\theta_{xy}^\delta(dr dt) = \eta_{xy}^\delta(dr; t) dt$, with $\eta_{xy}^\delta([0, \infty); t) = \mu_x^\delta(t)$. Also, since $\int_0^\infty \int_0^\infty \ell(r) \theta_{xy}^{n,\delta}(dr dt)$ is uniformly bounded and ℓ is superlinear, we have convergence of the first moments of the first marginal, i.e.,

$$\int_{\mathbb{R}} f(t) r \theta_{xy}^{n,\delta}(dr dt) \rightarrow \int_{\mathbb{R}} f(t) r \theta_{xy}^\delta(dt), \quad \forall f \in C_b(\mathbb{R}).$$

Hence for q^δ defined by $\mu_x^\delta(t) q_{xy}^\delta(t) = \int_0^\infty r \eta_{xy}^\delta(dr; t)$, we get that for all $(x, y) \in \mathcal{Z}$

$$\int_0^\infty f(t) \mu_x^{n,\delta}(t) q_{xy}^{n,\delta}(t) dt \rightarrow \int_0^\infty f(t) \mu_x^\delta(t) q_{xy}^\delta(t) dt, \quad \forall f \in C_b(\mathbb{R}). \quad (82)$$

Using the fact that $d(\boldsymbol{\mu}^{n,\delta}, \tilde{\boldsymbol{\mu}}^{n,\delta}) \rightarrow 0$ and (74), we get

$$\left| \boldsymbol{\mu}^{n,\delta}(t) - \mathbf{m}^n - \sum_{(x,y) \in \mathcal{Z}} \mathbf{v}_{xy} \int_0^{T^{n,\delta} \wedge t} \mu_x^{n,\delta}(s) q_{xy}^{n,\delta}(s) ds \right| = \left| \boldsymbol{\mu}^{n,\delta}(t) - \tilde{\boldsymbol{\mu}}^{n,\delta}(t) \right| \rightarrow 0, \quad (83)$$

for a.e. t . Applying (82) for suitable choices of f and using (83),

$$\boldsymbol{\mu}^\delta(t) = \mathbf{m} + \sum_{(x,y) \in \mathcal{Z}} \mathbf{v}_{xy} \int_0^{T^\delta \wedge t} \mu_x^\delta(s) q_{xy}^\delta(s) ds$$

for a.e. t , and since the left side is cadlag and the right side is continuous in the last display, equality holds for $t \geq 0$. We conclude that \mathbf{q}^δ is the control that generates $\boldsymbol{\mu}^\delta$, and we also already noticed that $\mu_x^\delta(t) q_{xy}^\delta(t) = \int_0^\infty r \eta_{xy}^\delta(dr; t)$. Finally, since $\boldsymbol{\mu}^{n,\delta}(T^{n,\delta}) \in K_\delta$ and $d(\boldsymbol{\mu}^{n,\delta}, \boldsymbol{\mu}^\delta) \rightarrow 0$, by continuity of $\boldsymbol{\mu}^\delta$ we get $\boldsymbol{\mu}^\delta(T^\delta) \in K_\delta$. As discussed below (73), this concludes the lower bound proof.

6 Upper bound

Before we proceed with the proof of the upper bound

$$\limsup_{n \rightarrow \infty} V_K^n(\mathbf{m}^n) \leq V_K(\mathbf{m}),$$

we establish some preliminary lemmas. In the following lemmas, we make use of \mathcal{T}_m , \mathcal{U}_m and \hat{F}_{xy}^∞ , defined in (52), (53), and (57) respectively. For the properties of \hat{F}_{xy}^∞ , see Lemma 4.

Lemma 19 *Let $\mathbf{m} \in \mathcal{P}_*(\mathcal{X})$, and $\mathbf{q} \in \mathcal{U}_m$ be such that $(\boldsymbol{\mu}, \mathbf{q}) \in \mathcal{T}_m$. Given $T < \infty$ and $\epsilon > 0$, we can find $a_1, a_2, a_3 \in (0, \infty)$ and $\tilde{\mathbf{q}} \in \mathcal{U}_m$, with $(\tilde{\boldsymbol{\mu}}, \tilde{\mathbf{q}}) \in \mathcal{T}_m$, such that*

$$a_1 \leq \inf_{(x,y) \in \mathcal{Z}, t \in [0, T]} \tilde{q}_{xy}(t) \leq \sup_{(x,y) \in \mathcal{Z}, t \in [0, T]} \tilde{q}_{xy}(t) \leq a_2, \quad (84)$$

$$\inf_{x \in \mathcal{X}, t \in [0, T]} \tilde{\mu}_x(t) > a_3, \quad \sup_{t \in [0, T]} \|\boldsymbol{\mu}(t) - \tilde{\boldsymbol{\mu}}(t)\| < \epsilon, \quad \text{and} \quad (85)$$

$$\sum_{(x,y) \in \mathcal{Z}} \int_0^T \tilde{\mu}_x(t) \hat{F}_{xy}^\infty(\tilde{q}_{xy}(t)) dt \leq \sum_{(x,y) \in \mathcal{Z}} \int_0^T \mu_x(t) \hat{F}_{xy}^\infty(q_{xy}(t)) dt. \quad (86)$$

Proof. Recall that $\mathbf{m} \in \mathcal{P}_*(\mathcal{X})$ implies $\mathbf{m}_x > 0$ for all $x \in \mathcal{X}$. Let $\boldsymbol{\nu}(\mathbf{m}, t)$ be the solution to the equation $\dot{\boldsymbol{\nu}}(t) = \boldsymbol{\gamma}\boldsymbol{\nu}(t)$, with initial data \mathbf{m} . By Remark 16, we know that there exists $1 \geq a > 0$ such that $\boldsymbol{\nu}(\mathbf{m}, t) \in \mathcal{P}_a(\mathcal{X})$, for every $t \in [0, T]$. We can assume without loss that the right hand side of (86) is greater than zero, since if not true then the controlled rates are $\boldsymbol{\gamma}$ and the conclusion of the lemma is automatic. For $\frac{\epsilon}{2} \geq \delta > 0$, let

$$\boldsymbol{\mu}^\delta(\cdot) \doteq \delta \boldsymbol{\nu}(\mathbf{m}, \cdot) + (1 - \delta) \boldsymbol{\mu}(\cdot), \quad (87)$$

and note that $\mu_x^\delta(t) > 0$ for every $t \in [0, T]$ and $x \in \mathcal{X}$. Therefore, for δ as above and $(x, y) \in \mathcal{Z}$, we can define

$$q_{xy}^\delta(\cdot) = \gamma_{xy} \frac{\delta \nu_x(\mathbf{m}, \cdot)}{\mu_x^\delta(\cdot)} + q_{xy}(\cdot) \frac{(1 - \delta) \mu_x(\cdot)}{\mu_x^\delta(\cdot)}. \quad (88)$$

Then it is straightforward to check that $(\boldsymbol{\mu}^\delta, \mathbf{q}^\delta) \in \mathcal{T}_m$. Moreover, since $\frac{\delta \nu_x(\mathbf{m}, t)}{\mu_x^\delta(t)} + \frac{(1 - \delta) \mu_x(t)}{\mu_x^\delta(t)} = 1$ for all $t \in [0, T]$, by the convexity of \hat{F}^∞ we obtain

$$\begin{aligned} \sum_{(x,y) \in \mathcal{Z}} \int_0^T \mu_x^\delta(t) \hat{F}_{xy}^\infty(q_{xy}^\delta(t)) dt &= \sum_{(x,y) \in \mathcal{Z}} \int_0^T \mu_x^\delta(t) \hat{F}_{xy}^\infty \left(\gamma_{xy} \frac{\delta \nu_x(\mathbf{m}, t)}{\mu_x^\delta(t)} + q_{xy} \frac{(1 - \delta) \mu_x(t)}{\mu_x^\delta(t)} \right) dt \\ &\leq \sum_{(x,y) \in \mathcal{Z}} \int_0^T \mu_x^\delta(t) \frac{\delta \nu_x(\mathbf{m}, t)}{\mu_x^\delta(t)} \hat{F}_{xy}^\infty(\gamma_{xy}) dt + \sum_{(x,y) \in \mathcal{Z}} \int_0^T \mu_x^\delta(t) \frac{(1 - \delta) \mu_x(t)}{\mu_x^\delta(t)} \hat{F}_{xy}^\infty(q_{xy}(t)) dt \\ &\leq (1 - \delta) \sum_{(x,y) \in \mathcal{Z}} \int_0^T \mu_x(t) \hat{F}_{xy}^\infty(q_{xy}(t)) dt, \end{aligned} \quad (89)$$

where in the second inequality, we used the fact that $\hat{F}^\infty(\gamma_{xy}) = 0$ [see Lemma 4]. Therefore, we get a triplet $(\boldsymbol{\mu}^\delta, \mathbf{q}^\delta) \in \mathcal{T}_m$ with cost strictly less than the initial one, and with $\boldsymbol{\mu}^\delta$ that satisfies

$$\mu_x^\delta(t) \geq \delta a \quad \text{and} \quad \frac{(1-\delta)\mu_x(t)}{\mu_x^\delta(t)} \leq \frac{(1-\delta)}{\delta a + (1-\delta)} \equiv c < 1, \quad (90)$$

for all $t \in [0, T]$. However, since this triplet does not necessarily satisfy condition (84), we modify it even further. Specifically, we pick $M \in (2\gamma_{\max}, \infty)$ big enough such that

$$\sum_{(x,y) \in \mathcal{Z}} \int_0^T \mu_x^\delta(t) |\min\{q_{xy}^\delta(t), M\} - q_{xy}^\delta(t)| dt \leq \frac{a\delta(1-\sqrt{c})}{\sqrt{2}}, \quad (91)$$

and define

$$\boldsymbol{\mu}^{\delta, M}(t) = \int_0^t \sum_{(x,y) \in \mathcal{Z}} \mu_x^\delta(t) \min\{q_{xy}^\delta(t), M\} \mathbf{v}_{xy} dt. \quad (92)$$

Then

$$\begin{aligned} |\mu_x^{\delta, M}(t) - \mu_x^\delta(t)| &\leq \|\boldsymbol{\mu}^{\delta, M}(t) - \boldsymbol{\mu}^\delta(t)\| \stackrel{(92)}{=} \left\| \sum_{(x,y) \in \mathcal{Z}} \int_0^t (\mu_x^\delta(t) (q_{xy}^\delta(t) - \min\{q_{xy}^\delta(t), M\})) \mathbf{v}_{xy} dt \right\| \\ &\leq \sum_{(x,y) \in \mathcal{Z}} \int_0^T |\mu_x^\delta(t) (q_{xy}^\delta(t) - \min\{q_{xy}^\delta(t), M\})| \|\mathbf{v}_{xy}\| dt \\ &\leq \sqrt{2} \sum_{(x,y) \in \mathcal{Z}} \int_0^T (\mu_x^{\delta, M}(t) (q_{xy}^\delta(t) - \min\{q_{xy}^\delta(t), M\})) dt \stackrel{(91)}{\leq} a\delta(1-\sqrt{c}), \end{aligned} \quad (93)$$

and therefore for $t \in [0, T]$,

$$\mu_x^{\delta, M}(t) \geq \mu_x^\delta(t) - |\mu_x^{\delta, M}(t) - \mu_x^\delta(t)| \stackrel{(90)}{\geq} a\delta - |\mu_x^{\delta, M}(t) - \mu_x^\delta(t)| \stackrel{(93)}{\geq} a\delta\sqrt{c}. \quad (94)$$

We also get

$$\left| 1 - \frac{\mu_x^{\delta, M}(t)}{\mu_x^\delta(t)} \right| \stackrel{(93)}{\leq} \frac{a\delta(1-\sqrt{c})}{\min_x \mu_x^\delta} \stackrel{(90)}{\leq} (1-\sqrt{c}) \quad (95)$$

or

$$\frac{\mu_x^\delta(t)}{\mu_x^{\delta, M}(t)} \geq \frac{1}{2-\sqrt{c}} \quad \text{and} \quad \frac{\mu_x^\delta(t)}{\mu_x^{\delta, M}(t)} \leq \frac{1}{\sqrt{c}} = \frac{\sqrt{c}}{c}. \quad (96)$$

We deduce that $\boldsymbol{\mu}^{\delta, M}(t) \in \mathcal{P}_*(\mathcal{X})$, for all $t \in [0, T]$, and therefore can define

$$q_{xy}^{\delta, M}(t) = \frac{\min\{q_{xy}^\delta(t), M\} \mu_x^\delta(t)}{\mu_x^{\delta, M}(t)}, \quad (97)$$

which will give $(\boldsymbol{\mu}^{\delta, M}, \mathbf{q}^{\delta, M}) \in \mathcal{T}_m$. We can see that (84) is satisfied, since by (88) and the LHS of (96) for the bound from below and the RHS of (96) for the bound from above we have

$$\frac{\gamma_{xy} \delta \nu_x(\mathbf{m}, \cdot)}{2} \leq q_{xy}^{\delta, M}(t) \leq M \frac{\sqrt{c}}{c} \quad (98)$$

It is worth mentioning at this point that trying to get an estimate for the cost of $(\boldsymbol{\mu}^{\delta, M}, \mathbf{q}^{\delta, M})$, with respect to the cost of $(\boldsymbol{\mu}^\delta, \mathbf{q}^\delta)$, would require some extra properties of \hat{F}^∞ . However, we can obtain an estimate of the cost $(\boldsymbol{\mu}^{\delta, M}, \mathbf{q}^{\delta, M})$ with respect to the cost of the initial triplet $(\boldsymbol{\mu}, \mathbf{q})$, by utilizing only the convexity of \hat{F}_{xy}^∞ , and choosing the right parameters. Using the fact that \hat{F}_{xy}^∞ is increasing on $[\gamma_{xy}, \infty)$ in the first inequality, and that $M\mu_x^\delta(t)/\mu_x^{\delta, M}(t) \geq \gamma_{xy}$ by (96) and $M \geq 2\gamma_{xy}$,

$$\begin{aligned} \hat{F}_{xy}^\infty(q_{xy}^{\delta, M}(t)) &\stackrel{(97)}{=} \hat{F}_{xy}^\infty\left(\frac{\min\{q_{xy}^\delta(t), M\} \mu_x^\delta(t)}{\mu_x^{\delta, M}(t)}\right) \leq \hat{F}_{xy}^\infty\left(\frac{q_{xy}^\delta(t) \mu_x^\delta(t)}{\mu_x^{\delta, M}(t)}\right) \\ &\stackrel{(88)}{=} \hat{F}_{xy}^\infty\left(\frac{\mu_x^\delta(t)}{\mu_x^{\delta, M}(t)} \left(\gamma_{xy} \frac{\delta \nu_x(\mathbf{m}, t)}{\mu_x^\delta(t)} + q_{xy}(t) \frac{(1-\delta)\mu_x(t)}{\mu_x^\delta(t)}\right)\right) \\ &= \hat{F}_{xy}^\infty\left(\gamma_{xy} \frac{\delta \nu_x(\mathbf{m}, t)}{\mu_x^{\delta, M}(t)} + q_{xy}(t) \frac{(1-\delta)\mu_x(t)}{\mu_x^{\delta, M}(t)}\right). \end{aligned} \quad (99)$$

However, from (90) and (96), we have

$$\frac{(1-\delta)\mu_x(t)}{\mu_x^{\delta,M}(t)} = \frac{(1-\delta)\mu_x(t)}{\mu_x^\delta(t)} \frac{\mu_x^\delta(t)}{\mu_x^{\delta,M}(t)} \leq c \frac{\sqrt{c}}{c} = \sqrt{c} < 1.$$

Therefore using the convexity of \hat{F}^∞ we have

$$\begin{aligned} \hat{F}_{xy}^\infty \left(\gamma_{xy} \frac{\delta\nu_x(\mathbf{m}, t)}{\mu_x^{\delta,M}(t)} + q_{xy}(t) \frac{(1-\delta)\mu_x(t)}{\mu_x^{\delta,M}(t)} \right) &= \hat{F}_{xy}^\infty \left(\frac{\left(1 - \frac{(1-\delta)\mu_x(t)}{\mu_x^{\delta,M}(t)}\right)}{\left(1 - \frac{(1-\delta)\mu_x(t)}{\mu_x^{\delta,M}(t)}\right)} \gamma_{xy} \frac{\delta\nu_x(\mathbf{m}, t)}{\mu_x^{\delta,M}(t)} + q_{xy}(t) \frac{(1-\delta)\mu_x(t)}{\mu_x^{\delta,M}(t)} \right) \\ &\leq \left(1 - \frac{(1-\delta)\mu_x(t)}{\mu_x^{\delta,M}(t)}\right) \hat{F}_{xy}^\infty \left(\gamma_{xy} \frac{\delta\nu_x(\mathbf{m}, t)}{\mu_x^{\delta,M}(t) - (1-\delta)\mu_x(t)} \right) + \frac{(1-\delta)\mu_x(t)}{\mu_x^{\delta,M}(t)} \hat{F}_{xy}^\infty(q_{xy}(t)). \end{aligned} \quad (100)$$

Combining (99) and (100) and then using (87), we obtain

$$\begin{aligned} \mu_x^{\delta,M}(t) \hat{F}_{xy}^\infty(q_{xy}^{\delta,M}(t)) &\leq (\mu_x^{\delta,M}(t) - (1-\delta)\mu_x(t)) \hat{F}_{xy}^\infty \left(\gamma_{xy} \frac{\delta\nu_x(\mathbf{m}, t)}{\mu_x^{\delta,M}(t) - (1-\delta)\mu_x(t)} \right) + (1-\delta)\mu_x(t) \hat{F}_{xy}^\infty(q_{xy}(t)) \\ &= (\mu_x^{\delta,M}(t) - \mu_x^\delta(t) + \delta\nu_x(\mathbf{m}, t)) \hat{F}_{xy}^\infty \left(\gamma_{xy} \frac{\delta\nu_x(\mathbf{m}, t)}{\mu_x^{\delta,M}(t) - \mu_x^\delta(t) + \delta\nu_x(\mathbf{m}, t)} \right) + (1-\delta)\mu_x(t) \hat{F}_{xy}^\infty(q_{xy}(t)). \end{aligned} \quad (101)$$

We can make $|\mu_x^{\delta,M}(t) - \mu_x^\delta(t)|$ uniformly as close to zero as desired and therefore we can make the quantity $\frac{\gamma_{xy} \frac{\delta\nu_x(\mathbf{m}, t)}{\mu_x^{\delta,M}(t) - \mu_x^\delta(t) + \delta\nu_x(\mathbf{m}, t)}}{\mu_x^{\delta,M}(t) - \mu_x^\delta(t) + \delta\nu_x(\mathbf{m}, t)}$ as close to γ_{xy} as desired by picking M sufficiently large. Since $\hat{F}_{xy}^\infty(\gamma_{xy}) = 0$ and $\hat{F}_{xy}^\infty(\cdot)$ is continuous on $(0, \infty)$ by Lemma 15, we can pick $M < \infty$ such that for every $t \in [0, T]$,

$$\hat{F}_{xy}^\infty \left(\gamma_{xy} \frac{\delta\nu_x(\mathbf{m}, t)}{\mu_x^{\delta,M}(t) - \mu_x^\delta(t) + \delta\nu_x(\mathbf{m}, t)} \right) \leq \frac{1}{2T} \int_0^T \mu_x(s) \hat{F}_{xy}^\infty(q_{xy}(s)) ds. \quad (102)$$

Then from (101) and (102) and the fact that $\nu_x(\mathbf{m}, t) \leq 1$ and (93), for $t \in [0, T]$

$$\begin{aligned} \sum_{(x,y) \in \mathcal{Z}} \int_0^T \mu_x^{\delta,M}(t) \hat{F}_{xy}^\infty(q_{xy}^{\delta,M}(t)) dt &\leq \sum_{(x,y) \in \mathcal{Z}} \int_0^T (2\delta) \left(\frac{1}{2T} \int_0^T \mu_x(s) \hat{F}_{xy}^\infty(q_{xy}(s)) ds \right) dt + \\ &\int_0^T (1-\delta)\mu_x(t) \hat{F}_{xy}^\infty(q_{xy}(t)) dt = \sum_{(x,y) \in \mathcal{Z}} \int_0^T \mu_x(t) \hat{F}_{xy}^\infty(q_{xy}(t)) dt. \end{aligned} \quad (103)$$

■

Next, we are going to prove the following result.

Lemma 20 (Law of large numbers) *Let $T \in (0, \infty)$ be given. There exists a constant $c < \infty$ such that if $(\mu^n, \gamma) \in \mathcal{T}_m^n$ (see (21)), and $(\nu, \gamma) \in \mathcal{T}_m$, then*

$$\mathbb{P} \left(\sup_{t \in [0, T]} \|\mu^n(t) - \nu(\mathbf{m}, t)\| \geq \epsilon \right) \leq \frac{c}{\epsilon \sqrt{n}}. \quad (104)$$

Proof. We have

$$\begin{aligned} \|\mu^n(t) - \nu(\mathbf{m}, t)\| &\leq \sum_{(x,y)} \left| \int_0^t \int_0^\infty 1_{[0, \mu_x^n(s)\gamma_{xy}]}(r) \frac{1}{n} N_{xy}^n(dsdr) - \int_0^t \int_0^\infty 1_{[0, \nu_x(\mathbf{m}, s)\gamma_{xy}]}(r) dsdr \right| \\ &\leq \sum_{(x,y)} \left| \int_0^t \int_0^\infty 1_{[0, \mu_x^n(s)\gamma_{xy}]}(r) \frac{1}{n} N_{xy}^n(dsdr) - \int_0^t \int_0^\infty 1_{[0, \mu_x^n(s)\gamma_{xy}]}(r) dsdr \right| \\ &\quad + \sum_{(x,y)} \left| \int_0^t \int_0^\infty 1_{[0, \mu_x^n(s)\gamma_{xy}]}(r) dsdr - \int_0^t \int_0^\infty 1_{[0, \nu_x(\mathbf{m}, s)\gamma_{xy}]}(r) dsdr \right|. \end{aligned}$$

For a constant K that depends on d and the maximum of γ_{xy} ,

$$\sum_{(x,y)} \left| \int_0^t \int_0^\infty 1_{[0, \mu_x^n(s)\gamma_{xy}]}(r) ds dr - \int_0^t \int_0^\infty 1_{[0, \nu_x(m,s)\gamma_{xy}]}(r) ds \right| \leq K \sup_{0 \leq s \leq t} \|\mu^n(s) - \nu(m, s)\|.$$

Hence by Gronwall's inequality, for $r \in [0, T]$

$$\|\mu^n(r) - \nu(m, r)\| \leq e^{KT} \sup_{0 \leq t \leq r} \sum_{(x,y)} \left| \int_0^t \int_0^\infty 1_{[0, \mu_x^n(s)\gamma_{xy}]}(r) \frac{1}{n} N_{xy}^n(ds dr) - \int_0^t \int_0^\infty 1_{[0, \mu_x^n(s)\gamma_{xy}]}(r) ds dr \right|.$$

Using the Burkholder-Gundy-Davis inequality as was done to obtain (81),

$$P \left(\sup_{t \in [0, T]} \left| \int_0^t \int_0^\infty 1_{[0, \mu_x^n(s)\gamma_{xy}]}(r) \frac{1}{n} N_{xy}^n(ds dr) - \int_0^t \int_0^\infty 1_{[0, \mu_x^n(s)\gamma_{xy}]}(r) ds dr \right| \geq \epsilon \right) \leq \frac{\bar{c}}{\epsilon \sqrt{n}},$$

and hence

$$P \left(\sup_{t \in [0, T]} \|\mu^n(t) - \nu(m, t)\| \geq \epsilon \right) \leq d^2 \frac{e^{KT} \bar{c}}{\epsilon \sqrt{n}},$$

which is (104).
■

We now obtain the following result.

Lemma 21 *The sequence $V^n(m)$ is bounded, uniformly in n and $m \in \mathcal{P}(X)$.*

Proof. Let $\tau = \text{diameter}(\mathcal{P}(X))$. By Remark 16, there exists $\bar{a} > 0$ such that $\nu(m, \tau) \in \mathcal{P}_{2\bar{a}}(X)$ regardless of the initial data m . We can further assume that $\mathcal{P}_{\bar{a}}(X) \cap K^\circ \neq \emptyset$, and in particular that there exists an element \tilde{m} such that $B(\tilde{m}, \bar{a}/2) \subset \mathcal{P}_{\bar{a}}(X) \cap K^\circ$. Without loss of generality, we replace a by $a \wedge \bar{a}$.

Since $\tilde{m} \in \mathcal{P}_a$, the first part of Theorem 17 implies that for every point m in $\mathcal{P}_a(X)$ we can find a control q_m with the following properties: there is a unique μ such that $(\mu, q_m) \in \mathcal{T}_m$; μ is a constant speed parametrization of the straight line that connects m to \tilde{m} in time $T_{\{\tilde{m}\}} = \|m - \tilde{m}\|$; and the control q_m satisfies

$$\gamma_{xy} \leq q_{m,xy}(t) \leq \frac{c_1}{a}, \quad (105)$$

for $t \in [0, T_{\{\tilde{m}\}}]$, $(x, y) \in \mathcal{Z}$, where $c_1 > 0$ is a constant that does not depend on a . For every m , we let

$$q_{xy}(m, t) = \begin{cases} q_{m,xy}(t) & t \leq \|m - \tilde{m}\|, \\ \gamma_{xy} & t > \|m - \tilde{m}\|, \end{cases} \quad (106)$$

denote the control that takes m to \tilde{m} in time $\|m - \tilde{m}\|$, in the sense that it was described above, and after that time is equal to the original rates.

For $i \in \mathbb{N}$ we define a control for the interval $i\tau \leq t < (i+1)\tau$ as follows. Let $f(t-)$ denote the limit of $f(s)$ from the left at time t , and recall that $\mu(m, \cdot)$ is the straight line that connects m to \tilde{m} in time $T_{\{\tilde{m}\}}$, where \tilde{m} is fixed and we explicitly indicate the dependence on m . Then set

$$q_{xy}^n(t) = \begin{cases} q_{xy}(m, t - i\tau) \frac{\mu_x^n(t-)}{\mu_x(m, t - i\tau)}, & \text{if } \left(\sup_{s \in [i\tau, t]} \|\mu(m, s) - \mu^n(s)\| \leq \frac{a}{2} \right) \text{ and } (\mu^n(i\tau) = m \in \mathcal{P}_a(X)) \\ \gamma_{xy}, & \text{otherwise.} \end{cases} \quad (107)$$

The idea with these controls is that, within each time interval with length τ , the control considers the starting point m , and then if $m \in \mathcal{P}_a(X)$, it attempts to force the process to follow the straight line to \tilde{m} . If $m \notin \mathcal{P}_a(X)$ or the process goes close to the boundary of the simplex $\mathcal{P}(X) \setminus \mathcal{P}_*(X)$, then we just use original rates to push the process inside $\mathcal{P}_a(X)$. Since all controls used are bounded from above and below, the total cost is a multiple of $\mathbb{E}[T^n]$. Thus we need only show this expected exit time is uniformly bounded.

By using (81), we can find constant $c < \infty$ such that

$$\mathbb{P} \left(\sup_{t \in [i\tau, (i+1)\tau]} \|\mu^n(t) - \mu(m, t)\| \geq \frac{a}{2} \mid \mu^n(i\tau) = m \in \mathcal{P}_a(X) \right) \leq c \frac{2}{\sqrt{na}}, \quad (108)$$

from which we get

$$\begin{aligned} P(T^n > (i+1)\tau | \boldsymbol{\mu}^n(i\tau) \in \mathcal{P}_a(\mathcal{X})) &\leq \inf_{\mathbf{m} \in \mathcal{P}_a(\mathcal{X})} \mathbb{P} \left(\sup_{t \in [i\tau, (i+1)\tau]} \|\boldsymbol{\mu}^n(t) - \boldsymbol{\mu}(\mathbf{m}, t)\| \geq \frac{a}{2} \middle| \boldsymbol{\mu}^n(i\tau) = \mathbf{m} \right) \\ &\leq c \frac{2}{\sqrt{na}}. \end{aligned} \quad (109)$$

By Lemma 20, we have that for some $c' < \infty$

$$\mathbb{P} \left(\sup_{t \in [i\tau, (i+1)\tau]} \|\boldsymbol{\mu}^n(t) - \boldsymbol{\nu}(\mathbf{m}, t)\| \geq \frac{a}{2} \middle| \boldsymbol{\mu}^n(i\tau) = \mathbf{m} \notin \mathcal{P}_a(\mathcal{X}) \right) \leq c' \frac{2}{a\sqrt{n}}, \quad (110)$$

which implies that

$$\begin{aligned} &\mathbb{P} \left(\boldsymbol{\mu}^n((i+1)\tau) \notin \mathcal{P}_a(\mathcal{X}) \middle| \boldsymbol{\mu}^n(i\tau) \notin \mathcal{P}_a(\mathcal{X}) \right) \\ &\leq \inf_{\mathbf{m} \notin \mathcal{P}_a(\mathcal{X})} \mathbb{P} \left(\sup_{t \in [i\tau, (i+1)\tau]} \|\boldsymbol{\mu}^n(t) - \boldsymbol{\nu}(\mathbf{m}, t)\| \geq \frac{a}{2} \middle| \boldsymbol{\mu}^n(i\tau) = \mathbf{m} \right) \leq c' \frac{2}{a\sqrt{n}}. \end{aligned} \quad (111)$$

Thus the probability to escape in the next 2τ units of time has a positive lower bound that is independent of n and the starting position. This implies the uniform upper bound on the mean escape time. ■

Now we proceed with the proof of the upper bound.

Proof of upper bound. We will initially assume that \mathbf{m} is in $\mathcal{P}_a(\mathcal{X})$, for some $a > 0$. Recall that $V_K(\mathbf{m}) < \infty$. Let $\epsilon > 0$. By the definition of $V_K(\mathbf{m})$, we can find a triplet $(\boldsymbol{\mu}, \mathbf{q}) \in \mathcal{T}_{\mathbf{m}}$ and a $T \in [0, \infty]$, such that

$$\int_0^T \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(t) \hat{F}_{xy}^\infty(q_{xy}(t)) + R^\infty(\boldsymbol{\mu}(t)) \right) dt \leq V_K(\mathbf{m}) + \epsilon. \quad (112)$$

Since we assumed that R^∞ is bounded from below by a positive constant for every compact subset of K^c , we can furthermore find a δ such that for finite time $T^\delta \in [0, \infty)$ we have

$$\int_0^{T^\delta} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x(t) \hat{F}_{xy}^\infty(q_{xy}(t)) + R^\infty(\boldsymbol{\mu}(t)) \right) dt \leq V_K(\mathbf{m}) + \epsilon, \quad (113)$$

and $d(\boldsymbol{\mu}(T^\delta), K) \leq \delta$. By the second part of Theorem 17, we can extend the path so it can reach a point $\tilde{\mathbf{m}}$ of K , with extra cost less than ϵ . Since $K = \overline{(K^\circ)}$, by a second application of Theorem 17, we can assume that $\tilde{\mathbf{m}}$ is an internal point of K , by again adding an extra cost less than ϵ .

Let $r > 0$ be such that $B(\tilde{\mathbf{m}}, r) \subset K^\circ$. From Lemma 19, without any loss of generality, we can assume that there exist $a_1, a_2, a_3 \in (0, \infty)$ such that

$$a_1 \leq \inf_{(x,y) \in \mathcal{Z}, t \in [0, T]} q_{xy}(t) \leq \sup_{(x,y) \in \mathcal{Z}, t \in [0, T]} q_{xy}(t) \leq a_2, \quad (114)$$

and

$$\inf_{x \in \mathcal{X}, t \in [0, T]} \mu_x(t) > a_3, \quad \|\boldsymbol{\mu}(T) - \tilde{\mathbf{m}}\| < \frac{r}{2}, \quad (115)$$

where the T used above is the one obtained by starting with T^δ and adding bits as just described, and not the T introduced at the beginning of the proof. Finally, by applying the first part of Theorem 17, we can assume the existence of a $r_1 > 0$ such that for every point $\tilde{\mathbf{m}}$ in the neighborhood $B(\tilde{\mathbf{m}}, r_1)$, we can find a path like the one described above, by connecting $\tilde{\mathbf{m}}$ with a straight line to \mathbf{m} . Of course this could generate different, though universal, a_1, a_2, a_3 from the initial ones.

Now let \mathbf{m}^n be a sequence that converges to \mathbf{m} . For big enough n , we can assume that $\mathbf{m}^n \in B(\mathbf{m}, r_1)$. By the continuity of \hat{F}^∞ on compact subsets of $(0, \infty)$, we can find $r_2 > 0$ such that if $\mathbf{m}_1, \mathbf{m}_2 \in \mathcal{P}_{\frac{a_3}{2}}(\mathcal{X})$ and $\|\mathbf{m}_1 - \mathbf{m}_2\| \leq r_2$, then for every \mathbf{q} that satisfies (114), we have

$$\sum_{(x,y) \in \mathcal{Z}} \left| m_{1,x} \hat{F}_{xy}^\infty(q_{xy}) - m_{2,x} \hat{F}_{xy}^\infty \left(q_{xy} \frac{m_{1,x}}{m_{2,x}} \right) \right| \leq \frac{\epsilon}{T}. \quad (116)$$

Now for every $n \in \mathbb{N}$, we define the following control for the time interval $[0, T]$,

$$q_{xy}^n(t) = \begin{cases} q_{xy}(t) \frac{\mu_x^n(t-)}{\mu_x(t)}, & \text{if } \sup_{s \in [0,t]} \|\boldsymbol{\mu}(s) - \boldsymbol{\mu}^n(s)\| \leq r_2 \\ \gamma_{xy}, & \text{otherwise.} \end{cases} \quad (117)$$

Note that either $\boldsymbol{\mu}^n$ enters K by time T , or the control has switch to γ_{xy} before T . For every n , we define an auxiliary stopping time $\bar{T}^n = \inf\{t \in [0, T] : \|\boldsymbol{\mu}^n(t) - \boldsymbol{\mu}(t)\| > r_2\}$, and also we define $R_{max} = \sup_{n \in \mathbb{N}, \mathbf{m} \in \mathcal{P}(\mathcal{X})} R^n(\mathbf{m})$. For sufficiently large n , by uniform convergence of F^n to \hat{F}^∞ on compact subsets of $(0, \infty)$, and the uniform convergence of R^n to R^∞ , we can get an estimate of the cost accumulated up to time T , for the triple $(\boldsymbol{\mu}^n, \mathbf{q}^n) \in \mathcal{T}_{\mathbf{m}^n}^n$. Specifically,

$$\begin{aligned} & \mathbb{E} \left[\int_0^T \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x^n(t) F_{xy}^n(q_{xy}^n(t)) + R^n(\boldsymbol{\mu}^n(t)) \right) dt \right] \\ & \leq \mathbb{E} \left[\int_0^T \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x^n(t) F_{xy}^n(q_{xy}^n(t)) + R^n(\boldsymbol{\mu}^n(t)) \right) dt \cdot \mathbf{1}_{\{\sup_{t \in [0,T]} \|\boldsymbol{\mu}(t) - \boldsymbol{\mu}^n(t)\| \leq r_2\}} \right] \\ & + \mathbb{P} \left(\sup_{t \in [0,T]} \|\boldsymbol{\mu}^n(t) - \boldsymbol{\mu}(t)\| > r_2 \right) \times \\ & \left(\mathbb{E} \left[\int_0^{\bar{T}^n} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x^n(t) F_{xy}^n(q_{xy}^n(t)) + R^n(\boldsymbol{\mu}^n(t)) \right) dt \middle| \sup_{t \in [0,T]} \|\boldsymbol{\mu}^n(t) - \boldsymbol{\mu}(t)\| > r_2 \right] + TR_{max} \right) \end{aligned} \quad (118)$$

Now by (117) the last quantity is equal to

$$\begin{aligned} & \mathbb{E} \left[\int_0^T \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x^n(t) F_{xy}^n \left(q_{xy}(t) \frac{\mu_x^n(t-)}{\mu_x(t)} \right) + R^n(\boldsymbol{\mu}^n(t)) \right) dt \cdot \mathbf{1}_{\{\sup_{t \in [0,T]} \|\boldsymbol{\mu}(t) - \boldsymbol{\mu}^n(t)\| \leq r_2\}} \right] \\ & + \mathbb{P} \left(\sup_{t \in [0,T]} \|\boldsymbol{\mu}^n(t) - \boldsymbol{\mu}(t)\| > r_2 \right) \times \\ & \left(\mathbb{E} \left[\int_0^{\bar{T}^n} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x^n(t) F_{xy}^n \left(q_{xy}(t) \frac{\mu_x^n(t-)}{\mu_x(t)} \right) + R^n(\boldsymbol{\mu}^n(t)) \right) dt \middle| \sup_{t \in [0,T]} \|\boldsymbol{\mu}(t) - \boldsymbol{\mu}^n(t)\| > r_2 \right] + TR_{max} \right) \\ & \leq \mathbb{E} \left[\int_0^T \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x^n(t) \hat{F}_{xy}^\infty \left(q_{xy}(t) \frac{\mu_x^n(t-)}{\mu_x(t)} \right) + R^\infty(\boldsymbol{\mu}^n(t)) \right) dt \cdot \mathbf{1}_{\{\sup_{t \in [0,T]} \|\boldsymbol{\mu}(t) - \boldsymbol{\mu}^n(t)\| \leq r_2\}} \right] + \epsilon \\ & + \mathbb{P} \left(\sup_{t \in [0,T]} \|\boldsymbol{\mu}^n(t) - \boldsymbol{\mu}(t)\| > r_2 \right) \times \\ & \left(\mathbb{E} \left[\int_0^{\bar{T}^n} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x^n(t) \hat{F}_{xy}^\infty \left(q_{xy}(t) \frac{\mu_x^n(t-)}{\mu_x(t)} \right) + R^\infty(\boldsymbol{\mu}^n(t)) \right) dt \middle| \sup_{t \in [0,T]} \|\boldsymbol{\mu}(t) - \boldsymbol{\mu}^n(t)\| > r_2 \right] + \epsilon + TR_{max} \right). \end{aligned} \quad (119)$$

Then using (116) with $m_{1,x} = \mu_x(t)$, $m_{2,x} = \mu_x^n(t-)$, for big enough n we can bound

$$\mathbb{E} \left[\int_0^T \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x^n(t) F_{xy}^n(q_{xy}^n(t)) + R^n(\boldsymbol{\mu}^n(t)) \right) dt \right]$$

above by

$$V_K(\mathbf{m}) + 2\epsilon + \mathbb{P}\left(\sup_{t \in [0, T]} \|\boldsymbol{\mu}^n(t) - \boldsymbol{\mu}(t)\| > r_2\right) (V_K(\mathbf{m}) + TR_{\max} + 2\epsilon). \quad (120)$$

By using (81), the probability that there was no exit in the time interval $[0, T]$ is

$$\mathbb{P}(T^n \geq T) \leq \mathbb{P}\left(\sup_{t \in [0, T]} \|\boldsymbol{\mu}^n(t) - \boldsymbol{\mu}(t)\| > r_2\right) \leq c \frac{1}{\sqrt{nr_2}}.$$

Letting V_{\max} be the upper bound identified in Lemma 21 for the given $a > 0$, the total cost satisfies

$$\begin{aligned} V_K^n(\mathbf{m}^n) &\leq \mathbb{E}\left[\int_0^T \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x^n(t) F_{xy}^n(q_{xy}^n(t)) + R^n(\boldsymbol{\mu}^n(t))\right) dt + V(\boldsymbol{\mu}^n(T \wedge T^n))\right] \leq \\ &V_K(\mathbf{m}) + 2\epsilon + \mathbb{P}\left(\sup_{t \in [0, T]} \|\boldsymbol{\mu}^n(t) - \boldsymbol{\mu}(t)\| > r_2\right) (V_K(\mathbf{m}) + TR_{\max} + 2\epsilon) + \mathbb{P}(T^n \geq T) V_{\max} \leq \\ &V_K(\mathbf{m}) + 2\epsilon + 2(TR_{\max} + V_{\max} + 2\epsilon) \frac{c}{\sqrt{nr_2}}. \end{aligned} \quad (121)$$

By sending n to infinity we get the upper bound if $\mathbf{m} \in \mathcal{P}_a(\mathcal{X})$ for some $a > 0$. Next let $\mathbf{m} \in \mathcal{P}(\mathcal{X}) \setminus \mathcal{P}_*(\mathcal{X})$. Let $t_0 \leq \epsilon$ be such that $V_K(\boldsymbol{\nu}(\mathbf{m}, t_0)) \leq V_K(\mathbf{m}) + \epsilon$, where $\boldsymbol{\nu}(\mathbf{m}, t)$ is the solution to the original equation after time t . We can find a $r > 0$ such that for every $\tilde{\mathbf{m}} \in B(\boldsymbol{\nu}(\mathbf{m}, t_0), r)$, $V_K(\tilde{\mathbf{m}}) \leq V_K(\mathbf{m}) + 2\epsilon$. If $q^n(\tilde{\mathbf{m}}, t)$ is an ϵ -optimal control that corresponds to each initial condition $\tilde{\mathbf{m}}$, we define the control

$$q_{xy}^n(t) = \begin{cases} \gamma_{xy}, & t \leq t_0, \\ q_{xy}^n(\boldsymbol{\mu}^n(t_0), t - t_0), & t > t_0, \end{cases}$$

which gives

$$\begin{aligned} V_K^n(\mathbf{m}^n) &\leq \mathbb{E}\left[\int_0^{T^n} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x^n(s) F_{xy}^n(q_{xy}^n(s)) + R^n(\boldsymbol{\mu}^n(s))\right) dt\right] \\ &\leq \mathbb{E}\left[\int_0^{t_0} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x^n(s) F_{xy}^n(q_{xy}^n(s)) + R^n(\boldsymbol{\mu}^n(s))\right) dt\right] \\ &+ \mathbb{E}\left[\int_{t_0}^{T^n} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x^n(s) F_{xy}^n(q_{xy}^n(s)) + R^n(\boldsymbol{\mu}^n(s))\right) dt\right] \\ &\leq \mathbb{E}\left[\int_0^{t_0} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x^n(s) F_{xy}^n(\gamma_{xy}^n(s)) + R^n(\boldsymbol{\mu}^n(s))\right) dt\right] \\ &+ \mathbb{E}\left[\int_{t_0}^{T^n} \left(\sum_{(x,y) \in \mathcal{Z}} \mu_x^n(s) F_{xy}^n(q_{xy}^n(\boldsymbol{\mu}^n(t_0), s - t_0)) + R^n(\boldsymbol{\mu}^n(s))\right) dt\right] \leq t_0 R_{\max} + \mathbb{E}[V(\boldsymbol{\mu}^n(t_0))] \\ &\stackrel{\text{Lemma 21}}{\leq} \epsilon R_{\max} + P(\boldsymbol{\mu}^n(t_0) \in B(\boldsymbol{\nu}(\mathbf{m}, t_0), r)) (V_K(\mathbf{m}) + 2\epsilon) + P(\boldsymbol{\mu}^n(t_0) \notin B(\boldsymbol{\nu}(\mathbf{m}, t_0), r)) V_{\max} \\ &\leq V_K(\mathbf{m}) + (2 + R_{\max}^n)\epsilon + P(\boldsymbol{\mu}^n(t_0) \notin B(\boldsymbol{\nu}(\mathbf{m}, t_0), r)) V_{\max}. \end{aligned} \quad (122)$$

Now by an application of Lemma 20, we get that the last term goes to zero as n goes to ∞ , and since ϵ is arbitrary, we get that

$$\limsup V_K^n(\mathbf{m}^n) \leq V_K(\mathbf{m}).$$

■

7 Appendix: Properties of F_{xy}^n

Proof of Lemma 4. (1) We have

$$F_{xy}^n(q) = \sup_{u \in (0, \infty)} \left\{ u \ell \left(\frac{q}{u} \right) - \gamma_{xy} C_{xy}^n \left(\frac{u}{\gamma_{xy}} \right) \right\} \geq \gamma_{xy} \ell \left(\frac{q}{\gamma_{xy}} \right) - \gamma_{xy} C_{xy}^n \left(\frac{\gamma_{xy}}{\gamma_{xy}} \right) \geq \gamma_{xy} \ell \left(\frac{q}{\gamma_{xy}} \right) \geq 0.$$

(2) We have

$$\begin{aligned} F_{xy}^n(\gamma_{xy}) &= \sup_{u \in (0, \infty)} G_{xy}^n(u, \gamma_{xy}) = \sup_{u \in (0, \infty)} \left\{ u \ell \left(\frac{\gamma_{xy}}{u} \right) - \gamma_{xy} C_{xy}^n \left(\frac{u}{\gamma_{xy}} \right) \right\} \\ &= \sup_{u \in (0, \infty)} \left\{ \gamma_{xy} \log \gamma_{xy} - \gamma_{xy} \log u - \gamma_{xy} + u - \gamma_{xy} C_{xy}^n \left(\frac{u}{\gamma_{xy}} \right) \right\}, \end{aligned}$$

and by applying part 2 of Lemma 3

$$\gamma_{xy} C_{xy}^n \left(\frac{u}{\gamma_{xy}} \right) \geq \gamma_{xy} \log \gamma_{xy} - \gamma_{xy} \log u - \gamma_{xy} + u.$$

Therefore $F_{xy}^n(\gamma_{xy}) \leq 0$. However, by part (1) of this lemma $F_{xy}^n(\gamma_{xy}) \geq 0$, and therefore the equality follows.

(3) By definition $F_{xy}^n(q) = \sup_{u \in (0, \infty)} G_{xy}^n(u, q)$. Let $a \in (0, 1)$ and $0 \leq q_1 < q_2 < \infty$, and let $q = aq_1 + (1-a)q_2$. Using the convexity of $G_{xy}^n(u, q)$ for fixed u as a function of q , we have

$$\begin{aligned} F_{xy}^n(aq_1 + (1-a)q_2) &= \sup_{u \in (0, \infty)} G_{xy}^n(u, aq_1 + (1-a)q_2) \\ &\leq \sup_{u \in (0, \infty)} \left\{ aG_{xy}^n(u, q_1) + (1-a)G_{xy}^n(u, q_2) \right\} \\ &\leq a \sup_{u \in (0, \infty)} G_{xy}^n(u, q_1) + (1-a) \sup_{u \in (0, \infty)} G_{xy}^n(u, q_2) \\ &\leq aF_{xy}^n(q_1) + (1-a)F_{xy}^n(q_2). \end{aligned}$$

■

For the proof of Lemma 15 that is given below, we will use the following auxiliary lemma. Recall the definition of C_{xy}^n in (9).

Lemma 22 *If $\{C^n\}$ satisfies Assumption 13, then the following hold for every $(x, y) \in \mathcal{Z}$.*

1 *There exists a positive real number M , that does not depend on (x, y) , such that for the decreasing function $M_{xy}^1 : (0, \infty) \rightarrow [0, \infty)$, given by*

$$M_{xy}^1(q) \doteq \min \left\{ \gamma_{xy} \left(\frac{\gamma_{xy}}{q} \right)^{1/p}, M \right\},$$

we have that $G_{xy}^n(u, q)$ is increasing as a function of u on the interval $(0, M_{xy}^1(q)]$.

2 *There exists a decreasing function $M_{xy}^2 : (0, \infty) \rightarrow [0, \infty)$, with $M_{xy}^2(q) \geq M_{xy}^1(q)$, such that $G_{xy}^n(u, q)$ is decreasing as a function of u on the interval $[M_{xy}^2(q), \infty)$.*

Proof. By taking the derivative with respect to u in the definition (9) we get

$$-\frac{q}{u} - (C_{xy}^n)' \left(\frac{u}{\gamma_{xy}} \right) + 1. \quad (123)$$

(1) By part 2 of Assumption 13 there exists $M \in (0, \infty)$ such that if $u < M$, then

$$-\frac{q}{u} - (C_{xy}^n)' \left(\frac{u}{\gamma_{xy}} \right) + 1 \geq -\frac{q}{u} + \left(\frac{\gamma_{xy}}{u} \right)^{p+1} + 1,$$

and by taking $u \leq \gamma_{xy} (\gamma_{xy}/q)^{1/p}$ we get

$$-\frac{q}{u} + \left(\frac{\gamma_{xy}}{u}\right)^{p+1} + 1 \geq -\frac{q}{u} + \frac{q}{u} + 1 > 0.$$

Therefore for

$$M_{xy}^1(q) = \min \left\{ \gamma_{xy} \left(\frac{\gamma_{xy}}{q}\right)^{1/p}, M \right\},$$

we have $-\frac{q}{u} - (C_{xy}^n)' \left(\frac{u}{\gamma_{xy}}\right) + 1 \geq 0$ on the interval $(0, M_{xy}^1(q)]$.

(2) By applying part 3 of Assumption 13, we get that there exists decreasing $\tilde{M}_{xy}^2(q) < \infty$, such that if $u > \tilde{M}_{xy}^2(q)$ then

$$\frac{u}{\gamma_{xy}} (C_{xy}^n)' \left(\frac{u}{\gamma_{xy}}\right) - \frac{u}{\gamma_{xy}} \geq -\frac{q}{\gamma_{xy}}. \quad (124)$$

Then $M_{xy}^2(q) \doteq \max\{M_{xy}^1(q), \tilde{M}_{xy}^2(q)\}$, is decreasing and bigger than M_{xy}^1 , and using (124) we get

$$-\frac{q}{u} - (C_{xy}^n)' \left(\frac{u}{\gamma_{xy}}\right) + 1 = -\frac{q}{u} - \frac{\gamma_{xy}}{u} \left(\frac{u}{\gamma_{xy}} (C_{xy}^n)' \left(\frac{u}{\gamma_{xy}}\right) - \frac{u}{\gamma_{xy}}\right) \leq 0$$

on the interval $[M_{xy}^2(q), \infty)$. ■

Proof of Lemma 15. (1) Let $\epsilon > 0$, and $q \geq \epsilon$. By Lemma 22, we have that $G_{xy}^n(u, q)$, as a function of u , is increasing on the interval $(0, M_{xy}^1(q)]$. Therefore for all $u \in (0, M_{xy}^1(q)]$ we have

$$\begin{aligned} u\ell\left(\frac{q}{u}\right) - \gamma_{xy} C_{xy}^n\left(\frac{u}{\gamma_{xy}}\right) &\leq M_{xy}^1(q)\ell\left(\frac{q}{M_{xy}^1(q)}\right) - \gamma_{xy} C_{xy}^n\left(\frac{M_{xy}^1(q)}{\gamma_{xy}}\right) \leq M_{xy}^1(q)\ell\left(\frac{q}{M_{xy}^1(q)}\right) \\ &\leq q \log\left(\frac{q}{M_{xy}^1(q)}\right) + M_{xy}^1(q) \leq q \log\left(\frac{q}{M_{xy}^1(q)}\right) + M_{xy}^1(\epsilon) \\ &\leq q \log(q) - q \log(M_{xy}^1(q)) + M_{xy}^1(\epsilon) \\ &\stackrel{M_{xy}^1(\epsilon) \leq M_{xy}^2(\epsilon)}{\leq} q \log(q) - q \log(M_{xy}^1(q)) + M_{xy}^2(\epsilon). \end{aligned}$$

By the second part of Lemma 22, we have that $G_{xy}^n(u, q)$ is decreasing on the interval $(M_{xy}^2(\epsilon), \infty)$. Therefore for all $u \in (M_{xy}^2(\epsilon), \infty)$

$$\begin{aligned} u\ell\left(\frac{q}{u}\right) - \gamma_{xy} C_{xy}^n\left(\frac{u}{\gamma_{xy}}\right) &\leq M_{xy}^2(\epsilon)\ell\left(\frac{q}{M_{xy}^2(\epsilon)}\right) - \gamma_{xy} C_{xy}^n\left(\frac{M_{xy}^2(\epsilon)}{\gamma_{xy}}\right) \leq M_{xy}^2(\epsilon)\ell\left(\frac{q}{M_{xy}^2(\epsilon)}\right) \\ &\leq q \log\left(\frac{q}{M_{xy}^2(\epsilon)}\right) + M_{xy}^2(\epsilon) \\ &\stackrel{M_{xy}^2(q) \leq M_{xy}^2(\epsilon)}{\leq} q \log(q) - q \log(M_{xy}^2(q)) + M_{xy}^2(\epsilon) \\ &\stackrel{M_{xy}^1(q) \leq M_{xy}^2(q)}{\leq} q \log(q) - q \log(M_{xy}^1(q)) + M_{xy}^2(\epsilon). \end{aligned}$$

Finally for the interval $[M_{xy}^1(q), M_{xy}^2(\epsilon)]$ we have

$$\begin{aligned} u\ell\left(\frac{q}{u}\right) - \gamma_{xy} C_{xy}^n\left(\frac{u}{\gamma_{xy}}\right) &\leq u\ell\left(\frac{q}{u}\right) = q \log q - q \log u - q + u \\ &\leq q \log q - q \log(M_{xy}^1(q)) + M_{xy}^2(\epsilon). \end{aligned}$$

Now if we recall the definition of M_{xy}^1 given in Lemma 22 and set $\bar{M}(q) \doteq \max\{M_{xy}^2(q) : (x, y) \in \mathcal{Z}\}$, then

$$G_{xy}^n(u, q) \leq q \log \frac{q}{\min \left\{ \gamma_{xy} \left(\frac{\gamma_{xy}}{q}\right)^{1/p}, M \right\}} + \bar{M}(\epsilon),$$

and by taking supremum over u we end up with $F_{xy}^n(q)$ satisfying the same bound.

(2) This is straightforward since F_{xy}^n is finite on the interval $(0, \infty)$, and convex.

(3) Let $1 > \epsilon > 0$. For every $q \in [\epsilon, \frac{1}{\epsilon}]$ and $n \in \mathbb{N}$, we have that $u\ell\left(\frac{q}{u}\right) - \gamma_{xy}C_{xy}^n\left(\frac{u}{\gamma_{xy}}\right)$ is increasing on the interval $(0, M_{xy}^2(\epsilon)]$ and decreasing on $[M_{xy}^1(\frac{1}{\epsilon}), \infty)$ and since C_{xy}^n converges locally uniformly to C_{xy} , then the same conclusion holds for $u\ell\left(\frac{q}{u}\right) - \gamma_{xy}C_{xy}\left(\frac{u}{\gamma_{xy}}\right)$. It is straight forward to conclude that in all cases, the supremum is achieved on the interval $[M_{xy}^2(\epsilon), M_{xy}^1(\frac{1}{\epsilon})]$. We now define the sets

$$A_{\epsilon, xy}^{a, i} = \left\{ u : \gamma_{xy}C_{xy}^i\left(\frac{u}{\gamma_{xy}}\right) \leq a \sup_{\epsilon \leq q \leq \frac{1}{\epsilon}, M_{xy}^2(\epsilon) \leq \tilde{u} \leq M_{xy}^1(\frac{1}{\epsilon})} \tilde{u}\ell\left(\frac{q}{\tilde{u}}\right) \right\},$$

with $i \in \mathbb{N} \cup \{\infty\}$. By uniform convergence of C_{xy}^n in C_{xy}^∞ in compact subsets of $(u_{1, xy}, u_{2, xy})$, and the monotonicity properties of C_{xy}^n , we get that for large enough n , we have $A_{\epsilon, xy}^{1, n} \subset A_{\epsilon, xy}^{2, \infty} \subset A_{\epsilon, xy}^{3, n}$. For every $q \in [\epsilon, \frac{1}{\epsilon}]$, we have

$$\begin{aligned} F_{xy}^n(q) &= \sup_{u \in [M_{xy}^2(\epsilon), M_{xy}^1(\frac{1}{\epsilon})]} \left\{ u\ell\left(\frac{q}{u}\right) - \gamma_{xy}C_{xy}^n\left(\frac{u}{\gamma_{xy}}\right) \right\} \\ &= \sup_{u \in [M_{xy}^2(\epsilon), M_{xy}^1(\frac{1}{\epsilon})] \cap A_{\epsilon, xy}^{1, n}} \left\{ u\ell\left(\frac{q}{u}\right) - \gamma_{xy}C_{xy}^n\left(\frac{u}{\gamma_{xy}}\right) \right\} \\ &\leq \sup_{u \in [M_{xy}^2(\epsilon), M_{xy}^1(\frac{1}{\epsilon})] \cap A_{\epsilon, xy}^{2, \infty}} \left\{ u\ell\left(\frac{q}{u}\right) - \gamma_{xy}C_{xy}^\infty\left(\frac{u}{\gamma_{xy}}\right) \right\} \\ &\quad + \sup_{u \in [M_{xy}^2(\epsilon), M_{xy}^1(\frac{1}{\epsilon})] \cap A_{\epsilon, xy}^{2, \infty}} \left\{ \left| \gamma_{xy}C_{xy}^n\left(\frac{u}{\gamma_{xy}}\right) - \gamma_{xy}C_{xy}^\infty\left(\frac{u}{\gamma_{xy}}\right) \right| \right\} \\ &\leq \sup_{u \in [M_{xy}^2(\epsilon), M_{xy}^1(\frac{1}{\epsilon})]} \left\{ u\ell\left(\frac{q}{u}\right) - \gamma_{xy}C_{xy}^\infty\left(\frac{u}{\gamma_{xy}}\right) \right\} \\ &\quad + \sup_{u \in [M_{xy}^2(\epsilon), M_{xy}^1(\frac{1}{\epsilon})] \cap A_{\epsilon, xy}^{2, \infty}} \left\{ \left| \gamma_{xy}C_{xy}^n\left(\frac{u}{\gamma_{xy}}\right) - \gamma_{xy}C_{xy}^\infty\left(\frac{u}{\gamma_{xy}}\right) \right| \right\}. \end{aligned}$$

By doing the same for \bar{F}_{xy}^∞ , we get

$$\sup_{q \in [\epsilon, \frac{1}{\epsilon}]} |F_{xy}^n(q) - \bar{F}_{xy}^\infty(q)| \leq \sup_{u \in [M_{xy}^2(\epsilon), M_{xy}^1(\frac{1}{\epsilon})] \cap A_{\epsilon, xy}^{1, \infty}} \left\{ \left| \gamma_{xy}C_{xy}^n\left(\frac{u}{\gamma_{xy}}\right) - \gamma_{xy}C_{xy}^\infty\left(\frac{u}{\gamma_{xy}}\right) \right| \right\},$$

and therefore F_{xy}^n converges locally uniformly to \bar{F}_{xy}^∞ on $A_{\epsilon, xy}^{1, \infty}$. Now by the definition of F_{xy}^∞ , i.e.

$$F_{xy}^\infty(q) = \inf \left\{ \liminf_{n \rightarrow \infty} F_{xy}^n(q^n) : \{q^n\} \text{ sequence in } \mathbb{R} \text{ with } q^n \rightarrow q \right\},$$

it is easy to see that since F_{xy}^n converges locally uniformly to \bar{F}_{xy}^∞ on $(0, \infty)$, and so $F_{xy}^\infty = \bar{F}_{xy}^\infty$ on $(0, \infty)$. Also F_{xy}^∞ as a pointwise limit of convex functions on $(0, \infty)$, is also a convex function there. It remains to investigate what happens with F_{xy}^∞ at zero. Let $q^n \rightarrow 0$ with $F_{xy}^n(q^n) \rightarrow F_{xy}^\infty(0)$. For $a \in (0, 1)$, we have

$$\begin{aligned} aF_{xy}^\infty(0) + (1-a)F_{xy}^\infty(q) &= aF_{xy}^n(q^n) + (1-a)F_{xy}^n(q) + a(F_{xy}^n(q^n) - F_{xy}^\infty(0)) + (1-a)(F_{xy}^\infty(q) - F_{xy}^n(q)) \\ &\geq F_{xy}^n(aq^n + (1-a)q) + a(F_{xy}^n(q^n) - F_{xy}^\infty(0)) + (1-a)(F_{xy}^\infty(q) - F_{xy}^n(q)) \end{aligned}$$

Now if we take the limit, then by continuity of each F_{xy}^n on $(0, \infty)$ and the uniform convergence on every compact subset (also on $[(1-a)q, q]$), we have $aF_{xy}^\infty(0) + (1-a)F_{xy}^\infty(q) \geq F_{xy}^\infty((1-a)q)$. ■

8 Appendix: Tightness functionals

Proof of Lemma 18. Let $c_2 > 0$ and $\{(\mu^n, T^n)\}$ be a deterministic sequence in S with μ^n absolutely continuous such that

$$\int_0^{T^n} \ell(|\dot{\mu}^n(t)|) dt + c_1 T^n \leq c_2$$

and $|\dot{\mu}^n(t)| = 0$ for $t > T^n$. We need to show that H has level sets with compact closure. Since all elements are positive, we have that $T^n \leq c_2/c_1$. Let $\bar{\mu}^n$ denote the restriction of μ^n to $[0, c_2/c_1]$. If we prove that $\bar{\mu}^n$ converges along some subsequence then we are done. Using the inequality $ab \leq e^{ca} + \ell(b)/c$, which is valid for $a, b \geq 0$, and $c \geq 1$, we have that

$$|\mu^n(t) - \mu^n(s)| \leq \int_t^s |\dot{\mu}^n(r)| dr \leq (t-s)e^c + \frac{c_2}{c}.$$

This shows that $\{\bar{\mu}^n\}$ are equicontinuous. Since $\bar{\mu}^n(t)$ takes values in the compact set $\mathcal{P}(\mathcal{X})$, by the Arzela-Ascoli theorem there is a convergent subsequence. ■

9 Appendix: Properties of Hamiltonians

Lemma 23 *Under Assumption 2 the Isaac condition is satisfied, i.e.,*

$$H^{-,n}(\mathbf{m}, \xi) = \sup_{\mathbf{u} \in (0, \infty)^{\otimes \mathcal{Z}}} \inf_{\mathbf{q} \in [0, \infty)^{\otimes \mathcal{Z}}} \sum_{(x,y) \in \mathcal{Z}} m_x \{q_{xy} \xi_{xy} + G_{xy}^n(u_{xy}, q_{xy})\} = \quad (125)$$

$$\inf_{\mathbf{q} \in [0, \infty)^{\otimes \mathcal{Z}}} \sup_{\mathbf{u} \in (0, \infty)^{\otimes \mathcal{Z}}} \sum_{(x,y) \in \mathcal{Z}} m_x \{q_{xy} \xi_{xy} + G_{xy}^n(u_{xy}, q_{xy})\} = H^{+,n}(\mathbf{m}, \xi). \quad (126)$$

Proof. We have

$$H^{-,n}(\mathbf{m}, \xi) = \sup_{\mathbf{u} \in (0, \infty)^{\otimes \mathcal{Z}}} \inf_{\mathbf{q} \in [0, \infty)^{\otimes \mathcal{Z}}} \sum_{(x,y) \in \mathcal{Z}} m_x \{q_{xy} \xi_{xy} + G_{xy}^n(u_{xy}, q_{xy})\} =$$

$$\sum_{(x,y) \in \mathcal{Z}} m_x \sup_{u_{xy} \in (0, \infty)} \inf_{q_{xy} \in [0, \infty)} \left\{ q_{xy} \xi_{xy} + u_{xy} \ell \left(\frac{q_{xy}}{u_{xy}} \right) - \gamma_{xy} C_{xy}^m \left(\frac{u_{xy}}{\gamma_{xy}} \right) \right\}.$$

If we prove the exchange of sup and inf for each $(x, y) \in \mathcal{Z}$, then we are done.

Since ℓ is convex

$$L_{xy}^n(u, q) = q\xi + u\ell \left(\frac{q}{u} \right) - \gamma_{xy} C_{xy}^n \left(\frac{u}{\gamma_{xy}} \right)$$

is convex with respect to q . It is easy to see that $L_{xy}^n(u, q)$ is not concave with respect to u , however under Assumption 2, we can show that L_{xy}^n is quasi-concave with respect to u (i.e., $\{u : L_{xy}^n(u, q) \geq c\}$ is convex for every $q \in [0, \infty)$, and $c \in \mathbb{R}$.)

By differentiating with respect to u we get

$$\partial_u L_{xy}^n(u, q) = -\frac{q}{u} + 1 - (C_{xy}^n)' \left(\frac{u}{\gamma_{xy}} \right).$$

If we prove that for each q the set of roots for $\partial_u L_{xy}^n(u, q)$ is an interval or a point we are done, because a real function that changes monotonicity from increasing to decreasing at most once is quasi-concave. However $\partial_u L_{xy}^n(u, q)$ has the same roots as $u(C_{xy}^n)' \left(\frac{u}{\gamma_{xy}} \right) - u + q$. By part 1 of Assumption 1 $u(C_{xy}^n)' \left(\frac{u}{\gamma_{xy}} \right) - u + q$ is increasing, which gives what is needed.

To prove the exchange between supremum and infimum, we apply Sion's Theorem (Corollary 3.3 in [26]), which states that if a continuous $F(u, q)$ is quasi-concave for every u in some convex set U and quasi-convex for every q in some convex set Q , and if one of the two sets is compact, then we can exchange the supremum with the infimum. In our case both sets are non-compact, and so this result cannot be applied directly, but it can be applied using the fact that $\lim_{q \rightarrow \infty} L_{xy}^n(q, 1) = \infty$, as we now explain.

If we prove that

$$\inf_{q \in [0, \infty)} \sup_{u \in (0, \infty)} L_{xy}^n(u, q) = \lim_{r \rightarrow \infty} \inf_{q \in [0, \infty)} \sup_{u \in [r, \frac{1}{r}]} L_{xy}^n(u, q),$$

then we are done, since by Corollary 3.3 in [26]

$$\inf_{q \in [0, \infty)} \sup_{u \in (0, \infty)} L_{xy}^n(u, q) = \lim_{r \rightarrow \infty} \inf_{q \in [0, \infty)} \sup_{u \in [r, \frac{1}{r}]} L_{xy}^n(u, q) =$$

$$\lim_{r \rightarrow \infty} \sup_{u \in [r, \frac{1}{r}]} \inf_{q \in [0, \infty)} L_{xy}^n(u, q) = \sup_{u \in (0, \infty)} \inf_{q \in [0, \infty)} L_{xy}^n(u, q).$$

Let $M \doteq \inf_{q \in [0, \infty)} \sup_{u \in (0, \infty)} L_{xy}^n(u, q)$. We will assume that $M < \infty$, and note that the case $M = \infty$ is treated similarly. Since $\lim_{q \rightarrow \infty} L_{xy}^n(q, 1) = \infty$, we can find \tilde{q} such that $L_{xy}^n(q, 1) > 2M$ for every $q \geq \tilde{q}$. Now we have

$$\inf_{q \in [0, \infty)} \sup_{u \in (0, \infty)} L_{xy}^n(u, q) = \inf_{q \in [0, \tilde{q}]} \sup_{u \in (0, \infty)} L_{xy}^n(u, q),$$

and

$$\inf_{q \in [0, \tilde{q}]} \sup_{u \in [r, \frac{1}{r}]} L_{xy}^n(u, q) = \inf_{q \in [0, \infty)} \sup_{u \in [r, \frac{1}{r}]} L_{xy}^n(u, q),$$

which gives

$$\begin{aligned} \inf_{q \in [0, \infty)} \sup_{u \in (0, \infty)} L_{xy}^n(u, q) &= \inf_{q \in [0, \tilde{q}]} \sup_{u \in (0, \infty)} L_{xy}^n(u, q) = \sup_{u \in (0, \infty)} \inf_{q \in [0, \tilde{q}]} L_{xy}^n(u, q) = \\ \lim_{r \rightarrow \infty} \sup_{u \in [r, \frac{1}{r}]} \inf_{q \in [0, \tilde{q}]} L_{xy}^n(u, q) &= \lim_{r \rightarrow \infty} \inf_{q \in [0, \tilde{q}]} \sup_{u \in [r, \frac{1}{r}]} L_{xy}^n(u, q) = \lim_{r \rightarrow \infty} \inf_{q \in [0, \infty)} \sup_{u \in [r, \frac{1}{r}]} L_{xy}^n(u, q). \end{aligned}$$

■

Theorem 24 Let $H^{+,n}(m, \xi)$ and $H^{-,n}(m, \xi)$ as in (125). If C_{xy}^n is twice differentiable on an interval $(u_{1,xy}^n, u_{2,xy}^n)$ and the equality holds, then the first part of Assumption 2 holds true on that interval.

Proof. Since each term in the sum that generates $H^{+,n}$ is bigger than the corresponding one in the sum of $H^{-,n}$, we get equality for all of them. By the theory of Legendre transformations we know that $\inf_{q \in [0, \infty)} \sup_{u \in (0, \infty)} \{q\xi_{xy} + G_{xy}^n(u, q)\}$ is actually a concave function. Since we can exchange the order between supremum and infimum, $\sup_{u \in (0, \infty)} \inf_{q \in [0, \infty)} \{q\xi_{xy} + G_{xy}^n(u, q)\}$ must be a concave function as well. By using the formula

$$\sup_{u \in (0, \infty)} \inf_{q \in [0, \infty)} \{q\xi + G_{xy}^n(u, q)\} = \sum_{(x,y) \in \mathcal{Z}} m_x \gamma_{xy} (C_{xy}^n)^* (-\ell^*(-\xi_{xy}))$$

we have that $(C_{xy}^n)^* (-\ell^*(\xi)) = (C_{xy}^n)^* (1 - e^\xi)$ must also be concave. By differentiating with respect to ξ we get, $e^{2\xi} \left((C_{xy}^n)^* \right)'' (1 - e^\xi) - e^\xi \left((C_{xy}^n)^* \right)' (1 - e^\xi) \leq 0$, from which, by using the identity $(f^*)' = (f')^{-1}$, we get

$$e^{2\xi} \left(\left((C_{xy}^n)' \right)^{-1} \right)' (1 - e^\xi) - e^\xi \left((C_{xy}^n)' \right)^{-1} (1 - e^\xi) \leq 0.$$

By substituting $\tilde{u} = 1 - e^\xi$ we get

$$\begin{aligned} (1 - \tilde{u}) \left(\left((C_{xy}^n)' \right)^{-1} \right)' (\tilde{u}) - \left((C_{xy}^n)' \right)^{-1} (\tilde{u}) &\leq 0, & \text{with } \tilde{u} \leq 1 \\ (1 - \tilde{u}) \frac{1}{(C_{xy}^n)'' \left(\left((C_{xy}^n)' \right)^{-1} (\tilde{u}) \right)} - \left((C_{xy}^n)' \right)^{-1} (\tilde{u}) &\leq 0, & \text{with } \tilde{u} \leq 1 \\ \left(1 - (C_{xy}^n)'(r) \right) \frac{1}{(C_{xy}^n)''(r)} - r &\leq 0, & \text{with } (C_{xy}^n)'(r) \leq 1 \\ 1 - (C_{xy}^n)'(r) - r (C_{xy}^n)''(r) &\leq 0, & \text{with } (C_{xy}^n)'(r) \leq 1 \\ r (C_{xy}^n)''(r) + (C_{xy}^n)'(r) - 1 &\geq 0, & \text{with } (C_{xy}^n)'(r) \leq 1. \end{aligned}$$

Now the last inequality implies that either $(C_{xy}^n)'(u) \geq 1$ or that $u(C_{xy}^n)'(u) - u$ is locally increasing and even more that if $(C_{xy}^n)'(u_0) \geq 1$ for some u_0 , then it must remain like that for every $u \geq u_0$. If that was not the case then we can find $u_1 > u_0$ such that $u_1(C_{xy}^n)'(u_1) - u_1 < \hat{q}$ for some negative \hat{q} , while $u_0(C_{xy}^n)'(u_0) - u_0 \geq 0$. By a suitable application of the mean value theorem we will get the existence of an r that the last inequality fails. If we set $\tilde{u}_{xy}^n = \inf\{u : (C_{xy}^n)'(u) \geq 1\}$, then the Assumption 2 is recovered. ■

References

- [1] A. Arapostathis, V. S. Borkar, E. Fernández-Gaucherand, M. K. Ghosh, and S. I. Marcus. Discrete-Time Controlled Markov Processes with Average Cost Criterion: A Survey. *SIAM Journal on Control and Optimization*, 31(2):282–344, mar 1993.

- [2] G. Avila-Godoy and E. Fernández-Gaucherand. Controlled Markov chains with exponential risk-sensitive criteria: modularity, structured policies and applications. In *Decision and Control, 1998. Proceedings of the 37th IEEE Conference on*, volume 1, pages 778–783. IEEE, 1998.
- [3] D. P. Bertsekas. *Dynamic Programming and Optimal Control: Approximate Dynamic Programming*, volume 2. Athena Scientific, 2012.
- [4] P. Billingsley. *Probability and measure*. Wiley Series in Probability and Mathematical Statistics. John Wiley & Sons, Inc., New York, third edition, 1995.
- [5] V. S. Borkar and S. P. Meyn. Risk-Sensitive Optimal Control for Markov Decision Processes with Monotone Cost. *Mathematics of Operations Research*, 27(1):192–209, feb 2002.
- [6] R. Cavazos-Cadena. Optimality equations and inequalities in a class of risk-sensitive average cost Markov decision chains. *Mathematical Methods of Operations Research*, 71(1):47–84, feb 2010.
- [7] K.-J. Chung and M. J. Sobel. Discounted MDP's: Distribution Functions and Exponential Utility Maximization. *SIAM Journal on Control and Optimization*, 25(1):49–62, jan 1987.
- [8] G. B. Di Masi and Ł. Stettner. Infinite Horizon Risk Sensitive Control of Discrete Time Markov Processes under Minorization Property. *SIAM Journal on Control and Optimization*, 46(1):231–252, jan 2007.
- [9] P. Dupuis, M. James, and I. Petersen. Robust properties of risk-sensitive control. In *Proceedings of the 37th IEEE Conference on Decision and Control (Cat. No.98CH36171)*, volume 2, pages 2365–2370. IEEE.
- [10] P. Dupuis, M. R. James, and I. R. Petersen. Robust properties of risk-sensitive control. *Math. Control Signals Systems*, 13:318–332, 2000.
- [11] P. Dupuis and W. McEneaney. Risk-sensitive and robust escape criteria. *SIAM journal on control and optimization*, 35(6):2021–2049, 1997.
- [12] P. Dupuis, K. Ramanan, and W. Wu. Large Deviation Principle For Finite-State Mean Field Interacting Particle Systems. *arXiv:1601.06219*, page 62, jan 2016.
- [13] W. H. Fleming and D. Hernández-Hernández. Risk-Sensitive Control of Finite State Machines on an Infinite Horizon I. *SIAM Journal on Control and Optimization*, 35(5):1790–1810, sep 1997.
- [14] W. H. Fleming and H. M. Soner. Asymptotic expansions for Markov processes with Levy generators. *Appl. Math. Optimization*, 19:203–223, 1989.
- [15] M. K. Ghosh and S. Saha. Risk-sensitive control of continuous time Markov chains. *Stochastics: An International Journal of Probability and Stochastic Processes*, (October 2014):37–41, 2014.
- [16] D. Hernandez-Hernández and S. I. Marcus. Risk sensitive control of Markov processes in countable state space. *Systems & Control Letters*, 29(3):147–155, nov 1996.
- [17] O. Hernández-Lerma and J. B. Lasserre. *Further Topics in Discrete Time Markov Control Processes*. Springer, 1999.
- [18] R. Howard and J. Matheson. Risk-sensitive Markov decision processes. *Management science*, 1972.
- [19] N. Ikeda and S. Watanabe. *Stochastic differential equations and diffusion processes*, volume 24 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam; Kodansha, Ltd., Tokyo, second edition, 1989.
- [20] A. Jaskiewicz. Average optimality for risk-sensitive control with general state space. *Annals of Applied Probability*, 17(2):654–675, apr 2007.
- [21] S. I. Marcus, E. Fernandez-Gaucherand, D. Hernandez-Hernandez, S. P. Coraluppi, and P. Fard. Risk Sensitive Markov Decision Processes. In *Systems and Control in the 21st Century*, page 17. Birkhäuser Boston, Boston, MA, 1997.
- [22] S. N. Ethier and T. G. Kurtz. Markov Processes. In *SpringerReference*, page 544. 1986.
- [23] I. Petersen, M. James, and P. Dupuis. Minimax optimal control of stochastic uncertain systems with relative entropy constraints. *Automatic Control, IEEE Transactions on*, 45(3):398–412, 2000.

- [24] M. Puterman. Markov decision processes. In D. Heyman and M. Sobel, editors, *Stochastic Models, Volume 2*, chapter 8. North Holland, Amsterdam, 1991.
- [25] A. Ruszczyński. Risk-averse dynamic programming for Markov decision processes. *Mathematical Programming*, 125(2):235–261, oct 2010.
- [26] M. Sion. On general minimax theorems. *Pacific J. Math*, 1958.
- [27] P. Whittle. Optimal Control: Basics and Beyond. *Wiley-Interscience series in systems and optimization*, page 464, 1996.
- [28] G. Xianping and O. Hernández-Lerma. *Continuous-time Markov decision processes : theory and applications*. Springer, 2009.