Weierstraß–Institut für Angewandte Analysis und Stochastik

im Forschungsverbund Berlin e.V.

Approximate wavelets and the approximation of pseudodifferential operators

Vladimir Maz'ya¹ , Gunther Schmidt²

submitted: 13rd June 1996

 Department of Mathematics Linköping University
 S – 581 83 Linköping
 Sweden
 e-mail: vlmaz@math.liu.se ² Weierstrass Institute for Applied Analysis and Stochastics Mohrenstraße 39 D - 10117 Berlin Germany e-mail: schmidt@wias-berlin.de

Preprint No. 249 Berlin 1996

1991 Mathematics Subject Classification. 41A30, 41A63; Secondary 65D30.

Key words and phrases. Cubature formulas, approximate multiresolution, multivariate wavelets.

Edited by Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS) Mohrenstraße 39 D — 10117 Berlin Germany

Fax: + 49 30 2044975 e-mail (X.400): c=de;a=d400-gw;p=WIAS-BERLIN;s=preprint e-mail (Internet): preprint@wias-berlin.de ABSTRACT. The paper studies an approximate multiresolution analysis for spaces generated by smooth functions which provide high order cubature formulas for integral operators of mathematical physics. Since these functions satisfy refinement equations with any prescribed accuracy methods of the wavelet theory can be applied. We obtain a decomposition of the finest scale space into almost orthogonal wavelet spaces. For one example we study some properties of the analytic prewavelets, describe the projection operators onto the wavelet spaces and consider some applications to the cubature of integral operators.

1. INTRODUCTION

In this paper we introduce the so called *approximate* wavelet decompositions of spaces of approximating functions which appeared to be very useful for constructing high order semi-analytic cubature formulas for important classes of pseudodifferential and other integral operators of mathematical physics. The application of wavelet based methods to the representation of integral and differential operators is one of the actual research topics in the numerical analysis of solution methods for the corresponding operator equations.

Let us describe a usual setting. Starting from a finite sequence of nested closed subspaces

(1.1)
$$V_0 \subset V_1 \subset \ldots \subset V_n \subset L_2(\mathbf{R}^d)$$

the space of approximating functions V_n , corresponding to the finest grid, is decomposed into the orthogonal sum

(1.2)
$$V_n = V_0 \bigoplus_{j=0}^{n-1} W_j ,$$

where the wavelet space W_j is the orthogonal complement $W_j = V_{j+1} \ominus V_j$. The chain (1.1) is called a multiresolution analysis of V_n if the spaces V_j have the properties

- (i) $f(\mathbf{x}) \in V_0$ if and only if $f(\mathbf{x} \mathbf{m}) \in V_0$ for any $\mathbf{m} \in \mathbf{Z}^d$;
- (ii) $f(\mathbf{x}) \in V_j$ if and only if $f(2\mathbf{x}) \in V_{j+1}$ for any $j = 0, \ldots, n-1$;
- (iii) there exists a function ϕ such that $\{\phi(\cdot \mathbf{m})\}_{\mathbf{m}\in\mathbb{Z}^d}$ is an L_2 -stable basis in V_0 , i.e. there exist constants $c_2 > c_1 > 0$ such that

(1.3)
$$c_1 \|\{a_{\mathbf{m}}\}\|_{\ell^2} \le \|\sum_{\mathbf{m}\in\mathbf{Z}^d} a_{\mathbf{m}}\phi(\cdot-\mathbf{m})\|_2 \le c_2 \|\{a_{\mathbf{m}}\}\|_{\ell_2}, \quad \forall \{a_{\mathbf{m}}\} \in \ell_2(\mathbf{Z}^d)$$

Then obviously the spaces V_j are spanned by the dilated shifts $\phi(2^j \cdot -\mathbf{m})$, $\mathbf{m} \in \mathbf{Z}^d$, of the scaling function ϕ . The main goal of the multiresolution is to determine a new basis of the space V_n , which is used in numerical procedures. It is well known that there exist $2^d - 1$ functions $\psi_v \in W_0$, called prewavelets, such that the shifts $\{\psi_v(2^j \cdot -\mathbf{m}), \mathbf{m} \in \mathbf{Z}^n, v \in \mathcal{V}'\}$ form an L_2 -stable basis in the space W_j ([6], [12]). Here we index the prewavelets ψ_v by the set $\mathcal{V}' = \mathcal{V} \setminus \{\mathbf{0}\} \{\mathbf{0}\}$ with V denoting the set of vertices of the cube $[0, 1/2]^d$. Thus one obtains an L_2 -stable basis of the space V_n consisting of

$$\{\phi(\cdot-\mathbf{m}),\mathbf{m}\in\mathbf{Z}^d\} \quad ext{and} \quad \{\psi_v(2^j\cdot-\mathbf{m}),\mathbf{m}\in\mathbf{Z}^d, v\in\mathcal{V}', j=0,\ldots,n-1\}\;.$$

Similar to other transform methods elements of V_n and operators are now expanded into the new basis and the computations take place in this system of coordinates, where one hopes to achieve that the computation is faster than in the original system. Additionally, some features of wavelets, as the localization in both space and frequency domains and vanishing moment properties lead to a number of new and interesting properties of wavelet based numerical methods. The multiresolution structure of the wavelet expansion leads to an effective organisation of transformations. expansion Furthermore, the vanishing moments of wavelets imply that within a prescribed accuracy pseudodifferential operators admit sparse matrix representations, which allows to design fast numerical algorithms for these operators. There exists a series of papers on the application of wavelet methods to the computation of integral operators and the solution of integral equations, where different types of scaling functions and wavelets are used (see [1], [2], [5] and [13] the references therein). Since the scaling function ϕ has to satisfy the so called refinement equation

(1.4)
$$\phi(\mathbf{x}) = \sum_{\mathbf{m}\in\mathbf{Z}^n} a_{\mathbf{m}} \phi(2\mathbf{x} - \mathbf{m})$$

as a rule these functions are piecewise polynomials satisfying some smoothness and vanishing moment conditions. Many interesting examples can be found in [6], [12], [4] and the above mentioned papers. However, one drawback of these functions is that it is practically impossible to derive analytic formulas for the action of important integral operators of mathematical physics on these functions, especially in the multidimensional case. Thus it is necessary to use cubatures for integral operators with singular kernel functions applied to piecewise polynomials.

A different point of view to the cubature of integral operators of mathematical physics

$$Ku(\mathbf{x}) = \int\limits_{\mathbf{R}^d} k(\mathbf{x}, \mathbf{y}) u(\mathbf{y}) \, d\mathbf{y}$$

was developed in [7], [8], [9] and [11], where the function u is approximated by linear combinations of the form

(1.5)
$$u_h(\mathbf{x}) := \mathcal{D}^{-n/2} \sum_{\mathbf{m} \in \mathbf{Z}^d} u_{\mathbf{m}} \eta \left(\frac{\mathbf{x} - h\mathbf{m}}{\sqrt{\mathcal{D}}h} \right) \,.$$

Here the generating function η decays together with its Fourier transform $\mathcal{F}\eta$ rapidly at infinity and has the property that the integral $K\eta$ can be evaluated very effectively, either analytically or by simple one-dimensional quadrature. For example, the Newton potential of the function $\eta(\mathbf{x}) = (5/2 - |\mathbf{x}|^2) \exp(-|\mathbf{x}|^2)$, $\mathbf{x} \in \mathbf{R}^3$, is given by

(1.6)
$$\mathcal{N}\eta(\mathbf{x}) := \frac{1}{4\pi} \int_{\mathbf{R}^3} \frac{(5/2 - |\mathbf{y}|^2) e^{-|\mathbf{y}|^2}}{|\mathbf{x} - \mathbf{y}|} d\mathbf{y} = \frac{1}{2\pi^{3/2}} \left(\frac{1}{|\mathbf{x}|} \int_{0}^{|\mathbf{x}|} e^{-\tau^2} d\tau + e^{-|\mathbf{x}|^2} \right)$$

Hence, if the density can be represented as the sum $u_h(\mathbf{x})$ one obtains analytic formulas for the Newton potential and its derivatives, for example.

The crucial problem of this approach lies in the approximation properties of the spaces $\left\{\eta\left(\frac{\mathbf{x}-h\mathbf{m}}{\sqrt{\mathcal{D}}h}\right), \mathbf{m} \in \mathbf{Z}^d\right\}$, where the parameter $\mathcal{D} > 0$ is fixed and $h \to 0$. In [10] we proved that if the generating function η satisfies the moment conditions

(1.7)
$$\int_{\mathbf{R}^d} \eta(\mathbf{x}) \, d\mathbf{x} = 1 , \quad \int_{\mathbf{R}^d} \mathbf{x}^{\boldsymbol{\alpha}} \eta(\mathbf{x}) \, d\mathbf{x} = 0, \quad \forall \, \boldsymbol{\alpha} \, , \, 1 \le |\boldsymbol{\alpha}| < N \, ,$$

(here we use standard multiindex notations), then the quasi-interpolant

(1.8)
$$u_h(\mathbf{x}) := \mathcal{D}^{-d/2} \sum_{\mathbf{m} \in \mathbf{Z}^d} u(h\mathbf{m}) \, \eta \left(\frac{\mathbf{x} - h\mathbf{m}}{\sqrt{\mathcal{D}}h} \right) \,,$$

provides the following representation: For any $u \in C^N(\mathbf{R}^d) \cap W^N_{\infty}(\mathbf{R}^d)$ there holds

(1.9)
$$u_h(\mathbf{x}) = u(\mathbf{x}) + R_h(\mathbf{x})$$

where

(1.10)
$$|R_h(\mathbf{x})| \le c_\eta (\sqrt{\mathcal{D}}h)^N ||\nabla_N u||_{L_\infty(\mathbf{R}^d)} + |u(\mathbf{x})| O\Big(\sum_{\boldsymbol{\nu}\in\mathbf{Z}^d\setminus\{\mathbf{0}\}} |\mathcal{F}\eta(\sqrt{\mathcal{D}}\boldsymbol{\nu})|\Big)$$

Due to the second term the quasi-interpolant u_h does not converge to u. However, the rapid decay of $\mathcal{F}\eta$ ensures that one can choose \mathcal{D} such that this saturation error can be made arbitrarily small, for example less than the needed accuracy or the machine precision. Then u_h behaves in numerical computations like a usual approximant of the order $O(h^N)$. This behaviour was studied in the framework of "approximate approximations" in [8],[9] and [10], where the approximation error was estimated in different norms and the construction of suitable generating functions for quasi-interpolation formulas and other simple approximants were described.

Applied to the example (1.6) concerning the Newton potential this approach leads to the following result:

If $u \in W_p^6(\mathbf{R}^3)$, 1 , then the cubature formula

(1.11)
$$\mathcal{N}_{h}u = \frac{h^{2}}{\sqrt{\mathcal{D}}} \sum_{\mathbf{m}\in\mathbf{Z}^{3}} u(h\mathbf{m}) \,\mathcal{N}\eta\left(\frac{\mathbf{x}-h\mathbf{m}}{\sqrt{\mathcal{D}}h}\right)$$

provides the estimate

(1.12)
$$\|\mathcal{N}u - \mathcal{N}_h u\|_{L_q(\mathbf{R}^3)} \le \left(c_1 \left(\sqrt{\mathcal{D}}h\right)^4 + c_2 h^2 e^{-\pi^2 \mathcal{D}}\right) \|u\|_{W_p^6(\mathbf{R}^3)}$$

with positive constants c_1 , c_2 and q = 3p/(3-2p) (for a proof see [11], where also cubature formulas of arbitrary order for different potentials are given). In view of $\exp(-\pi^2) =$ $0.51723...10^{-4}$ we see, that for sufficiently large \mathcal{D} , say $\mathcal{D} \ge 4$, the formula (1.11) behaves in numerical computations like a fourth order cubature. Thus giving up to require the convergence of the approximants (1.5) as $h \to 0$ does not imply any serious restrictions, as far as numerical computations are concerned and η and \mathcal{D} are suitable chosen. But one obtains an essentially greater flexibility in the choice of basic functions resulting in simple multidimensional approximation formulas and cubatures of important classes of integral operators.

It is clear that the basic functions we have in mind do not satisfy a refinement equation of the form (1.4). But it turns out that for a wide class of interesting functions refinement equations are valid in some approximate sense. For example, for $\mathbf{x} \in \mathbf{R}^d$ we have

$$\left| e^{-|\mathbf{x}|^2/\mathcal{D}} - \frac{2^d}{(3\pi\mathcal{D})^{d/2}} \sum_{\mathbf{m}\in\mathbf{Z}^d} e^{-|\mathbf{m}|^2/3\mathcal{D}} e^{-|2\mathbf{x}-\mathbf{m}|^2/\mathcal{D}} \right| \le (2+\epsilon) d \, e^{-3\pi^2\mathcal{D}/4} \, e^{-|\mathbf{x}|^2/\mathcal{D}}$$

with $\epsilon \ll 1$, such that the Gaussian function $\phi_{\mathcal{D}}(\mathbf{x}) := e^{-|\mathbf{x}|^2/\mathcal{D}}$ satisfies a refinement equation within any prescribed tolerance if \mathcal{D} is chosen sufficiently large. This leads to the idea to perform an approximate multiresolution analysis and wavelet construction similar to the case, where one has an exact refinement equation. This is the goal of our paper.

In section 2 we prove that there exists a large class of basic functions satisfying approximate refinement equations and study some approximation properties of the Gaussian radial function $\phi_{\mathcal{D}}$. For this example we provide in section 3 an approximate multiresolution analysis. We show that any element of the L_2 -closure of the linear span

(1.13)
$$\mathbf{V}_n := \left\{ \phi_{\mathcal{D}}(2^n \cdot -\mathbf{m}), \mathbf{m} \in \mathbf{Z}^d \right\}$$

can be approximated by elements of the direct sum

(1.14)
$$\mathbf{V}_n := \mathbf{V}_0 \dotplus \mathbf{W}_0 \dotplus \dots \dotplus \mathbf{W}_{n-1} ,$$

with some small relative error of the form $\varepsilon = (4+\epsilon) dn e^{-3\pi^2 \mathcal{D}/4}$, $\epsilon \ll 1$. Here the wavelet spaces \mathbf{W}_j are almost orthogonal such that the approximate decomposition (1.14) of \mathbf{V}_n can be performed using the orthogonal projections P_0 onto \mathbf{V}_0 and Q_j onto \mathbf{W}_j .

The univariate wavelet construction and some properties of the prewavelet are discussed in section 4. The univariate wavelet spaces W_j are spanned by rapidly decaying analytic prewavelets, plotted in fig. 1, which belong to V_{j+1} and are orthogonal to all elements of V_j .



FIGURE 1. Graph of the normed wavelet $\psi_3(x)$.

It is interesting that there exist simple analytic formulas for a perturbation of the wavelet with a relative error ε^2 . More precisely, the space W_0 can be defined within the assumed tolerance ε as the span of the integer translates of the function

(1.15)
$$\psi_{\mathcal{D}}(x) = e^{-(2x-1)^2/6\mathcal{D}} \cos \frac{5\pi}{6} (2x-1) .$$

Since the values of many integral operators applied to the wavelets can be given analytically one can use approximants from $\tilde{\mathbf{V}}_n$ to derive the cubature of these operators. Here one assumes that the transformation to the basis in $\tilde{\mathbf{V}}_n$ leads to some data compression, at least for sufficiently smooth integrands u. Additionally, the power moments of the prewavelets are very small and can be controlled by the parameter \mathcal{D} . This implies a fast decay of the integrals $K\psi$ if the kernel $k(\mathbf{x}, \mathbf{y})$ satisfies

$$|\partial_{\mathbf{y}}^{\boldsymbol{\alpha}}k(\mathbf{x},\mathbf{y})| \leq c_{\boldsymbol{\alpha}} |\mathbf{x}-\mathbf{y}|^{-(\gamma+|\boldsymbol{\alpha}|)} \quad ext{for some } \gamma > 0 \; .$$

The effect of the nearly vanishing moments can be seen by the example of the Hilbert transform of $\psi_{\mathcal{D}}$

$$\mathcal{H}\psi_\mathcal{D}(x):=rac{1}{\pi}\int\limits_{-\infty}^\inftyrac{\psi_\mathcal{D}(y)}{y-x}\,dy$$

shown in fig. 2. We see, that the essential supports of $\psi_{\mathcal{D}}$ and $\mathcal{H}\psi_{\mathcal{D}}$ are very close, which leads to a high compression rate for the matrix representation of the Hilbert transform in the basis

$$\{\phi_\mathcal{D}(\cdot-k), k\in \mathbf{Z}\} \hspace{0.1in} ext{and} \hspace{0.1in} \{\psi_\mathcal{D}(2^j\cdot-k), k\in \mathbf{Z}, j=0,\ldots,n-1\} \;.$$



FIGURE 2. Graph of the Hilbert transform of the normed wavelet $\psi_3(x)$.

In section 5 we consider multivariate approximate wavelets and the multiresolution structure of the spaces spanned by the Gaussian radial function. We construct a wavelet basis with the property that important pseudodifferential operators admit semi-analytic representations. Further we give explicit formulas for the orthogonal projection P_0 onto \mathbf{V}_0 and almost orthogonal projections \tilde{Q}_j onto decomposition (1.14), \mathbf{W}_j , which are proved in the final section 6, such that for any $\varphi_n \in \mathbf{V}_n$ the estimate

(1.16)
$$\left\|\varphi_n - P_0\varphi_n - \sum_{j=0}^{n-1} \widetilde{Q}_j\varphi_n\right\|_2 \le c \varepsilon \|\varphi_n\|_2$$

holds with some constant not depending on φ_n and \mathcal{D} .

2. Approximate refinement equations

Suppose that $u(\mathbf{x})$, $\mathbf{x} = (x_1, \ldots, x_d) \in \mathbf{R}^d$, is the restriction of an entire function $u(\mathbf{z})$, $\mathbf{z} = (z_1, \ldots, z_d) \in \mathbf{C}^d$, such that for some fixed positive parameters h and \mathcal{D} and any $\mathbf{x} \in \mathbf{R}^d$ the estimate

(2.1)
$$|u(\mathbf{x} + i\pi\mathcal{D}h\mathbf{y}) \exp(-\pi^2\mathcal{D}|\mathbf{y}|^2)| \le A(1+|\mathbf{y}|)^{-d-\delta}, \quad \mathbf{y} \in \mathbf{R}^d,$$

holds. Then for any $\mathbf{m} \in \mathbf{Z}^d$ there exists the number

(2.2)
$$u_{\mathbf{m}} := \int_{\mathbf{R}^d} \exp(-\pi^2 \mathcal{D}|\mathbf{y}|^2) u(h\mathbf{m} + i\pi \mathcal{D}h\mathbf{y}) d\mathbf{y} .$$

In the following we denote by $\langle \mathbf{x}, \mathbf{y} \rangle$ the usual scalar product in \mathbf{R}^d , $|\mathbf{x}| = \langle \mathbf{x}, \mathbf{x} \rangle^{1/2}$, and the Fourier transform is defined as

$$\mathcal{F}\varphi(\lambda) = \int\limits_{\mathbf{R}^d} \varphi(\mathbf{x}) \, e_{\lambda}(-\mathbf{x}) \, d\mathbf{x} \quad ext{with} \quad e_{\lambda}(\mathbf{x}) := e^{2\pi i \langle \mathbf{x}, \lambda
angle} \, .$$

Lemma 2.1. If the entire function u satisfies (2.1) and for given $x \in \mathbb{R}^d$ the semi-discrete convolution

$$u_h(\mathbf{x}) := \sum_{\mathbf{m} \in \mathbf{Z}^d} u_{\mathbf{m}} \exp\left(-\frac{|\mathbf{x} - h\mathbf{m}|^2}{\mathcal{D}h^2}\right) = \sum_{\mathbf{m} \in \mathbf{Z}^d} u_{\mathbf{m}} \phi_{\mathcal{D}}(\mathbf{x}/h - \mathbf{m})$$

converges absolutely then the equality

$$u_h(\mathbf{x}) = \sum_{\boldsymbol{
u} \in \mathbf{Z}^d} u(\mathbf{x} + i\pi \mathcal{D}h \boldsymbol{
u}) \, \exp(-\pi^2 \mathcal{D}|\boldsymbol{
u}|^2) \, e_{\boldsymbol{
u}}(\mathbf{x}/h)$$

holds.

Proof. By repeated application of Cauchy's Theorem one gets

$$\begin{split} u_{\mathbf{m}} \phi_{\mathcal{D}}(\mathbf{x}/h - \mathbf{m}) &= (\pi \mathcal{D}h)^{-n} \phi_{\mathcal{D}}(\mathbf{x}/h - \mathbf{m}) \int_{\mathbf{R}^{d}} u(h\mathbf{m} + i\mathbf{y}) \phi_{\mathcal{D}}(\mathbf{y}/h) \, d\mathbf{y} \\ &= (\pi \mathcal{D}h)^{-n} \phi_{\mathcal{D}}(\mathbf{x}/h) \int_{\mathbf{R}^{d}} u(i\mathbf{y}) \phi_{\mathcal{D}}(\mathbf{y}/h) e_{\mathbf{m}} \left(-\frac{2i}{\mathcal{D}h}(\mathbf{y} + i\mathbf{x})\right) d\mathbf{y} \\ &= (\pi \mathcal{D}h)^{-n} \int_{\mathbf{R}^{d}} u(\mathbf{x} + i\mathbf{y}) \phi_{\mathcal{D}}(\mathbf{y}/h) \, e_{\mathbf{y}} \left(\frac{2i}{\mathcal{D}h}(\mathbf{x}/h - \mathbf{m})\right) d\mathbf{y} \\ &= \int_{\mathbf{R}^{d}} u(\mathbf{x} + i\pi \mathcal{D}h\mathbf{y}) \, \exp(-\pi^{2} \mathcal{D}|\mathbf{y}|^{2}) \, e_{\mathbf{y}}(\mathbf{x}/h) \, e_{\mathbf{m}}(-\mathbf{y}) \, d\mathbf{y} \, . \end{split}$$

Denoting by

$$f_{\mathbf{x}}(\mathbf{y}) := u(\mathbf{x} + i\pi \mathcal{D}h\mathbf{y}) \exp(-\pi^2 \mathcal{D}|\mathbf{y}|^2) e_{\mathbf{y}}(\mathbf{x}/h)$$

we see that

$$u_{\mathbf{m}} \, \phi_{\mathcal{D}}(\mathbf{x}/h - \mathbf{m}) = \mathcal{F} f_{\mathbf{x}}(\mathbf{m})$$
 .

Now we have only to apply Poisson's summation formula (see [14], Th. VII. 2.4),

$$\sum_{\mathbf{m}\in\mathbf{Z}^d}\mathcal{F}f_{\mathbf{x}}(\mathbf{m}) = \sum_{\boldsymbol{\nu}\in\mathbf{Z}^d} f_{\mathbf{x}}(\boldsymbol{\nu}) = \sum_{\boldsymbol{\nu}\in\mathbf{Z}^d} u(\mathbf{x} + i\pi\mathcal{D}h\boldsymbol{\nu}) \, \exp(-\pi^2\mathcal{D}|\boldsymbol{\nu}|^2) \, e_{\boldsymbol{\nu}}(\mathbf{x}/h) \; ,$$

which is valid due to the absolute convergence of both series.

Thus one obtains the relation

(2.3)
$$u_h(\mathbf{x}) - u(\mathbf{x}) = \sum_{\boldsymbol{\nu} \in \mathbf{Z}^d \setminus \{\mathbf{0}\}} u(\mathbf{x} + i\pi \mathcal{D}h\boldsymbol{\nu}) \exp(-\pi^2 \mathcal{D}|\boldsymbol{\nu}|^2) e_{\boldsymbol{\nu}}(\mathbf{x}/h) ,$$

showing that if \mathcal{D} is suitable chosen then u_h is for relative large h a very precise approximant to analytic functions u of first order of growth. In that case u has a compactly supported Fourier transform and we obtain an equivalent formula for the coefficients

$$u_{\mathbf{m}} = (\pi \mathcal{D})^{-d/2} \int\limits_{\mathbf{R}^d} \mathcal{F}u(\boldsymbol{\lambda}) \, \exp(\pi^2 \mathcal{D}h^2 |\boldsymbol{\lambda}|^2) \, e_{\mathbf{m}}(h\boldsymbol{\lambda}) \, d\boldsymbol{\lambda} \; .$$

Let us give some examples. First the case of a polynomial u. It can be easily seen that

$$\int_{-\infty}^{\infty} (hm + i\pi \mathcal{D}hy)^j \exp(-\pi^2 \mathcal{D}y^2) \, dy = \frac{1}{\sqrt{\pi \mathcal{D}}} \Big(\frac{\sqrt{\mathcal{D}}h}{2}\Big)^j H_j\Big(\frac{m}{\sqrt{\mathcal{D}}}\Big) \,,$$

with the Hermite polynomials

$$H_j(y) := (-1)^j e^{y^2} \left(\frac{d}{dy}\right)^j e^{-y^2}.$$

In particular we obtain from Lemma 2.1 the estimate

$$\left|x^{j} - \frac{1}{\sqrt{\pi \mathcal{D}}} \left(\frac{\sqrt{\mathcal{D}}}{2}\right)^{j} \sum_{m=-\infty}^{\infty} H_{j}\left(\frac{m}{\sqrt{\mathcal{D}}}\right) e^{-(x-m)^{2}/\mathcal{D}} \right| \leq \sum_{k \in \mathbf{Z} \setminus \{0\}} |x + i\pi \mathcal{D}k|^{j} e^{-\pi^{2} \mathcal{D}k^{2}}$$

Hence, any polynomial $p(\mathbf{x})$ can be approximated by linear combinations of the functions $\phi_{\mathcal{D}}(\mathbf{x}-\mathbf{m}), \mathbf{m} \in \mathbf{Z}^d$, with an arbitrary relative error $\epsilon > 0$ if \mathcal{D} is chosen sufficiently large. Next we consider the example of the exponential function $u(\mathbf{x}) = e^{\langle \mathbf{x}, \mathbf{a} \rangle}, \mathbf{a} \in \mathbf{C}^d$. Here Lemma 2.1 leads to

$$egin{aligned} u_h(\mathbf{x}) &= (\pi \mathcal{D})^{-d/2} \, e^{-\mathcal{D}h^2 \mathbf{a}^2/4} \sum_{\mathbf{m} \in \mathbf{Z}^d} e^{\langle h\mathbf{m},\mathbf{a}
angle} \, \phi_\mathcal{D}(\mathbf{x}/h-\mathbf{m}) \ &= e^{\langle \mathbf{x},\mathbf{a}
angle} \Big(1 + \sum_{oldsymbol{
u} \in \mathbf{Z}^d \setminus \{\mathbf{0}\}} e^{-\pi^2 \mathcal{D}|oldsymbol{
u}|^2} \, e_{oldsymbol{
u}} \Big(rac{\mathbf{x}}{h} + rac{\mathcal{D}h\mathbf{a}}{2} \Big) \Big) \,. \end{aligned}$$

Thus, if $\mathbf{a} \in \mathbf{R}^d$ then for any h > 0 the series u_h approximates the exponential function with the relative error

$$\sum_{oldsymbol{
u}\in \mathbf{Z}^d\setminus\{\mathbf{0}\}}e^{-\pi^2\mathcal{D}|oldsymbol{
u}|^2}$$
 .

If $\mathbf{a} = \mathbf{u} + i\mathbf{v}$, $\mathbf{u}, \mathbf{v} \in \mathbf{R}^d$ with $\mathbf{v} \neq \mathbf{0}$, and $h \leq \pi/2|\mathbf{v}|$ then u_h approximates $e^{\langle \mathbf{x}, \mathbf{a} \rangle}$ with the relative error less or equal to

$$\sum_{oldsymbol{
u}\in \mathbf{Z}^d\setminus\{\mathbf{0}\}}e^{-\pi^2\mathcal{D}|oldsymbol{
u}|^2/2}$$
 .

If d = 1 then we get some special cases of the well-known transformation formula for Jacobi's elliptic theta functions

(2.4)
$$\sum_{k \in \mathbb{Z}} e^{-k^2 \pi a + 2\pi i k z} = a^{-1/2} \sum_{m \in \mathbb{Z}} e^{-\pi (z-m)^2/a} , \quad z \in \mathbb{C} , \text{ Re } a > 0 ,$$

in particular

(2.5)

$$\cos 2\pi x = \frac{e^{\pi^2 \mathcal{D}/4}}{2\sqrt{\pi \mathcal{D}}} \sum_{m=-\infty}^{\infty} (-1)^m e^{-(2x-m)^2/\mathcal{D}} - R(x)$$

where $R(x) = e^{\pi^2 \mathcal{D}/4} \sum_{k=1}^{\infty} \cos\left(2\pi (2k+1)x\right) e^{-\pi^2 \mathcal{D}(2k+1)^2/4} = O(e^{-2\pi^2 \mathcal{D}})$.

Finally we apply Lemma 2.1 to $\varphi_{\mathcal{D}}(\mathbf{x})$. If h < 1 then we obtain the equality

(2.6)
$$\varphi_{\mathcal{D}}(\mathbf{x}) = (\pi \mathcal{D}(1-h^2))^{-d/2} \sum_{\mathbf{m} \in \mathbf{Z}^d} e^{-h^2 |\mathbf{m}|^2 / \mathcal{D}(1-h^2)} \varphi_{\mathcal{D}}(\mathbf{x}/h-\mathbf{m}) -\varphi_{\mathcal{D}}(\mathbf{x}) \sum_{\boldsymbol{\nu} \in \mathbf{Z}^d \setminus \{\mathbf{0}\}} e_{\boldsymbol{\nu}/h}((1-h^2)\mathbf{x}) e^{-\pi^2 \mathcal{D}(1-h^2)|\boldsymbol{\nu}|^2}.$$

Therefore the Gaussian function satisfies the approximate refinement equation

(2.7)
$$\varphi_{\mathcal{D}}(\mathbf{x}) \doteq (\pi \mathcal{D}(1-h^2))^{-d/2} \sum_{\mathbf{m} \in \mathbf{Z}^d} e^{-h^2 |\mathbf{m}|^2 / \mathcal{D}(1-h^2)} \varphi_{\mathcal{D}}(\mathbf{x}/h-\mathbf{m}) ,$$

with the accuracy $\varphi_{\mathcal{D}}(\mathbf{x}) \sum_{\boldsymbol{\nu} \in \mathbf{Z}^d \setminus \{\mathbf{0}\}} e^{-\pi^2 \mathcal{D}(1-h^2)|\boldsymbol{\nu}|^2}.$

Approximate refinement equations of the from (2.7) are valid for a large class of basic functions η as shown by the following assertion.

Lemma 2.2. Let η belong to the Schwartz space, $\eta \in S(\mathbb{R}^d)$, with positive Fourier transform $\mathcal{F}\eta > 0$ and such that for given h < 1 the function $g \in S(\mathbb{R}^d)$, which is defined via its Fourier transform by $\mathcal{F}g(\lambda) := \mathcal{F}\eta(\lambda)/\mathcal{F}\eta(h\lambda)$. Then for any $\varepsilon > 0$ there exists $\mathcal{D} > 0$ such that for all $\mathbf{x} \in \mathbb{R}^n$

(2.8)
$$\left|\eta\left(\frac{\mathbf{x}}{\sqrt{\mathcal{D}}}\right) - \mathcal{D}^{-d/2} \sum_{\mathbf{m} \in \mathbf{Z}^d} g\left(\frac{h\mathbf{m}}{\sqrt{\mathcal{D}}}\right) \eta\left(\frac{\mathbf{x} - h\mathbf{m}}{\sqrt{\mathcal{D}}h}\right)\right| < \varepsilon .$$

Proof. Again we may apply the Poisson summation formula to

$$\mathcal{D}^{-d/2} \sum_{\mathbf{m} \in \mathbf{Z}^d} g\left(\frac{h\mathbf{m}}{\sqrt{\mathcal{D}}}\right) \eta\left(\frac{\mathbf{x} - h\mathbf{m}}{\sqrt{\mathcal{D}}h}\right) = \sum_{\boldsymbol{\nu} \in \mathbf{Z}^d} \int_{\mathbf{R}^d} g(h\mathbf{y}) \eta\left(\frac{\mathbf{x}}{\sqrt{\mathcal{D}}h} - \mathbf{y}\right) e_{\boldsymbol{\nu}}(\sqrt{\mathcal{D}}\mathbf{y}) d\mathbf{y}$$
$$= \sum_{\boldsymbol{\nu} \in \mathbf{Z}^d} \int_{\mathbf{R}^d} \mathcal{F}g(\boldsymbol{\lambda}) \int_{\mathbf{R}^d} \eta(\mathbf{y}) e_{\sqrt{\mathcal{D}}\boldsymbol{\nu} - h\boldsymbol{\lambda}}\left(\frac{\mathbf{x}}{\sqrt{\mathcal{D}}}\right) d\mathbf{y} d\boldsymbol{\lambda}$$
$$= \sum_{\boldsymbol{\nu} \in \mathbf{Z}^d} e_{\boldsymbol{\nu}}\left(\frac{x}{h}\right) \int_{\mathbf{R}^d} \frac{\mathcal{F}\eta(\boldsymbol{\lambda})}{\mathcal{F}\eta(h\boldsymbol{\lambda})} \mathcal{F}\eta(h\boldsymbol{\lambda} + \sqrt{\mathcal{D}}\boldsymbol{\nu}) e_{\boldsymbol{\lambda}}\left(\frac{\mathbf{x}}{\sqrt{\mathcal{D}}}\right) d\boldsymbol{\lambda}$$
$$= \eta\left(\frac{\mathbf{x}}{\sqrt{\mathcal{D}}}\right) + \sum_{\boldsymbol{\nu} \in \mathbf{Z}^d \setminus \{\mathbf{0}\}} e_{\boldsymbol{\nu}}\left(\frac{x}{h}\right) \int_{\mathbf{R}^d} \frac{\mathcal{F}\eta(\boldsymbol{\lambda})}{\mathcal{F}\eta(h\boldsymbol{\lambda})} \mathcal{F}\eta(h\boldsymbol{\lambda} + \sqrt{\mathcal{D}}\boldsymbol{\nu}) e_{\boldsymbol{\lambda}}\left(\frac{\mathbf{x}}{\sqrt{\mathcal{D}}}\right) d\boldsymbol{\lambda}$$

Now it is easy to see that the function of $\mathbf{y} \in \mathbf{R}^d$

$$\int\limits_{\mathbf{R}^d} rac{\mathcal{F}\eta(oldsymbol{\lambda})}{\mathcal{F}\eta(holdsymbol{\lambda})} \, \mathcal{F}\eta(holdsymbol{\lambda}+\mathbf{y}) \, doldsymbol{\lambda} \, \in \, \mathcal{S}(\mathbf{R}^d) \, ,$$

hence the sum

$$\sum_{\boldsymbol{\nu}\in\mathbf{Z}^d\setminus\{\mathbf{0}\}} \Big| \int\limits_{\mathbf{R}^d} \frac{\mathcal{F}\eta(\boldsymbol{\lambda})}{\mathcal{F}\eta(h\boldsymbol{\lambda})} \, \mathcal{F}\eta(h\boldsymbol{\lambda}+\sqrt{\mathcal{D}}\boldsymbol{\nu}) \, e_{\boldsymbol{\lambda}}\Big(\frac{\mathbf{x}}{\sqrt{\mathcal{D}}}\Big) d\boldsymbol{\lambda} \Big|$$

can be made arbitrarily small by choosing \mathcal{D} large enough.

We note a special case of basic functions giving high order semi-analytic quasi-interpolation formulas of the form (1.8). Using the generalized Laguerre polynomials $L_j^{(\alpha)}$ we define the basic functions (see [8], [9])

$$\eta_{2M}(\mathbf{x}) := \pi^{-d/2} L_{M-1}^{(d/2)}(|\mathbf{x}|^2) e^{-|\mathbf{x}|^2}, \quad M = 1, 2, \dots,$$

having the Fourier transform

$$\mathcal{F}\eta_{2M}(oldsymbol{\lambda}) = e^{-\pi^2|oldsymbol{\lambda}|^2}\sum_{j=0}^{M-1}rac{(\pi^2|oldsymbol{\lambda}|^2)^j}{j!}$$

and thus satisfying the moment conditions (1.7) with N = 2M. Note that the basic function η , considered in (1.6), corresponds to the case N = 4 and d = 3. For the functions η_{2M} one can estimate the error bound of Lemma 2.2 by

$$arepsilon \leq \sum_{oldsymbol{
u}\in \mathbf{Z}^d\setminus\{\mathbf{0}\}} q(\sqrt{\mathcal{D}}oldsymbol{
u}) \, e^{-\pi^2 \mathcal{D}(1-h^2)|oldsymbol{
u}|^2}$$

with some polynomial q of degree 2M.

_

It is clear that the mask values $g(hm/\sqrt{D})$ in (2.8) can be computed by simple onedimensional quadratures if η is a the radial function. Some of the above mentioned basic functions allow to derive analytic formulas for the function g. For example, for the function η considered in (1.6) one has

$$g(\mathbf{x}) = \frac{e^{-|\mathbf{x}|^2/(1-h^2)}}{h^2\sqrt{\pi \mathcal{D}(1-h^2)^3}} -\frac{1-h^2}{h^4} \frac{e^{(1-h^2)/h^2}}{2\pi |\mathbf{x}|\sqrt{\mathcal{D}^3}} \left(e^{-2|\mathbf{x}|/h} \operatorname{erfc}\left(\frac{1-h|\mathbf{x}|-h^2}{h\sqrt{1-h^2}}\right) - e^{2|\mathbf{x}|/h} \operatorname{erfc}\left(\frac{1+h|\mathbf{x}|-h^2}{h\sqrt{1-h^2}}\right)\right),$$

where erfc denotes the complementary error function.

We remark that the knowledge of g is very useful also for another interesting application of the approximate refinement equation, which allows to derive quasi-interpolation formulas for nonuniformly distributed mesh points.

3. Approximate multiresolution

In this section we provide for the example of the Gaussian function $\phi_{\mathcal{D}}$ an approximate multiresolution analysis. We introduce the closed linear subspaces of $L_2(\mathbf{R}^d)$

$$\mathbf{V}_j := \left\{ \sum_{\mathbf{m} \in \mathbf{Z}^d} a_{\mathbf{m}} \phi_{\mathcal{D}}(2^j \cdot -\mathbf{m}), \ \{a_{\mathbf{m}}\} \in \ell_2(\mathbf{Z}^d) \right\}.$$

Since

$$\begin{split} \Big\| \sum_{\mathbf{m}\in\mathbf{Z}^d} a_{\mathbf{m}} \,\phi_{\mathcal{D}}(\cdot-\mathbf{m}) \Big\|_2^2 &= (\pi\mathcal{D})^d \int_{\mathbf{R}^d} e^{-2\pi^2\mathcal{D}|\boldsymbol{\lambda}|^2} \,\Big| \sum_{\mathbf{m}\in\mathbf{Z}^d} a_{\mathbf{m}} \,e_{\mathbf{m}}(\boldsymbol{\lambda}) \Big|^2 d\boldsymbol{\lambda} \\ &= (\pi\mathcal{D})^d \int_{[0,1]^d} \sum_{\mathbf{k}\in\mathbf{Z}^d} e^{-2\pi^2\mathcal{D}|\boldsymbol{\lambda}-\mathbf{k}|^2} \Big| \sum_{\mathbf{m}\in\mathbf{Z}^d} a_{\mathbf{m}} \,e_{\mathbf{m}}(\boldsymbol{\lambda}) \Big|^2 d\boldsymbol{\lambda} , \end{split}$$

the set $\{\phi_{\mathcal{D}}((\cdot - \mathbf{m})\}_{\mathbf{m}\in\mathbf{Z}^d}$ is an L_2 -stable basis in \mathbf{V}_0 . From the approximate refinement equation (2.6) for the Gaussian function it is clear that for any l < j the space \mathbf{V}_l is almost included in \mathbf{V}_j . In particular, if h = 1/2 then we have

$$\phi_{\mathcal{D}}(\mathbf{x}) = \frac{2^d}{(3\pi\mathcal{D})^{d/2}} \sum_{\mathbf{m}\in\mathbf{Z}^d} e^{-|\mathbf{m}|^2/3\mathcal{D}} \phi_{\mathcal{D}}(2\mathbf{x}-\mathbf{m}) - \phi_{\mathcal{D}}(\mathbf{x}) \sum_{\boldsymbol{\nu}\in\mathbf{Z}^d\setminus\{\mathbf{0}\}} e^{-3\pi^2\mathcal{D}|\boldsymbol{\nu}|^2/4} e^{3\pi i \langle \boldsymbol{x}, \boldsymbol{\nu} \rangle}$$

hence for any $\varphi_j \in \mathbf{V}_j$ the small perturbation

(3.1)
$$\varphi_j(\mathbf{x}) \left(1 + \sum_{\boldsymbol{\nu} \in \mathbf{Z}^d \setminus \{\mathbf{0}\}} e^{-3\pi^2 \mathcal{D}|\boldsymbol{\nu}|^2/4} e^{3\pi i \langle \mathbf{x}, \boldsymbol{\nu} \rangle} \right) \in \mathbf{V}_{j+1}$$

Consequently, for any $\varphi_j \in \mathbf{V}_j$ we can find $\varphi_{j+1} \in \mathbf{V}_{j+1}$ such that

$$\|\varphi_j - \varphi_{j+1}\|_2 \le \varepsilon \, \|\varphi_j\|_2$$

with

$$arepsilon = arepsilon_{\mathcal{D}} := \sum_{oldsymbol{
u} \in \mathbf{Z}^d \setminus \{\mathbf{0}\}} e^{-3\pi^2 \mathcal{D} |oldsymbol{
u}|^2/4} = (2+\epsilon) \, d \, e^{-3\pi^2 \mathcal{D}/4} \quad ext{with } \epsilon \ll 1 \; .$$

Furthermore, if we introduce the closed subspace $\mathbf{W}_j \subset \mathbf{V}_{j+1}$ of all functions which are orthogonal to \mathbf{V}_j , then it can be easily seen that

$$|(\varphi_j, \varphi_l)_2| \le \left((1+\varepsilon)^{|j-l|-1} - 1 \right) ||\varphi_j||_2 ||\varphi_l||_2$$

for $\varphi_j \in \mathbf{W}_j$. Thus the situation is very similar to the case when exact refinement equations are valid, which was mentioned in the introduction.

Let us fix some integer n > 0, which determines the grid for the approximating functions. In the following we show that any element of \mathbf{V}_n can be represented within some prescribed tolerance as an element of the multiresolution structure

$$\mathbf{V}_n := \mathbf{V}_0 + \mathbf{W}_0 + \ldots + \mathbf{W}_{n-1}$$

To this end introduce the orthogonal projections $P_j : L_2(\mathbf{R}^d) \to \mathbf{V}_j, \ j = 0, \ldots, n$, and $Q_j : L_2(\mathbf{R}^d) \to \mathbf{W}_j, \ j = 0, \ldots, n-1$, and denote $Q_{-1} = P_0$.

Theorem 3.1. Any $\varphi_n \in \mathbf{V}_n$ can be approximatively represented as an element of the multiresolution structure (3.3) and there holds

$$\left\|\varphi_n - \sum_{j=-1}^{n-1} Q_j \varphi_n\right\|_2 \le n \frac{2\varepsilon - \varepsilon^2}{1 - \varepsilon} \|\varphi_n\|_2.$$

Proof. Use the telescopic series

$$\varphi_n = P_n \varphi_n = \sum_{j=1}^n (P_j - P_{j-1})\varphi_n + P_0 \varphi_n$$
$$= \sum_{j=1}^n (P_j - P_{j-1})\varphi_n + Q_{-1}\varphi_n = \sum_{j=-1}^{n-1} Q_j \varphi_n + \sum_{j=0}^{n-1} (P_{j+1} - P_j - Q_j)\varphi_n$$

 and

Lemma 3.1. For j = 0, ..., n - 1 we have

$$||P_j + Q_j - P_{j+1}||_2 \le \frac{2\varepsilon - \varepsilon^2}{1 - \varepsilon}.$$

Proof. Note that (3.2) implies the inequality

(3.4)
$$\|\varphi_j - P_{j+1}\varphi_j\|_2 \le \varepsilon \, \|\varphi_j\|_2 \,, \quad \forall \, \varphi_j \in \mathbf{V}_j \,.$$

Since $P_{j+1}(\mathbf{V}_j) = \mathbf{V}_{j+1} \ominus \mathbf{W}_j$ any $\varphi_{j+1} \in \mathbf{V}_{j+1}$ can be written in the form

$$\varphi_{j+1} = P_{j+1}\varphi_j + Q_j\varphi_{j+1}$$

with some $\varphi_j \in \mathbf{V}_j$. From (3.4) we derive therefore

$$\|\varphi_{j+1} - (\varphi_j + Q_j \varphi_{j+1})\|_2 = \|P_{j+1}\varphi_j - \varphi_j\|_2 \le \varepsilon \|\varphi_j\|_2$$

and

$$\|\varphi_j\|_2 \leq \frac{1}{1-\varepsilon} \|P_{j+1}\varphi_j\|_2 = \frac{1}{1-\varepsilon} \|\varphi_{j+1} - Q_j\varphi_{j+1}\|_2 \leq \frac{\|\varphi_{j+1}\|_2}{1-\varepsilon} .$$

Now we use that the sum $P_j + Q_j$ is the orthogonal projection onto $\mathbf{V}_j \oplus \mathbf{W}_j$. Hence for any $u \in L_2(\mathbf{R}^d)$ we obtain the estimate

$$\|(I - (P_j + Q_j))P_{j+1}u\|_2 = \inf_{v \in \mathbf{V}_j \oplus \mathbf{W}_j} \|P_{j+1}u - v\|_2 \le \frac{\varepsilon}{1 - \varepsilon} \|u\|_2,$$

leading together with (3.4) to

$$\|(P_j + Q_j - P_{j+1})u\|_2 \le \|(I - P_{j+1})(P_j + Q_j)u\|_2 + \|P_{j+1}(I - P_j - Q_j)u\|_2$$

$$\le \inf_{v \in \mathbf{V}_{j+1}} \|(P_j + Q_j)u - v\|_2 + \|(I - P_j - Q_j)P_{j+1}\|_2 \|u\|_2 \le (\varepsilon + \frac{\varepsilon}{1 - \varepsilon})\|u\|_2.$$

4. Approximate univariate wavelets

Now we apply some well-known constructions from wavelet theory to the univariate case. Here $\phi_{\mathcal{D}}(x) = e^{-x^2/\mathcal{D}}$, we denote the corresponding scaled and wavelet spaces by V_j and W_j , respectively. Since the function

$$\sum_{n \in \mathbf{Z}} (-1)^{m-1} \mu_{m-1} \phi_{\mathcal{D}}(2 \cdot -m) \quad \text{with} \quad \mu_m = \int_{\mathbf{R}} \phi_{\mathcal{D}}(x) \phi_{\mathcal{D}}(2x+m) \, dx$$

is orthogonal to all integer shifts of the scaling function $\phi_{\mathcal{D}}$ we obtain a first element of the wavelet space

$$\sum_{m \in \mathbf{Z}} (-1)^{m-1} e^{-(m-1)^2/5\mathcal{D}} \phi_{\mathcal{D}}(2x-m) \in W_0 .$$

Using (2.5) we derive the formula

$$\begin{split} \sum_{n \in \mathbf{Z}} (-1)^m e^{-m^2/5\mathcal{D}} \,\phi_{\mathcal{D}}(2x-1-m) \\ &= e^{-(2x-1)^2/6\mathcal{D}} \sum_{m \in \mathbf{Z}} (-1)^m \exp\left(-\frac{6}{5\mathcal{D}} \left(\frac{5}{6}(2x-1)-m\right)^2\right) \\ &= \sqrt{\frac{10\pi\mathcal{D}}{3}} \, e^{-5\pi^2\mathcal{D}/24} \, e^{-(2x-1)^2/6\mathcal{D}} \left(\cos\frac{5\pi}{6}(2x-1)+R_{\mathcal{D}}(x)\right) \,, \end{split}$$

with

$$R_{\mathcal{D}}(x) = \sum_{k=1}^{\infty} \cos \frac{5\pi}{6} (2k+1)(2x-1) e^{-5\pi^2 \mathcal{D}(k^2+k)/6} = O(e^{-5\pi^2 \mathcal{D}/3})$$

So we introduce the univariate prewavelet

(4.1)
$$\psi_{\mathcal{D}}(x) := \sqrt{\frac{3}{10\pi\mathcal{D}}} e^{5\pi^2\mathcal{D}/24} \sum_{m \in \mathbf{Z}} (-1)^m e^{-m^2/5\mathcal{D}} \phi_{\mathcal{D}}(2x-1-m)$$

and its perturbation, the approximate prewavelet

(4.2)
$$\tilde{\psi}_{\mathcal{D}}(x) := e^{-(2x-1)^2/6\mathcal{D}} \cos \frac{5\pi}{6} (2x-1) ,$$

differing by

$$|\psi_{\mathcal{D}}(x) - \tilde{\psi}_{\mathcal{D}}(x)| \le (1+\epsilon) e^{-(2x-1)^2/6\mathcal{D}} e^{-5\pi^2 \mathcal{D}/3}, \quad \epsilon \ll 1,$$

such that for any $\varphi = \sum a_m \psi_{\mathcal{D}}(x-m)$ the estimate

(4.3)
$$\left\|\varphi - \sum a_m \tilde{\psi}_{\mathcal{D}}(x-m)\right\|_2 \leq \frac{\varepsilon_{\mathcal{D}}^2}{2d} \|\varphi\|_2$$

holds. Consequently, in numerical computations with the precision $\varepsilon_{\mathcal{D}}$ one can use both formulas of the prewavelet with the same rights.

In order to define multivariate wavelet bases we formulate the following assertions which can be easily checked by using the Fourier transform of the prewavelet

(4.4)
$$\mathcal{F}\psi_{\mathcal{D}}(\lambda) = \sqrt{\frac{3\pi\mathcal{D}}{8}} e^{5\pi^2\mathcal{D}/24} e^{-\pi i\lambda} e^{-\pi^2\mathcal{D}\lambda^2/4} \sigma_{5\mathcal{D}}\left(\frac{\lambda+1}{2}\right),$$

where σ_{α} denotes the positive and 1-periodic function

$$\sigma_{\alpha}(\lambda) = \frac{1}{\sqrt{\alpha\pi}} \sum_{m \in \mathbf{Z}} e^{-m^2/\alpha} e^{2\pi i m \lambda} = \sum_{j \in \mathbf{Z}} e^{-\alpha \pi^2 (\lambda+j)^2} .$$

Lemma 4.1.

- 1. The half-shifts of the prewavelet $\{\psi_{\mathcal{D}}(\cdot m/2)\}_{m \in \mathbb{Z}}$ are an L_2 -stable basis in the scaled space V_1 .
- 2. The integer shifts of the prewavelet $\psi_{\mathcal{D}}$ and of the function

(4.5)
$$\tilde{\phi}_{\mathcal{D}}(x) := \frac{2}{\sqrt{3\pi\mathcal{D}}} \sum_{m \in \mathbf{Z}} e^{-m^2/3\mathcal{D}} \phi_{\mathcal{D}}(2x-m) \in V_1$$

form an L_2 -stable basis in V_1 .

Besides the fast decay of $\psi_{\mathcal{D}}$ we are interested in the moments of the prewavelet. Since this function is orthogonal to the integer shifts of the Gaussian $\phi_{\mathcal{D}}$, which approximate polynomials very accurate as seen in section 2, one can expect that even higher moments are very small and decrease if \mathcal{D} increases. Using the Fourier transform (4.4) we have computed the first eight moments of $\psi_{\mathcal{D}}$, which are contained in the following table.

moment	$\mathcal{D}=2$	$\mathcal{D}=3$	$\mathcal{D}=4$	$\mathcal{D}=5$
0	$5.103\cdot10^{-9}$	$2.143 \cdot 10^{-13}$	$8.482 \cdot 10^{-18}$	$3.251 \cdot 10^{-22}$
1	$-2.551 \cdot 10^{-9}$	$-1.071 \cdot 10^{-13}$	$-4.241 \cdot 10^{-18}$	$-1.626 \cdot 10^{-22}$
2	$-3.059 \cdot 10^{-7}$	$-2.920 \cdot 10^{-11}$	$-2.065 \cdot 10^{-15}$	$-1.240 \cdot 10^{-19}$
3	$4.601 \cdot 10^{-7}$	$4.386 \cdot 10^{-11}$	$3.100 \cdot 10^{-15}$	$1.862 \cdot 10^{-19}$
4	$1.616 \cdot 10^{-6}$	$3.685 \cdot 10^{-9}$	$4.759 \cdot 10^{-13}$	$4.533 \cdot 10^{-17}$
5	$-4.116 \cdot 10^{-6}$	$-9.287 \cdot 10^{-9}$	$-1.195 \cdot 10^{-12}$	$-1.134 \cdot 10^{-16}$
6	$-7.304 \cdot 10^{-4}$	$-4.263 \cdot 10^{-7}$	$-1.032 \cdot 10^{-10}$	$-1.582 \cdot 10^{-14}$
7	$2.701\cdot 10^{-3}$	$1.525\cdot 10^{-6}$	$3.655 \cdot 10^{-10}$	$5.577 \cdot 10^{-14}$
8	$2.768 \cdot 10^{-2}$	$4.460 \cdot 10^{-5}$	$2.096 \cdot 10^{-8}$	$5.255 \cdot 10^{-12}$

5. Approximate multivariate wavelet decomposition

Now we are in the position to discuss the approximate wavelet decomposition of \mathbf{V}_n in the multivariate case. First we introduce an L_2 -stable basis in the wavelet space \mathbf{W}_0 . Consider the function

(5.1)
$$\Psi_{\mathcal{D}}(\mathbf{x}) = \left(\frac{3}{10\pi\mathcal{D}}\right)^{d/2} e^{5\pi^2 d\mathcal{D}/24} \sum_{\mathbf{m}\in\mathbf{Z}^d} (-1)^{|\mathbf{m}|} e^{-|\mathbf{m}|^2/5\mathcal{D}} e^{-|2\mathbf{x}-\mathbf{m}|^2/\mathcal{D}},$$

which represents the tensor product of the univariate functions $\psi_{\mathcal{D}}(x_j-1/2), j=1,\ldots,d$, and its perturbation

(5.2)
$$\widetilde{\Psi}_{\mathcal{D}}(\mathbf{x}) = \prod_{j=1}^{d} \widetilde{\psi}_{\mathcal{D}}(x_j - 1/2) = \phi_{3\mathcal{D}}(\mathbf{x}) \prod_{j=1}^{d} \cos \frac{5\pi}{3} x_j$$

¿From (4.3) it is clear, that within the precision $\varepsilon_{\mathcal{D}}$ these two functions can be identified. By Lemma 4.1, the scaled space \mathbf{V}_1 is spanned by the half-shifts $\{\Psi_{\mathcal{D}}(\cdot - \mathbf{m}/2)\}_{\mathbf{m}\in\mathbf{Z}^d}$ and consequently the set

$$ig \{ \Psi_\mathcal{D}(\cdot - \mathbf{v} - \mathbf{m}): \; \mathbf{m} \in \mathbf{Z}^d, \; \mathbf{v} \in \mathcal{V}' ig \}$$

is an L_2 -stable basis in \mathbf{W}_0 . Here again \mathcal{V} denotes the set of vertices of the cube $[0, 1/2]^d$ and $\mathcal{V}' = \mathcal{V} \setminus \{\mathbf{0}\}$. Thus we obtain the L_2 -stable basis

$$\left\{\phi_{\mathcal{D}}(\cdot-\mathbf{m}): \mathbf{m} \in \mathbf{Z}^{d}\right\}, \left\{\Psi_{\mathcal{D}}(2^{j}\cdot-\mathbf{v}-\mathbf{m}): \mathbf{m} \in \mathbf{Z}^{d}, \mathbf{v} \in \mathcal{V}', j=0,\ldots,n-1\right\}$$

in $\widetilde{\mathbf{V}}_n$, which is by Theorem 3.1 almost the space \mathbf{V}_n .

The elements of this basis have the property that the application of important integral operators can be given analytically. Besides series representations obtained from (5.1) efficient expressions can be derived by using the perturbed basis function (5.2). For example,

$$\mathcal{N}\widetilde{\Psi}_{\mathcal{D}}(\mathbf{x}) = \frac{1}{4\pi} \int_{\mathbf{R}^3} \frac{e^{-|\mathbf{y}|^2/3\mathcal{D}}}{|\mathbf{x} - \mathbf{y}|} \prod_{j=1}^3 \cos\frac{5\pi}{3} y_j \, d\mathbf{y} = \frac{1}{8} \sum_{|\mu_j|=1} \frac{1}{4\pi} \int_{\mathbf{R}^3} \frac{e^{-|\mathbf{y}|^2/3\mathcal{D}}}{|\mathbf{x} - \mathbf{y}|} \, e_{\mathbf{y}}\left(\frac{5\mu}{6}\right) \, d\mathbf{y}$$

with $\mu = (\mu_1, \mu_2, \mu_3) \in \mathbf{R}^3$. Using the Fourier transforms one obtains

$$\begin{split} &\frac{1}{4\pi} \int\limits_{\mathbf{R}^3} \frac{e^{-|\mathbf{y}|^2/3\mathcal{D}}}{|\mathbf{x}-\mathbf{y}|} \, e_{\mathbf{y}} \left(\frac{5\mu}{6}\right) d\mathbf{y} = (3\pi\mathcal{D})^{3/2} \int\limits_{\mathbf{R}^3} \frac{e^{-3\pi^2\mathcal{D}|\boldsymbol{\lambda}-5\boldsymbol{\mu}/6|^2}}{4\pi^2|\boldsymbol{\lambda}|^2} \, e_{\boldsymbol{\lambda}}(\mathbf{x}) \, d\boldsymbol{\lambda} \\ &= 3\mathcal{D}\pi^{3/2} \, e^{-25\pi^2\mathcal{D}/4} \int\limits_{\mathbf{R}^3} \frac{e^{-\pi^2|\boldsymbol{\lambda}|^2}}{4\pi^2|\boldsymbol{\lambda}|^2} \, e_{\boldsymbol{\lambda}/\sqrt{3\mathcal{D}}} \left(\mathbf{x} - \frac{5\pi\mathcal{D}i}{2}\boldsymbol{\mu}\right) d\boldsymbol{\lambda} \, . \end{split}$$

Since the integral

$$\pi^{3/2}\int\limits_{\mathbf{R}^3}rac{e^{-\pi^2|oldsymbol{\lambda}|^2}}{4\pi^2|oldsymbol{\lambda}|^2}e_{oldsymbol{\lambda}}(\mathbf{x})\,doldsymbol{\lambda}=rac{1}{2|\mathbf{x}|}\int\limits_{0}^{|\mathbf{x}|}e^{-t^2}\,dt\;,\quad\mathbf{x}\in\mathbf{R}^3\;,$$

(see [11]) is analytic we derive

$$\frac{1}{4\pi} \int_{\mathbf{R}^3} \frac{e^{-|\mathbf{y}|^2/3\mathcal{D}}}{|\mathbf{x} - \mathbf{y}|} \, e_{\mathbf{y}}\left(\frac{5\mu}{6}\right) \, d\mathbf{y} = \frac{3\mathcal{D} \, e^{-25\pi^2 \mathcal{D}/4}}{2} \int_{0}^{1} e^{-zt^2} \, dt \; ,$$

where

$$z = rac{1}{3\mathcal{D}}\sum_{j=1}^3 \left(x_j - rac{5\pi\mathcal{D}i}{2}\mu_j
ight)^2 = rac{|\mathbf{x}|^2}{3\mathcal{D}} - rac{25\pi^2\mathcal{D}}{4} - rac{5\pi i}{3}\langle\mathbf{x},oldsymbol{\mu}
angle \,.$$

Thus we get the one-dimensional integral

$$\mathcal{N}\widetilde{\Psi}_{\mathcal{D}}(\mathbf{x}) = \frac{3\mathcal{D}}{2} \int_{0}^{1} \exp\left(-\frac{|\mathbf{x}|^{2}t^{2}}{3\mathcal{D}} + \frac{25\pi^{2}\mathcal{D}(t^{2}-1)}{4}\right) \prod_{j=1}^{3} \cos\frac{5\pi}{3} y_{j}t^{2} dt ,$$

which can be expressed also by the error function of complex arguments.

Now we consider the problem to find the approximate wavelet decomposition of a given element belonging to V_n and to prove estimate (1.16). Following Theorem 3.1 one has to determine the orthogonal projections onto V_0 and W_j .

Since $\phi_{\mathcal{D}}(\mathbf{x}) = \prod \phi_{\mathcal{D}}(x_j)$ the orthogonal projections P_0 onto \mathbf{V}_0 can be given as the tensor product

$$(5.3) P_0 = R_0 \otimes \cdots \otimes R_0$$

of the univariate L_2 -projections R_0 onto V_0 . This mapping is described in the following lemma, which will be proved in the final section.

Lemma 5.1. The orthogonal projection R_0 onto V_0 has the form

(5.4)
$$R_0 f = \sum_{k \in \mathbf{Z}} (f, \check{\phi}_{\mathcal{D}}(\cdot - k))_2 \phi_{\mathcal{D}}(\cdot - k) ,$$

where the function $\check{\phi}_{\mathcal{D}} \in V_0$ is given by the formula

$$\check{\phi}_{\mathcal{D}}(x) = \sum_{k \in \mathbf{Z}} a_k(\mathcal{D}) \, \phi_{\mathcal{D}}(x-k)$$

with the coefficients

$$a_k(\mathcal{D}) = (-1)^k c(\mathcal{D}) e^{k^2/2\mathcal{D}} \sum_{r=|k|}^{\infty} (-1)^r e^{-(r+1/2)^2/2\mathcal{D}}$$

and the constant

$$c(\mathcal{D}) = \frac{1}{2\pi^2 \mathcal{D}^2} \Big(\sum_{j \in \mathbf{Z}} (-1)^j j \, e^{-\pi^2 \mathcal{D}(2j+1)^2/2} \Big)^{-1} \, .$$

Let us turn to the orthogonal projections Q_j onto \mathbf{W}_j . Since these spaces are derived by scaling from \mathbf{W}_0 it suffices to consider the operator Q_0 . We will construct a mapping \widetilde{Q}_0 onto \mathbf{W}_0 such that $\|\widetilde{Q}_0 - Q_0\|_2 \leq c \varepsilon_{\mathcal{D}}$ with some constant c depending only on the dimension d. To this end we consider the basis in \mathbf{W}_0 obtained by the simple tensor product approach for the construction of multivariate prewavelets (see [12]). Denoting

(5.5)
$$w_0(x) := \tilde{\phi}_{\mathcal{D}}(x) , \quad w_{1/2}(x) := \psi_{\mathcal{D}}(x)$$

collection (see Lemma 4.1), we introduce the collection of functions

$$\Phi_{\mathbf{v}}(\mathbf{x}) = w_{v_1}(x_1) \cdots w_{v_d}(x_d) \in \mathbf{W_0} \;, \quad \mathbf{v} \in \mathcal{V}' \;.$$

Since the function w_0 is the right-hand side of the approximate refinement equation (2.7) for $\phi_{\mathcal{D}}$ it follows from Lemma 4.1 that the set $\{\Phi_{\mathbf{v}}(\cdot - \mathbf{m})\}_{\mathbf{m}\in\mathbf{Z}^d,\mathbf{v}\in\mathcal{V}'}$ is an L_2 -stable basis in \mathbf{W}_0 . Moreover, if in the definition of $\Phi_{\mathbf{v}}$ the function w_0 is replaced by the Gaussian function $\phi_{\mathcal{D}}(x)$, then the corresponding principal shift invariant spaces

$$X_{\mathbf{v}} := \{ \Phi_{\mathbf{v}}(\cdot - \mathbf{m}), \ \mathbf{m} \in \mathbf{Z}^d \}$$

are orthogonal, $X_{\mathbf{v}} \perp X_{\mathbf{v}'}$ for $\mathbf{v} \neq \mathbf{v}'$. Thus the orthogonal projection onto $\bigoplus_{\mathbf{v} \in \mathcal{V}'} X_{\mathbf{v}}$ is

(5.6)
$$\overline{Q}_0 := \sum_{\mathbf{v} \in \mathcal{V}'} R_{v_1} \otimes \cdots \otimes R_{v_d}$$

with the univariate projections $R_0: L_2(\mathbf{R}) \to V_0$ and $R_{1/2}: L_2(\mathbf{R}) \to W_0$. It is evident that

(5.7)
$$\|\overline{Q}_0 - Q_0\| \le c \varepsilon_{\mathcal{D}} ,$$

where c depends on d only. To derive a mapping into \mathbf{W}_0 we introduce the small perturbation of the operator \overline{Q}_0

(5.8)
$$\widetilde{Q}_0 := \sum_{\mathbf{v}\in\mathcal{V}'} \widetilde{R}_{v_1}\otimes\cdots\otimes\widetilde{R}_{v_d},$$

where we define (see (5.4) and (5.5))

$$\widetilde{R}_0 f = \sum_{k \in \mathbf{Z}} (f, \check{\phi}_{\mathcal{D}}(\cdot - k))_2 w_0(\cdot - k) ,$$

and the mapping $\widetilde{R}_{1/2}$ is described in the following lemma, which will be proved in the next section.

Lemma 5.2. The orthogonal projection $R_{1/2}$ onto W_0 can be approximatively represented in the form

$$\widetilde{R}_{1/2}f = \sum_{k \in \mathbf{Z}} (f, \check{\psi}_{\mathcal{D}}(\cdot - k))_2 \ \psi_{\mathcal{D}}(\cdot - k) \ ,$$

where the function $\check{\psi}_{\mathcal{D}}$ is given as the sum

$$\check{\psi}_{\mathcal{D}}(x):=\sum_{m\in\mathbf{Z}}a_k\,\psi_{\mathcal{D}}(x-k)\in W_0\;,$$

with the coefficients

$$a_{k} = e^{k^{2}/3\mathcal{D}} \left(c_{0} \sum_{j=|k|}^{\infty} (-1)^{j} e^{-3(j+1/2)^{2}/\mathcal{D}} \right)$$

(5.9)

$$+ (-1)^k c_1 \sum_{j=|k|}^{\infty} \left(e^{-3(j+1/4)^2/\mathcal{D}} - e^{-3(j+3/4)^2/\mathcal{D}} \right)
ight)$$

and the numbers c_0 and c_1 equal to

$$c_{0} = \frac{4}{3\pi^{2}\mathcal{D}^{2}} \left(\sum_{j \in \mathbf{Z}} (-1)^{j} (6j+1) e^{-\pi^{2}\mathcal{D}(6j+1)^{2}/12} \right)^{-1},$$

$$c_{1} = \frac{2}{3\pi^{2}\mathcal{D}^{2}} \left(\sum_{j \in \mathbf{Z}} (-1)^{j} (6j+1) e^{-\pi^{2}\mathcal{D}(6j+1)^{2}/3} \right)^{-1}.$$

There exists a constant c such that

(5.10)
$$||R_{1/2}f - \widetilde{R}_{1/2}f||_2 \le c e^{-\pi^2 \mathcal{D}} ||f||_2 , \forall f \in L_2(\mathbf{R}).$$

; From Lemma 5.2, the refinement equation (2.7) and (5.7) it is clear that we have

(5.11)
$$\|\widetilde{Q}_0 - Q_0\| \le c \varepsilon_{\mathcal{D}} ,$$

with some constant c depending only on d. Thus we obtain the following approximate wavelet decomposition of the space V_n .

Theorem 5.1. There exists a constant c depending on the space dimension d and on n such that for any $\varphi_n \in \mathbf{V}_n$ the estimate

$$\|\varphi_n - \sum_{j=-1}^{n-1} \widetilde{Q}_j \varphi_n \varphi_n\|_2 \le c \varepsilon_{\mathcal{D}} \|\varphi_n\|_2$$

holds, where $\tilde{Q}_{-1} = P_0$ is defined in (5.3) and the mappings \tilde{Q}_j onto \mathbf{W}_j are obtained by scaling from \tilde{Q}_0 given in (5.8).

6. The orthogonal projections R_0 and $R_{1/2}$

The construction of the orthogonal projections onto V_0 and W_0 uses some well-known facts about principal shift invariant spaces (see [3]), which we recall briefly. Denote by $S(\eta)$ the L_2 -closure of finite linear combinations of the shifts $\eta(\cdot - m)$, $m \in \mathbb{Z}$, of a generating function η and suppose that the shifts form an L_2 -stable basis of $S(\eta)$. Note that for any

$$\phi(x) = \sum_{m \in \mathbf{Z}} a_m \, \eta(x - m) \in S(\eta)$$

we have

$$\mathcal{F}\phi(\lambda) = \tau(\lambda) \mathcal{F}\eta(\boldsymbol{\lambda}) \quad \text{with} \ \tau(\lambda) := \sum_{m \in \mathbf{Z}} a_m e_{-m}(\lambda) \in L_2(0,1) \ .$$

Furthermore if $\mathcal{F}\phi = \tau \mathcal{F}\eta \in L_2(\mathbf{R})$ with a 1-periodic function τ , then

$$\|\phi\|_{L_2(\mathbf{R})} = \|\tau[\mathcal{F}\eta, \mathcal{F}\eta]^{1/2}\|_{L_2(0,1)}$$

where the bracket product stands for the 1-periodic function

$$[f,g] := \sum_{k \in \mathbb{Z}} f(\cdot - k) \overline{g(\cdot - k)} .$$

For each $f \in L_2(\mathbf{R})$ the L_2 -projection $Pf \in S(\eta)$ is given by

$$\mathcal{F}(Pf) = \frac{[\mathcal{F}f, \mathcal{F}\eta]}{[\mathcal{F}\eta, \mathcal{F}\eta]} \, \mathcal{F}\eta$$

Hence the shifts of the function $\check{\eta} \in S(\eta)$ defined via

(6.1)
$$\mathcal{F}\check{\eta} = \frac{\mathcal{F}\eta}{[\mathcal{F}\eta, \mathcal{F}\eta]}$$

form the corresponding biorthogonal basis, i.e. $(\eta(\cdot - k), \check{\eta})_2 = \delta_{0k}, k \in \mathbb{Z}$.

Applied to R_0 we have to determine the Fourier coefficients of the inverse function to

$$[\mathcal{F}\phi_{\mathcal{D}}, \mathcal{F}\phi_{\mathcal{D}}] = \pi \mathcal{D} \sum_{m \in \mathbf{Z}} e^{-\pi^2 \mathcal{D}(\lambda - m)^2} = \sqrt{\frac{\pi \mathcal{D}}{2}} \sum_{k \in \mathbf{Z}} e^{-k^2/2\mathcal{D}} e^{2\pi i k x} ,$$

which is closely connected with the theta function.

Lemma 6.1. Let a > 0. Then

$$\left(\sum_{k\in\mathbf{Z}}e^{-ak^2}e^{2\pi ikx}\right)^{-1}=\sum_{k\in\mathbf{Z}}a_k\,e^{2\pi ikx}\;,$$

where

$$a_k = (-1)^k c e^{ak^2} \sum_{r=|k|}^{\infty} (-1)^r e^{-a(r+1/2)^2}$$
,

and

$$c^{-1} = \sum_{r=0}^{\infty} (-1)^r (2r+1) e^{-a(r+1/2)^2}$$

Setting $a^{-1} = 2\mathcal{D}$ we obtain immediately the assertion of Lemma 5.1.

Note that the cardinal interpolant $\eta^* \in S(\eta)$ which satisfies $\eta^*(k) = \delta_{0k}$, $k \in \mathbb{Z}$, has the Fourier transform

$$\mathcal{F}\eta^*:=rac{\mathcal{F}\eta}{[\mathcal{F}\eta,1]} \ .$$

Hence Lemma 6.1 gives an explicit formula for the interpolant $Q_h f$ composed of the scaled shifts $\phi_{\mathcal{D}}(\mathbf{x}/h - \mathbf{k})$ with $Q_h f(h\mathbf{m}) = f(h\mathbf{m})$ for all $\mathbf{m} \in \mathbf{Z}^d$,

$$Q_h f(\mathbf{x}) = \sum_{\mathbf{m} \in \mathbf{Z}^d} f(h\mathbf{m}) q_{\mathcal{D}}(\mathbf{x}/h - \mathbf{m}) ,$$

where

$$q_\mathcal{D}(\mathbf{x}) = \sum_{\mathbf{k} \in \mathbf{Z}^d} A_\mathbf{k} \, \phi_\mathcal{D}(\mathbf{x} - \mathbf{k}) \quad ext{with} \quad A_\mathbf{k} = a_{k_1} \cdots a_{k_d}$$

and

$$a_{k} = (-1)^{k} c \sum_{r=0}^{\infty} (-1)^{r} e^{-(r+1/2)^{2}/\mathcal{D}} e^{-|k|(2r+1)/\mathcal{D}} ,$$
$$c^{-1} = \sum_{r=0}^{\infty} (-1)^{r} (2r+1) e^{-(r+1/2)^{2}/\mathcal{D}} .$$

It was proved in [10] that this interpolant approximates with exponential rate up to a saturation error

$$|f(\mathbf{x}) - Q_h f(\mathbf{x})| \leq \left((2+4d) \, e^{-a/2h} + 4 \, d \, e^{-\pi^2 \mathcal{D}}
ight) \int\limits_{\mathbf{R}^d} |\mathcal{F}f(oldsymbol{\lambda})| \, \exp(a|oldsymbol{\lambda}|) \, doldsymbol{\lambda} \; ,$$

if the last integral is finite.

Now we come to the proof of Lemma 5.2. We will construct an almost biorthogonal basis to $\{\psi(\cdot - m)\}_{m \in \mathbb{Z}}$ in the space $W_0 = S(\psi_D)$. Following (6.1) one has to find the inverse function of

(6.2)
$$\sum_{m \in \mathbf{Z}} |\mathcal{F}\psi_{\mathcal{D}}(\lambda + m)|^2$$

Using several times the transformation formula (2.4) it can be shown that this function is approximated with a relative error less than $3e^{-5\pi^2 D/2}$ by the function

$$g(\lambda) := \frac{3\pi\mathcal{D}}{4} \sum_{k \in \mathbf{Z}} \left(e^{-3\pi^2\mathcal{D}(k+\lambda+1/6)^2} + e^{-3\pi^2\mathcal{D}(k-\lambda+1/6)^2} \right) \,.$$

Hence is suffices to determine the Fourier coefficients of the 1-periodic function g^{-1} . Note that g(z) is analytic and quasi doubly-periodic with

(6.3)
$$g(z+1) = g(z), \quad g\left(z + \frac{i}{\pi D}\right) = -e^{3/D}e^{6\pi i z}g(z),$$

therefore in any rectangle $[z, z + 1) \times [z, z + i/\pi D)$ there are three zeros of g(z). It can be easily seen that in $[-1/4, 3/4) \times [-1/4, -1/4 + i/\pi D)$ the zeros are at $z_0 = (0, i/2\pi D)$ and near the points $(1/2, i/4\pi D)$ and $(1/2, 3i/4\pi D)$. More precisely, we have

$$g(1/2+iy) = \frac{3\pi\mathcal{D}}{2}e^{3\pi^2\mathcal{D}y^2} \sum_{k=0}^{\infty} \left(\cos 2\pi^2\mathcal{D}y(3k+1)e^{-\pi^2\mathcal{D}(3k+1)^2/3} + \cos 2\pi^2\mathcal{D}y(3k+2)e^{-\pi^2\mathcal{D}(3k+2)^2/3}\right),$$

such that g(1/2 + iy) = 0 at the points

(6.4)
$$z_1 = \frac{1}{2} + \frac{i}{4\pi \mathcal{D}} - i\epsilon$$
 and $z_2 = \frac{1}{2} + \frac{3i}{4\pi \mathcal{D}} + i\epsilon$ with $0 < \epsilon < e^{-\pi^2 \mathcal{D}}$.

To determine the Fourier coefficients we apply (6.3) and the Residue Theorem to the integral

$$a_k = \int_{-1/4}^{3/4} \frac{e^{-2\pi i k x}}{g(x)} \, dx$$

and obtain the recurrence relation

(6.5)
$$a_k = -e^{(2k-3)/\mathcal{D}}a_{k-3} + (r_0 + r_1 + r_2)$$

where

$$r_j = 2\pi i \operatorname{Res}_{z=z_j} rac{e^{-2\pi i k z}}{g(z)}$$

It is easy to see that

(6.6)
$$r_0 = c_0 e^{k/\mathcal{D}} e^{-3/4\mathcal{D}} , \ c_0 = \frac{4}{3\pi^2 \mathcal{D}^2} \Big(\sum_{j \in \mathbf{Z}} (-1)^j (6j+1) e^{-\pi^2 \mathcal{D}(6j+1)^2/12} \Big)^{-1}$$

Further, the residues r_1 and r_2 are

$$\begin{aligned} r_1 &= -(-1)^k c_1 e^{k/2\mathcal{D}} e^{-3/16\mathcal{D}} e^{-\pi\epsilon(2k-3/2)} e^{-3\pi^2\mathcal{D}\epsilon^2} ,\\ r_2 &= (-1)^k c_1 e^{3k/2\mathcal{D}} e^{-27/16\mathcal{D}} e^{\pi\epsilon(2k-3/2)} e^{-3\pi^2\mathcal{D}\epsilon^2} , \end{aligned}$$

with the constant

(6.7)
$$c_1 = \frac{2}{3\pi^2 \mathcal{D}^2} \left(\sum_{j \in \mathbf{Z}} (3j+1) \sin \frac{\pi}{2} (1 - 4\pi \mathcal{D}\epsilon) (3j+1) e^{-\pi^2 \mathcal{D}(3j+1)^2/3} \right)^{-1}$$

It turns out, that (6.5) has a unique solution $\{a_k\} \in \ell_2(\mathbb{Z})$ which is given by

(6.8)
$$a_{k} = e^{k^{2}/3\mathcal{D}} \left(c_{0} \sum_{j=|k|}^{\infty} (-1)^{j} e^{-3(j+1/2)^{2}/\mathcal{D}} + (-1)^{k} c_{1} \sum_{j=|k|}^{\infty} \left(e^{3\pi\epsilon/2} e^{-3(j+1/4-\pi\mathcal{D}\epsilon)^{2}/\mathcal{D}} - e^{-3\pi\epsilon/2} e^{-3(j+3/4+\pi\mathcal{D}\epsilon)^{2}/\mathcal{D}} \right) \right).$$

So we have derived the Fourier coefficients of g^{-1} , i.e. up to a relative error less than $3 e^{-5\pi^2 \mathcal{D}/2}$ the coefficients of the biorthogonal basis function to the wavelet basis $\psi_{\mathcal{D}}(\cdot -k)$. Furthermore, since $0 < \epsilon < e^{-4\pi^2 \mathcal{D}/3}$ (see (6.4)) we may set in (6.7) and (6.8) the small number $\epsilon = 0$ and obtain together with (6.6) formula (5.9) of Lemma 5.2. In doing so we make an error for the Fourier coefficients, which is less than $e^{-\pi^2 \mathcal{D}} |a_k|$. In view of the rapid decay of the wavelet the estimate (5.10) follows immediately, which completes the proof of Lemma 5.2.

References

- Alpert, B., Beylkin, G., Coifman, R., Rokhlin, V., Waveletlike bases for the fast solution of secondkind integral equations, SIAM J. Sci. Statist. Comp. 14 (1993), 159–184.
- [2] Beylkin, G., Wavelets and fast numerical algorithms, Proc. Symp. Appl. Math. 47 (1993), 89-117.
- [3] de Boor, C., DeVore, R. A., Ron, A. Approximation from Shift-invariant Subspaces of L₂(R^d), Trans. AMS 341 (1994), 787-806.
- [4] Chui, C. K., Stöcker, J., Ward, J. D., Compactly supported box spline wavelets, Proc. Amer. Math. Soc. 113 (1992), 785–793.
- [5] Dahmen, W., Prössdorf, S., Schneider, R., Wavelet approximation methods for pseudodifferential equations II, Adv. Comp. Math. 1 (1993), 259–335.
- [6] Daubechies, I., Ten lectures on wavelets, CBMS-NSF Regional Conf. Series in Appl. Math. 61, 1992.
- [7] Maz'ya, V., A New Approximation Method and its Applications to the Calculation of Volume Potentials. Boundary Point Method, in: 3. DFG-Kolloqium des DFG-Forschungsschwerpunktes "Randelementmethoden", 1991.
- [8] Maz'ya, V., Approximate Approximations, in: The Mathematics of Finite Elements and Applications. Highlights 1993, J. R. Whiteman (ed.), 77-104, Wiley & Sons, Chichester 1994.
- [9] Maz'ya, V., Schmidt, G., On Approximate Approximations, Preprint LiTH-MAT-R-94-12, Linköping University 1994.
- [10] V. Maz'ya, G. Schmidt, On Approximate Approximations using Gaussian Kernels, IMA J. Num. Anal. 16 (1996), 13-29.
- [11] Maz'ya, V., Schmidt, G., "Approximate Approximations" and the cubature of potentials, Rend. Mat. Acc. Lincei, s. 9, 6 (1995), 161-184.
- [12] Meyer, Y., Wavelets and operators, University Press, Cambridge 1992.
- [13] von Petersdorf, T., Schneider, R., Schwab, C., Multi wavelet approximation for the double layer potential equation, Preprint TH Darmstadt, 1995.
- [14] E. M. Stein, G. Weiss, Introduction to Fourier Analysis on Euclidian Spaces, University Press, Princeton 1971.

Recent publications of the Weierstraß–Institut für Angewandte Analysis und Stochastik

Preprints 1996

- 220. Boris N. Khoromskij, Siegfried Prößdorf: Fast computations with the harmonic Poincaré-Steklov operators on nested refined meshes.
- 221. Anton Bovier, Véronique Gayrard: Distribution of overlap profiles in the one-dimensional Kac-Hopfield model.
- 222. Jürgen Sprekels, Dan Tiba: A duality-type method for the design of beams.
- 223. Wolfgang Dahmen, Bernd Kleemann, Siegfried Prößdorf, Reinhold Schneider: Multiscale methods for the solution of the Helmholtz and Laplace equation.
- 224. Herbert Gajewski, Annegret Glitzky, Jens Griepentrog, Rolf Hünlich, Hans-Christoph Kaiser, Joachim Rehberg, Holger Stephan, Wilfried Röpke, Hans Wenzel: Modellierung und Simulation von Bauelementen der Nano- und Optoelektronik.
- **225.** Andreas Rathsfeld: A wavelet algorithm for the boundary element solution of a geodetic boundary value problem.
- **226.** Sergej Rjasanow, Wolfgang Wagner: Numerical study of a stochastic weighted particle method for a model kinetic equation.
- **227.** Alexander A. Gushchin: On an information-type inequality for the Hellinger process.
- **228.** Dietmar Horn: Entwicklung einer Schnittstelle für einen DAE-Solver in der chemischen Verfahrenstechnik.
- **229.** Oleg V. Lepski, Vladimir G. Spokoiny: Optimal pointwise adaptive methods in nonparametric estimation.
- 230. Bernd Kleemann, Andreas Rathsfeld, Reinhold Schneider: Multiscale methods for boundary integral equations and their application to boundary value problems in scattering theory and geodesy.
- 231. Jürgen Borchardt, Ludger Bruell, Friedrich Grund, Dietmar Horn, Frank Hubbuch, Tino Michael, Horst Sandmann, Robert Zeller: Numerische Lösung großer strukturierter DAE-Systeme der chemischen Prozeßsimulation.
- **232.** Herbert Gajewski, Klaus Zacharias: Global behaviour of a reaction-diffusion system modelling chemotaxis.

- 233. Fréderic Guyard, Reiner Lauterbach: Forced symmetry breaking perturbations for periodic solutions.
- **234.** Vladimir G. Spokoiny: Adaptive and spatially adaptive testing of a nonparametric hypothesis.
- 235. Georg Hebermehl, Rainer Schlundt, Horst Zscheile, Wolfgang Heinrich: Simulation of monolithic microwave integrated circuits.
- 236. Georg Hebermehl, Rainer Schlundt, Horst Zscheile, Wolfgang Heinrich: Improved numerical solutions for the simulation of monolithic microwave integrated circuits.
- 237. Pavel Krejčí, Jürgen Sprekels: Global solutions to a coupled parabolichyperbolic system with hysteresis in 1-d magnetoelasticity.
- 238. Georg Hebermehl, Friedrich-Karl Hübner: Portabilität und Adaption von Software der linearen Algebra für Distributed Memory Systeme.
- **239.** Michael H. Neumann: Multivariate wavelet thresholding: a remedy against the curse of dimensionality?
- 240. Anton Bovier, Miloš Zahradník: The low-temperature phase of Kac-Ising models.
- 241. Klaus Zacharias: A special system of reaction equations.
- 242. Susumu Okada, Siegfried Prößdorf: On the solution of the generalized airfoil equation.
- **243.** Alexey K. Lopatin: Oscillations and dynamical systems: Normalization procedures and averaging.
- **244.** Grigori N. Milstein: Stability index for invariant manifolds of stochastic systems.
- 245. Luis Barreira, Yakov Pesin, Jörg Schmeling: Dimension of hyperbolic measures – A proof of the Eckmann–Ruelle conjecture.
- 246. Leonid M. Fridman, Rainer J. Rumpel: On the asymptotic analysis of singularly perturbed systems with sliding mode.
- 247. Björn Sandstede: Instability of localised buckling modes in a one-dimensional strut model.
- 248. Björn Sandstede, Christopher K.R.T. Jones, James C. Alexander: Existence and stability of N-pulses on optical fibers with phase-sensitive amplifiers.