

## **On forward and inverse uncertainty quantification for models involving hysteresis operators**

Olaf Klein<sup>1</sup>, Daniele Davino<sup>2</sup>, Ciro Visone<sup>2</sup>

submitted: December 18, 2018

<sup>1</sup> Weierstrass Institute  
Mohrenstr. 39  
10117 Berlin  
Germany  
E-Mail: [olaf.klein@wias-berlin.de](mailto:olaf.klein@wias-berlin.de)

<sup>2</sup> Università degli Studi del Sannio  
P.zza Roma  
21 - 82100  
Benevento  
Italy  
E-Mail: [davino@unisannio.it](mailto:davino@unisannio.it)  
[visone@unisannio.it](mailto:visone@unisannio.it)

No. 2561  
Berlin 2018



Edited by  
Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS)  
Leibniz-Institut im Forschungsverbund Berlin e. V.  
Mohrenstraße 39  
10117 Berlin  
Germany

Fax: +49 30 20372-303  
E-Mail: [preprint@wias-berlin.de](mailto:preprint@wias-berlin.de)  
World Wide Web: <http://www.wias-berlin.de/>

# On forward and inverse uncertainty quantification for models involving hysteresis operators

Olaf Klein, Daniele Davino, Ciro Visone

## Abstract

Parameters within hysteresis operators modeling real world objects have to be identified from measurements and are therefore subject to corresponding errors. To investigate the influence of these errors, the methods of *Uncertainty Quantification (UQ)* are applied.

## 1 Uncertainties in models with hysteresis operators and uncertainty quantification

Considering, e.g., magnetization, piezo-electric effects, elasto-plastic behavior, or magnetostrictive materials, one has to take into account hysteresis effects. Many models involve therefore *hysteresis operators*. The parameter in the models are identified using results from *measurements*, sometimes performed only for some *sample specimens* but also used also for other specimens.

The parameter in the hysteresis operators are therefore also subject to uncertainties. We apply the methods of *Uncertainty Quantification (UQ)*, see, e.g., [11, 12], to deal with theses uncertainties, i.e., we describe them by introducing appropriate random variables modeling the corresponding information/assumptions/beliefs and use probability theory to describe and determine the influence of the uncertainties.

In this paper, we present results of elementary *Forward Uncertainty Quantification*. During forward UQ, one starts from the random variable representing the parameter value and considers the resulting model output as random variable. This allows to compute properties like expected value, variation, probabilities for outputs entering some interval, credible intervals, and other *Quantities of Interest (QoI)*. Moreover, we will also present a brief example of *Inverse UQ*, i.e., of using (further) data and measurements, to determine (reduce/adapt) the uncertainty of the parameter, i.e., to determine a (new) random variable taking into account the (new) information, and use the (new) random variable to represent the parameter afterwards.

## 2 The play operator with deterministic data

Considering a final time  $T > 0$ , some *yield limit*  $r \geq 0$  and some *initial state*  $z_0$ , the *play operator*  $\mathcal{P}_r[z_0, \cdot]$  maps  $u \in C[0, T]$  being piecewise monotone to  $\mathcal{P}_r[z_0, u] \in C[0, T]$  being piecewise

monotone and it holds, see, e.g., [3, 8, 9, 13],

$$\mathcal{P}_r[z_0, u](0) = \max(u(0) - r, \min(u(0) + r, z_0)), \quad (2.1a)$$

$$\mathcal{P}_r[z_0, u](t) = \begin{cases} \max(\mathcal{P}_r[z_0, u](t_0), u(t) - r), & \text{if } u \text{ is} \\ & \text{increasing on } [t_0, t], \\ \min(\mathcal{P}_r[z_0, u](t_0), u(t) + r), & \text{if } u \text{ is} \\ & \text{decreasing on } [t_0, t], \end{cases} \quad (2.1b)$$

for all  $t_0, t \in [0, T]$  with  $t_0 < t$  such that  $u$  is **monotone** on  $[t_0, t]$ . An example for an input function and the output of the play operator is shown in Figure 1.




Figure 1: Left-hand side: evolution of input function  $u$ , of  $u+2$ , of  $u-2$  and of  $\mathcal{P}_2[0, u]$ ; right-hand side: input-output diagram, showing evolution of  $(u(t), \mathcal{P}_2[0, u](t))$ , sizes of marking circles decreases with time

In Figure 2 the outputs of the play for one input function and several yield limits are shown. Therein, we observe that even if  $\mathcal{P}_r[0, u]$  and  $\mathcal{P}_{r'}[0, u]$  with  $r \neq r'$  coincide on some time interval, they may not coincide in the future.




Figure 2: One input function and the output functions  $\mathcal{P}_r[0, u]$  for the play-operator with yield limits  $r \in \{1, 1.5, 2, 2.5, 3\}$ . Top: evolution; bottom, left-hand side: input-output diagram for  $r \in \{1, 2, 2.5\}$ ; bottom, right-hand side: input-output diagram for  $r \in \{1.5, 3\}$ .

In the next section, we need to deal with the evolutions of  $\mathcal{P}_r[r, u](\cdot)$  for all  $r \geq 0$  simultaneously. The method in use is inspired by consideration for the Preisach operator and the representation of the Preisach operator by using the play operator and a *memory sequence*.

Adapting the construction of the memory sequence as in [9, Text from (II.2.11) up to Prop. II.2.5] to a play operator with an initial configuration being equal to some constant value  $w \in \mathbb{R}$  on  $[0, +\infty[$ , we deal with piecewise monotone input functions  $u \in C[0, T]$ , and  $t \in [0, T]$ , and consider the following construction:

■ Let

$$r_{w,u,t,0} := \max\{|u(\tau) - w| : \tau \in [0, t]\}, \quad (2.2a)$$

$$s_{w,u,t,0} := \max\{\tau \in [0, t] : |u(\tau) - w| = r_{w,u,t,0}(t)\}, \quad (2.2b)$$

$$z_{w,u,t,0} := u(s_{w,u,t,0}). \quad (2.2c)$$

■ If  $r_{w,u,t,0} > 0$  then it follows that  $z_{w,u,t,0} \neq w$ . We define

$$p_{w,u,t} := \begin{cases} 1, & \text{if } z_{w,u,t,0} < w, \\ -1, & \text{if } z_{w,u,t,0} > w. \end{cases} \quad (2.3)$$

For  $k = 1, 2, \dots$ , we recursively define

$$z_{w,u,t,k} := \begin{cases} \max\{u(\tau) : \tau \in [s_{w,u,t,k-1}, t]\}, & \text{if } k \text{ is odd and } p_{w,u,t} > 0, \\ \max\{u(\tau) : \tau \in [s_{w,u,t,k-1}, t]\}, & \text{if } k \text{ is even and } p_{w,u,t}(t) < 0, \\ \min\{u(\tau) : \tau \in [s_{w,u,t,k-1}, t]\}, & \text{otherwise,} \end{cases} \quad (2.4)$$

$$r_{w,u,t,k} := \frac{1}{2} |z_{w,u,t,k} - z_{w,u,t,k-1}|, \quad (2.5)$$

$$s_{w,u,t,k} := \max\{\tau \in [s_{w,u,t,k-1}, t] : u(\tau) = z_{w,u,t,k}\}, \quad (2.6)$$

until it holds that  $s_{w,u,t,k}(t) = t$ . Then we define  $K_{w,u,t} := k$  and  $r_{w,u,t,K_{w,u,t}+1} = 0$ . We have  $s_{w,u,t,k} = t$  for some finite index  $k$  since  $u$  restricted to  $[0, t]$  has different local extrema at  $s_{w,u,t,0}, s_{w,u,t,1}, \dots$ , and the number of them is limited since  $u$  is piecewise monotone. Moreover, we see that the construction yields that

$$0 = r_{w,u,t,K_{w,u,t}+1} < r_{w,u,t,K_{w,u,t}} < \dots < r_{w,u,t,1} < r_{w,u,t,0} < \infty. \quad (2.7)$$

**Lemma 1.** *In the considered situation, it holds:*

a)  $\mathcal{P}_r[w, u](t) = w$  for all  $r \geq r_{w,u,t,0}$ .

b) If  $r_{w,u,t,0} > 0$ , we have

$$\mathcal{P}_r[w, u](t) = z_{w,u,t,k} + (-1)^k p_{w,u,t} r, \quad \forall r \in [r_{w,u,t,k+1}, r_{w,u,t,k}], \quad k \in \{0, \dots, K_{w,u,t}\}. \quad (2.8)$$

*Proof.* a) Let  $r \geq r_{w,u,t,0}$  be given. Recalling (2.2a), we deduce that  $u(\tau) - r \leq w \leq u(\tau) + r$  for all  $\tau \in [0, t]$ . In view of (2.1a), we conclude that  $\mathcal{P}_r[w, u](0) = w$ . Since  $u$  is piecewise monotone, we can perform an induction argument over the monotonicity intervals and observe by using (2.1b) that  $\mathcal{P}_r[w, u](\tau) = w$  for all  $\tau \in [0, t]$ .

- b) Assume that  $r_{w,u,t,0} > 0$ . Let  $\lambda : [0, \infty) \rightarrow \mathbb{R}$  be defined by  $\lambda(r) := 0$  for all  $r \geq 0$  if  $w = 0$ ; otherwise let

$$\lambda(r) := \begin{cases} w, & \text{if } r \leq r_{w,u,t,0}, \\ w \left(1 - \frac{1}{|w|} (r - r_{w,u,t,0})\right), & \text{if } r_{w,u,t,0} \leq r \leq r_{w,u,t,0} + |w|, \\ 0, & \text{if } r \geq r_{w,u,t,0} + |w|. \end{cases} \quad (2.9)$$

For both definitions it holds that  $\lambda \in W^{1,\infty}(0, \infty)$ , that  $\left|\frac{d\lambda(r)}{dr}\right| \leq 1$  for all  $r \geq 0$  and that  $\lambda(s) = 0$  for all  $s \geq r_{w,u,t,0} + |w|$ . This yield that  $\lambda$  belongs to the subset  $\Lambda_0$  of *memory configurations* as defined in [9, II.(2.5)].

Following [9, II.(2.11)], we define  $m_\lambda : \mathbb{R} \rightarrow [0, +\infty[$  by

$$m_\lambda(v) := \min\{r \geq 0 : |v - \lambda(r)| = r\}, \quad \forall v \in \mathbb{R}. \quad (2.10)$$

Then it follows that  $m_\lambda(v) = |v - w|$  for all  $v \in \mathbb{R}$  with  $|v - w| \leq r_{w,u,t,0}$ . Recalling (2.2a), we see that the equation holds for  $v = u(\tau)$  for all  $\tau \in [0, t]$ . Considering  $\bar{r}$  and  $\bar{t}$  as in [9, II.(2.14)], we deduce by recalling [9, II.(2.12)], (2.2a), (2.2b), and (2.3) that

$$\begin{aligned} \bar{r} &:= \max\{m_\lambda(u(\tau)) : \tau \in [0, t]\} \\ &= \max\{|u(\tau) - w| : \tau \in [0, t]\} = r_{w,u,t,0}, \end{aligned} \quad (2.11)$$

$$\begin{aligned} \bar{t} &:= \max\{\tau \in [0, t] : m_\lambda(u(\tau)) = \bar{r}\} \\ &= \max\{\tau \in [0, t] : |u(\tau) - w| = r_{w,u,t,0}(t)\} = s_{w,u,t,0}, \end{aligned} \quad (2.12)$$

$$\begin{aligned} u(\bar{t}) - \lambda(\bar{r}) &= u(s_{w,u,t,0}) - w = -p_{w,u,t} |u(s_{w,u,t,0}) - w| \\ &= -p_{w,u,t} r_{w,u,t,0} = -p_{w,u,t} \bar{r}. \end{aligned} \quad (2.13)$$

- If  $p_{w,u,t} = 1$  then it follows from (2.13) that  $u(\bar{t}) = \lambda(\bar{r}) - \bar{r}$ . We define  $r_k := r_{w,u,t,k}$  for all  $k \in \{0, \dots, K_{w,u,t} + 1\}$  and  $t_k := s_{w,u,t,k}$  for all  $k \in \{0, \dots, K_{w,u,t}\}$ . Defining moreover  $t_{K_{w,u,t}+1} = t$ ,  $r_k := 0$  and  $t_k := t$  for all  $k \in \mathbb{N}$  with  $k > K_{w,u,t} + 1$  and recalling the definition of a *memory sequence* as in [9, text before Prop. II.2.5], it follows that  $\{(t_j, r_j)\}_{j=0}^\infty$  is the memory sequence of  $u$  at the point  $t$  with respect to the initial configuration  $\lambda$ . Moreover, it holds that  $(-1)^j = p_{w,u,t}(-1)^j$  for all  $j = 0, 1, 2, \dots$ .
- If  $p_{w,u,t} \neq 1$  then it follows from (2.3) and (2.13) that  $u(\bar{t}) = \lambda(\bar{r}) + \bar{r}$ . We define  $r_k := r_{w,u,t,k-1}$  for all  $k \in \{1, \dots, K_{w,u,t} + 2\}$  and  $t_k := s_{w,u,t,k-1}$  for all  $k \in \{1, \dots, K_{w,u,t} + 1\}$ . Defining moreover  $t_{K_{w,u,t}+2} = t$ ,  $r_k := 0$  and  $t_k := t$  for all  $k \in \mathbb{N}$  with  $k > K_{w,u,t} + 2$  and recalling the definition of a *memory sequence* as in [9, Text before Prop. II.2.5] it follows that  $\{(t_j, r_j)\}_{j=1}^\infty$  is the memory sequence of  $u$  at the point  $t$  with respect to the initial configuration  $\lambda$ . Moreover, it holds that  $(-1)^j = p_{w,u,t}(-1)^{j-1}$  for all  $j = 1, 2, \dots$ .

Recalling now [9, II.(2.17)] in Prop. II.2.5] and considering the value of  $p_{w,u,t}$ , we see that the assertion is proved. □

### 3 The play operator with stochastic yield limit

Now, we want to consider a situation, wherein the true value of the yield limit is not known exactly, but we think that its value is near to 2. Hence, we interpret the yield limit  $r \geq 0$  as value of the random variable  $R$  generated from the normal distribution  $N(2, 0.5^2)$  with mean 2 and standard deviation 0.5 by ignoring  $(-\infty, 0]$  and rescaling, leading to the following *probability density function*  $\rho_R$  of  $R$ , see also Figure 3:

$$\rho_R(r) = \begin{cases} \frac{1}{C} e^{-\frac{(r-2)^2}{2 \cdot 0.5^2}}, & \text{if } r \geq 0 \\ 0, & \text{if } r < 0, \end{cases} \quad \text{with } C = \int_0^\infty e^{-\frac{(r-2)^2}{2 \cdot 0.5^2}} dr. \quad (3.1)$$




Figure 3: Probability density  $\rho_R$  and expected value  $\mathbb{E}(R) \approx 2.00007$  of  $R$  defined in (3.1)

The mapping  $[0, \infty) \ni r \mapsto \mathcal{P}_r[w, u](t)$  is continuous, see e.g., [9, Coro. II.2.2]. Hence, it follows that the composition of this mapping with  $R$  generates a random variable, denoted by  $\mathcal{P}_R[w, u](t)$ . The collection  $(\mathcal{P}_R[w, u](t))_{t \in [0, T]}$  is a stochastic process.

For an initial state  $w \in \mathbb{R}$ , an input function  $u$  and a time  $t \in [0, T]$ , we denote by  $\mu_{w, u, t}$  the probability measure on  $\mathbb{R}$  for  $\mathcal{P}_R[w, u](t)$  and observe that for the expected value  $\mathbb{E}(\mathcal{P}_R[w, u](t))$  it holds that

$$\mathbb{E}(\mathcal{P}_R[w, u](t)) = \int_0^\infty \mathcal{P}_r[w, u](t) \rho_R(r) dr = \int_{-\infty}^\infty s d\mu_{w, u, t}(s). \quad (3.2)$$

Hence, we see that the expected value is equal to the output of a *Prandtl-Ishlinskii operator* with weight function  $\rho_R$  and an initial state function that is equal to  $w$  on  $[0, \infty)$ .

To compute  $\mathcal{P}_R[w, u](t)$  or the measure  $\mu_{w, u, t}$  by using  $(\mathcal{P}_r[w, u](t))_{r \geq 0}$ , we consider the representation as in Lemma 1.

Let  $r_{w, u, t, 0}$  be defined as in (2.2a). Then it follows from Lemma 1.a that  $\mathcal{P}_r[w, u](t) = w$  for all  $r \geq r_{w, u, t, 0}$ .

If  $r_{w, u, t, 0} > 0$ , we consider  $(r_{w, u, t, k})_{k=1}^{K_{w, u, t}+1}$ ,  $(z_{w, u, t, k})_{k=0}^{K_{w, u, t}}$  and  $p_{w, u, t}$  as defined in (2.2c)–(2.5). Using Lemma 1.b, we see that  $[r_{w, u, t, k+1}, r_{w, u, t, k}] \ni r \rightarrow \mathcal{P}_r[w, u](t) \in I_{w, u, t, k}$  with

$$I_{w, u, t, k} := [z_{w, u, t, k} + \min \{ (-1)^k p_{w, u, t} r_{w, u, t, k}, (-1)^k p_{w, u, t} r_{w, u, t, k+1} \}, z_{w, u, t, k} + \max \{ (-1)^k p_{w, u, t} r_{w, u, t, k}, (-1)^k p_{w, u, t} r_{w, u, t, k+1} \}] \quad (3.3)$$

is a one-to-one mapping to with derivative  $(-1)^k p_{w, u, t} \in \{1, -1\}$  and it holds for  $r \in [r_{w, u, t, k+1}, r_{w, u, t, k}]$  and  $z \in I_k$  that  $\mathcal{P}_r[w, u](t) = z$  if and only if  $r = \frac{z - z_{w, u, t, k}}{(-1)^k p_{w, u, t}}$ . Using this properties, we can show that

$t$	$u(t)$	$\mathcal{P}_2[0, u](t)$	$\mathbb{E}(\mathcal{P}_R[0, u](t))$	$\max_{\tau \in [0, t]}  u(\tau) $	$\int_{\max\{ u(\tau)  : \tau \in [0, t]\}} \phi_R(r) dr$
0	0	0	0	0	1
0.5	2	0	0.199411	2	0.500016
1	4	2	1.99994	4	0.0000316722
2	0.5	2.5	1.60112	4	0.0000316722
3	3.5	1.5	1.7988	4	0.0000316722
$3\frac{13}{32}$	0.25	2.25	1.66774	4	0.0000316722
4	-4.5	-2.5	-2.49993	4.5	$2.8661 \cdot 10^{-7}$

Table 1: Values for the input function,  $u(t)$ ,  $\mathcal{P}_2[0, u](t)$ ,  $\mathbb{E}(\mathcal{P}_R[0, u](t))$ ,  $\max\{|u(\tau)| : \tau \in [0, t]\}$ , and  $\int_{\max\{|u(\tau)| : \tau \in [0, t]\}} \rho_R(r) dr$  for some values for  $t$  also considered in Figures 4 and 5.

the probability measure  $\mu_{w,u,t}$  for the random variable  $\mathcal{P}_R[0, u](t)$  is the sum of a Dirac measure at  $w$  weighted by  $\int_{r_{w,u,t,0}} \rho_R(r) dr = \int_{\max\{|u(\tau)-w| : \tau \in [0, t]\}} \rho_R(r) dr$  and of a measure with a density on  $\mathbb{R}$  being equal to 0 if  $r_{w,u,t,0} = 0$  and being equal to

$$\mathbb{R} \ni z \mapsto \sum_{i=0}^{K_{w,u,t}} \begin{cases} 0, & \text{if } z \notin I_{w,u,t,k}, \\ \rho_R\left(\frac{z - z_{w,u,t,k}}{(-1)^k p_{w,u,t}}\right) & \text{if } z \in I_{w,u,t,k}. \end{cases} \quad (3.4)$$

otherwise.

As example, we consider the piecewise monotone input function  $u \in C[0, T]$  such that the value of the function increases linearly on  $[0, 1]$  from 0 to 4, afterwards decreases linearly on  $[1, 2]$  to 0, afterwards increases linearly on  $[2, 3]$  to 3.5 and afterwards decreases linearly on  $[3, 4]$  to -4.5.

The probability measure  $\mu_{0,u,t}$  for the random variable  $\mathcal{P}_R[0, u](t)$  is the sum of a Dirac measure at 0 weighted by  $\int_{\max\{|u(\tau)| : \tau \in [0, t]\}} \rho_R(r) dr$  and of a measure with a density, denoted by  $\phi_t$ . For  $t \in \{0, 0.5, 1, 2, 3, 3\frac{13}{32}, 4\}$ , the values of  $u(t)$ , of the output  $\mathcal{P}_2[0, u](t)$  of the play operator with yield limit 2, of the expected value  $\mathbb{E}(\mathcal{P}_R[0, u](t))$  of  $(\mathcal{P}_R[0, u](t))$ , of  $\max\{|u(\tau)| : \tau \in [0, t]\}$  and of the weight  $\int_{\max\{|u(\tau)| : \tau \in [0, t]\}} \rho_R(r) dr$  are presented in Table 1. Moreover, the density  $\phi_t$ , the expected value and the weight are also shown in Figures 4 and 5.

To compute  $\mathcal{P}_r[0, u]\left(\frac{13}{32}\right)$  as function of  $r$  for  $t = 3\frac{13}{32} = \frac{109}{32}$ , we use Lemma 1. Using (2.2a)–(2.6) and the text afterwards, we get the values shown in Table 2, such that Lemma 1 yield that for  $[0, \infty) \ni r \rightarrow \mathcal{P}_r[0, u]\left(\frac{13}{32}\right)$  it holds that the function increases linearly on  $[0, 1.625]$  from 0.25 to 1.875, decreases linearly on  $[1.625, 1.75]$  to 1.75, increases linearly on  $[1.75, 2]$  to 2 and decreases linearly on  $[2, 4]$  to 0 and is equal to 0 on  $[4, \infty)$ , see also Figure 6. For computing the corresponding density  $\phi_{\frac{109}{32}}$ , shown in the top, right-hand side of Figure 5 by using (3.4), we have to sum up several values for  $\rho_R(r)$ . In Figure 6 the corresponding values are marked for  $z \in \{0.1, 1.3, 1.8, 1.9\}$ . In view of the  $I_k$  shown in Table 2, we see that for  $z \in [0, 0.25)$  we need to evaluate  $\rho_R(r)$  for one value of  $r$  to evaluate (3.4). In the corresponding computation for  $z \in [0.25, 1.75) \cup (1.875, 2]$ , we have to sum up two values of  $\rho_R$ , while for  $z \in [1.75, 1.875]$  we have to sum up four values.

As one can see in Figure 7, showing evolutions of  $\mathcal{P}_2[0, u](t)$  and of the expected value  $\mathbb{E}(\mathcal{P}_R[0, u](t))$ , it holds that these evolutions differ. Since [9, Coro. II.2.2] yields that

$$|\mathcal{P}_2[0, u](t) - \mathcal{P}_{\mathbb{E}(R)}[0, u](t)| \leq |\mathbb{E}(R) - 2| = 0.00007,$$

such that in the scale used in this figure the graphs of  $\mathcal{P}_2[0, u]$  and  $\mathcal{P}_{\mathbb{E}(R)}[0, u]$  would coincide, we deduce that also the evolutions of  $\mathcal{P}_{\mathbb{E}(R)}[0, u](t)$  and of the expected value  $\mathbb{E}(\mathcal{P}_R[0, u](t))$  differ.






Figure 4: Density function  $\phi_t$  and weight  $\int_{\max\{[u(\tau)]: \tau \in [0, t]\}} \phi_R(r) dr$  for Dirac measure concentrated at 0 leading to the probability measure  $\mu_{0,u,t}$  of  $\mathcal{P}_R[0, u](t)$  for some values of  $t$ , see also Table 1. Also  $\mathbb{E}(\mathcal{P}_R[0, u](t))$  is shown, marked by  $\mathbb{E}$ . It holds top, left-hand side:  $t = 0$ ; top, right-hand side:  $t = 0.5$ ; bottom, left-hand side:  $t = 1$ ; bottom, right-hand side:  $t = 2$ .

$k$	$r_{0,u,3\frac{13}{32},k}$	$s_{0,u,3\frac{13}{32},k}$	$z_{0,u,3\frac{13}{32},k}$	$I_{0,u,3\frac{13}{32},k}$
0	4	1	4	$[0, 2]$
1	2	2	0	$[1.75, 2]$
2	1.75	3	3.5	$[1.75, 1.875]$
3	1.625	$3\frac{13}{32}$	0.25	$[0.25, 1.875]$
4	0	—	—	

Table 2: Values for  $r_{0,u,3\frac{13}{32},k}$ ,  $s_{0,u,3\frac{13}{32},k}$ ,  $z_{0,u,3\frac{13}{32},k}$  and the interval  $I_{0,u,3\frac{13}{32},k}$  for the considered input function. It holds  $K_{0,u,3\frac{13}{32}} = 3$  and  $p_{0,u,3\frac{13}{32}} = -1$ .




Figure 5: Density function  $\phi_t$  and weight  $\int_{\max\{[u(\tau)]: \tau \in [0, t]\}} \phi_R(r) dr$  for Dirac measure concentrated at 0 leading to the probability measure  $\mu_{0,u,t}$  of  $\mathcal{P}_R[0, u](t)$  for some values of  $t$ , see also Table 1. Also  $\mathbb{E}(\mathcal{P}_R[0, u](t))$  is shown, marked by  $\mathbb{E}$ . It holds top, left-hand side:  $t = 3$ ; top, right-hand side:  $t = 3\frac{13}{32} = \frac{109}{32}$ , bottom  $t = 4$ .




Figure 6: Output of play operator  $\mathcal{P}_r[0, u](t)$  as function of  $r$  for  $t = 3\frac{13}{32} = \frac{109}{32}$  and evaluating value for  $\rho_R$  during computing the sum as in (3.4) for  $z \in \{0.1, 1.3, 1.8, 1.9\}$ .




Figure 7: Left-hand side: evolution of  $\mathcal{P}_2[0, u](t)$  (indistinguishable from evolution of  $\mathcal{P}_{\mathbb{E}(R)}[0, u](t)$  in the considered scale) and of the expected value  $\mathbb{E}(\mathcal{P}_R[0, u](t))$ ; right-hand side: corresponding input-output diagram

## 4 Generalized Prandtl-Ishlinskii operator

Following [6, 7, 14], we consider *generalized Prandtl-Ishlinskii operators* that are defined by combining a Prandtl-Ishlinskii operators and a given function  $g : \mathbb{R} \rightarrow \mathbb{R}$  by mapping an input function  $u : [0, T] \rightarrow [0, T]$  to the output of the Prandtl-Ishlinskii considered with the composition  $g \circ u$  used as input function. These kind of operators can be inverted analytically and can model effects like saturation, asymptotic behaviors, etc, see, e.g., discussions in [1, 5, 14].

In [4, Sec. 3 and 5] the following generalized Prandtl-Ishlinskii operator has been used to model the *magnetization* of Galfenol for an applied magnetic field  $H$ . This operator is defined by mapping  $H \in C[0, T]$  to  $\mathcal{G}_{c_1, c_2, c_3}[H] \in C[0, T]$  defined by

$$\mathcal{G}_{c_1, c_2, c_3}[H](t) = \mathcal{PI}_{c_1, c_2}[\tanh(c_3 H)](t), \quad \forall t \in [0, T], \quad (4.1)$$

with the Prandtl-Ishlinskii operator  $\mathcal{PI}_{c_1, c_2} : C[0, T] \rightarrow C[0, T]$  being defined by

$$\mathcal{PI}_{c_1, c_2}[u](t) = \int_0^\infty C_1 e^{-r/C_2} \mathcal{P}_r[\lambda_0(r), u](t) dr, \quad \forall u \in C[0, T], t \in [0, T], \quad (4.2)$$

$c_1, c_2, c_3$  being positive parameters and  $\lambda_0 : [0, \infty) \rightarrow \mathbb{R}$  being an appropriate function.

Related to a Prandtl-Ishlinskii operator is the *initial loading curve*, whose value at  $x \in \mathbb{R}$  can be determined by letting the input monotonically increase or decrease linearly from 0 to  $x$  on  $[0, T]$  with an initial configuration  $\lambda$  being constant 0, and evaluating the output of the Prandtl-Ishlinskii operator at  $T$ .

In applications to plasticity modeled by a Prandtl-Ishlinskii operator this curve can be measured by considering a material with no previous memory, e.g., if the memory was erased for instance by heating above the Curie temperature and cooling again or applying an input function with decreasing oscillations.

The initial loading curve  $\psi_{c_1, c_2}$  for  $\mathcal{PI}_{c_1, c_2}$  satisfies

$$\begin{aligned} \psi_{c_1, c_2}(s) &= \int_0^\infty C_1 e^{-r/C_2} \begin{cases} (s-r), & \text{if } r \leq s, \\ 0, & \text{if } -r \leq s \leq r, \\ (s+r), & \text{if } r \geq -s, \end{cases} dr, \\ &= s c_1 c_2 + c_1 c_2^2 \begin{cases} e^{-\frac{s}{c_2}} - 1, & \text{if } s \geq 0, \\ 1 - e^{\frac{s}{c_2}}, & \text{if } s \leq 0, \end{cases} \end{aligned} \quad (4.3)$$

for all  $s \in \mathbb{R}$ .




Figure 8: FORC and measurements during creation of these curves.,

Following, [5, 6, 10], we consider  $u \in C([0, T])$  and  $0 \leq t_a < t_b < t_c \leq T$  with  $u$  being monotone on  $[t_a, t_b]$  and on  $[t_b, t_c]$  and  $u(t_a) = u(t_c)$ , we have for all  $w_0 \in \mathbb{R}$  that

$$\mathcal{P}_r[\beta_0(r), u](t_c) - \mathcal{P}_r[\beta_0(r), u](t_b) = 2 \begin{cases} 0, & \text{if } \left| \frac{u(t_c) - u(t_b)}{2} \right| \leq r, \\ \frac{u(t_c) - u(t_b)}{2} - r, & \text{if } \frac{u(t_c) - u(t_b)}{2} > r, \\ \frac{u(t_c) - u(t_b)}{2} + r, & \text{if } -\frac{u(t_c) - u(t_b)}{2} > r. \end{cases} \quad (4.4)$$

Therefore, invoking (4.3) and (4.2), we get

$$\begin{aligned} & \psi_{c_1, c_1} \left( \left| \frac{u(t_c) - u(t_b)}{2} \right| \right) \\ &= \frac{1}{2} \begin{cases} \mathcal{PI}_{c_1, c_2}[\lambda_0, u](t_c) - \mathcal{PI}_{c_1, c_2}[\lambda_0, u](t_b), & \text{if } u(t_c) \geq u(t_b), \\ \mathcal{PI}_{c_1, c_2}[\lambda_0, u](t_b) - \mathcal{PI}_{c_1, c_2}[\lambda_0, u](t_c), & \text{if } u(t_b) \leq u(t_c). \end{cases} \end{aligned} \quad (4.5)$$

Considering an input cycle and differences for corresponding measurements for a process that one would like to represent by a Prandtl-Ishlinskii operator, one can derive an *approximation for the initial loading curve* on  $[0, |(u(t_2) - u(t_b))/2|]$  from the measurements during  $[t_1, t_2]$ .

## 5 Identification

### 5.1 Considered situation

In [2], a magnetostrictive Terfenol-actuator is investigated and the hysteresis between the reference signal determining the current generating the magnetic field and the resulting displacement is considered, and the data of the *First-Order-Reversal-Curves (FORC)* are used to determine a Preisach operator and a generalized Prandtl-Ishlinskii operator.

We will now use the data measured during the preparation of the FORC-diagram in [2], considering now the measured current (corrente) generated from the reference signal as input data, such that we

get Figure 8 from the measurements, quite similar to the FORC diagram in [2, Fig. 4.(a)], except for the scale of the input.

Now, we will approximate these data by the initial loading curve corresponding to  $\mathcal{G}_{c_1, c_2, c_3}[H](t)$ , i.e., we consider the differences derived from the function  $\tanh(c_3 I(t))$  as input for  $\psi_{c_1, c_1}$ . This will be used to determine an appropriate value for  $c_3$  and to derive information about  $c_1$  and  $c_2$  and their uncertainty.

## 5.2 Identification of $c_3$

To determine  $c_3$ , we consider the generated approximation of the initial loading curve. For  $j \in \{30, \dots, 58\}$ , we have sets of times  $t_{i,0} < t_{i,1} < \dots < t_{i,K_i}$  such that the current  $I$  as function of time has a discrete local maximum at  $t_{i,0}$  and it holds  $I(t_{i,0}) > I(t_{i,1}) > \dots > I(t_{i,K_i})$ . For  $j \in \{1, \dots, 29\}$  it holds that the current  $I$  has a discrete local minima at  $t_{i,0}$  and it holds  $I(t_{i,0}) < I(t_{i,1}) < \dots < I(t_{i,K_i})$ .

Let  $L$  denote the length change measured by the length sensor as function of time. Then it follows that the data sets  $(I(t_{i,k}), L(t_{i,k}))_{k=0}^{K_i}$  for  $i = 30, \dots, 58$  represent the FORCs.

To derive the initial loading curve  $\zeta_{i,c_3}$  for  $i \in \{1, 2, \dots, 58\}$  and for  $c_3 > 0$  by considering the data set  $(I(t_{i,k}), L(t_{i,k}))_{k=0}^{K_i}$  and the transformation  $x \mapsto \tanh(c_3 x)$ , we consider (4.5) with  $\psi_{c_1, c_1}$  replaced by  $\zeta_{i,c_3}$ ,  $\mathcal{PI}_{c_1, c_2}[\lambda_0, u]$  replaced by  $L$ ,  $u(t) := \tanh(c_3 I(t))$ ,  $t_b = t_{i,0}$ , and  $t_c = t_{i,k}$  for all  $k = 0, \dots, K_i$ , leading to

$$\begin{aligned} & \zeta_{i,c_3} \left( \frac{|\tanh(c_3 I(t_{i,k})) - \tanh(c_3 I(t_{i,0}))|}{2} \right) \\ &= \frac{1}{2} \begin{cases} L(t_{i,k}) - L(t_{i,0}), & \text{if } i \leq 29, \\ L(t_{i,0}) - L(t_{i,k}), & \text{if } i > 29, \end{cases} \quad \forall k \in \{0, 1, \dots, K_i\}, i \in \{1, \dots, 58\}. \end{aligned} \quad (5.1)$$

By interpolation, we get a function  $\zeta_{i,c_3}$  defined on  $[0, \frac{1}{2} |\tanh(c_3 I(t_{i,k})) - \tanh(c_3 I(t_{i,0}))|]$ .

Now,  $c_3$  is determined by requesting that the sum over the squared  $L^2$  difference between these generated approximations is minimized. We end up with the optimal value  $c_3 = c_{\tanh} = 0.682138$ . The corresponding  $(\zeta_{i,c_{\tanh}})_{i=1}^{58}$  set of initial loading curves is shown in Figure 9.

## 5.3 Identification of $c_{1,i}$ and $c_{2,i}$

For each approximation  $\zeta_{i,c_{\tanh}}$  of the initial loading curve derived from the measurements by using interpolation, we determine  $(c_{1,i}, c_{2,i}) \in (0, 250) \times (0.0000001, 25)$  by minimizing the  $L^2$  difference between this approximations for the initial loading and the  $\psi_{c_{1,i}, c_{2,i}}$  loading curve corresponding to the Prandtl-Ishlinskii-operator  $\mathcal{PI}_{c_{1,i}, c_{2,i}}$ , see an example in Figure 10.

Moreover, a data pair  $(c_1, c_2)$  has been computed by minimizing the sum over the  $L^2$ -difference between  $\zeta_{i,c_{\tanh}}$  and  $\psi_{c_{1,i}, c_{2,i}}$  for all  $i \in \{1, \dots, 58\}$ , leading to  $c_1 = 11.7463$  and  $c_2 = 0.1636$ .

If only the computed parameter pairs for the curves generated for decreasing inputs are considered, and also the pairs belonging to the three shortest definition intervals are ignored, we get the *appropriate subset*  $((c_{1,i}, c_{2,i}))_{i=30}^{55}$  of 26 parameter pairs. Computing for this subset the *mean* and the *standard deviation* of the corresponding values for  $c_{1,i}$  and  $c_{2,i}$  leads to  $\text{mean}_i(c_{1,i}) = 13.3218$ ,  $\text{mean}_i(c_{2,i}) = 0.116617$ ,  $\text{std}(c_{1,i}) = 2.67912$ , and  $\text{std}(c_{2,i}) = 0.043755$ .




Figure 9: Derived initial loading curves



Figure 10: The first measured initial loading curve (solid line) and  $\psi_{c_1, i, c_2, i}$  (dashed line)

Figure 11: Input-output plot for  $\mathcal{P}\mathcal{I}_{\text{mean}_i(c_1, i), \text{mean}_i(c_2, i), c_3}[u]$ ,  $\mathcal{P}\mathcal{I}_{\text{mean}_i(c_1, i), \text{mean}_i(c_2, i) \pm \text{std}_i(c_2, i), c_3}[u]$ , and  $\mathcal{P}\mathcal{I}_{\text{mean}_i(c_1, i), \text{mean}_i(c_2, i) \pm 2\text{std}_i(c_2, i), c_3}[u]$ , for the initial state  $\lambda_0 \equiv 0$  and an input function  $u$  increasing linearly from 0 to 1.8, decreasing linearly afterwards to 0 and increasing linearly to 1.8 afterwards, Left-hand side: evolution of 0 — input  $u$ , 1 —  $\mathcal{P}\mathcal{I}_{\text{mean}_i(c_1, i), \text{mean}_i(c_2, i) - 2\text{std}_i(c_2, i), c_3}[u]$ , 2 —  $\mathcal{P}\mathcal{I}_{\text{mean}_i(c_1, i), \text{mean}_i(c_2, i) - \text{std}_i(c_2, i), c_3}[u]$ , 3 —  $\mathcal{P}\mathcal{I}_{\text{mean}_i(c_1, i), \text{mean}_i(c_2, i), c_3}[u]$ , 4 —  $\mathcal{P}\mathcal{I}_{\text{mean}_i(c_1, i), \text{mean}_i(c_2, i) + \text{std}_i(c_2, i), c_3}[u]$ , 5 —  $\mathcal{P}\mathcal{I}_{\text{mean}_i(c_1, i), \text{mean}_i(c_2, i) + 2\text{std}_i(c_2, i), c_3}[u]$ , right-hand side: input-output plots for 1–5

One could now use this data to start further investigations. As simple example, we consider  $\mathcal{PI}_{\text{mean}_i(c_{1,i}), \text{mean}_i(c_{2,i}), c_3}[u]$  and  $\mathcal{PI}_{\text{mean}_i(c_{1,i}), \text{mean}_i(c_{2,i}) \pm 2 \text{std}_i(c_{2,i}), c_3}[u]$ , for the initial state  $\lambda_0 \equiv 0$  and an input function  $u$  increasing linearly from 0 to 1.8, decreasing afterwards linearly to 0 and increasing linearly to 1.8 leading to the outputs shown in Figure 11.

As an example for forward UQ, we now assume that the value of  $c_2$  can be represented by a random variable  $C_2$  with density  $\rho_{C_2}$  that is derived from the distribution  $N(\text{mean}_i(c_{1,2}), (\text{std}_i(c_{2,2}))^2) = N(0.116617, (0.043755)^2)$  by ignoring  $]-\infty, 0.0000001]$  and rescaling, see left-hand side of Figure 12. In right-hand side of this figure, an input function  $u$ , the graph of  $\mathcal{PI}_{\text{mean}_i(c_1), \text{mean}_i(c_2)}[u]$  (graph output does not change in this scale, if  $\text{mean}_i(c_2) = 0.116617$  is replaced by  $\mathbb{E}(C_2) = 0.11712$ ) and the graph of  $[0, T] \ni t \mapsto \mathbb{E}(\mathcal{PI}_{\text{mean}_i(c_1), C_2}[u](t))$  with  $\lambda_0 \equiv 0$  are shown. In this example there is only a small difference between these graphs.




Figure 12: Left-hand side: Density  $\rho_2$  of random variable  $C_2$  describing  $c_2$  derived from  $N(\text{mean}_i(c_{1,2}), (\text{std}_i(c_{2,2}))^2) = N(0.116617, (0.043755)^2)$  by ignoring  $]-\infty, 0.0000001]$  and rescaling, expected value  $\mathbb{E}(C_2) = 0.11712$  is marked by  $\mathbb{E}$ . Right-hand side: an Input function  $u$  (marked by 0), graph of  $\mathcal{PI}_{\text{mean}_i(c_1), \text{mean}_i(c_2)}[u]$  (marked by 1) and graph of  $\mathbb{E}(\mathcal{PI}_{\text{mean}_i(c_1), C_2}[u])$  (marked by 2)

If all computed parameter pairs are considered, computing the mean and the standard deviation for all values for  $c_1$  and  $c_2$ , we get  $\text{mean}_{i,\text{all}}(c_{1,i}) = 16.0156$ ,  $\text{mean}_{i,\text{all}}(c_{2,i}) = 4.71742$ ,  $\text{std}_{i,\text{all}}(c_{1,i}) = 13.3218$ , and  $\text{std}_{i,\text{all}}(c_{2,i}) = 7.94603$ . In view of these values, it is obvious, that one can not use them to derive a satisfying assumption for the random variables representing the positive parameter  $c_1$  and  $c_2$ .

We could consider  $N(m_1, \delta_1^2)$  and  $N(m_2, \delta_2^2)$  for some positive constants  $m_1, m_2, \delta_1, \delta_2$  and could perform some truncation, to derive random variables  $C_1$  and  $C_2$  to be used in UQ to represent an appropriate approximations for  $c_1$  and  $c_2$ . But, this could only be justified if the pairs  $((c_{1,i}, c_{2,i}))_{i=1}^{58}$  or the pairs  $((c_{1,i}, c_{2,i}))_{i=30}^{55}$  could be considered as typical independent samples for  $N(m_1, \delta_1^2)$  and  $N(m_2, \delta_2^2)$ . A typical example for the cloud generated by this kind of samples is shown in Figure 13 for  $(N(1, 1^2), N(2, (\frac{1}{2})^2))$ .

Considering the computed parameter pairs  $((c_{1,i}, c_{2,i}))_{i=1}^{58}$ , see Figure 14, it is obvious that we can not get these set of pairs as samples if we assume that the parameter can be represented by two independent normal distributed random variables, even if the correlation between  $(c_{1,i})_i$  and  $(c_{2,i})_i$  is  $-0.0737148$ . Also the 26 data pairs  $((c_{1,i}, c_{2,i}))_{i=30}^{55}$  in the appropriate subset introduced on page 5.3, see Figure 15, with a corresponding correlation of  $-0.532375$ , can not be derived as samples in this way.




Figure 13: 58 independent samples for  $(A_1, A_2) = \left( N\left(1, (1)^2\right), N\left(2, \left(\frac{1}{2}\right)^2\right) \right)$ .




Figure 14: Sample pairs  $(c_{1,1}, c_{2,1}), \dots, (c_{1,58}, c_{2,58})$ . Left-hand side: plot of all pairs, right-hand side: zoom plot






Figure 15: Appropriate subset  $(c_{1,30}, c_{2,30}), \dots, (c_{1,55}, c_{2,55})$  of 26 parameter pairs introduced on page 11

Hence, we are interested in finding a more general random variable  $(C_1^*, C_2^*)$  with values in  $(0, \infty) \times (0, \infty)$  such that either  $((c_{1,i}, c_{2,i}))_{i=1}^{58}$  or  $((c_{1,i}, c_{2,i}))_{i=30}^{55}$  could be considered as typical samples of this random variable.

## 5.4 Applications of Bayes' Theorem

Now, we will use Bayes' Theorem to derive a probability density on  $(0, \infty) \times (0, \infty)$  by considering the measured data.

Let  $v^{\text{obs}}$  be a vector containing the sets of the non-trivial discrete values for the initial loading curve used to derived their approximations in Section 5.2 and 5.3, i.e.

$$v^{\text{obs}} = \left( \begin{cases} \left( \frac{1}{2} (L(t_{i,k}) - L(t_{i,0})) \right)_{k=1}^{K_i} & \text{if } i \leq 29, \\ \left( \frac{1}{2} (L(t_{i,0}) - L(t_{i,k})) \right)_{k=1}^{K_i} & \text{if } i > 29, \end{cases} \right)_{i=1}^{58} \in \mathbb{R}^{K_1} \times \mathbb{R}^{K_2} \times \dots \times \mathbb{R}^{K_{58}}. \quad (5.2)$$

We consider on  $(0, \infty) \times (0, \infty)$  the uniform probability density  $\pi_0$  for  $(0, 250) \times (0.0000001, 25)$  as *priori* density. Updating this priori density by following Bayes' Theorem of Inverse Problems, see, e.g., [11, Sec. 8.1], [12, Sec. 2.8 and Sec. 6.2], we get the *posterior* probability density  $\pi_{\text{new}}$  on  $(0, \infty) \times (0, \infty)$  defined by

$$\pi_{\text{new}}(c_1, c_2) = \frac{L(c_1, c_2 | v^{\text{obs}}) \pi_0(c_1, c_2)}{\int_0^\infty \int_0^\infty L(c'_1, c'_2 | v^{\text{obs}}) \pi_0(c'_1, c'_2) dc'_2 dc'_1} \quad (5.3)$$

with  $L((c'_1, c'_2) | v^{\text{obs}})$  being the *likelihood* that the parameter pair has the value  $(c'_1, c'_2)$  if the output  $v^{\text{obs}}$  is observed. To derive the *forward function*  $\Psi_{c_1, c_2} : (0, \infty) \times (0, \infty)$  allowing to formulate the likelihood function, we use a method similar to the derivation of (5.1). We consider (4.5) with  $u(t) := \tanh(c_3 I(t))$ ,  $t_b = t_{i,0}$ ,  $t_b = t_{i,0}$ , and  $t_c = t_{i,k}$  for all  $k = 0, \dots, K_i$ . Using that  $L(t_{i,k})$  and  $L(t_{i,0})$  are approximations of  $\mathcal{PI}_{c_1, c_2}[u](t_{i,k})$  and  $\mathcal{PI}_{c_1, c_2}[u](t_{i,0})$ , we end up with the assumption that there is a sample  $\gamma = ((\gamma_{i,k})_{k=0}^{K_i})_{i=1}^{58}$  of a random variable  $\Gamma = ((\Gamma_{i,k})_{k=0}^{K_i})_{i=1}^{58}$  on  $\mathbb{R}^{K_1+1} \times \mathbb{R}^{K_2+1} \times \dots \times \mathbb{R}^{K_{58}+1}$  such that

$$\begin{aligned} & \psi_{c_1, c_2} \left( \frac{|\tanh(c_3 I(t_{i,k})) - \tanh(c_3 I(t_{i,0}))|}{2} \right) \\ &= \frac{1}{2} \begin{cases} L(t_{i,k}) + \gamma_{i,k} - (L(t_{i,0}) + \gamma_{i,0}), & \text{if } i \leq 29, \\ L(t_{i,0}) + \gamma_{i,0} - (L(t_{i,k}) + \gamma_{i,k}), & \text{if } i > 29, \end{cases} \quad \forall k \in \{0, 1, \dots, K_i\}. \end{aligned} \quad (5.4)$$

Now, defining the forward function  $\Psi : (0, \infty) \times (0, \infty) \rightarrow \mathbb{R}^{K_1} \times \mathbb{R}^{K_2} \times \dots \times \mathbb{R}^{K_{58}}$  by

$$\Psi(c_1, c_2) := \left( \left( \psi_{c_1, c_2} \left( \frac{|\tanh(c_3 I(t_{i,k})) - \tanh(c_3 I(t_{i,0}))|}{2} \right) \right)_{k=1}^{K_i} \right)_{i=1}^{58}. \quad (5.5)$$

and

$$\eta := \left( \left\{ \begin{array}{ll} \left( \frac{1}{2} (\gamma_{i,0} - \gamma_{i,k}) \right)_{k=1}^{K_i} & \text{if } i \leq 29, \\ \left( \frac{1}{2} (\gamma_{i,k} - \gamma_{i,0}) \right)_{k=1}^{K_i} & \text{if } i > 29, \end{array} \right\} \right)_{i=1}^{58} \in \mathbb{R}^{K_1} \times \mathbb{R}^{K_2} \times \dots \times \mathbb{R}^{K_{58}}, \quad (5.6)$$

we have

$$\Psi(c_1, c_2) + \eta = v^{\text{obs}} \quad (5.7)$$

and that  $\eta$  is a sample for  $H := \left( \left\{ \begin{array}{ll} \left( \frac{1}{2} (\Gamma_{i,0} - \Gamma_{i,k}) \right)_{k=1}^{K_i} & \text{if } i \leq 29, \\ \left( \frac{1}{2} (\Gamma_{i,k} - \Gamma_{i,0}) \right)_{k=1}^{K_i} & \text{if } i > 29, \end{array} \right\} \right)_{i=1}^{58}$  and that  $v^{\text{obs}}$  is a sample for  $\Psi(c_1, c_2) + H$ . Hence, we have to consider the likelihood function  $L$  associated to function mapping  $(c'_1, c'_2) \in (0, \infty) \times (0, \infty)$  to the random variable  $\Psi(c'_1, c'_2) + H$ .

In the following, we assume that the components

$$\Gamma_{1,0}, \Gamma_{1,1}, \dots, \Gamma_{1,K_1}, \Gamma_{2,0}, \Gamma_{2,1}, \dots, \Gamma_{2,K_2}, \dots, \Gamma_{58,0}, \Gamma_{58,1}, \dots, \Gamma_{58,K_{58}}, \text{ of } \Gamma$$

are independent random variables. Moreover, we assume that we have some  $\sigma > 0$  such that  $\Gamma_{1,1}, \Gamma_{1,1}, \dots, \Gamma_{1,K_1}, \Gamma_{2,1}, \Gamma_{2,1}, \dots, \Gamma_{2,K_2}, \dots, \Gamma_{58,1}, \Gamma_{58,1}, \dots, \Gamma_{58,K_{58}}$  have the distribution  $N(0, \sigma^2)$  representing the measurement error, i.e. the error between  $\mathcal{PT}_{c_1, c_2}[u](t_{i,k})$  and  $L(t_{i,k})$ .

For  $i \in \{1, \dots, 58\}$ , it is assumed that  $\gamma_{i,0}$  contains in addition to the measurement error between  $\mathcal{PT}_{c_1, c_2}[u](t_{i,0})$  and  $L(t_{i,0})$  also the error generated by evaluating (4.5) for  $t_b = t_{i,0}$  and not for  $t_b$  being the time  $t_i^*$  near to  $t_{i,0}$  such that the continuous function  $I$  has the local extremum in  $t_i^*$ . Since  $t_i^*$  may be different from  $t_{i,0}$ , (4.5) only holds with some additional error term. Hence, we assume that  $\Gamma_{1,0}, \Gamma_{2,0}, \dots, \Gamma_{58,0}$  have the distribution  $N(0, (2\sigma)^2)$ .

Now, one can show that  $H$  has the density  $\rho^* : \mathbb{R}^{K_1} \times \mathbb{R}^{K_2} \times \dots \times \mathbb{R}^{K_{58}} \rightarrow [0, \infty)$  satisfying

$$\rho^* \left( \left( (w_{i,k})_{k=1}^{K_i} \right)_{i=1}^{58} \right) = \left( \frac{2}{\sqrt{2\pi\sigma^2}} \right)^{\sum_{i=1}^{58} K_i} \left( \prod_{i=1}^{58} \sqrt{\frac{1}{4K_i + 1}} \right) \exp \left( -\frac{2}{\sigma^2} \sum_{i=1}^{58} \left( \sum_{k=1}^{K_i} w_{i,k}^2 - \left( \frac{1}{4} + K_i \right)^{-1} \left( \sum_{k=1}^{K_i} w_{i,k} \right)^2 \right) \right). \quad (5.8)$$

For  $(c'_1, c'_2) \in (0, \infty) \times (0, \infty)$  it holds that  $\Psi(c'_1, c'_2) + H$  has the density

$$\mathbb{R}^{K_1} \times \mathbb{R}^{K_2} \times \dots \times \mathbb{R}^{K_{58}} \ni \left( (w_{i,k})_{k=1}^{K_i} \right)_{i=1}^{58} \mapsto \rho^* \left( \left( (w_{i,k})_{k=1}^{K_i} \right)_{i=1}^{58} - \Psi(c'_1, c'_2) \right),$$

such that we can define  $L \left( (c'_1, c'_2) \left| \left( (w_{i,k})_{k=1}^{K_i} \right)_{i=1}^{58} \right. \right)$  to be the formulae on the right-hand side.

The *normalized likelihood function*  $L_{\text{norm}}$  is defined by

$$L_{\text{norm}} \left( (c'_1, c'_2) \left| \left( (w_{i,k})_{k=1}^{K_i} \right)_{i=1}^{58} \right. \right) = \frac{L \left( (c'_1, c'_2) \left| \left( (w_{i,k})_{k=1}^{K_i} \right)_{i=1}^{58} \right. \right)}{\max_{(c''_1, c''_2) \in (0, \infty)^2} L \left( (c''_1, c''_2) \left| \left( (w_{i,k})_{k=1}^{K_i} \right)_{i=1}^{58} \right. \right)}. \quad (5.9)$$




Figure 16: Normalized likelihood for the standard deviation 0.01 and 0.02 for the noise in the measured length. Left-hand side: 3D plot, right-hand side contour plot with isoline

We can replace the likelihood in (5.3) by the normalized likelihood without changing the result, also if we use a numerical approximation for the maximum instead of true maximum in the evaluation.

For  $\sigma = 0.01$  corresponding functions plots for  $L_{norm}((c'_1, c'_2)|v^{obs})$  can be found in Figure 16, the position for numerical maximum for the likelihood is  $(c_1, c_2) = (12.2253, 0.1519)$ . For  $\sigma = 0.05$  the corresponding functions plots are shown in figure 17, the position for numerical maximum for the likelihood is the same as for  $\sigma = 0.01$ .

Hence, we see that this likelihood function will generate a posterior probability density which is numerical equal to 0 except for a small region, such that the 58 computed pairs  $(c_{1,i}, c_{2,i})$  are obviously no typical samples for this probability density.

Recalling the derivation of the likelihood function and [11, 12], one realizes that this function is derived from a probability density for a noise disturbed output with using **one** parameter pair  $(c_1, c_2)$  for all possible initial loading curves. Therefore, the Bayes' theorem will generated a density on  $(0, \infty) \times (0, \infty)$  represents somehow the informations about the position of this **one** parameter pair that can be extracted by investigating the observed values in  $v^{obs}$  and does not contain information for dealing with different parameter pairs for different subsets of the observation.

## 5.5 Further research for UQ for magneto-mechanical component

### 5.5.1 UQ-issues for further research

We would like to find a probability density on  $(0, \infty) \times (0, \infty)$  such that the 58 computed pairs of parameter values or the appropriate subset of 26 parameter pairs introduced on page 11 are typical samples for for this probability density. A further investigation of Bayes' theorem as discussed for example [11, 12] yields that applying this theorem would require to use these probability densities as parameter instead of the numbers pairs  $(c_1, c_2)$  considered in Section 5.4, and that Bayes' theorem would return a probability density over a set of probability density on  $(0, \infty) \times (0, \infty)$  as result.

If one should apply now Bayes' theorem or try to determine the probability density on  $(0, \infty) \times (0, \infty)$  maximizing the likelihood will be subject of further research.




Figure 17: Normalized likelihood for the standard deviation 0.05 and 0.1 for the noise in the measured length. Left-hand side: 3D plot, right-hand side contour plot with isoline, top: input data range as in Figure 16, bottom: input data range adapted to considered likelihood function

### 5.5.2 Modeling issues

The considered data are derived from length change over magnetic field data. Hence, to be compatible with the thermodynamical consistent model derived in [4], one would need to replace the generalized Prandtl-Ishlinskii operator  $\mathcal{G}_{c_1, c_2, c_3}[H](t)$  by a function evaluation involving also the counter clockwise potential operator of an hysteresis operator appropriate to model the magnetization, e.g., the counter clockwise potential operator to  $\mathcal{G}_{c_1, c_2, c_3}$ . Corresponding investigations will be subject of further research.

## 6 Conclusion

The output of hysteresis operators dependent on parameter vales, but their values may not exactly known when modeling real world processes.

These parameter can be considered as random variables and the methods of uncertainty quantification (UQ) allow to investigate the influence of the uncertainty on the output of the model and to determine the random variable representing the information on the parameter.

## Acknowledgment

O. Klein would like to thank Prof. C. Schillings, University of Mannheim, for fruitful discussions.

## References

- [1] M. Al Janaideh, D. Davino, P. Krejčí, and C. Visone. Comparison of Prandtl-Ishlinskii and Preisach modeling for smart devices applications. *Physica B: Condensed Matter*, 486:155–159, apr 2016.
- [2] M. Al Janaideh, C. Visone, D. Davino, and P. Krejčí. The generalized Prandtl-Ishlinskii model: relation with the Preisach nonlinearity and inverse compensation error. In *2014 American Control Conference (ACC) June 4-6, 2014. Portland, Oregon, USA, 2014*.
- [3] M. Brokate and J. Sprekels. *Hysteresis and phase transitions*. Springer, New York, 1996.
- [4] D. Davino, P. Krejčí, and C. Visone. Fully coupled modeling of magneto-mechanical hysteresis through ‘thermodynamic’ compatibility. *Smart Mater. Struct.*, 22(9), SEP 2013.
- [5] D. Davino and C. Visone. Rate-independent memory in magneto-elastic materials. *Discrete and Continuous Dynamical Systems - Series S*, 8(4):649–691, Oct 2015.
- [6] O. Klein and P. Krejčí. Outwards pointing hysteresis operators and asymptotic behaviour of evolution equations. *Nonlinear Anal. Real World Appl.*, 4(5):755–785, Dezember 2003.
- [7] O. Klein and P. Krejčí. Asymptotic behaviour of evolution equations involving outwards pointing hysteresis operators. *Phys. B*, 343:53–58, 2004. Proceedings of the Fourth International Symposium on Hysteresis and Micromagnetic Modeling (HMM-2003).

- [8] M. Krasnosel'skiĭ and A. Pokrovskii. *Systems with Hysteresis*. Springer-Verlag, Heidelberg, 1989. Russian edition: Nauka, Moscow, 1983.
- [9] P. Krejčí. *Hysteresis, Convexity and Dissipation in Hyperbolic Equations*, volume 8 of *Gakuto Int. Series Math. Sci. & Appl.* Gakkōtoshō, Tokyo, 1996.
- [10] I. D. Mayergoyz. *Mathematical Models of Hysteresis and their Applications*. Elsevier, second edition edition, 2003.
- [11] R. C. Smith. *Uncertainty quantification: theory, implementation, and applications*, volume 12 of *Computational Science & Engineering*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2014. Theory, implementation, and applications.
- [12] T. J. Sullivan. *Introduction to uncertainty quantification*, volume 63 of *Texts in Applied Mathematics*. Springer, Cham, 2015.
- [13] A. Visintin. *Differential Models of Hysteresis*, volume 111 of *Applied Mathematical Sciences*. Springer, New York, 1994.
- [14] C. Visone and M. Sjöström. Exact invertible hysteresis models based on play operators. *Physica B: Condensed Matter*, 343(1):148 – 152, 2004. Proceedings of the Fourth Intional Conference on Hysteresis and Micromagnetic Modeling.