

**Essential boundedness for solutions of the  
Neumann problem on general domains**

A.F.M. ter Elst<sup>1</sup>, Hannes Meinlschmidt<sup>2</sup>, Joachim Rehberg<sup>3</sup>

submitted: February 15, 2019

<sup>1</sup> Department of Mathematics  
University of Auckland  
Private bag 92019  
Auckland 1142  
New Zealand  
E-Mail: terelst@math.auckland.ac.nz

<sup>2</sup> Johann Radon Institute for Computational  
and Applied Mathematics (RICAM)  
Altenberger Str. 69  
4040 Linz  
Austria  
E-Mail: hannes.meinlschmidt@ricam.oeaw.ac.at

<sup>3</sup> Weierstrass Institute  
Mohrenstr. 39  
10117 Berlin  
Germany  
E-Mail: joachim.rehberg@wias-berlin.de

No. 2574  
Berlin 2019



---

2010 *Mathematics Subject Classification.* 35D10, 46E35.

*Key words and phrases.* Regularity, Neumann problem, elliptic operators.

The first and third named authors are grateful for a most stimulating stay at the RICAM. Part of this work is supported by the Marsden Fund Council from Government funding, administered by the Royal Society of New Zealand.

Edited by  
Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS)  
Leibniz-Institut im Forschungsverbund Berlin e. V.  
Mohrenstraße 39  
10117 Berlin  
Germany

Fax: +49 30 20372-303  
E-Mail: [preprint@wias-berlin.de](mailto:preprint@wias-berlin.de)  
World Wide Web: <http://www.wias-berlin.de/>

# Essential boundedness for solutions of the Neumann problem on general domains

A.F.M. ter Elst, Hannes Meinschmidt, Joachim Rehberg

## Abstract

Let the domain under consideration be bounded. Under the suppositions of very weak Sobolev embeddings we prove that the solutions of the Neumann problem for an elliptic, second order divergence operator are essentially bounded, if the right hand sides are taken from the dual of a Sobolev space which is adapted to the above embedding.

## 1 Introduction

If  $A$  is a pure second-order elliptic operator with Dirichlet boundary conditions and real measurable coefficients on a bounded connected open set  $\Omega \subset \mathbb{R}^d$ , then it is not too hard to show that the resolvent operator  $A^{-1}$  maps  $L^q(\Omega)$  into  $L^\infty(\Omega)$  if  $q > d$ . It is a famous result of Stampacchia ([Sta65] Theorem 4.4) that  $A^{-1}$  extends to a continuous operator from  $W^{-1,q}(\Omega)$  into  $L^\infty(\Omega)$  if  $q > d$ . Inspecting the proof, it becomes clear that this result extends without difficulties from the pure Dirichlet case to the case of mixed boundary conditions, as long as the Dirichlet part of the boundary is large enough to imply a Poincaré inequality. It is also possible to extend the result to merely Neumann boundary conditions if a positive scalar is added to the operator, and hence the resulting operator is coercive. The idea how to do this can be found in the book of Tröltzsch ([Trö10] Section 7.2.2). What remains open is the pure divergence form operator with pure Neumann boundary condition. It is clear that a ‘naive’ generalisation cannot work, since one can add to any solution of such a Neumann problem an arbitrary constant and again obtains a solution.

The main theorem of this paper is as follows.

**Theorem 1.1.** *Let  $\Omega \subset \mathbb{R}^d$  be a bounded connected open set. Let  $r \in (2, \infty)$  and suppose that  $W^{1,2}(\Omega) \subset L^r(\Omega)$ . Let  $\mu: \Omega \rightarrow \mathbb{R}^{d \times d}$  be a bounded measurable function. Suppose there exists a  $\nu > 0$  such that*

$$\operatorname{Re} \sum_{k,\ell=1}^d \mu(x) \xi_k \bar{\xi}_\ell \geq \nu |\xi|^2$$

for all  $\xi \in \mathbb{C}^d$  and almost all  $x \in \Omega$ . Define  $\mathcal{A}: W^{1,2}(\Omega) \rightarrow (W^{1,2}(\Omega))^*$  by

$$\mathcal{A}(u, v) = \int_{\Omega} \mu \nabla u \cdot \bar{\nabla} v.$$

Let  $q \in (d, \infty)$  and suppose that  $\frac{1}{2} - \frac{1}{q} > \frac{1}{r}$ . If  $u \in W^{1,2}(\Omega)$  with  $\mathcal{A}u \in (W^{1,p}(\Omega))^*$ , where  $p$  is the dual exponent of  $q$ , then  $u \in L^\infty(\Omega)$ .

More precisely, for all  $T \in (W^{1,p}(\Omega))^*$  with  $T(\mathbb{1}) = 0$  there is a unique  $u \in W^{1,2}(\Omega)$  with  $\int_{\Omega} u = 0$  satisfying  $\mathcal{A}u = T$ . Moreover, there exists a  $c > 0$  independent of  $T$  such that  $\|u\|_{L^\infty(\Omega)} \leq c \|T\|_{(W^{1,p}(\Omega))^*}$ .

We emphasise that the Sobolev embedding  $W^{1,2}(\Omega) \subset L^r(\Omega)$  assumption is very weak. If  $2^*$  is the first Sobolev exponent, that is  $\frac{1}{2^*} = \frac{1}{2} - \frac{1}{d}$ , then it follows from scaling that  $r \leq 2^*$ . The assumption

$\frac{1}{2} - \frac{1}{q} > \frac{1}{r}$  implies that  $q > d$ . It is well known that there is a connection between Sobolev embeddings and solvability of Neumann problems. We exemplarily refer to Maz'ya and Poborchii [MP07, MP09] and [Maz11], Section 6.10.

If  $d \geq 3$ , then the optimal case in our assumption is  $r = 2^*$ , the first Sobolev exponent. Then the condition  $\frac{1}{2} - \frac{1}{q} > \frac{1}{r}$  is merely the condition  $q > d$ , as in the Stampacchia theorem for the Dirichlet boundary condition. This optimal assumption is satisfied for example by any open bounded set which is the finite union of connected  $W^{1,2}$ -extension domains, such as for example Lipschitz domains. Another example is that of a connected John domain ([Boj88], Section 6). If the domain has cusps, then the full Sobolev embedding is usually not available, but the embedding  $W^{1,2}(\Omega) \subset L^r(\Omega)$  still holds for some  $r \in (2, 2^*)$  if the cusps are of polynomial type by [AF03], Theorem 4.51. We also refer to Maz'ya [Maz11], Section 6.9 for more geometric conditions. It is also known that the embedding cannot hold true for any  $r > 2$  if the boundary of  $\Omega$  has cusps of exponential sharpness, see [AF03] Theorem 4.48. Note that in the case of Dirichlet boundary conditions, one always has the optimal embeddings  $W_0^{1,2}(\Omega) \subset L^{2^*}(\Omega)$  if  $d \geq 3$ , and  $W_0^{1,2}(\Omega) \subset L^r(\Omega)$  for all  $r \in (2, \infty)$  if  $d = 2$ .

The proof of Theorem 1.1 follows the ideas of Stampacchia and uses truncations of Sobolev functions. It relies on the Stampacchia lemma ([KS80] Chapter II, Appendix B, Lemma 2.1) and at its heart lies a uniform estimation of the Poincaré constants of the truncations of mean value free Sobolev functions, Lemma 3.3 below.

We also prove that the pure Neumann operator  $\mathcal{A}$  admits optimal Sobolev regularity in the setting of Theorem 1.1 for  $q$  sufficiently close to 2. This means that the domain of the part of the operator  $\mathcal{A}$  in  $(W^{1,p}(\Omega))^*$  coincides with  $W_{\perp}^{1,q}(\Omega)$ , the mean value free functions in  $W^{1,q}(\Omega)$ , where again  $p$  is the dual exponent to  $q$ . The result relies on interpolation and the Šneĭberg stability theorem. We refer to Theorem 4.3 below.

The outline of this paper is as follows. In Section 2 we show that a Sobolev embedding implies a Poincaré inequality on any  $L^p$ -space. We use this in Section 3 to adapt the argument of Stampacchia to deduce the boundedness as stated in Theorem 1.1. In Section 4 we derive optimal Sobolev regularity results for  $\mathcal{A}$  and some consequences of these based on the results in Section 2.

We conclude with an example. We formally attach the following boundary value problem to the equation  $\mathcal{A}u = T$  with  $T \in (W^{1,p}(\Omega))^*$  as in Theorem 1.1:

$$\begin{aligned} -\operatorname{div}(\mu \nabla u) &= f \quad \text{in } \Omega, \\ -n \cdot \mu \nabla u &= g \quad \text{on } \partial\Omega, \end{aligned}$$

where  $f \in L^s(\Omega)$  and  $g \in L^t(\partial\Omega; \mathcal{H}_{d-1})$  for appropriate values of  $s$  and  $t$ , where  $n$  is the normal. Since  $T$  is only supposed to be a functional on  $W^{1,p}(\Omega)$ , inhomogeneous boundary data is allowed. For the foregoing boundary value problem,  $T$  takes the form

$$T(v) = \int_{\Omega} f \bar{v} + \int_{\partial\Omega} g \bar{\tau v} \, d\mathcal{H}_{d-1},$$

where  $\tau$  is the trace operator onto  $\partial\Omega$ . If the domain  $\Omega$  is sufficiently regular to allow the application of the divergence theorem and to admit a suitable trace operator, this formulation and its connection to  $\mathcal{A}u = T$  can be made rigorous, see Ciarlet ([Cia78], Chapter 1.2) or [GGZ74], Chapter 2.2. A particular case would be that of a Lipschitz graph domain  $\Omega$ .

## 2 Sobolev and Poincaré

We first show that a Sobolev type embedding extrapolates to compactness of the inclusion map  $W^{1,p}(\Omega) \subset L^p(\Omega)$ .

**Lemma 2.1.** *Let  $\Omega \subset \mathbb{R}^d$  be open and bounded. Let  $q \in (1, \infty)$  and suppose there exists a  $\delta > 0$  such that  $W^{1,q}(\Omega) \subset L^{q+\delta}(\Omega)$ . Let  $p \in (1, \infty)$ . Then the inclusion  $W^{1,p}(\Omega) \subset L^p(\Omega)$  is compact. Moreover, there exists a  $\delta' > 0$  such that  $W^{1,p}(\Omega) \subset L^{p+\delta'}(\Omega)$ .*

*Proof.* We show that there exists an  $s > p$  such that  $W^{1,p}(\Omega) \subset L^s(\Omega)$ . Then the compactness of the inclusion  $W^{1,p}(\Omega) \subset L^p(\Omega)$  follows as in [Dan02] Lemma 7.1. Suppose that  $p \in (1, q)$  (the case  $p \in (q, \infty)$  is similar). Fix  $r \in (1, p)$ . It follows from Liu–Tai [LT97] Theorem 9 that the real interpolation space  $(W^{1,1}(\Omega), W^{1,\infty}(\Omega))_{1-\frac{1}{r}, r} = W^{1,t}(\Omega)$  for all  $t \in (1, \infty)$ . Here  $W^{1,\infty}(\Omega)$  is the Sobolev space of all  $L^\infty(\Omega)$  functions whose weak partial derivatives are also  $L^\infty(\Omega)$  functions. Let  $\theta \in (0, 1)$  be such that  $\frac{1}{p} = \frac{1-\theta}{r} + \frac{\theta}{q}$ . Then by complex interpolation

$$\begin{aligned} [W^{1,r}(\Omega), W^{1,q}(\Omega)]_\theta &= \left[ (W^{1,1}(\Omega), W^{1,\infty}(\Omega))_{1-\frac{1}{r}, r}, (W^{1,1}(\Omega), W^{1,\infty}(\Omega))_{1-\frac{1}{q}, q} \right]_\theta \\ &= (W^{1,1}(\Omega), W^{1,\infty}(\Omega))_{1-\frac{1}{p}, p} = W^{1,p}(\Omega), \end{aligned} \quad (1)$$

where we used the reiteration theorem [BL76] Theorem 4.7.2 in the second step. The inclusions  $W^{1,r}(\Omega) \rightarrow L^r(\Omega)$  and  $W^{1,q}(\Omega) \rightarrow L^{q+\delta}(\Omega)$  are continuous. Hence by complex interpolation one deduces that  $W^{1,p}(\Omega) \subset L^s(\Omega)$ , where  $\frac{1}{s} = \frac{1-\theta}{r} + \frac{\theta}{q+\delta} < \frac{1-\theta}{r} + \frac{\theta}{q} = \frac{1}{p}$ . Note that  $s > p$  as required.  $\square$

Arguing as in Ziemer [Zie89] Theorem 4.4.2 one obtains a Poincaré inequality from the compact inclusion  $W^{1,p}(\Omega) \subset L^p(\Omega)$ .

**Proposition 2.2.** *Let  $\Omega \subset \mathbb{R}^d$  be open, bounded and connected. Let  $p \in (1, \infty)$  and suppose that the inclusion  $W^{1,p}(\Omega) \subset L^p(\Omega)$  is compact. Let  $\Omega_0 \subset \Omega$  be measurable and suppose that the Lebesgue measure  $|\Omega_0| > 0$ . Then there exists a  $c > 0$  such that*

$$\|u\|_p \leq c \|\nabla u\|_p$$

for all  $u \in W^{1,p}(\Omega)$  with  $\int_{\Omega_0} u = 0$ .

*Proof.* Suppose not. Then for all  $n \in \mathbb{N}$  there exists a  $u_n \in W^{1,p}(\Omega)$  such that  $\|u_n\|_p > n \|\nabla u_n\|_p$  and  $\int_{\Omega_0} u_n = 0$ . Without loss of generality  $\|u_n\|_p = 1$  for all  $n \in \mathbb{N}$ . Then  $\|\nabla u_n\|_p \leq \frac{1}{n}$ . So the sequence  $(u_n)_{n \in \mathbb{N}}$  is bounded in  $W^{1,p}(\Omega)$ . Passing to a subsequence if necessary there exists a  $u \in W^{1,p}(\Omega)$  such that  $\lim u_n = u$  weakly in  $W^{1,p}(\Omega)$ . Then  $\lim u_n = u$  strongly in  $L^p(\Omega)$  and  $\int_{\Omega_0} u = 0$ . Moreover  $\|u\|_p = 1$  and  $u \neq 0$ . Next  $\|\nabla u\|_p \leq \liminf_{n \rightarrow \infty} \|\nabla u_n\|_p = 0$ . Since  $\Omega$  is connected it follows that  $u$  is constant by [Zie89] Corollary 2.1.9. Because  $\int_{\Omega_0} u = 0$  and  $|\Omega_0| > 0$  one deduces that  $u = 0$ . This is a contradiction.  $\square$

If  $\Omega \subset \mathbb{R}^d$  is a bounded open set and  $p \in (1, \infty)$ , then we define

$$W_{\perp}^{1,p}(\Omega) = \left\{ u \in W^{1,p}(\Omega) : \int_{\Omega} u = 0 \right\}.$$

It follows from Proposition 2.2 that  $W_{\perp}^{1,p}(\Omega)$  equipped with the norm  $u \mapsto \|\nabla u\|_p$  is a Banach space.

**Corollary 2.3.** *Let  $\Omega \subset \mathbb{R}^d$  be open, bounded and connected. Let  $p \in (1, \infty)$  and suppose that the inclusion  $W^{1,p}(\Omega) \subset L^p(\Omega)$  is compact. Define  $\|\cdot\| : W^{1,p}(\Omega) \rightarrow [0, \infty)$  by  $\|u\| = \|\nabla u\|_p + \left| \int_{\Omega} u \right|$ . Then one has the following.*

(a) *The function  $\|\cdot\|$  is a norm on  $W^{1,p}(\Omega)$  which is equivalent to  $\|\cdot\|_{W^{1,p}(\Omega)}$ .*

(b) The map

$$P: u \mapsto u - \frac{1}{|\Omega|} \int_{\Omega} u$$

is a projection from  $W^{1,p}(\Omega)$  onto  $W_{\perp}^{1,p}(\Omega)$ . In particular,

$$u \mapsto \left( \frac{1}{|\Omega|} \int_{\Omega} u, u - \frac{1}{|\Omega|} \int_{\Omega} u \right)$$

is a topological isomorphism from  $W^{1,p}(\Omega)$  onto  $\mathbb{C} \oplus W_{\perp}^{1,p}(\Omega)$ .

*Proof.* By Proposition 2.2 there exists a  $c > 0$  such that  $\|u\|_p \leq c \|\nabla u\|_p$  for all  $u \in W_{\perp}^{1,p}(\Omega)$ . If  $u \in W^{1,p}(\Omega)$ , then

$$\begin{aligned} \|u\|_p &\leq \left\| u - \frac{1}{|\Omega|} \int_{\Omega} u \right\|_p + \left\| \frac{1}{|\Omega|} \int_{\Omega} u \right\|_p \\ &\leq c \left\| \nabla \left( u - \frac{1}{|\Omega|} \int_{\Omega} u \right) \right\|_p + |\Omega|^{-1+\frac{1}{p}} \left| \int_{\Omega} u \right| \\ &= c \|\nabla u\|_p + |\Omega|^{-1+\frac{1}{p}} \left| \int_{\Omega} u \right| \end{aligned}$$

and the lemma follows easily.  $\square$

**Proposition 2.4.** Let  $\Omega \subset \mathbb{R}^d$  be open, bounded and connected. Let  $p \in (1, \infty)$  and suppose that the inclusion  $W^{1,p}(\Omega) \subset L^p(\Omega)$  is compact. Then for all  $T \in (W^{1,p}(\Omega))^*$  there exist  $\kappa \in \mathbb{C}$  and  $f_1, \dots, f_d \in L^q(\Omega)$  such that

$$\langle T, u \rangle_{(W^{1,p}(\Omega))^* \times W^{1,p}(\Omega)} = \kappa \int_{\Omega} \bar{u} + \sum_{j=1}^d \int_{\Omega} f_j \bar{\partial}_j u$$

for all  $u \in W^{1,p}(\Omega)$ , where  $q$  is the dual exponent of  $p$ .

*Proof.* Using Corollary 2.3(b) it suffices to show that for all  $S \in (W_{\perp}^{1,p}(\Omega))^*$  there exist  $f_1, \dots, f_d \in L^q(\Omega)$  such that

$$\langle S, u \rangle_{(W_{\perp}^{1,p}(\Omega))^* \times W_{\perp}^{1,p}(\Omega)} = \sum_{j=1}^d \int_{\Omega} f_j \bar{\partial}_j u$$

for all  $u \in W_{\perp}^{1,p}(\Omega)$ , where  $u \mapsto \|\nabla u\|_p$  is the norm on  $W_{\perp}^{1,p}(\Omega)$ . Consider the subspace  $M = \{\nabla u: u \in W_{\perp}^{1,p}(\Omega)\}$  in  $L^p(\Omega)^d$ . Define  $F: M \rightarrow \mathbb{C}$  by  $F(\nabla u) = Su$ . Then  $F$  is well-defined and continuous. Therefore by Hahn–Banach there exists an extension  $\tilde{F} \in (L^p(\Omega)^d)^*$  of  $F$ . The rest of the proof is straight forward.  $\square$

### 3 Proof of Theorem 1.1

In this section we prove Theorem 1.1. Let  $\Omega \subset \mathbb{R}^d$  be a bounded connected open set. Let  $\mu: \Omega \rightarrow \mathbb{R}^{d \times d}$  be a bounded measurable function. We suppose that  $\mu$  is **elliptic**, that is there exists a  $\nu > 0$  such that

$$\operatorname{Re} \sum_{k,\ell=1}^d \mu(x) \xi_k \bar{\xi}_{\ell} \geq \nu |\xi|^2$$

for all  $\xi \in \mathbb{C}^d$  and almost all  $x \in \Omega$ . Let  $r > 2$  and suppose that  $W^{1,2}(\Omega) \subset L^r(\Omega)$ . Define  $\mathcal{A}: W^{1,2}(\Omega) \rightarrow (W^{1,2}(\Omega))^*$  by

$$\mathcal{A}(u, v) = \int_{\Omega} \mu \nabla u \cdot \overline{\nabla v}.$$

Recall that  $W_{\perp}^{1,p}(\Omega) = \{u \in W^{1,p}(\Omega) : \int_{\Omega} u = 0\}$  for all  $p \in (1, \infty)$ . If  $q \in (1, \infty)$  then we define

$$W_{\emptyset}^{-1,q}(\Omega) = (W^{1,p}(\Omega))^*,$$

where  $p$  is the dual exponent of  $q$ . Moreover, we define

$$W_{\perp}^{-1,q}(\Omega) = \{T \in W_{\emptyset}^{-1,q}(\Omega) : T(\mathbb{1}) = 0\}.$$

Clearly  $\mathcal{A}u \in W_{\perp}^{-1,2}(\Omega)$  for all  $u \in W^{1,2}(\Omega)$  and  $\ker \mathcal{A} = \mathbb{C}\mathbb{1}$  since  $\Omega$  is connected. Define  $\mathcal{A}_{\perp}: W_{\perp}^{1,2}(\Omega) \rightarrow W_{\perp}^{-1,2}(\Omega)$  by  $\mathcal{A}_{\perp}u = \mathcal{A}u$ . Then  $\mathcal{A}_{\perp}$  is injective. We next show that it is also surjective and  $W_{\perp}^{-1,2}(\Omega) = (W_{\perp}^{1,2}(\Omega))^*$ , up to isomorphism.

**Proposition 3.1.** *The map  $\mathcal{A}_{\perp}$  is a topological isomorphism.*

*Proof.* Define the form  $\mathfrak{b}: W_{\perp}^{1,2}(\Omega) \times W_{\perp}^{1,2}(\Omega) \rightarrow \mathbb{C}$  by

$$\mathfrak{b}(u, v) = \int_{\Omega} \mu \nabla u \cdot \overline{\nabla v}.$$

Then  $\mathfrak{b}$  is a continuous coercive sesquilinear form by Lemma 2.1 and Proposition 2.2. Let  $\mathcal{B}: W_{\perp}^{1,2}(\Omega) \rightarrow (W_{\perp}^{1,2}(\Omega))^*$  be such that  $\mathfrak{b}(u, v) = \langle \mathcal{B}u, v \rangle_{(W_{\perp}^{1,2}(\Omega))^* \times W_{\perp}^{1,2}(\Omega)}$  for all  $u, v \in W_{\perp}^{1,2}(\Omega)$ . Then  $\mathcal{B}$  is surjective by the Lax–Milgram theorem. Let  $T \in W_{\perp}^{-1,2}(\Omega)$ . Then  $T \in W_{\emptyset}^{-1,2}(\Omega) = (W^{1,2}(\Omega))^*$ . Let  $\tilde{T} = T|_{W_{\perp}^{1,2}(\Omega)}$ . Then  $\tilde{T} \in (W_{\perp}^{1,2}(\Omega))^*$ . Hence there is a  $u \in W_{\perp}^{1,2}(\Omega)$  such that  $\mathcal{B}u = \tilde{T}$ . If  $v \in W_{\perp}^{1,2}(\Omega)$ , then

$$\langle \mathcal{A}u, v \rangle_{W_{\emptyset}^{-1,2}(\Omega) \times W^{1,2}(\Omega)} = \mathfrak{b}(u, v) = \langle \mathcal{B}u, v \rangle_{(W_{\perp}^{1,2}(\Omega))^* \times W_{\perp}^{1,2}(\Omega)} = \tilde{T}(v) = T(v).$$

Since  $\langle \mathcal{A}u, \mathbb{1} \rangle_{W_{\emptyset}^{-1,2}(\Omega) \times W^{1,2}(\Omega)} = 0 = T(\mathbb{1})$  it follows by linearity and Corollary 2.3(b) that  $\mathcal{A}_{\perp}u = \mathcal{A}u = T$ .  $\square$

As a main tool for the proof of Theorem 1.1 we need truncations of Sobolev functions, which we consider next.

For all  $u \in W^{1,2}(\Omega, \mathbb{R})$  and  $k \in [0, \infty)$  define  $\zeta_{u,k} = (\operatorname{sgn} u) (|u| - k)^+$ . If no confusion is possible then we write  $\zeta_k = \zeta_{u,k}$ . Moreover, define  $A_k = \{x \in \Omega : |u(x)| > k\} = [|u| > k]$ .

**Lemma 3.2.** *Let  $u \in W^{1,2}(\Omega, \mathbb{R})$ . Then one has the following.*

- (a)  $\zeta_k \in W^{1,2}(\Omega)$  for all  $k \in [0, \infty)$ .
- (b)  $\mathbb{1}_{A_k} D_j u = \mathbb{1}_{A_k} D_j \zeta_k$  for all  $j \in \{1, \dots, d\}$  and  $k \in [0, \infty)$ .
- (c) The map  $k \mapsto \zeta_k$  is continuous from  $[0, \infty)$  into  $W^{1,2}(\Omega)$ .
- (d) If  $k \in [0, \infty)$ , then the map  $v \mapsto \zeta_{v,k}$  is continuous from  $W^{1,2}(\Omega, \mathbb{R})$  into  $W^{1,2}(\Omega)$ .

*Proof.* ‘(a)’ and ‘(b)’. Note that  $\zeta_k = (u^+ - k)^+ - (u^- - k)^+$ . Then the statements follow from [GT83] Lemma 7.6.

‘(c)’. This follows from the Lebesgue dominated convergence theorem.

‘(d)’. This follows from (a) and [MM79] Theorem 1.  $\square$

A key estimate for the proof of Theorem 1.1 is the next lemma.

**Lemma 3.3.** *Let  $u \in W_{\perp}^{1,2}(\Omega, \mathbb{R})$ . Then there exists a  $\gamma \geq 0$  such that  $\|\zeta_k\|_2 \leq \gamma \|\nabla \zeta_k\|_2$  for all  $k \in [0, \infty)$ .*

*Proof.* We split the proof into two cases depending whether  $u$  is bounded or not.

**Case 1.** Suppose  $u$  is unbounded. If  $k \in [0, \infty)$  and  $\|\nabla \zeta_k\|_2 = 0$ , then  $\zeta_k$  is constant and consequently  $u$  is bounded, which is a contradiction. Hence  $\|\nabla \zeta_k\|_2 \neq 0$  for all  $k \in [0, \infty)$ . Since both  $k \mapsto \|\zeta_k\|_2$  and  $k \mapsto \|\nabla \zeta_k\|_2$  are continuous on  $[0, \infty)$  by Lemma 3.2(c), it suffices to show that

$$\limsup_{k \rightarrow \infty} \frac{\|\zeta_k\|_2}{\|\nabla \zeta_k\|_2} \leq 1. \quad (2)$$

Suppose that (2) is false. Then there exists a sequence  $(k_n)_{n \in \mathbb{N}}$  in  $\mathbb{R}$  such that  $k_n \geq n$  for all  $n \in \mathbb{N}$  and  $\|\zeta_{k_n}\|_2 > \|\nabla \zeta_{k_n}\|_2$  for all  $n \in \mathbb{N}$ . Define  $v_n = \|\zeta_{k_n}\|_2^{-1} \zeta_{k_n}$  for all  $n \in \mathbb{N}$ . Then  $v_n \in W^{1,2}(\Omega)$ ,  $\|v_n\|_2 = 1$  and  $\|\nabla v_n\|_2 \leq 1$  for all  $n \in \mathbb{N}$ . So the sequence  $(v_n)_{n \in \mathbb{N}}$  is bounded in  $W^{1,2}(\Omega)$ . Passing to a subsequence if necessary we may assume that there is a  $v \in W^{1,2}(\Omega)$  such that  $\lim v_n = v$  weakly in  $W^{1,2}(\Omega)$ . Then  $\lim v_n = v$  in  $L^2(\Omega)$ . So  $\|v\|_2 = 1$  and in particular  $v \neq 0$ . But  $v(x) = \lim_{n \rightarrow \infty} v_n(x) = 0$  for almost every  $x \in \Omega$ . This is a contradiction.

**Case 2.** Suppose  $u$  is bounded. Without loss of generality we may assume that  $u \neq 0$ . Let  $k \in [0, \|u\|_{\infty})$  and suppose that  $\|\nabla \zeta_k\|_2 = 0$ . Then  $\zeta_k$  is constant, say  $\delta$ . If  $\delta = 0$ , then  $|u| \leq k$  a.e., which is not possible since  $k < \|u\|_{\infty}$ . Suppose  $\delta > 0$ . Note that  $\zeta_k(x) \leq 0 < \delta$  for all  $x \in \Omega$  with  $u(x) \leq k$ . So  $u(x) = k + \delta$  for all  $x \in \Omega$ . But then  $\int_{\Omega} u \neq 0$ . Similarly  $\delta < 0$  gives a contradiction. Hence  $\|\nabla \zeta_k\|_2 \neq 0$  for all  $k \in [0, \|u\|_{\infty})$ .

Arguing as in Case 1 and using Lemma 3.2(c) it follows that for all  $k_1 \in (0, \|u\|_{\infty})$  there exists a  $c_1 > 0$  such that  $\|\zeta_k\|_2 \leq c_1 \|\nabla \zeta_k\|_2$  for all  $k \in [0, k_1]$ .

Finally we show that there exist  $k_0 \in (0, \|u\|_{\infty})$  and  $c_0 > 0$  such that  $\|\zeta_k\|_2 \leq c_0 \|\nabla \zeta_k\|_2$  for all  $k \in (k_0, \infty)$ . If  $|u| = \|u\|_{\infty}$  a.e., then  $[|u| = \|u\|_{\infty}] = \frac{1}{2} |\Omega| > 0$ , where we use that  $\int_{\Omega} u = 0$ . Then  $w = \mathbf{1}_{[|u| = \|u\|_{\infty}]} u = u \vee 0 \in W^{1,2}(\Omega)$ . Using [GT83] Lemma 7.7 we deduce that  $\nabla w = 0$  a.e. and this implies that  $[|u| = \|u\|_{\infty}] \in \{0, |\Omega|\}$ , which is a contradiction. Hence there is a  $k_0 \in (0, \|u\|_{\infty})$  such that  $[|u| \leq k_0] > 0$ . Write  $\Omega_0 = [|u| \leq k_0]$ . By Lemma 2.1 and Proposition 2.2 there exists a  $c_0 > 0$  such that  $\|v\|_2 \leq c_0 \|\nabla v\|_2$  for all  $v \in W^{1,2}(\Omega)$  with  $\int_{\Omega_0} v = 0$ . If  $k \in (k_0, \infty)$ , then  $\zeta_k(x) = 0$  for all  $x \in \Omega_0$ , so  $\int_{\Omega_0} \zeta_k = 0$ . Hence  $\|\zeta_k\|_2 \leq c_0 \|\nabla \zeta_k\|_2$ .  $\square$

For all  $u \in W_{\perp}^{1,2}(\Omega, \mathbb{R})$  define  $\gamma_u \in [0, \infty)$  to be the minimum of all  $\gamma \geq 0$  such that  $\|\zeta_k\|_2 \leq \gamma \|\nabla \zeta_k\|_2$  for all  $k \in [0, \infty)$ . Recall that  $r > 2$  is such that  $W^{1,2}(\Omega) \subset L^r(\Omega)$ .

**Proposition 3.4.** *Let  $u \in W_{\perp}^{1,2}(\Omega, \mathbb{R})$  and  $q > d$  with  $\frac{1}{2} - \frac{1}{q} > \frac{1}{r}$ . Further let  $f_1, \dots, f_d \in L^q(\Omega)$  and suppose that  $\langle \mathcal{A}u, v \rangle_{W_0^{-1,2}(\Omega) \times W^{1,2}(\Omega)} = \sum_{j=1}^d (f_j, \partial_j v)_2$  for all  $v \in W^{1,2}(\Omega)$ . Then  $u \in L^{\infty}(\Omega)$ . Moreover*

$$\|u\|_{\infty} \leq 2^{(\frac{1}{2} - \frac{1}{q})/\delta} \frac{E}{\nu} \sqrt{(1 + \gamma_u^2) |\Omega|^{\delta}} \left( \sum_{j=1}^d \|f_j\|_q^2 \right)^{\frac{1}{2}},$$

where  $\delta = \frac{1}{2} - \frac{1}{q} - \frac{1}{r} > 0$  and  $\nu$  is the ellipticity constant of  $\mu$ . Finally,  $E > 0$  is such that  $\|v\|_r \leq E \|v\|_{W^{1,2}(\Omega)}$  for all  $v \in W^{1,2}(\Omega)$ .

*Proof.* For all  $k \in [0, \infty)$  define  $\zeta_k = (\text{sgn } u) (|u| - k)^+ \in W^{1,2}(\Omega)$  and  $A_k = [|u| > k]$  as before. Let  $k \in [0, \infty)$ . Then

$$\nu \|\nabla \zeta_k\|_2^2 \leq \int_{\Omega} \mu \nabla \zeta_k \cdot \nabla \zeta_k = \int_{\Omega} \mu \nabla u \cdot \nabla \zeta_k = \sum_{j=1}^d \int_{A_k} f_j \partial_j \zeta_k$$



$$\begin{aligned} &\leq \left( \sum_{j=1}^d \int_{A_k} |f_j|^2 \right)^{1/2} \|\nabla \zeta_k\|_2 \\ &\leq \frac{\nu}{2} \|\nabla \zeta_k\|_2^2 + \frac{1}{2\nu} \sum_{j=1}^d \int_{A_k} |f_j|^2. \end{aligned}$$

Hence

$$\|\nabla \zeta_k\|_2^2 \leq \frac{1}{\nu^2} \sum_{j=1}^d \int_{A_k} |f_j|^2 \leq \frac{|A_k|^{1-\frac{2}{q}}}{\nu^2} \sum_{j=1}^d \|f_j\|_q^2.$$

By assumption  $W^{1,2}(\Omega) \subset L^r(\Omega)$ . Then

$$\begin{aligned} \left( \int_{A_k} (|u| - k)^r \right)^{\frac{2}{r}} &= \|\zeta_k\|_{L^r(\Omega)}^2 \\ &\leq E^2 \|\zeta_k\|_{W^{1,2}(\Omega)}^2 = E^2 \left( \|\zeta_k\|_2^2 + \|\nabla \zeta_k\|_2^2 \right) \\ &\leq E^2 (1 + \gamma_u^2) \frac{|A_k|^{1-\frac{2}{q}}}{\nu^2} \sum_{j=1}^d \|f_j\|_q^2. \end{aligned}$$

Next let  $h, k \in [0, \infty)$  with  $h > k$ . Then  $A_h \subset A_k$  and

$$(h - k)^2 |A_h|^{\frac{2}{r}} \leq \left( \int_{A_h} (|u| - k)^r \right)^{\frac{2}{r}} \leq \left( \int_{A_k} (|u| - k)^r \right)^{\frac{2}{r}} \leq E^2 (1 + \gamma_u^2) \frac{|A_k|^{1-\frac{2}{q}}}{\nu^2} \sum_{j=1}^d \|f_j\|_q^2.$$

Equivalently

$$|A_h| \leq \frac{1}{(h - k)^r} \left( \frac{E}{\nu} \right)^r (1 + \gamma_u^2)^{\frac{r}{2}} \left( \sum_{j=1}^d \|f_j\|_q^2 \right)^{\frac{r}{2}} |A_k|^{(1-\frac{2}{q})\frac{r}{2}} \dots\dots$$

Due to  $(1 - \frac{2}{q})\frac{r}{2} = (\frac{1}{2} - \frac{1}{q})r > 1$  by assumption, it now follows from the Stampacchia lemma ([KS80] Chapter II, Appendix B, Lemma 2.1) that  $u \in L^\infty(\Omega)$  and

$$\|u\|_\infty \leq 2^{(\frac{1}{2}-\frac{1}{q})/\delta} \frac{E}{\nu} \sqrt{(1 + \gamma_u^2)|\Omega|^\delta} \left( \sum_{j=1}^d \|f_j\|_q^2 \right)^{\frac{1}{2}}.$$

This completes the proof of the proposition.  $\square$

**Proof of Theorem 1.1.** Let  $u \in W^{1,2}(\Omega)$  be such that  $\mathcal{A}u \in (W^{1,p}(\Omega))^*$ , where  $p$  is the dual exponent of  $q$ . By Lemma 2.1 the inclusion  $W^{1,p}(\Omega) \subset L^p(\Omega)$  is compact. Hence by Proposition 2.4 there exist  $\kappa \in \mathbb{C}$  and  $f_1, \dots, f_d \in L^q(\Omega)$  such that

$$\langle \mathcal{A}u, v \rangle_{(W^{1,p}(\Omega))^* \times W^{1,p}(\Omega)} = \kappa \int_{\Omega} \bar{v} + \sum_{j=1}^d \int_{\Omega} f_j \bar{\partial}_j v$$

for all  $v \in W^{1,p}(\Omega)$ . Choosing  $v = \mathbb{1}$  one deduces that  $\kappa = 0$  and  $\mathcal{A}u \in W_{\perp}^{-1,2}(\Omega)$ . Without loss of generality we may assume that  $u \in W_{\perp}^{1,2}(\Omega)$ . Moreover, we may also assume that  $u$  is real valued. Now apply Proposition 3.4 to obtain  $u \in L^\infty(\Omega)$ .

If we start with  $T \in W_{\perp}^{-1,q}(\Omega)$ , then there exists a unique  $u \in W_{\perp}^{1,2}(\Omega)$  such that  $\mathcal{A}u = T$  by Proposition 3.1. Then  $\mathcal{A}u \in W_{\perp}^{-1,q}(\Omega) \subset (W^{1,p}(\Omega))^*$ , so  $u \in L^{\infty}(\Omega)$  by the above.

For the estimate it suffices to show that the map  $T \mapsto u$  has closed graph in the space  $W_{\perp}^{-1,q}(\Omega) \times L^{\infty}(\Omega)$ . Let  $T, T_1, T_2, \dots \in W_{\perp}^{-1,q}(\Omega)$  and  $u \in L^{\infty}(\Omega)$ . Suppose that  $\lim T_n = T$  in  $W_{\perp}^{-1,q}(\Omega)$  and  $\lim(\mathcal{A}_{\perp})^{-1}T_n = u$  in  $L^{\infty}(\Omega)$ . Then  $\lim T_n = T$  in  $W_{\perp}^{-1,2}(\Omega)$ , so  $\lim(\mathcal{A}_{\perp})^{-1}T_n = (\mathcal{A}_{\perp})^{-1}T$  in  $W_{\perp}^{1,2}(\Omega)$  and hence also in  $L^2(\Omega)$ . But  $\lim(\mathcal{A}_{\perp})^{-1}T_n = u$  in  $L^{\infty}(\Omega)$  and therefore also in  $L^2(\Omega)$ . Consequently  $(\mathcal{A}_{\perp})^{-1}T = u$  as required.  $\square$

## 4 Interpolation and maximal Sobolev regularity

In this section, we use the structure of  $W_{\perp}^{1,p}(\Omega)$  as a complemented subspace of  $W^{1,p}(\Omega)$  to establish interpolation results. Optimal Sobolev regularity for the pure Neumann operator  $\mathcal{A}_{\perp}$  for  $p$  close to 2 also follows. This is particularly interesting for space dimension  $d = 2$ . The first step is to show that  $W_{\perp}^{1,p}(\Omega)$  and  $W_{\perp}^{-1,p}(\Omega)$  form an interpolation scale with respect to  $p$ .

**Proposition 4.1.** *Let  $\Omega \subset \mathbb{R}^d$  be open and bounded. Let  $p_0, p_1 \in (1, \infty)$ ,  $\theta \in (0, 1)$  and set  $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$ . Then*

$$[W_{\perp}^{1,p_0}(\Omega), W_{\perp}^{1,p_1}(\Omega)]_{\theta} = (W_{\perp}^{1,p_0}(\Omega), W_{\perp}^{1,p_1}(\Omega))_{\theta,p} = W_{\perp}^{1,p}(\Omega)$$

and

$$[W_{\perp}^{-1,p_0}(\Omega), W_{\perp}^{-1,p_1}(\Omega)]_{\theta} = (W_{\perp}^{-1,p_0}(\Omega), W_{\perp}^{-1,p_1}(\Omega))_{\theta,p} = W_{\perp}^{-1,p}(\Omega).$$

*Proof.* It follows from (1) that

$$[W^{1,p_0}(\Omega), W^{1,p_1}(\Omega)]_{\theta} = W^{1,p}(\Omega).$$

Arguing as in (1), but using the reiteration theorem for real interpolation [BL76], Theorem 3.5.3 one deduces similarly

$$(W^{1,p_0}(\Omega), W^{1,p_1}(\Omega))_{\theta,p} = W^{1,p}(\Omega).$$

Note that for all  $r \in (1, \infty)$  the projection  $P$  in Corollary 2.3(b) maps  $W^{1,r}(\Omega)$  onto  $W_{\perp}^{1,r}(\Omega)$ , so  $W_{\perp}^{1,r}(\Omega)$  is a complemented subspace of  $W^{1,r}(\Omega)$ . We further observe that  $W_{\perp}^{1,p_i}(\Omega) = W^{1,p_i}(\Omega) \cap W_{\perp}^{1,\min(p_0,p_1)}(\Omega)$  for  $i = 1, 2$ . Thus, interpolation theory for complemented subspaces ([Tri78] Theorem 1.17.1.1) shows that

$$[W_{\perp}^{1,p_0}(\Omega), W_{\perp}^{1,p_1}(\Omega)]_{\theta} = (W_{\perp}^{1,p_0}(\Omega), W_{\perp}^{1,p_1}(\Omega))_{\theta,p} = W^{1,p}(\Omega) \cap W_{\perp}^{1,\min(p_0,p_1)}(\Omega) = W_{\perp}^{1,p}(\Omega).$$

Concerning the dual spaces, it is easy to see that for all  $q \in (1, \infty)$  the operator  $T \mapsto T - \frac{1}{|\Omega|} \langle T, \mathbb{1} \rangle \mathbb{1}$  is a projection from  $W^{-1,q}(\Omega)$  onto  $W_{\perp}^{-1,q}(\Omega)$ . Hence the assertion follows with the same argument and the duality properties of the real and complex interpolation functors, see [Tri78] Subsections 1.11.2 and 1.11.3.  $\square$

The first result derived from Proposition 4.1 together with Theorem 1.1 is the following mapping property for  $\mathcal{A}_{\perp}^{-1}$  on the  $W_{\perp}^{-1,p}(\Omega)$  spaces for all  $p > 2$ . Note that we do not require that  $p > d$ .

**Corollary 4.2.** *Let  $\Omega \subset \mathbb{R}^d$  be a bounded connected open set. Let  $r \in (2, \infty)$  and suppose that  $W^{1,2}(\Omega) \subset L^r(\Omega)$ . Let further  $q \in (d, \infty)$  and suppose that  $\frac{1}{2} - \frac{1}{q} > \frac{1}{r}$ . Let  $p \in (2, q)$ . Let  $\mu: \Omega \rightarrow \mathbb{R}^{d \times d}$  be a bounded measurable elliptic function and let  $\mathcal{A}: W^{1,2}(\Omega) \rightarrow (W^{1,2}(\Omega))^*$  be the associated operator. Then  $\mathcal{A}_{\perp}^{-1}$  maps  $W_{\perp}^{-1,p}(\Omega)$  into  $L^s(\Omega)$ , where  $\frac{1}{s} = \frac{1-\theta}{r}$  and  $\theta \in (0, 1)$  is such that  $\frac{1}{p} = \frac{1-\theta}{2} + \frac{\theta}{q}$ .*

*Proof.* The operator  $\mathcal{A}_\perp^{-1}$  maps  $W_\perp^{-1,2}(\Omega)$  continuously into  $W_\perp^{1,2}(\Omega) \subset L^r(\Omega)$  by Proposition 3.1. Moreover,  $\mathcal{A}_\perp^{-1}$  maps  $W_\perp^{-1,q}(\Omega)$  continuously into  $L^\infty(\Omega)$  by Theorem 1.1. Now use complex interpolation and Proposition 4.1.  $\square$

Due to Proposition 4.1 and the work from the previous sections, a maximal Sobolev regularity result for  $p$  close to 2 follows by an application of the Šneĭberg stability theorem.

**Theorem 4.3.** *Let  $\Omega \subset \mathbb{R}^d$  be a bounded connected open set. Let  $r \in (2, \infty)$  and suppose that  $W^{1,2}(\Omega) \subset L^r(\Omega)$ . Let  $\mu: \Omega \rightarrow \mathbb{R}^{d \times d}$  be a bounded measurable elliptic function and let  $\mathcal{A}: W^{1,2}(\Omega) \rightarrow (W^{1,2}(\Omega))^*$  be the associated operator. Then there exists a  $\delta > 0$  such that  $\mathcal{A}_\perp$  is a topological isomorphism between  $W_\perp^{1,p}(\Omega)$  and  $W_\perp^{-1,p}(\Omega)$  for all  $p \in (2 - \delta, 2 + \delta)$ .*

*Proof.* Under the assumptions,  $\mathcal{A}_\perp$  is a topological isomorphism between  $W_\perp^{1,2}(\Omega)$  and  $W_\perp^{-1,2}(\Omega)$  by Proposition 3.1. Proposition 4.1 shows that these spaces are simultaneous interpolation spaces in the  $W_\perp^{1,p}(\Omega)$  and  $W_\perp^{-1,p}(\Omega)$  scale. The Šneĭberg stability theorem [Sne74] implies that there is a  $\delta > 0$  such that  $\mathcal{A}_\perp$  remains an isomorphism between  $W_\perp^{1,p}(\Omega)$  and  $W_\perp^{-1,p}(\Omega)$  for all  $p \in (2 - \delta, 2 + \delta)$ .  $\square$

There exist quantitative results on the size of  $\delta$  derived from the Šneĭberg result in Theorem 4.3. We refer to [ABES19], Appendix A. The most crucial information is that one can choose  $\delta$  to depend only on the ellipticity constant and the upper bound  $\|\mu\|_\infty$  of the coefficient function  $\mu$  of  $\mathcal{A}$ . Moreover, for all  $p \in (2 - \delta, 2 + \delta)$ , the operator norm  $\|\mathcal{A}_\perp^{-1}\|_{W_\perp^{-1,p}(\Omega) \rightarrow W_\perp^{1,p}(\Omega)}$  can be estimated by a multiple of  $\|\mathcal{A}_\perp^{-1}\|_{W_\perp^{-1,2}(\Omega) \rightarrow W_\perp^{1,2}(\Omega)}$ . By Lax-Milgram, the latter can be estimated by  $1/\nu$ , where  $\nu$  is the ellipticity constant of  $\mu$ .

Theorem 4.3 yields further corollaries for  $d = 2$ .

**Corollary 4.4.** *Adopt the notation and assumptions of Theorem 4.3. Let  $d = 2$ . Let  $q \in (2, 2 + \delta)$  and suppose that  $\frac{1}{2} - \frac{1}{q} > \frac{1}{r}$ . Then  $W^{1,s}(\Omega) \subset L^\infty(\Omega)$  for all  $s \geq q$ .*

*Proof.* It follows from Theorem 1.1 that  $\mathcal{A}_\perp^{-1}W_\perp^{-1,q}(\Omega) \subset L^\infty(\Omega)$ . But  $\mathcal{A}_\perp^{-1}W_\perp^{-1,q}(\Omega) = W_\perp^{1,q}(\Omega)$  by Theorem 4.3. Since  $W^{1,q}(\Omega) = W_\perp^{1,q}(\Omega) + \mathbb{C}\mathbb{1}$  the corollary follows.  $\square$

The parameter  $\delta$  in the previous corollary depends on the coefficient function  $\mu$  via the Šneĭberg theorem. If  $\Omega$  is smooth enough so that the full Sobolev embedding for  $W^{1,2}(\Omega)$  is available, then no coefficient function is needed (at least in the formulation of the corollary).

**Corollary 4.5.** *Let  $\Omega \subset \mathbb{R}^2$  be a bounded connected open set. Suppose that  $W^{1,2}(\Omega) \subset L^r(\Omega)$  for all  $r \in (2, \infty)$ . Then  $W^{1,s}(\Omega) \subset L^\infty(\Omega)$  for all  $s \in (2, \infty)$ .*

*Proof.* Choose  $\mu = I$ . Let  $\delta > 0$  be as in Theorem 4.3. Let  $s \in (2, \infty)$ . Then there exists a  $q \in (2, 2 + \delta) \cap (2, s]$ . Now apply Corollary 4.4.  $\square$

The third corollary concerns Hölder regularity of solutions  $u$  of  $\mathcal{A}_\perp u = T$  with  $T \in W_\perp^{-1,q}(\Omega)$  for  $q > 2$  and a uniform estimate. We do not pass through Theorem 1.1 for this result. The price to pay is a Sobolev embedding assumption for the Hölder space similar to the one in Theorem 1.1.

**Corollary 4.6.** *Let  $\Omega \subset \mathbb{R}^2$  be a bounded connected open set. Suppose that for all  $q \in (2, \infty)$  there exists an  $\alpha \in (0, 1)$  such that  $W^{1,q}(\Omega) \subset C^\alpha(\overline{\Omega})$ . Let  $\mu: \Omega \rightarrow \mathbb{R}^{d \times d}$  be a bounded measurable elliptic function and let  $\mathcal{A}: W^{1,2}(\Omega) \rightarrow (W^{1,2}(\Omega))^*$  be the associated operator. Then one has the following.*

(a) *For all  $q \in (2, \infty)$  there exists an  $\alpha \in (0, 1)$  such that  $\mathcal{A}_\perp^{-1}W_\perp^{-1,q}(\Omega) \subset C^\alpha(\overline{\Omega})$ .*

(b) For all  $q \in (2, \infty)$  and  $R > 0$  the set

$$\{\mathcal{A}_{\perp}^{-1}(T) : T \in W_{\perp}^{-1,q}(\Omega) \text{ and } \|T\|_{W_{\perp}^{-1,q}(\Omega)} \leq R\}$$

is compact in  $C(\overline{\Omega})$ .

*Proof.* ‘(a)’. Since  $C^{\alpha}(\overline{\Omega}) \subset L^{\infty}(\Omega) \subset L^r(\Omega)$  for all  $\alpha \in (0, 1)$  and  $r \in (1, \infty)$ , it follows from Lemma 2.1 that there exists an  $r \in (2, \infty)$  such that  $W^{1,2}(\Omega) \subset L^r(\Omega)$ . Let  $\delta > 0$  be as in Theorem 4.3. Let  $s \in (2, 2+\delta) \cap (2, q]$ . By assumption there exists an  $\alpha \in (0, 1)$  such that  $W^{1,s}(\Omega) \subset C^{\alpha}(\overline{\Omega})$ . Then  $\mathcal{A}_{\perp}^{-1}W_{\perp}^{-1,q}(\Omega) \subset \mathcal{A}_{\perp}^{-1}W_{\perp}^{-1,s}(\Omega) = W_{\perp}^{1,s}(\Omega) \subset C^{\alpha}(\overline{\Omega})$ .

‘(b)’. This follows from statement (a) and the Arzelà–Ascoli theorem.  $\square$

The situation for the Hölder-Sobolev embedding assumption in Corollary 4.6 is similar to the assumption on the Sobolev embedding in Theorem 1.1. It is satisfied for example when for all  $q \in (2, \infty)$  the domain  $\Omega$  is a connected  $W^{1,q}$ -extension domain and then one can choose  $\alpha = 1 - 2/q$ , but there are also examples of (non-extension) domains with sufficiently regular cusps where the assumption is satisfied in the weaker form, see [AF03] Theorem 4.53. Note however that the optimal embedding for  $W^{1,q}(\Omega)$  into the Hölder space of order  $1 - 2/q$  implies the  $W^{1,r}$ -extension property for all  $r > q$ , see [Kos98] Theorem A.

## References

- [ABES19] AUSCHER, P., BORTZ, S., EGERT, M. and SAARI, O., Nonlocal self-improving properties: a functional analytic approach. *Tunisian J. Math.* **1** (2019), 151–183.
- [AF03] ADAMS, R. A. and FOURNIER, J. J. F., *Sobolev spaces*. Second edition, Pure and Applied Mathematics 140. Elsevier/Academic Press, Amsterdam, 2003.
- [BL76] BERGH, J. and LÖFSTRÖM, J., *Interpolation spaces. An introduction*. Grundlehren der mathematischen Wissenschaften 223. Springer-Verlag, Berlin etc., 1976.
- [Boj88] BOJARSKI, B., Remarks on Sobolev imbedding inequalities. In LAINE, L., RICKMAN, S. and SORVALI, T., eds., *Complex analysis, Joensuu 1987*, Lecture Notes in Math. 1351, 52–68. Springer, Berlin, 1988.
- [Cia78] CIARLET, P. G., *The finite element method for elliptic problems*. Studies in Mathematics and its Applications 4. North-Holland, Amsterdam, 1978.
- [Dan02] DANERS, D., A priori estimates for solutions to elliptic equations on non-smooth domains. *Proc. Roy. Soc. Edinburgh Sect. A* **132** (2002), 793–813.
- [GGZ74] GAJEWSKI, H., GRÖGER, K. and ZACHARIAS, K., *Nichtlineare Operatorgleichungen und Operatordifferentialgleichungen*. Mathematische Lehrbücher und Monographien, II. Abteilung Mathematische Monographien 38. Akademie-Verlag, Berlin, 1974.
- [GT83] GILBARG, D. and TRUDINGER, N. S., *Elliptic partial differential equations of second order*. Second edition, Grundlehren der mathematischen Wissenschaften 224. Springer-Verlag, Berlin etc., 1983.
- [Kos98] KOSKELA, P., Extensions and imbeddings. *J. Funct. Anal.* **159** (1998), 369–383.
- [KS80] KINDERLEHRER, D. and STAMPACCHIA, G., *An introduction to variational inequalities and their applications*. Pure and Applied Mathematics 88. Academic Press, New York, 1980.

- [LT97] LIU, F.-C. and TAI, W.-S., Lusin properties and interpolation of Sobolev spaces. *Topol. Methods Nonlinear Anal.* **9** (1997), 163–177.
- [Maz11] MAZ'YA, V. G., *Sobolev spaces with applications to elliptic partial differential equations*. Second edition, Grundlehren der mathematischen Wissenschaften 342. Springer-Verlag, Berlin etc., 2011.
- [MM79] MARCUS, M. and MIZEL, V. J., Every superposition operator mapping one Sobolev space into another is continuous. *J. Funct. Anal.* **33** (1979), 217–229.
- [MP07] MAZ'YA, V. G. and POBORCHIĬ, S., Imbedding theorems for Sobolev spaces on domains with peak and on Hölder domains. *St. Petersburg Math. J.* **18** (2007), 583–605.
- [MP09] ———, On the solvability of the Neumann problem in domains with peak. *St. Petersburg Math. J.* **20** (2009), 757–790.
- [Sne74] SNEIBERG, I. J., Spectral properties of linear operators in interpolation families of Banach spaces. *Mat. Issled.* **9**, No. 2 (1974), 214–229.
- [Sta65] STAMPACCHIA, G., Le problème de Dirichlet pour les équations elliptiques du second ordre à coefficients discontinus. *Ann. Inst. Fourier, Grenoble* **15** (1965), 189–258.
- [Tri78] TRIEBEL, H., *Interpolation theory, function spaces, differential operators*. North-Holland, Amsterdam, 1978.
- [Trö10] TRÖLTZSCH, F., *Optimal control of partial differential equations. Theory, methods and applications*. Graduate Studies in Mathematics 112. American Mathematical Society, Providence, RI, 2010.
- [Zie89] ZIEMER, W. P., *Weakly differentiable functions*. Graduate Texts in Mathematics 120. Springer-Verlag, New York, 1989.