

Weierstraß–Institut für Angewandte Analysis und Stochastik

im Forschungsverbund Berlin e.V.

Convergence criteria for waveform iteration methods applied to partitioned DAE systems in chemical process simulation

Tino Michael, Jürgen Borchardt

submitted: 28th August 1996

Weierstrass Institute
for Applied Analysis
and Stochastics
Mohrenstraße 39
D – 10117 Berlin
Germany

Preprint No. 262
Berlin 1996

1991 Mathematics Subject Classification. Primary 65L05, 65H10, 65F50, 65Y05; Secondary 80A30, 92E20.

Key words and phrases. Algebraic–differential equations, waveform iteration, partitioned systems, parallelization of numerical methods, chemical process simulation.

Edited by
Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS)
Mohrenstraße 39
D — 10117 Berlin
Germany

Fax: + 49 30 2044975
e-mail (X.400): c=de;a=d400-gw;p=WIAS-BERLIN;s=preprint
e-mail (Internet): preprint@wias-berlin.de

1 Introduction

During the last two decades dynamic processes simulation has become an indispensable tool for the design and operation of complex chemical plants. The rigorous modeling of the chemical processes results in large-scale systems of differential and algebraic equations (DAE). Using an appropriate modeling the differential index of the resulting DAE systems can be restricted to index 1 in many cases [BP1]. For details about the numerical solution of DAE systems and the index problem we refer to [BC1, GH1].

For large-scale systems cost-effective numerical methods are needed. Under special modeling assumptions waveform iteration methods can be used for the concurrent dynamic process simulation [SM1]. These methods, also called dynamic iteration methods or Picard-type methods are well suited for the implementation on parallel computers.

Since the beginning of the 80'th the main application field for waveform iteration methods has been the circuit simulation of MOS digital integrated circuits [LRS]. Investigations of waveform iteration methods for systems of ordinary differential equations (ODE) [WS1, JP1, Br1, BB1] as well as for systems of differential algebraic equations [LRS, Mi1, SL1, SM1] are known. Convergence under certain conditions on the mathematical problem has been proven in [Br1] for explicit ODE systems and in [Mi1] for linear DAE systems of arbitrary index. Lelarsmee et al. [LRS] and Schneider [Sc1] proved convergence for a semiexplicit type of waveform iteration for semiexplicit DAEs of Index 1.

In this paper we consider the application of block waveform iteration methods to initial value problems for implicit DAE systems of index 1 arising in chemical process simulation. Block waveform iteration methods permit the concurrent treatment of blocks of subsystems of the entire system with multirate integration techniques gaining a coarse granular parallelism. Their convergence properties strongly depend on the assignment of variables to equations and the partitioning of the system into subsystem blocks.

First we proof convergence for waveform iteration methods applied to semiexplicit DAE systems of index 1. The convergence conditions are given in a blocksystem oriented manner, i.e. only the blocksystems have to satisfy some corresponding conditions. Then we show that the convergence conditions are fulfilled for a simplified modeling of distillation columns. An assignment and partitioning algorithm is given, which takes the requirements of the convergence theorems into account.

Based on this investigations a prototype of a waveform-iteration code has been implemented and tested by means of examples included in the user package of the chemical process simulator SPEEDUP [PS1].

2 Waveformiteration

We consider the initial value problem (IVP) for a semiexplicit DAE system:

$$\begin{aligned} \dot{x}_1 &= \tilde{f}(x_1, x_2, t) \\ 0 &= \tilde{h}(x_1, x_2, t) \\ x(t_0) &= x_0, \quad t \in [t_0, t_e] \end{aligned} \quad (2.1)$$

with $x = (x_1, x_2)^T$, x_0 the consistent initial value, $\tilde{f} : \mathbb{R}^k \times \mathbb{R}^l \times [t_0, t_e] \rightarrow \mathbb{R}^k$, $\tilde{h} : \mathbb{R}^k \times \mathbb{R}^l \times [t_0, t_e] \rightarrow \mathbb{R}^l$ and $n := k + l$.

Let the time interval be denoted by $T := [t_0, t_e]$.

For index 1 problems (2.1) the inverse $(\tilde{h}_{x_2})^{-1}$ exists in a neighbourhood of the solution of the IVP. In the following we assume that $(\tilde{h}_{x_2})^{-1}$ exists in the whole domain.

Definition 1 Let $f : \mathbb{R}^k \times \mathbb{R}^l \times \mathbb{R}^k \times \mathbb{R}^l \times T \rightarrow \mathbb{R}^k$, $h : \mathbb{R}^k \times \mathbb{R}^l \times \mathbb{R}^k \times \mathbb{R}^l \times T \rightarrow \mathbb{R}^l$ be functions with $f(x_1, x_2, x_1, x_2, t) \equiv \tilde{f}(x_1, x_2, t)$, $h(x_1, x_2, x_1, x_2, t) \equiv \tilde{h}(x_1, x_2, t)$. The general form of the continuous waveform iteration for the problem (2.1) is then given by

$$\begin{aligned} \dot{x}_1^k &= f(x_1^k, x_2^k, x_1^{k-1}(t), x_2^{k-1}(t), t) \\ 0 &= h(x_1^k, x_2^k, x_1^{k-1}(t), x_2^{k-1}(t), t) \\ x^k(t_0) &= x_0 \end{aligned} \quad (2.2)$$

for $k = 1, 2, \dots$ with $x^0 : T \rightarrow \mathbb{R}^n$ be any arbitrary approximation for the beginning of the iteration which satisfies $x^0(t_0) = x_0$.

For the discrete waveform iteration on subintervals, the so called "windows", we refer to [Br1].

After partitioning of (2.1) into a set of r semiexplicit index 1 problems with $x_1 = (x_{11}, \dots, x_{1r})^T$, $x_{1i} \in \mathbb{R}^{k_i}$, $\sum k_i = k$, $x_2 = (x_{21}, \dots, x_{2r})^T$, $x_{2i} \in \mathbb{R}^{l_i}$, $\sum l_i = l$, $n_i := k_i + l_i$, $\sum n_i = n$, $x_{-i} := (x_{1,i}, x_{2,i})^T$, $x = (x_{-1}, \dots, x_{-r})^T$, one obtains the following block structured representation

$$\begin{aligned} \dot{x}_{1i} &= \tilde{f}_i(x_{11}, \dots, x_{1r}, x_{21}, \dots, x_{2r}, t) \\ 0 &= \tilde{h}_i(x_{11}, \dots, x_{1r}, x_{21}, \dots, x_{2r}, t) \\ x_{-i}(t_0) &= x_{-i,0}, \quad i = 1(1)r. \end{aligned} \quad (2.3)$$

The corresponding block waveform iteration is then given by

$$\begin{aligned} \dot{x}_{1i}^k &= f_i(x_1^k, x_2^k, x_1^{k-1}(t), x_2^{k-1}(t), t), \\ 0 &= h_i(x_1^k, x_2^k, x_1^{k-1}(t), x_2^{k-1}(t), t), \\ x_{-i}^k(t_0) &= x_{-i,0}, \quad i = 1(1)r. \end{aligned} \quad (2.4)$$

Two special cases are the block Gauss Seidel waveform iteration where f_i and g_i are given by

$$\begin{aligned} f_i(x_1, x_2, y_1, y_2, t) &:= \tilde{f}_i(x_{11}, x_{21}, \dots, x_{1i}, x_{2i}, y_{1i+1}, y_{2i+1}, \dots, y_{1r}, y_{2r}, t) \\ h_i(x_1, x_2, y_1, y_2, t) &:= \tilde{h}_i(x_{11}, x_{21}, \dots, x_{1i}, x_{2i}, y_{1i+1}, y_{2i+1}, \dots, y_{1r}, y_{2r}, t) \end{aligned} \quad (2.5)$$

and the block Jacobi waveform iteration with

$$\begin{aligned} f_i(x_1, x_2, y_1, y_2, t) &:= \tilde{f}_i(y_{11}, y_{21}, \dots, y_{1i-1}, y_{2i-1}, x_{1i}, x_{2i}, y_{1i+2}, y_{2i+2}, \dots, y_{1r}, y_{2r}, t) \\ h_i(x_1, x_2, y_1, y_2, t) &:= \tilde{h}_i(y_{11}, y_{21}, \dots, y_{1i-1}, y_{2i-1}, x_{1i}, x_{2i}, y_{1i+1}, y_{2i+1}, \dots, y_{1r}, y_{2r}, t). \end{aligned} \quad (2.6)$$

The block Jacobi waveform iteration is especially suitable for parallelization.

3 Convergence considerations

For problems (2.1) with index 1 the existence of an unique solution $x_2 = \tilde{g}(x_1, t)$ of the algebraic equations $0 = \tilde{h}(x_1, x_2, t)$ follows from the implicit function theorem. Because each blocksystem in (2.3) is assumed to have index 1, every \tilde{h}_i is solvable for x_{2i} as well. Hence in the cases of block Gauss Seidel- and of block Jacobi waveform iteration the h_i are also solvable for x_{2i} .

Then (2.4) can formally be rewritten as

$$\begin{aligned} \dot{x}_{1i}^k &= f_i(x_1^k, x_2^k, x_1^{k-1}(t), x_2^{k-1}(t), t), \\ x_{2i}^k &= g_i(x_1^k, x_2^k, x_1^{k-1}(t), x_2^{k-1}(t), t), \\ x_{-i}^k(t_0) &= x_{-i,0}, \quad i = 1(1)r, \end{aligned} \quad (3.1)$$

In the case of the block Jacobi waveform iteration the g_i are even independent of x_2 , so that 3.1 has the representation

$$\begin{aligned} \dot{x}_{1i}^k &= f_i(x_1^k, x_2^k, x_1^{k-1}(t), x_2^{k-1}(t), t), \\ x_{2i}^k &= g_i(x_1^k, x_1^{k-1}(t), x_2^{k-1}(t), t), \\ x_{-i}^k(t_0) &= x_{-i,0}, \quad i = 1(1)r. \end{aligned} \quad (3.2)$$

Let $C(T, \mathbb{R}^n)$ and $C^1(T, \mathbb{R}^n)$ be the spaces of the continuous and the continuous differentiable n dimensional functions defined on T .

The direct sum $\bar{Z} := C^1(T, \mathbb{R}^k) \oplus C(T, \mathbb{R}^l)$ is a set of ordered pairs defined by

$$\bar{Z} := \{(z_1, z_2)^T : z_1 \in C^1(T, \mathbb{R}^k), z_2 \in C(T, \mathbb{R}^l)\}.$$

Let $|\cdot|_m$ be any norm in \mathbb{R}^m , then for $z_2 \in C(T, \mathbb{R}^l)$ and $z_1 \in C^1(T, \mathbb{R}^k)$ exponentially weighted norms for arbitrary but fixed $\alpha > 0$ are defined by

$$\begin{aligned}\|z_2\|_{\alpha, l} &:= \sup_{t \in T} (e^{-\alpha(t-t_0)} |z_2(t)|_l) \\ \|z_1\|_k &:= |x(t_0)|_k + \frac{1 - e^{-\alpha(t_e-t_0)}}{\alpha} \|\dot{z}_1\|_{\alpha, k}.\end{aligned}$$

With this norms a norm for $z \in \bar{Z}$ is defined by

$$\|z\|_{\bar{Z}} := \max\{\|z_1\|_k, \|z_2\|_{\alpha, l}\}.$$

Lemma 1 $\{\bar{Z}, \|\cdot\|_{\bar{Z}}\}$ is a Banach-space.

Proof:

It has to be shown, that the limit $z = (z_1, z_2)^T$ of an arbitrary Cauchy sequence $(z^i)_{i=1}^{\infty}$ in $\{\bar{Z}, \|\cdot\|_{\bar{Z}}\}$ is an element of \bar{Z} .

$\forall \epsilon > 0$ there $\exists i_0$ such that $\|z^q - z^p\|_{\bar{Z}} < \epsilon$ holds $\forall q, p > i_0$.

We have that

$$\begin{aligned}\epsilon &> \|z^q - z^p\|_{\bar{Z}} = \|(z_1^q, z_2^q)^T - (z_1^p, z_2^p)^T\|_{\bar{Z}} = \max\{\|z_1^q - z_1^p\|_k, \|z_2^q - z_2^p\|_{\alpha, l}\} \\ &= \max\left\{|z_1^q(t_0) - z_1^p(t_0)|_k + \frac{1 - e^{-\alpha(t_e-t_0)}}{\alpha} \|\dot{z}_1^q - \dot{z}_1^p\|_{\alpha, k}, \|z_2^q - z_2^p\|_{\alpha, l}\right\} \\ &= \max\left\{|z_1^q(t_0) - z_1^p(t_0)|_k + \frac{1 - e^{-\alpha(t_e-t_0)}}{\alpha} \sup_{t \in T} (e^{-\alpha(t-t_0)} |z_1^q(t) - z_1^p(t)|_k), \right. \\ &\quad \left. \sup_{t \in T} (e^{-\alpha(t-t_0)} |z_2^q(t) - z_2^p(t)|_l)\right\} \\ &\geq \max\left\{|z_1^q(t_0) - z_1^p(t_0)|_k + \frac{1 - e^{-\alpha(t_e-t_0)}}{\alpha} e^{-\alpha(t_e-t_0)} \sup_{t \in T} |z_1^q(t) - z_1^p(t)|_k, \right. \\ &\quad \left. e^{-\alpha(t_e-t_0)} \sup_{t \in T} |z_2^q(t) - z_2^p(t)|_l\right\}.\end{aligned}$$

Since $\epsilon \rightarrow 0$ for growing q and p it follows that

$$|z_1^q(t_0) - z_1^p(t_0)|_k \rightarrow 0, \quad \sup_{t \in T} |z_1^q(t) - z_1^p(t)|_k \rightarrow 0 \quad \text{and} \quad \sup_{t \in T} |z_2^q(t) - z_2^p(t)|_l \rightarrow 0.$$

So we have uniform convergence for $\dot{z}_1^i \rightarrow \dot{z}_1$, $z_2^i \rightarrow z_2$ and thus $\dot{z}_1 \in C(T, \mathbb{R}^k)$, $z_2 \in C(T, \mathbb{R}^l)$. Because also $z_1^i(t_0) \rightarrow z_1(t_0)$ it follows the uniform convergence $z_1^i \rightarrow z_1$ and hence $z_1 \in C^1(T, \mathbb{R}^k)$.

□

With respect to the partitioned system (2.3) we have $z_1 = (z_{11}, \dots, z_{1r})^T$ and $z_2 = (z_{21}, \dots, z_{2r})^T$. With $z_{\cdot i} := (z_{1i}, z_{2i})^T$ we can define function sets Z_i corresponding to

\bar{Z} and norms $\|\cdot\|_{Z_i}$ corresponding to $\|\cdot\|_{\bar{Z}}$. Because of Lemma 1, the spaces $\{Z_i, \|\cdot\|_{Z_i}\}$, $i = 1, \dots, r$, are Banach-spaces. For $z = (z_1, \dots, z_r)^T \in Z$, $Z := Z_1 \oplus Z_2 \oplus \dots \oplus Z_r$, a norm $\|\cdot\|_Z$ is defined by

$$\|z\|_Z := \max_{i \in \{1, \dots, r\}} \{\|z_i\|_{Z_i}\}.$$

One easily can show that $\{Z, \|\cdot\|_Z\}$ is a Banach-space.

For $x, y \in Z$, an operator $O : Z \times Z \rightarrow Z$ is defined by

$$O(x, y)(t) = O(x_1, x_2, y_1, y_2)(t) := \begin{bmatrix} x_{1,0} + \int_{t_0}^t f(x_1, x_2, y_1, y_2, s) ds \\ g(x_1, x_2, y_1, y_2, t) \end{bmatrix}.$$

From the definition of the operator O we find that the block waveform iteration (3.1) has the formal representation

$$x^k = O(x^k, x^{k-1})$$

with $x^k(t_0) = x_0$ and an arbitrary starting function $x^0 \in Z$.

We now consider the operator equation

$$x = O(x, y).$$

If this equation is uniquely solvable for x , an operator

$$P : Z \rightarrow Z \tag{3.3}$$

is defined where x is determined by the equation $x = P(y)$.

In the following Lemma 2 the Banach space $\{Z, \|\cdot\|_Z\}$ and the operator O are more generally defined. The coinciding notation is chosen because the above defined Banach space $\{Z, \|\cdot\|_Z\}$ and operator O are applicable to this Lemma, what will be shown later.

The proof of the Lemma 2 also can be found in [Br1].

Lemma 2 *Let $\{Z, \|\cdot\|_Z\}$ be a Banach-space and let the operator $O : Z \times Z \rightarrow Z$ satisfy the following assumptions:*

a1 *O is globally Lipschitz continuous, i.e. there exist constants $K_x, K_y \geq 0$ such that for all x, y, \bar{x}, \bar{y}*

$$\|O(x, y) - O(\bar{x}, \bar{y})\|_Z \leq K_x \|x - \bar{x}\|_Z + K_y \|y - \bar{y}\|_Z,$$

a2 $0 < K_x + K_y < 1$.

Then

f1 For each $y \in Z$ the equation $x = O(x, y)$ is uniquely solvable for x . The operator P defined in (3.3) is contractive with a Lipschitz-constant $M_y = \frac{K_y}{1-K_x}$.

f2 The sequence (x^k) defined by

$$x^k = O(x^k, x^{k-1})$$

converges to a unique limit x for any starting function $x^0 \in Z$.

Proof:

Let $y \in Z$ be arbitrary but fixed. First it is shown, that the operator

$$O_y := O(\cdot, y), \quad O_y : Z \rightarrow Z$$

is contractive and has a unique fixpoint. From a1 it is known that O_y is Lipschitz continuous with the Lipschitz constant K_x and from a2 we have that $K_x < 1$. Thus the fixpoint theorem of Banach is applicable to O_y which proofs the assertion.

The operator $P : Z \rightarrow Z$ is defined in that way, that P appoints to every $y \in Z$ the fixpoint of O_y .

For arbitrary $y, \bar{y} \in Z$ with $x := P(y)$ and $\bar{x} := P(\bar{y})$ we have

$$\|x - \bar{x}\|_Z = \|O(x, y) - O(\bar{x}, \bar{y})\|_Z \leq K_x \|x - \bar{x}\|_Z + K_y \|y - \bar{y}\|_Z$$

and thus

$$\|x - \bar{x}\|_Z \leq \frac{K_y}{1 - K_x} \|y - \bar{y}\|_Z.$$

Since

$$\|x - \bar{x}\|_Z = \|P(y) - P(\bar{y})\|_Z,$$

P is Lipschitz continuous. From a2 it follows that the Lipschitz constant $\frac{K_y}{1-K_x}$ is smaller than 1, i.e. P is contractive. This proofs f1.

With $x := x^k$ and $y := x^{k-1}$, we have

$$x^k = O(x^k, x^{k-1})$$

and thus

$$x^k = P(x^{k-1}).$$

Because of f1 the fixpoint theorem of Banach is applicable to P what proves f2. □

If $\mathbb{R}^m = \mathbb{R}^{m_1} \oplus \mathbb{R}^{m_2} \oplus \dots \oplus \mathbb{R}^{m_q}$, then an appropriate norm for $w = (w_1, \dots, w_q)^T \in \mathbb{R}^m$ is defined by

$$|w| := \max_{i \in \{1, \dots, q\}} |w_i|_{m_i}$$

Lemma 3 *Let the waveform iteration (3.1) for the partitioned system (2.3) be considered. If the functions f_i and g_i , $i = 1, \dots, r$ fulfill Lipschitz conditions, i.e. for all $\xi = (\xi_1, \xi_2)^T, \bar{\xi} = (\bar{\xi}_1, \bar{\xi}_2)^T, \eta = (\eta_1, \eta_2)^T, \bar{\eta} = (\bar{\eta}_1, \bar{\eta}_2)^T \in \mathbb{R}^k \oplus \mathbb{R}^l, t \in T$ exist positive constants $f_{ix}, f_{iy}, g_{ix}, g_{iy}$ with*

$$|f_i(\xi, \eta, t) - f_i(\bar{\xi}, \bar{\eta}, t)|_{k_i} \leq f_{ix}|\xi - \bar{\xi}| + f_{iy}|\eta - \bar{\eta}| \quad (3.4)$$

$$|g_i(\xi, \eta, t) - g_i(\bar{\xi}, \bar{\eta}, t)|_{l_i} \leq g_{ix}|\xi - \bar{\xi}| + g_{iy}|\eta - \bar{\eta}| \quad (3.5)$$

then the following estimations hold for $x, y, \bar{x}, \bar{y} \in Z$:

$$\begin{aligned} & \|f_i(x_1(\cdot), x_2(\cdot), y_1(\cdot), y_2(\cdot), \cdot) - f_i(\bar{x}_1(\cdot), \bar{x}_2(\cdot), \bar{y}_1(\cdot), \bar{y}_2(\cdot), \cdot)\|_{\alpha, k_i} \\ & \leq f_{ix}\|x - \bar{x}\|_Z + f_{iy}\|y - \bar{y}\|_Z, \end{aligned} \quad (3.6)$$

$$\begin{aligned} & \|g_i(x_1(\cdot), x_2(\cdot), y_1(\cdot), y_2(\cdot), \cdot) - g_i(\bar{x}_1(\cdot), \bar{x}_2(\cdot), \bar{y}_1(\cdot), \bar{y}_2(\cdot), \cdot)\|_{\alpha, l_i} \\ & \leq g_{ix}\|x - \bar{x}\|_Z + g_{iy}\|y - \bar{y}\|_Z. \end{aligned} \quad (3.7)$$

Proof: With $x, y, \bar{x}, \bar{y} \in Z$ one obtains from (3.4)

$$\begin{aligned} & |f_i(x_1(t), x_2(t), y_1(t), y_2(t), t) - f_i(\bar{x}_1(t), \bar{x}_2(t), \bar{y}_1(t), \bar{y}_2(t), t)|_{k_i} \\ & \leq f_{ix} \max_{i \in \{1, \dots, r\}} \{ \max\{|x_{1i}(t) - \bar{x}_{1i}(t)|_{k_i}, |x_{2i}(t) - \bar{x}_{2i}(t)|_{l_i}\} \} + \\ & \quad f_{iy} \max_{i \in \{1, \dots, r\}} \{ \max\{|y_{1i}(t) - \bar{y}_{1i}(t)|_{k_i}, |y_{2i}(t) - \bar{y}_{2i}(t)|_{l_i}\} \} \end{aligned}$$

Let j be the index i which yields the maximum for the x -component. Then we have

$$\begin{aligned} & f_{ix} \max_{i \in \{1, \dots, r\}} \{ \max\{|x_{1i}(t) - \bar{x}_{1i}(t)|_{k_i}, |x_{2i}(t) - \bar{x}_{2i}(t)|_{l_i}\} \} \\ & \leq f_{ix} \max \left\{ |x_{1j}(t_0) - \bar{x}_{1j}(t_0) + \int_{t_0}^t (\dot{x}_{1j}(s) - \dot{\bar{x}}_{1j}(s)) ds|_{k_j}, |x_{2j}(t) - \bar{x}_{2j}(t)|_{l_j} \right\} \\ & \leq f_{ix} \max \left\{ |x_{1j}(t_0) - \bar{x}_{1j}(t_0)| + \int_{t_0}^t e^{\alpha(s-t_0)} e^{-\alpha(s-t_0)} |\dot{x}_{1j}(s) - \dot{\bar{x}}_{1j}(s)|_{k_j} ds, \right. \\ & \quad \left. e^{\alpha(t-t_0)} e^{-\alpha(t-t_0)} |x_{2j}(t) - \bar{x}_{2j}(t)|_{l_j} \right\} \\ & \leq f_{ix} \max \left\{ |x_{1j}(t_0) - \bar{x}_{1j}(t_0)|_{k_j} + \int_{t_0}^t e^{\alpha(s-t_0)} \sup_{s \in T} \{ e^{-\alpha(s-t_0)} |\dot{x}_{1j}(s) - \dot{\bar{x}}_{1j}(s)|_{k_j} \} ds, \right. \\ & \quad \left. e^{\alpha(t-t_0)} \sup_{t \in T} \{ e^{-\alpha(t-t_0)} |x_{2j}(t) - \bar{x}_{2j}(t)|_{l_j} \} \right\} \\ & = f_{ix} \max \left\{ |x_{1j}(t_0) - \bar{x}_{1j}(t_0)|_{k_j} + \|\dot{x}_{1j} - \dot{\bar{x}}_{1j}\|_{\alpha, k_j} \int_{t_0}^t e^{\alpha(s-t_0)} ds, e^{\alpha(t-t_0)} \|x_{2j} - \bar{x}_{2j}\|_{\alpha, l_j} \right\} \\ & = f_{ix} \max \left\{ |x_{1j}(t_0) - \bar{x}_{1j}(t_0)|_{k_j} + \frac{e^{\alpha(t-t_0)} - 1}{\alpha} \|\dot{x}_{1j} - \dot{\bar{x}}_{1j}\|_{\alpha, k_j}, e^{\alpha(t-t_0)} \|x_{2j} - \bar{x}_{2j}\|_{\alpha, l_j} \right\} \\ & \leq f_{ix} e^{\alpha(t-t_0)} \max \left\{ |x_{1j}(t_0) - \bar{x}_{1j}(t_0)|_{k_j} + \frac{1 - e^{-\alpha(t-t_0)}}{\alpha} \|\dot{x}_{1j} - \dot{\bar{x}}_{1j}\|_{\alpha, k_j}, \|x_{2j} - \bar{x}_{2j}\|_{\alpha, l_j} \right\} \\ & = e^{\alpha(t-t_0)} f_{ix} \max \{ \|x_{1j} - \bar{x}_{1j}\|_{k_j}, \|x_{2j} - \bar{x}_{2j}\|_{\alpha, l_j} \}. \end{aligned}$$

Because the same estimate holds for the y -component we obtain

$$\begin{aligned} & |f_i(x_1(t), x_2(t), y_1(t), y_2(t), t) - f_i(\bar{x}_1(t), \bar{x}_2(t), \bar{y}_1(t), \bar{y}_2(t), t)|_{k_i} \\ & \leq e^{\alpha(t-t_0)} (f_{ix} \|x - \bar{x}\|_Z + f_{iy} \|y - \bar{y}\|_Z). \end{aligned}$$

Using the norm definition of $\|\cdot\|_{\alpha, k_i}$ one obtains (3.6).

Going for g through the same procedure as for f one gets (3.7). □

Theorem 1 *Let the assumptions of Lemma 3 be satisfied. If further the Lipschitz constants g_{ix} , g_{iy} of g_i fulfill for every i the condition*

$$g_{ix} + g_{iy} < 1,$$

then the sequence (x^k) obtained by the block waveform iteration (3.1) converges in $\{Z, \|\cdot\|_Z\}$ to the solution of (2.1).

Proof:

We show that the operator O is globally Lipschitz continuous and the Lipschitz constants satisfy the condition **a2** of Lemma 2. Then **f1**, **f2** can be followed.

We have

$$\|O(x, y) - O(\bar{x}, \bar{y})\|_Z = \|z - \bar{z}\|_Z = \max_{i \in \{1, \dots, r\}} \{ \max \{ \|z_{1i} - \bar{z}_{1i}\|_{k_i}, \|z_{2i} - \bar{z}_{2i}\|_{\alpha, l_i} \} \}.$$

Let j be the maximal i . Using the result of Lemma 3, it follows that

$$\begin{aligned} & \|O(x, y) - O(\bar{x}, \bar{y})\|_Z = \\ & = \max \left\{ |z_{1j}(t_0) - \bar{z}_{1j}(t_0)| + \frac{1 - e^{-\alpha(t_e - t_0)}}{\alpha} \|\dot{z}_{1j} - \dot{\bar{z}}_{1j}\|_{\alpha, k_j}, \|z_{2j} - \bar{z}_{2j}\|_{\alpha, l_j} \right\} \\ & = \max \left\{ \frac{1 - e^{-\alpha(t_e - t_0)}}{\alpha} \|f_j(x_1(\cdot), x_2(\cdot), y_1(\cdot), y_2(\cdot), \cdot) - f_j(\bar{x}_1(\cdot), \bar{x}_2(\cdot), \bar{y}_1(\cdot), \bar{y}_2(\cdot), \cdot))\|_{\alpha, k_j}, \right. \\ & \quad \left. \|g_j(x_1(\cdot), x_2(\cdot), y_1(\cdot), y_2(\cdot), \cdot) - g_j(\bar{x}_1(\cdot), \bar{x}_2(\cdot), \bar{y}_1(\cdot), \bar{y}_2(\cdot), \cdot))\|_{\alpha, l_j} \right\} \\ & \leq \max \left\{ \frac{1 - e^{-\alpha(t_e - t_0)}}{\alpha} (f_{jx} \|x - \bar{x}\|_Z + f_{jy} \|y - \bar{y}\|_Z), g_{jx} \|x - \bar{x}\|_Z + g_{jy} \|y - \bar{y}\|_Z \right\}. \end{aligned}$$

Hence for large enough chosen α the Lipschitz constants of O are

$$K_x = g_{jx}, \quad K_y = g_{jy}.$$

Because the assumption $g_{jx} + g_{jy} < 1$ satisfies the condition **a2** of Lemma 2 the proof is completed. □

Let $z_{-i} = (z_{1i}, z_{2i})^T \in Z_i$. Then norms $|\cdot|_0, |\cdot|_1$ usually used for the continuous and continuous differentiable functions are defined by

$$\begin{aligned} |z_{2i}|_0 &:= \sup_{t \in T} \{|z_{2i}(t)|\}, \\ |z_{1i}|_1 &:= |z_{1i}|_0 + |\dot{z}_{1i}|_0. \end{aligned}$$

Further a norm $|\cdot|_{Z_i}$ for $z_{-i} \in Z_i$ is defined by

$$|z_{-i}|_{Z_i} := \max\{|z_{1i}|_1, |z_{2i}|_0\}.$$

Another norm $|\cdot|_Z$ for $z = (z_{-1}, \dots, z_{-r})^T \in Z$ is then defined by

$$|z|_Z := \max\{|z_{-i}|_{Z_i}\}.$$

With this definitions $\{Z, |\cdot|_Z\}$ is a Banach space.

Theorem 2 *With the assumptions of theorem 1 the sequence (x^k) obtained by the block waveform iteration (3.1) converges in $\{Z, |\cdot|_Z\}$ to the solution of (2.1).*

Proof:

Let the fixpoint of P be denoted by x^* . To show that the block waveform iteration (3.1) converges to $z^* \in \{Z, |\cdot|_Z\}$ for $k \rightarrow \infty$ we look for a constant $K > 0$ such that

$$|x^k - x^*|_Z \leq K \|x^k - x^*\|_Z.$$

For $z = (z_{-1}, \dots, z_{-r})^T \in Z$ it holds by definition

$$|z|_Z = \max_{i \in \{1, \dots, r\}} \{\max\{|z_{1i}|_1, |z_{2i}|_0\}\}.$$

With

$$z_{1i}(t) = z_{1i}(t_0) + \int_{t_0}^t \dot{z}_{1i}(s) ds$$

we have

$$\begin{aligned} |z_{1i}(t)|_{k_i} &\leq |z_{1i}(t_0)|_{k_i} + \int_{t_0}^t e^{-\alpha(s-t_0)} \|\dot{z}_{1i}\|_{\alpha, k_i} ds \\ &\leq e^{\alpha(t-t_0)} \left(|z_{1i}(t_0)|_{k_i} + \frac{1 - e^{-\alpha(t_e-t_0)}}{\alpha} \|\dot{z}_{1i}\|_{\alpha, k_i} \right) \end{aligned}$$

and hence

$$|z_{1i}|_0 \leq e^{\alpha(t_e-t_0)} \|\dot{z}_{1i}\|_{k_i}.$$

Further it is

$$\begin{aligned} |\dot{z}_{1i}(t)|_{k_i} &= e^{\alpha(t-t_0)} e^{-\alpha(t-t_0)} |\dot{z}_{1i}(t)|_{k_i} \\ &\leq e^{\alpha(t-t_0)} \|\dot{z}_{1i}\|_{\alpha, k_i} \\ &\leq \frac{\alpha}{1 - e^{-\alpha(t_e-t_0)}} e^{\alpha(t-t_0)} \left(|z_{1i}(t_0)|_{k_i} + \frac{1 - e^{-\alpha(t_e-t_0)}}{\alpha} \|\dot{z}_{1i}\|_{k_i} \right) \end{aligned}$$

and hence

$$|\dot{z}_{1i}|_0 \leq \frac{\alpha}{1 - e^{-\alpha(t_e - t_0)}} e^{\alpha(t_e - t_0)} \|z_{1i}\|_{k_i}.$$

Finally one has

$$|z_{2i}(t)|_{l_i} \leq e^{\alpha(t - t_0)} \|z_{2i}\|_{\alpha, l_i}$$

and thus

$$|z_{2i}|_0 \leq e^{\alpha(t_e - t_0)} \|z_{2i}\|_{\alpha, l_i}$$

what results in

$$|z|_Z \leq \left(1 + \frac{\alpha}{1 - e^{-\alpha(t_e - t_0)}}\right) e^{\alpha(t_e - t_0)} \|z\|_Z. \quad (3.8)$$

Then replacing z by $x^k - x^*$ in (3.8) leads to the assertion. \square

In summary, for the convergence of the block waveform iteration (3.1) for semiexplicit DAE systems of index 1 the Lipschitz continuity of f_i and g_i , $i = 1, \dots, r$, is required, where the Lipschitz constants of g_i satisfy the inequality

$$g_{ix} + g_{iy} < 1.$$

A local approximation for the Lipschitz constants g_{ix} , g_{iy} can be determined by means of the Jacobians $\frac{\partial g_i}{\partial x}$, $\frac{\partial g_i}{\partial y}$ evaluated in a fixed point $(x_1(t^*), x_2(t^*), y_1(t^*), y_2(t^*), t^*)$. In the following these arguments are left away.

We remember that the block systems h_i were solved for $x_{2i} = g_i(x_1, x_2, y_1, y_2)$ with g_i independent of x_{2i} . A deduction from the implicit function theorem yields

$$\begin{bmatrix} \frac{\partial g}{\partial x_1} & \frac{\partial g}{\partial x_2} & \frac{\partial g}{\partial y_1} & \frac{\partial g}{\partial y_2} \end{bmatrix} = \begin{bmatrix} \frac{\partial h}{\partial x_{2i}} \end{bmatrix}^{-1} \begin{bmatrix} \frac{\partial h}{\partial x_1} & \frac{\partial h}{\partial x_2} & \frac{\partial h}{\partial y_1} & \frac{\partial h}{\partial y_2} \end{bmatrix}. \quad (3.9)$$

The constants g_{ix} and g_{iy} are then evaluated for the block Jacobi waveform iteration by

$$g_{ix} = \left| \frac{\partial g_i}{\partial x_1} \right| = \left| \begin{bmatrix} \frac{\partial h_i}{\partial x_{2i}} \end{bmatrix}^{-1} \frac{\partial h_i}{\partial x_{1i}} \right|_{n_{1i}} \quad (3.10)$$

$$g_{iy} = \left| \frac{\partial g_i}{\partial y} \right| = \sum_{j \neq i} \left| \begin{bmatrix} \frac{\partial h_i}{\partial x_{2i}} \end{bmatrix}^{-1} \frac{\partial h_i}{\partial y_{1j}} \right|_{n_{1j}} + \sum_{j \neq i} \left| \begin{bmatrix} \frac{\partial h_i}{\partial x_{2i}} \end{bmatrix}^{-1} \frac{\partial h_i}{\partial y_{2j}} \right|_{n_{2j}}. \quad (3.11)$$

and for the block Gauss Seidel waveform iteration by

$$\begin{aligned} \bar{g}_{ix} &= \left| \frac{\partial g_i}{\partial x} \right| = \sum_{j=1}^i \left| \begin{bmatrix} \frac{\partial h_i}{\partial x_{2i}} \end{bmatrix}^{-1} \frac{\partial h_i}{\partial x_{1j}} \right|_{n_{1j}} + \sum_{j=1}^{i-1} \left| \begin{bmatrix} \frac{\partial h_i}{\partial x_{2i}} \end{bmatrix}^{-1} \frac{\partial h_i}{\partial x_{2j}} \right|_{n_{2j}} \\ \bar{g}_{iy} &= \left| \frac{\partial g_i}{\partial y} \right| = \sum_{j=i+1}^r \left| \begin{bmatrix} \frac{\partial h_i}{\partial x_{2i}} \end{bmatrix}^{-1} \frac{\partial h_i}{\partial y_{1j}} \right|_{n_{1j}} + \sum_{j=i+1}^r \left| \begin{bmatrix} \frac{\partial h_i}{\partial x_{2i}} \end{bmatrix}^{-1} \frac{\partial h_i}{\partial y_{2j}} \right|_{n_{2j}} \end{aligned}$$

Here the matrix norm $|\cdot|$ is induced by the vector norm $|\cdot|$. The coinciding notation for the norms is used for simplicity.

These conditions just require the Jacobian of the discretized problem (2.1) to be block diagonal dominant because the diagonal elements of the Jacobian in the hyperrows, belonging to the differential part $\dot{x}_i - \tilde{f}_i = 0$ of the DAE system, get arbitrary large with reduction of the stepsize. This comes from the stepsize in the denominator of the discretized \dot{x} .

Definition 2 A matrix $A \in \mathbb{R}^{n \times n}$, $A = (A_{ij})_{i,j=1}^M$, $A_{ij} \in \mathbb{R}^{n_i \times n_j}$, $\sum_{i=1}^M n_i = n$, is called block diagonal dominant if

$$\sum_{i \neq j} |A_{ii}^{-1} A_{ij}| \leq 1, \quad i = 1, \dots, M$$

holds.

For an implementation on a computer the evaluation of these Lipschitz constants is still to expansive. In chapter 5 we give a strategy for the partitioning of implicit index 1 problems, which is related to the results of this chapter but requires less computation time due to some heuristics.

4 Mathematical modeling of distillation columns

The modeling of distillation columns like it has been done in [Wo1], [Re1] leads to linear DAE systems of index 1:

$$\begin{aligned} D\dot{u} &= \tilde{f}(u, v, t) \\ 0 &= \tilde{h}(u, v, t) \\ (u(t_0), v(t_0))^T &= (u_0, v_0)^T \end{aligned}$$

where D is a diagonal matrix with constant coefficients.

If the modeling is done with the following restrictions,

- constant molar hold-up HU^L of the liquid phase,
- neglectable hold-up HU^V of the vapor phase,
- temporal constant pressure profile over column height,
- ideal mixing in liquid phase,
- phase equilibrium between vapor and liquid phase with constant tray efficiency coefficient,
- no excessive enthalpy,

one obtains for each tray of the distillation column, except for the bottom tray, the following system of equations:

Material balance for each component $i = 1(1)N$:

$$\begin{aligned} \frac{d(u_{i,j} H U_j^L)}{dt} &= (L_j + U_j + U_{j,n}^R) u_{i,j} - L_{j-1} u_{i,j-1} \\ &+ (V_j + W_j + W_{j,m}^R) v_{i,j} - V_{j+1} v_{i,j+1} - F_j^L u_{i,j}^F \\ &- F_j^V v_{i,j}^F - U_{k,j}^R u_{j,k} - W_{l,j}^R v_{i,l} \end{aligned}$$

Energy balance:

$$\begin{aligned} \frac{d(H_j^L H U_j^L)}{dt} &= (L_j + U_j + U_{j,n}^R) H_j^L - L_{j-1} H_{j-1}^L \\ &+ (V_j + W_j + W_{j,m}^R) H_j^L - V_{j+1} H_{j+1}^V - F_j^L H_{F,j}^L \\ &- F_j^V H_{F,j}^V - U_{k,j}^R H_k^L - W_{l,j}^R H_l^V + Q_j \end{aligned}$$

with

$$\begin{aligned} H_j^L &= \sum_{i=1}^N u_{i,j} h_{i,j}^L, & H_j^V &= \sum_{i=1}^N v_{i,j} h_{i,j}^V, \\ H_{F,j}^L &= \sum_{i=1}^N u_{i,j}^F h_{i,j}^{L,F}, & H_{F,j}^V &= \sum_{i=1}^N v_{i,j}^F h_{i,j}^{V,F}. \end{aligned}$$

Phase equilibrium relation for $i = 1(1)N$:

$$\eta_j k_{i,j} u_{i,j} - v_{i,j} + (1 - \eta_j) v_{i,j+1} = 0.$$

The tray efficiency coefficients η_j of tray j are smaller than 1 and calculated from

$$\eta_j = \frac{v_{i,j} - v_{i,j+1}}{v_{i,j}^* - v_{i,j+1}}.$$

The equilibrium constants $k_{i,j}$ are evaluated by

$$k_{i,j} = \gamma_{i,j} \frac{p_{i,j}^0}{p_j},$$

where the activity coefficients $\gamma_{i,j}$ are given by

$$v_{i,j}^* p_j = u_{i,j} \gamma_{i,j} p_{i,j}^0.$$

The following quantities are for the liquid and vapor phase respectively:

Trivially $\left[\frac{\partial h_j}{\partial x_{2,j}}\right]^{-1} = \text{diag}(-1)$ and hence

$$\begin{aligned} g_{jx_1} + g_{jy} &= \left[\left[\frac{\partial h_j}{\partial x_{2j}} \right]^{-1} \frac{\partial h_j}{\partial x_{1j}} \right]_{n_j} + \left[\left[\frac{\partial h_j}{\partial x_{2j}} \right]^{-1} \frac{\partial h_j}{\partial y_{2,j+1}} \right]_{n_j} \\ &= \left[\begin{array}{ccc} -\eta_j k_{1j} & & \\ & \ddots & \\ & & -\eta_j k_{Nj} \end{array} \right] + \left[\begin{array}{ccc} \eta_j - 1 & & \\ & \ddots & \\ & & \eta_j - 1 \end{array} \right] \\ &= \eta_j k_{-j} + 1 - \eta_j < 1, \end{aligned}$$

for $\max_{i \in \{1, \dots, N\}} \{k_{ij}\} =: k_{-j} < 1$.

Hence the following theorem is proved.

Theorem 3 *For the described mathematical model of a distillation column the block Jacobi waveform iteration (3.2) converges for any partitioning given above.*

5 Implementation

The mathematical modeling of chemical processes in chemical plants leads usually [AT1] to initial value problems for implicit DAE systems

$$\begin{aligned} F(t, y(t), \dot{y}(t), u(t)) &= 0, \quad y(t_0) = y_0, \\ F : \mathbb{R} \times \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^q &\rightarrow \mathbb{R}^n, \quad t \in [t_0, t_e], \end{aligned} \quad (5.1)$$

with $F = (f_1, \dots, f_n)^T$, a given parameter vector function $u(t)$ and the unknown vector function $y(t) = (v_1(t), \dots, v_n(t))^T$.

In many cases during the process of modeling it can be made sure that the index of the system (5.1) is 1. Usually the arising systems are stiff, and their discretization and linearization yield systems of equations with sparse nonsymmetric Jacobian matrices. The systems can comprise several 10 000 equations (e.g. distillation columns) and are structured into subsystems

$$\begin{aligned} F_i(t, y, \dot{y}, u) &= 0, \quad y(t_0) = y_0, \\ F_i : \mathbb{R} \times \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^q &\rightarrow \mathbb{R}^{n_i}, \quad \sum n_i = n, \quad i = 1, \dots, m \end{aligned} \quad (5.2)$$

in accordance with the functional units of the chemical plant.

To apply the block Jacobi waveform iteration there has to be done a one to one assignment of variables to the equations and a partitioning of the system (5.1) in such a way that the waveform iteration is convergent. Because (5.1) is a fully-implicit problem the convergence condition for semiexplicit problems is not applicable. For this we use some heuristic. Like observed before, in case of block Jacobi waveform iteration for semiexplicit index 1 problems the convergence condition $g_{(ix_i)} + g_{(iy)} < 1$

implies the block diagonal dominance of the Jacobian of the discretized problem. So our partitioning algorithm puts priority to getting the Jacobian of the partitioned discretized problem (5.1) block diagonal dominant.

We assume that the hyperrows of the Jacobian $\frac{\partial \tilde{F}_i}{\partial y} = \frac{\partial F_i}{\partial y} + c_i * \frac{\partial F_i}{\partial \tilde{y}}$ can be computed separately. \tilde{F}_i is a discretized subsystem from (5.2) and c_i the corresponding discretization constant dependent on the integration method and stepsize used.

Assigning variables to equations, system (5.2) can be rewritten to

$$F_i(t, x_i(t), \dot{x}_i(t), y_i(t), \dot{y}_i(t), u(t)) = 0, \quad x_i(t_0) = x_{i,0}, \quad i = 1(1)m, \quad (5.3)$$

where x_i is the vector of variables assigned to the subsystem F_i and y_i is the vector remaining if the components x_i are removed from y .

The system is then partitioned by merging subsystems F_i to blocksystems $\mathcal{F}_j = (F_{j_1}, \dots, F_{j_{m_j}})^T$, $\sum m_j = m$. With the notations $X_j = (x_{j_1}, \dots, x_{j_{m_j}})^T$ and the corresponding Y_j , the partitioning leads to

$$\mathcal{F}_j(t, X_j(t), \dot{X}_j(t), Y_j(t), \dot{Y}_j(t), u(t)) = 0, \quad X_j(t_0) = X_{j,0}, \quad j = 1(1)M.$$

The waveform iteration is done over time windows $[t_p, t_{p+1}] \subseteq [t_0, t_e]$, $p = 1, \dots, p_e$, $\cup [t_p, t_{p+1}] = [t_0, t_e]$ [GB1] to speed up the convergence. Here the blocksystems can be solved concurrently using general known methods. We use BDF, Newton's method and sparse matrix solver.

The algorithm reads as

```

do for  $p = 0, 1, 2, \dots$ 
  set  $Y_j^0$  on  $[t_p, t_{p+1}]$  for  $j = 1(1)M$ 
  do for  $k = 1, 2, \dots$ 
    do for  $j = 1(1)M$ 
      solve for  $t \in [t_p, t_{p+1}]$ 

$$\begin{aligned} \mathcal{F}_j(t, X_j^k(t), \dot{X}_j^k(t), Y_j^{k-1}(t), \dot{Y}_j^{k-1}(t), u(t)) &= 0 \\ Y_j^k(t_p) &= \tilde{Y}_j(t_p) \end{aligned}$$

    enddo
  until  $\|Y^k - Y^{k-1}\|_{n_j, [t_p, t_{p+1}]} < \epsilon$ 
enddo

```

For assigning variables to equations as well as merging strongly connected subsystems into blocks we define "weights" for the couplings between equations and subsystems respectively using the Jacobian matrix

$$A := \frac{\partial \tilde{F}(y)}{\partial y} \Big|_{y=\tilde{y}}, \quad A = (a_{pq}) \in \mathbb{R}^{n \times n}$$

of the nonlinear system

$$\tilde{F}(y) = 0, \quad \tilde{F} : \mathbb{R}^n \rightarrow \mathbb{R}^n,$$




Figure 1: DYNEVAP: Jacobian matrix before and after assignment

obtained by the discretization of the DAE system at the time point $t = \tilde{t}$, $\tilde{y} \sim y(\tilde{t})$. During the assignment process each variable v_q is assigned to one and only one equation f_p , such that the resulting assignment $x_i \rightarrow F_i$ is consistent with respect to the state variables for each subsystem i and the $n_i \times n_i$ Jacobians $\frac{\partial \tilde{F}_i}{\partial x_i}$ are at least non singular.

To treat this problem we consider the linear weighted matching problem

$$\sum_{p=1}^n \sum_{q=1}^n w_{pq} s_{pq} \rightarrow \max, \quad \sum_{k=1}^n s_{pk} = 1, \quad \sum_{k=1}^n s_{kq} = 1,$$

$$s_{pq} = \begin{cases} 1 & : \text{if variable } v_q \text{ is assigned to equation } f_p \\ 0 & : \text{else} \end{cases}$$

$$w_{pq} = \begin{cases} 0 & : f_p \text{ depends neither on } v_q \text{ nor on } \dot{v}_q \\ 1 + \frac{|a_{pq}|}{\sum_{r=1}^n |a_{pr}|} & : f_p \text{ depends on } v_q, \text{ but not on } \dot{v}_q \\ 3 + \frac{|a_{pq}|}{\sum_{r=1}^n |a_{pr}|} & : f_p \text{ depends on } \dot{v}_q \end{cases}$$

Starting from the original Jacobian we generate a parametrized directed graph and solve the matching problem with graph algorithms from the package LEDA [Näl]. In Figure 1 the structure of the Jacobi matrix of the example DYNEVAP is given before and after the assignment of the variables to the equations. To merge tightly coupled subsystems to blocks we define "strong" connections between equations and subsystems respectively. We call a row p of the matrix A with $f_p \in F_i$ *dominant with*

respect to subsystem i , if

$$\sum_{q \notin K_i} |a_{pq}| / |a_{pp}| < 1, \text{ with index set } K_i = \{r \mid f_r \in F_i\}.$$

Then the subsystem i is called *strong input to subsystem j* , if there exists a

$$p_j \in \{p \mid f_p \in F_j, \text{ row } p \text{ is not dominant with respect to subsystem } j\},$$

so that

$$\frac{\sum_{k \in K_i} |a_{p_j k}|}{|a_{p_j p_j}|} \geq \beta_{p_j}, \quad 0 < \beta_{p_j} \leq 1.$$

After determining strong inputs for the subsystems, we initialize blocks with one subsystem each and merge blocks containing strong input subsystems successively. In general the block partitioning is done only once before the iteration process starts, but there is an option to repeat it for some $t > t_0$ if convergence problems appear.

Codes of block waveform iteration methods have been tested on sequential machines. Currently the block waveform algorithm uses a modified DASSL code [BC1] including our linear sparse matrix package [Gr1] for numerical integration of the block systems. A program, automatically creating an interface for our code out of the data supplied by SPEEDUP when simulating a process [Hol], is used. The interface contains the DAE system in a structured representation, so it is possible to evaluate the function and the Jacobian-matrix subsystem-wise. At present the parallel case is simulated by these codes.

The numerical methods were run on two examples. The example DYNEVAP consisting of 87 equations within 13 subsystems represents a double effect evaporator. The second example BTX, a mathematical model of a Benzene-Toluene-Xylene distillation column, is made of 52 subsystems containing 1089 equations. For the example DYNEVAP an appropriate partitioning has been found, so that block waveform iteration method converges rapidly.

For BTX an appropriate partitioning, fulfilling all conditions for assignment and block generation, has not been found. Obviously the subsystems are strongly connected due to strong feedbacks between the stages (trays) of the column. For this reason possibilities to combine our iterative block methods with parallelizable direct block methods are investigated. A paper concerning this topic is in preparation.

References

- [AT1] Aspen Technology, SPEEDUP, User Manual, Library Manual, Aspen Technology, Inc., Cambridge, Massachusetts, USA, 1995
- [Br1] I. Bremer, Kurveniteration auf diskreten Zeitskalen, IWR Universität Heidelberg, Preprint 93 - 06, 1993

- [BB1] J. Borchardt, I. Bremer, Zur Analyse großer strukturierter chemischer Reaktionssysteme mit Waveform-Iterationsverfahren, IAAS Berlin, Preprint Nr. 65, 1993
- [BP1] L. Brüll, U. Pallaske, On Consistent Initialization of Differential-Algebraic Equations with Discontinuities, Proc. Fourth Conf. on Math. in Ind. (Hj. Wacker and W. Zulehner eds.), 1991
- [BC1] K.E. Brenan, S.L. Campbell, L.R. Petzold, Numerical Solution of Initial-Value Problem in Differential-Algebraic Equations, North-Holland, New York, 1989
- [Gr1] F. Grund, Numerische Lösung von hierarchisch strukturierten Systemen von Algebro-Differentialgleichungen, in Intern. Ser. of Num. Math., Vol. 117, Birkhäuser Verlag Basel, 1994, 17-31
- [GB1] F. Grund, J. Borchardt, D. Horn, T. Michael, H. Sandmann, Differential-algebraic systems in the chemical process simulation, Springer, Proceedings Workshop Scientific Computing in der Verfahrenstechnik, Hamburg, Juni 1995
- [Ho1] D. Horn, Entwicklung einer Schnittstelle für einen DAE Solver in der chemischen Verfahrenstechnik, Proceedings Workshop Software Engineering in Scientific Computing, Hamburg, to appear 1996
- [JP1] R. Jeltsch, B. Pohl, Waveform Relaxation with Overlapping Splittings, Seminar f. Angew. Math., Eidgenössische Techn. Hochschule Zürich, Research Rep. No. 91-02, 1991
- [LRS] E. Lelarsmee, A.E. Ruehli, A.L. Sangiovanni-Vincentelli, The Waveform Relaxation Method for Time-Domain Analysis of Large Scale Integrated Circuits, IEEE Trans. on CAD of IC and Syst., vol. CAD-1, No.3, 131-145, 1982
- [GH1] E. Griepentrog, M. Hanke, R. März, Toward a better understanding of differential algebraic equations (Introductory survey), Seminarberichte Nr 92-1, Humboldt-Universität zu Berlin, Fachbereich Mathematik, 1992
- [Mi1] U. Mikkala, Dynamic iteration methods applied to linear DAE systems, Journal of Comp. and Appl. Math. 25, 133-151, 1989
- [Nä1] St. Näher, LEDA Manual 3.0, Max-Planck-Institut für Informatik, MPI-I-93-109, Saarbrücken, 1993
- [PS1] Perkins, J.D., Sargent, R.W.H., SPEEDUP: A Computer Program for Steady State and Dynamic Simulation and Design of Chemical Processes, AIChE Symp. Ser. 214 (1982), 1-11

- [Re1] E.U. Reuter, Simulation und Optimierung einer chemischen Reaktion mit überlagerter Rektifikation, Dissertation, Institut f. Prozeß und Anlagentechnik, 1992
- [Sc1] K. R. Schneider, Existence and Approximation Results to the Cauchy Problem for a Class of Differential–Algebraic Equations, Zeitschrift f. Anal. u. ihre Anw., Vol.10, 3, 375–384, 1991
- [SL1] A. R. Secchi, F.S. Laganier, M. Morari, Dynamic Process Simulation Using a Concurrent Differential and Algebraic Solver, European Symposium on Computer Aided Process Engineering-2, 1991
- [SM1] A. R. Secchi, M. Morari, E. C. Biscaia Jr., The Waveform Relaxation Method in the Concurrent Dynamic Process Simulation, Computers and Chemical Engineering, Vol. 17, No. 7, 1993, 683–704
- [Wo1] G. Wozny, Simulation und Energetische Analyse thermischer Trennprozesse in Bodenkolonnen, Habilitationsschrift, Universität–Gesamthochschule Siegen, 1983
- [WS1] J.K. White, A.L. Sangiovanni–Vincentelli, Relaxation Techniques for the simulation of VLSI–Circuits. Kluwer Academic Press, New York, 1971

Recent publications of the Weierstraß-Institut für Angewandte Analysis und Stochastik

Preprints 1996

233. Frédéric Guyard, Reiner Lauterbach: Forced symmetry breaking perturbations for periodic solutions.
234. Vladimir G. Spokoiny: Adaptive and spatially adaptive testing of a nonparametric hypothesis.
235. Georg Hebermehl, Rainer Schlundt, Horst Zscheile, Wolfgang Heinrich: Simulation of monolithic microwave integrated circuits.
236. Georg Hebermehl, Rainer Schlundt, Horst Zscheile, Wolfgang Heinrich: Improved numerical solutions for the simulation of monolithic microwave integrated circuits.
237. Pavel Krejčí, Jürgen Sprekels: Global solutions to a coupled parabolic-hyperbolic system with hysteresis in 1-d magnetoelasticity.
238. Georg Hebermehl, Friedrich-Karl Hübner: Portabilität und Adaption von Software der linearen Algebra für Distributed Memory Systeme.
239. Michael H. Neumann: Multivariate wavelet thresholding: a remedy against the curse of dimensionality?
240. Anton Bovier, Miloš Zahradník: The low-temperature phase of Kac-Ising models.
241. Klaus Zacharias: A special system of reaction equations.
242. Susumu Okada, Siegfried Prößdorf: On the solution of the generalized airfoil equation.
243. Alexey K. Lopatin: Oscillations and dynamical systems: Normalization procedures and averaging.
244. Grigori N. Milstein: Stability index for invariant manifolds of stochastic systems.
245. Luis Barreira, Yakov Pesin, Jörg Schmeling: Dimension of hyperbolic measures – A proof of the Eckmann-Ruelle conjecture.
246. Leonid M. Fridman, Rainer J. Rumpel: On the asymptotic analysis of singularly perturbed systems with sliding mode.

247. Björn Sandstede: Instability of localised buckling modes in a one-dimensional strut model.
248. Björn Sandstede, Christopher K.R.T. Jones, James C. Alexander: Existence and stability of N -pulses on optical fibers with phase-sensitive amplifiers.
249. Vladimir Maz'ya, Gunther Schmidt: Approximate wavelets and the approximation of pseudodifferential operators.
250. Gottfried Bruckner, Sybille Handrock-Meyer, Hartmut Langmach: On the identification of soil transmissivity from measurements of the groundwater level.
251. Michael Schwarz: Phase transitions of shape memory alloys in soft and hard loading devices.
252. Gottfried Bruckner, Masahiro Yamamoto: On the determination of point sources by boundary observations: uniqueness, stability and reconstruction.
253. Anton Bovier, Véronique Gayrard: Hopfield models as generalized random mean field models.
254. Matthias Löwe: On the storage capacity of the Hopfield model.
255. Grigori N. Milstein: Random walk for elliptic equations and boundary layer.
256. Lutz Recke, Daniela Peterhof: Abstract forced symmetry breaking.
257. Lutz Recke, Daniela Peterhof: Forced frequency locking in S^1 -equivariant differential equations.
258. Udo Krause: Idealkristalle als Abelsche Varietäten.
259. Nikolaus Bubner, Jürgen Sprekels: Optimal control of martensitic phase transitions in a deformation-driven experiment on shape memory alloys.
260. Christof Külske: Metastates in disordered mean field models: random field and Hopfield models.
261. Donald A. Dawson, Klaus Fleischmann: Longtime behavior of a branching process controlled by branching catalysts.