Weierstraß-Institut für Angewandte Analysis und Stochastik Leibniz-Institut im Forschungsverbund Berlin e.V.

Preprint

ISSN 2198-5855

Limiting shape for first-passage percolation models on random geometric graphs

Cristian F. Coletti¹, Lucas R. de Lima¹, Alexander Hinsen², Benedikt Jahnel²,

Daniel R. Valesin³

submitted: September 27, 2021

 Center for Mathematics, Computation, and Cognition Federal University of ABC Av. dos Estados, 5001 09210-580 Santo André, São Paulo Brazil E-Mail: Cristian.Coletti@ufabc.edu.br Lucas.Roberto@ufabc.edu.br ² Weierstrass Institute Mohrenstr. 39 10117 Berlin Germany E-Mail: Alexander.Hinsen@wias-berlin.de Benedikt.Jahnel@wias-berlin.de

 ³ Department of Mathematics Bernoulli Institute University of Groningen Nijenborgh 9
 9747 AG Groningen The Netherlands E-Mail: D.Rodrigues.valesin@rug.nl

No. 2877

Berlin 2021



2020 Mathematics Subject Classification. 52A22, 60F15, 60K35.

Key words and phrases. Poisson-Gilbert graph, Richardson model, Bernoulli bond percolation, subergodicity, high-density limit, isotropic shapes.

We thank the team from Orange S.A. as well as W. König and A. Tóbiás for inspiring discussions. This work was funded by the German Research Foundation under Germany's Excellence Strategy MATH+: The Berlin Mathematics Research Center, EXC-2046/1 project ID: 390685689 as well as Orange Labs S.A.

This research was also supported by grants #2017/10555-0, #2019/19056-2, and #2020/12868-9, São Paulo Research Foundation (FAPESP). The article was written when L.R. de Lima was visiting the Department of Mathematics of the Bernoulli Institute at the University of Groningen. He is thankful for their hospitality.

Edited by Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS) Leibniz-Institut im Forschungsverbund Berlin e. V. Mohrenstraße 39 10117 Berlin Germany

Fax:+49 30 20372-303E-Mail:preprint@wias-berlin.deWorld Wide Web:http://www.wias-berlin.de/

Limiting shape for first-passage percolation models on random geometric graphs

Cristian F. Coletti, Lucas R. de Lima, Alexander Hinsen, Benedikt Jahnel, Daniel R. Valesin

Abstract

Let a random geometric graph be defined in the supercritical regime for the existence of a unique infinite connected component in Euclidean space. Consider the first-passage percolation model with independent and identically distributed random variables on the random infinite connected component. We provide sufficient conditions for the existence of the asymptotic shape and we show that the shape is an Euclidean ball. We give some examples exhibiting the result for Bernoulli percolation and the Richardson model. For the Richardson model we further show that it converges weakly to a branching process in the joint limit of large intensities and slow passing times.

1 Introduction, main results and examples

First-passage percolation (FPP) was initially introduced by Hammersley and Welsh [3] to study the spread of fluids through random medium. Since then, several variations of the percolation process have been extensively investigated (see Auffinger et al. [1] for an overview of FPP on \mathbb{Z}^d) due to their considerable amount of theoretical consequences and applications. It determines a random metric space by assigning random weights to the edges of a graph.

We consider the FPP model defined on a random geometric graph (RGG) in \mathbb{R}^d with $d \ge 2$. Here, the RGG is defined as in Penrose [9] by setting the vertices to be given by a homogeneous Poisson point process (PPP) with intensity $\lambda > 0$ and the edges are defined between any pair of vertices that are within an Euclidean distance smaller than a fixed threshold r > 0. This random graph is also known as the Poisson–Gilbert disk model. It is a graph associated to the Poisson–Boolean model in continuum percolation, and it can also be seen as a particular case of the random-connection model (see for instance Meester and Roy [7]).

The properties and other details regarding the structure and definition of the process will be given later in the text. We present here the basic definition in general lines. Let the parameters (λ, r) of the RGG be supercritical for the almost-sure existence of a random infinite connected component \mathcal{H} . Note that the infinite component \mathcal{H} is unique almost surely and we define the FPP model on \mathcal{H} with independent and identically distributed random variables on the joint probability space $(\Omega, \mathscr{A}, \mathbb{P})$.

The aim of this paper is to investigate the \mathbb{P} -*a.s.* existence of the limiting shape of the above defined process. In fact, we show that, under some conditions, the random balls of \mathcal{H} converge \mathbb{P} -*a.s.* to the deterministic shape of an Euclidean ball. The additional conditions refer to the distribution of zero passing time on the edges and the at-least linear growth of the process.

The model will be formally defined in the next sections, we give below a simplified description of the process to state the main result. Let τ be a random variable which defines the common distribution of the i.i.d. passage times τ_e along each edge $e \in E(\mathcal{H})$.

Set $r_c(\lambda) > 0$ to be the critical r for the existence of the infinite connected component \mathcal{H} of the RGG $\mathcal{G}_{\lambda,r}$. Let $B_s(x)$ stand for the open Euclidean ball of radius $s \ge 0$ centered at $x \in \mathbb{R}^d$ and denote by v_d be volume of the unit ball in d-dimensional Euclidean space. Denote by H_t the random subset of \mathbb{R}^d of points for which their closest point in \mathcal{H} is reached by the FPP model up to time t > 0. We let H_0 be the set of points that have the same closest point in \mathcal{H} as the origin. Here is our first main theorem.

Theorem 1.1 (Shape theorem for FPP on RGGs). Let $d \ge 2$ and $r > r_c(\lambda)$. Consider the FPP with i.i.d. random variables defined on the infinite connected component \mathcal{H} of $\mathcal{G}_{\lambda,r}$. Suppose that the following conditions are satisfied:

 (A_1) We have that

 $\mathbb{P}(\tau=0) < \frac{1}{\upsilon_d \, r^d \, \lambda}.$

(A₂) There exist $\eta > 2d + 1$ such that

 $\mathbb{E}[\tau^{\eta}] < +\infty.$

Then, there exists $\varphi > 0$ such that, for all $\varepsilon \in (0, 1)$, one has \mathbb{P} -a.s. that

$$(1-\varepsilon)B_{\varphi}(o) \subseteq \frac{1}{n}H_n \subseteq (1+\varepsilon)B_{\varphi}(o)$$
(1.1)

for sufficiently large $n \in \mathbb{N}$.

The existence of the limiting shape is particularly interesting because the RGG is a random graph which exhibits unbounded degrees. The interest of applications for this class of models has already been pointed out by Jahnel and König [5]. In particular, they suggested the theorem for the Richardson model on telecommunication networks. The example is naturally associated with the contact process by stochastic domination as studied by Ménard and Singh [8], and Riblet [11]. Another interesting application is a lower bound for the critical probability of bond percolation on the RGG, which is also presented below.

It is worth pointing out that one can find in the literature a bigger class of random geometric graphs studied by Hirsch et al. [4] where the graph distance was interpreted as a FPP model. It suggests that the class of RGGs could also be expanded in our case. We chose to focus our attention on the standard definition in this work due to the usage of intermediate results presented in the next section.

Before we state our second main result, let us present some examples.

Example 1 (Bond percolation). We define the bond percolation by considering the clusters of the Bernoulli FPP only at time zero, see Figure 1 for an illustration. For this, let us call $e \in E(\mathcal{H})$ an *open edge* when $\tau_e = 0$. Set $\tau_e \sim Ber(1-p)$ independently for every $e \in E(\mathcal{H})$ and observe that (A_2) is immediately satisfied.

Then, the open clusters are maximally connected components defined by sites with passage time zero between them. Let us define the critical probability p_c for the bond percolation on the *d*-dimensional RGG by

 $p_c := \inf\{p \in [0,1] : \mathbb{P}(\exists \text{ an infinite open cluster in } \mathcal{H}) > 0, \tau_e \sim Ber(1-p)\}.$

Note that by Theorem 1.1 the case $p < 1/(v_d r^d \lambda)$ implies the existence of the limiting shape.



Figure 1: Simulation of the open clusters for a bond percolation model on a 2-dimensional RGG with $p < 1/(v_d r^d \lambda)$.

Thus, an immediate consequence of the theorem is the following lower bound for the critical probability

$$p_c \ge 1/(\upsilon_d r^d \lambda),$$

and for p = 0 we recover \mathcal{H} . We observe that the same lower bound can also be obtained by exploration methods.

Example 2 (Richardson's growth model). Consider the interacting particle system known as the Richardson model defined on the infinite connected component \mathcal{H} of the RGG with parameter $\lambda_{\rm I} > 0$. It is a random growth process based on a model introduced by Richardson [12] and illustrated in Figure 2. It is commonly referred to as a model for the spread of an infection or for the growth of a population.

At each time $t \ge 0$, a site of \mathcal{H} is in either of two states, healthy (vacant) or infected (occupied). Let $\zeta_t : V(\mathcal{H}) \to \{0, 1\}$ indicate the state of the sites at time t assigning the values 0 and 1 for the healthy and infected states, respectively. The process evolves as follows:

- A healthy particle becomes infected at rate $\lambda_{I} \sum_{y \sim x} \zeta_{t}(y)$ and
- an infected particle remains infected forever.

It is easily seen that the process is determined by FPP with edge passage times $\tau_e \sim \text{Exp}(\lambda_I)$ independently for each $e \in E(\mathcal{H})$. In particular, this version of the Richardson model conventionally stochastically dominates the basic contact process.

Conditions (A₁) and (A₂) are straightforward since $\mathbb{P}(\tau = 0) = 0 < 1/(\upsilon_d r^d \lambda)$ and since $\mathbb{E}[\exp(\alpha \tau)] < +\infty$ for $\alpha \in (0, \lambda_I)$. Hence, Theorem 1.1 is valid for the Richardson model on \mathcal{H} for any supercritical $r > r_c(\lambda)$.

Futhermore, it is immediate to see that Theorem 1.1 still holds for any initial configuration $Z \subseteq \mathbb{R}^d$ of infected particles whenever $Z \subseteq B_{s'}(o)$ for some s' > 0. In that case, we simply replace H_t by $H_t^Z := \bigcup_{z \in Z} H_t^z$.



Figure 2: Simulation of the spread of an infection given by the Richardson model on a bidimensional RGG.

Our second main result concerns the asymptotic behavior of the Richardson model in the limit as α diverges to infinity in $\alpha\lambda$ and $\lambda_{\mathrm{I}}/\alpha$. In words, we consider a coupled limit of high densities and slow infection rates. The limiting process is a branching process $(\mathcal{T}_t^{\lambda,\lambda_{\mathrm{I}}})_{t\geq 0}$ defined as follows. At time zero, the process has a node only at the origin, i.e., $\mathcal{T}_0^{\lambda,\lambda_{\mathrm{I}}} = o$. Then, iteratively, each node $X_i \in \mathbb{R}^d$ of the process produces offsprings independently according to a Poisson process in time with intensity $v_d r^d \lambda \lambda_{\mathrm{I}}$ and the offsprings are placed independently and uniformly within $B_r(X_i)$. We will write subscript [0, t] in cases where we consider the whole path from time 0 to time t and $\mathcal{H}_t^{\lambda,\lambda_{\mathrm{I}}}$ for the set of points in \mathcal{H} reached by the FPP model up to time t > 0, where we also indicate system parameters. As above, we let $\mathcal{H}_0^{\lambda,\lambda_{\mathrm{I}}}$ be the closest point in \mathcal{H} to the origin and define $\mathcal{H}_0^{\lambda,\lambda_{\mathrm{I}}} = \emptyset$ if there is no infinite component. Let us note that this process has similarities with the growth process as presented in Deijfen [2]. Here is our second main result.

Theorem 1.2 (Time-space rescaling for Richardson models). Let $d \ge 2$. For the Richardson model with parameters $r, \lambda, \lambda_{\rm I}$ where $r > r_{\rm c}(\lambda)$, we have that for all t > 0

$$\mathcal{H}^{\alpha\lambda,\lambda_{\mathrm{I}}/\alpha}_{[0,t]}\longrightarrow \mathcal{T}^{\lambda,\lambda_{\mathrm{I}}}_{[0,t]},$$

weakly with respect to the Skorokhod topology based on the vague topology, as α tends to infinity.

The rest of the manuscript is organized as follows. In Section 2, we have compiled some basic facts about the RGG and show results on the asymptotic behaviour of the infinite component \mathcal{H} . The FPP model is defined in detail in Section 3 where we also present the proof of Theorem 1.1. Finally, in Section 4 we present the proof of Theorem 1.2.

2 On the random geometric graph

In this section we present the definition and parameters for the RGG and the existence of the infinite connected component. We also show some results about its geometry in order to study the asymptotic shape in the next section.

Let \mathcal{P}_{λ} be the random set of points determined by the homogeneous PPP on \mathbb{R}^d with intensity $\lambda > 0$. The RGG $\mathcal{G}_{\lambda,r} = (V, E)$ on \mathbb{R}^d is defined by

$$V = \mathcal{P}_{\lambda} \quad \text{and} \quad E = \left\{ \{u, v\} \subseteq V : \|u - v\| < r, \ u \neq v \right\},\$$

where $\|\cdot\|$ is the Euclidean norm. Since $\lambda^{-1/d} \mathcal{P}_{\lambda} \sim \mathcal{P}_1$, we may regard λ as fixed due to the homogeneity of the norm. We write $\mathcal{G}_r := \mathcal{G}_{1,r}$ and $\mathcal{P} = \mathcal{P}_1$. Set (Ξ, \mathscr{F}, μ) to be the probability space induced by the construction of \mathcal{P} . Let us now introduce the group action $\vartheta : \mathbb{R}^d \curvearrowright \Xi$ which is determined by the spatial translation as a shift operator. That is, $\mathcal{P} \circ \vartheta_z = \{v - z : v \in \mathcal{P}\}$. The following lemma is a classical result on PPPs, which can be found for example in Meester and Roy [7, Proposition 2.6].

Lemma 2.1. The homogeneous PPP is mixing on $(\Xi, \mathscr{F}, \mu, \vartheta)$.

Remark 2.1. Let $S : \mathbb{R}^d \to \mathbb{R}^d$ be an isometry. Then, it is known that S induces a μ -preserving ergodic function $\sigma : \Xi \to \Xi$ where $S[\mathcal{P}] = \mathcal{P} \circ \sigma$.

We are interested in studying the spread of an infection on an infinite connected component of \mathcal{G}_r . It is a well-known fact from continuum percolation theory (see Meester and Roy [7] or Penrose [9, Chapter 10] for details) that, for all $d \geq 2$, there exists a critical $r_c > 0$ such that \mathcal{G}_r has an infinite component $\mathcal{H} \mu$ -a.s. for all $r > r_c$. Moreover, \mathcal{H} is μ -a.s. unique. Since \mathcal{H} is a subgraph of \mathcal{G}_r , we denote by $V(\mathcal{H})$ and $E(\mathcal{H})$ its sets of vertices and edges, respectively.

From now on, write $(\Xi', \mathscr{F}', \mu)$ for the probability space of the PPP conditioned on the existence of \mathcal{H} when $r > r_c$. It suffices for our purposes to know that $r_c \ge 1/\upsilon_d^{1/d}$ where υ_d denotes the volume of the unit ball in the *d*-dimensional Euclidean space. Indeed, improved lower and upper bounds can be found in Torquato and Jiao [14] and r_c approximates to $1/\upsilon_d^{1/d}$ from above as $d \to +\infty$.

Let us write $\theta_r := \mu (B_r(o) \cap V(\mathcal{H}) \neq \emptyset)$ and denote the cardinality of a set by $|\cdot|$.

Proposition 2.2. (Weaker version of Penrose and Pisztora [10, Theorem 1]). Let $d \ge 2$, $r > r_c$ and $\varepsilon \in (0, 1/2)$. Then, there exists c > 0 and $s_0 > 0$ such that, for all $s \ge s_0$,

$$\mu\left((1-\varepsilon)\theta_r < \frac{|V(\mathcal{H})\cap [-s/2, s/2]^d|}{s^d} < (1+\varepsilon)\theta_r\right) \ge 1 - \exp(-cs^{d-1}).$$

As a consequence of the last result, we present the following lemma without proof (see Yao et al. [15, Lemma 3.3]).

Lemma 2.3. Let $r > r_c$. Then, there exists C, C' > 0 such that, for each $x \in \mathbb{R}^d$ and all s > 0,

$$\mu(B_s(x) \cap V(\mathcal{H}) = \emptyset) \le C \exp(-C' s^{d-1}).$$

Let $\mathscr{P}(x,y)$ denote the set of self-avoiding paths from x to y in \mathcal{H} . The simple length of a path $\gamma = (x = x_0, x_1, \dots, x_m = y) \in \mathscr{P}(x, y)$ is denoted by $|\gamma| = m$.

Write D(x, y) for the \mathcal{H} -distance between $x, y \in V(\mathcal{H})$ given by

$$D(x, y) = \inf\{|\gamma| : \gamma \in \mathscr{P}(x, y)\}.$$

Let $x \in \mathbb{R}^d$, then we define $q : \mathbb{R}^d \to V(\mathcal{H})$ by

$$q(x) := \underset{y \in V(\mathcal{H})}{\arg\min} \{ \|y - x\| \}$$
(2.1)

the closest point to x in the infinity cluster. Observe from (2.1) that q may be multivalued for some $x \in \mathbb{R}^d$. In that case, we assume that q(x) is uniquely defined by an arbitrarily fixed outcome of (2.1). Hence q induces a Voronoi partition of \mathbb{R}^d with respect to \mathcal{H} , see Figure 3 for an illustration.



Figure 3: A random geometric graph on \mathbb{R}^2 with the Voronoi partition generated by the infinite connected component \mathcal{H} (in blue).

We now extend the domain of the \mathcal{H} -distance by defining D(x, y) := D(q(x), q(y)) for every $x, y \in \mathbb{R}^d$. The proposition below can be immediately adapted from the proof of Yao et al. [15, Theorem 2.2] by applying properties of Palm calculus and Lemma 2.3.

Proposition 2.4. (Adapted from Yao et al. [15, Theorem 2.2]). Let $d \ge 2$ and $r > r_c$. Then there exists $\rho_r > 0$ depending on r such that, μ -a.s., for all $x \in \mathbb{R}^d$,

$$\lim_{\|y\|\uparrow+\infty}\frac{\mathrm{D}(x,y)}{\|y-x\|}=\rho_r.$$

The constant ρ_r is called *stretch factor* of \mathcal{H} . Observe that $\rho_r \ge 1/r$. Due to the subadditivity of the \mathcal{H} -distance, one can easily see that $\mathbb{E}_{\mu}[\mathbf{D}(o, z)]$ with ||z|| = 1 is an upper-bound for ρ_r .

We have the following result about the tail behaviour of D(o, z).

Lemma 2.5. Let $d \ge 2$ and $r > r_c$. Then, there exist $c_1, c_2 > 0$ and $\beta' > 1$ such that, for all $x \in \mathbb{R}^d$ and every $t > \beta' ||x||$,

$$\mu(\mathrm{D}(o, x) \ge t) \le c_1 \exp(-c_2 t).$$

Proof. Let us write $\mu_{v,w}$ for the Palm measure $\mu * \delta_v * \delta_w$ for any given $v, w \in \mathbb{R}^d$. Set \overline{D} to be the simple \mathcal{G}_r -distance. In what follows, $v \nleftrightarrow w$ stands for the existence of a path between v and w in \mathcal{G}_r . It is clear that $D(v, w) = \overline{D}(v, w)$ whenever $v, w \in V(\mathcal{H})$. By [15, Lemma 3.4], there exist $\overline{c}_1, \overline{c}_2 > 0$ and $\beta' > 1$ such that

$$\mu_{v,w}(v \nleftrightarrow w, \text{ and } \mathcal{D}(v,w) \ge t) \le \overline{c}_1 \exp(-\overline{c}_2 t)$$
(2.2)

for all $t \ge \beta' ||v - w||/2$. Consider now $\mathcal{B}_r(z) := B_r(z) \cap \mathcal{P}$. We apply Lemma 2.3, (2.2), and Campbell's theorem to obtain that there exist C, C' > 0 such that

$$\mu \left(\mathbf{D}(o, x) \ge t \right) \le \mu \left(\|q(o)\| \ge t/(2\beta') \right) + \mu \left(\|q(x) - x\| \ge t/(2\beta') \right) + \mu \left(\bigcup_{v \in \mathcal{B}_{t/(2\beta')}(o), w \in \mathcal{B}_{t/(2\beta')}(x)} \{v \nleftrightarrow w \text{ and } \overline{\mathbf{D}}(v, w) \ge t \} \right) \le 2C \exp \left(-C't/(2\beta') \right) + \overline{c}_1 \frac{v_d^2}{2^{2d}} t^{2d} \exp(-\overline{c}_2 t)$$
(2.3)

for all $t \ge \beta' ||x||$. Hence, we can conclude the proof of (2.3) by suitably choosing $c_1, c_2 > 0$. \Box

Next, let us define for every $x \in \mathbb{R}^d$ the quantity

 $W_n^x := \{ \text{self-avoiding paths of length } n \text{ in } \mathcal{G}_r \text{ starting at } x \}$

and note that using Jahnel and König [5, Theorem 4.6.11]), we have that

$$\mathbb{E}_{\mu}[|W_n^x|] = (\mathbf{v}_d R^d)^n$$

Consider now $|W_n^{q(o)}|$, the number of self-avoiding paths in \mathcal{H} starting at q(o). We apply the previous result to prove the following lemma.

Lemma 2.6. Let $d \ge 2$, $r > r_c$, and $\kappa > 1$. Then, one has μ -a.s. that for sufficiently large $n \in \mathbb{N}$,

$$\left|W_n^{q(o)}\right| < \left(\kappa \,\boldsymbol{\upsilon}_d \, r^d\right)^n.$$

Proof. Let $n \in \mathbb{N}$ and define the events

$$\mathsf{A}_n := \left\{ |W_n^{q(o)}| \ge (\kappa \mathfrak{v}_d \, r^d)^n \right\}.$$

Recall the notation $\mathcal{B}_n(o) := B_n(o) \cap \mathcal{P}$. Then

$$\mathsf{A}_n \subseteq \left(\bigcup_{v \in \mathcal{B}_n(o)} \left\{ |W_n^v| \ge (\kappa \mathfrak{v}_d r^d)^n, \|q(o)\| < n \right\} \right) \cup \{\|q(o)\| \ge n\}.$$

Hence, by Markov's inequality, one has for any given $x \in \mathbb{R}^d$ that

$$\mu(|W_n^x| \ge (\kappa \upsilon_d r^d)^n) \le \mathbb{E}_{\mu} \left[|W_n^x| \right] / (\kappa \upsilon_d r^d)^n = 1/\kappa^n.$$

Thus, by Lemma 2.3 and Campbell's theorem, there exist C, C' > 0 such that

$$\sum_{n=1}^{+\infty} \mu(\mathsf{A}_n) \le \mathfrak{v}_d \sum_{n=1}^{+\infty} \frac{n^d}{\kappa^n} + C \sum_{n=1}^{+\infty} e^{-C'n^{d-1}} < +\infty,$$

and hence, an application of the Borel-Cantelli lemma completes the proof.

DOI 10.20347/WIAS.PREPRINT.2877

Berlin 2021

3 First-passage percolation

We proceed to formally define our process. Let $\{\tau_e\}_{e \in E}$ be a family of independent and identically distributed random variables taking values in the time set $[0, +\infty)$. We say that τ_e is the passage time of the edge $e \in E(\mathcal{H})$.

Set $(\Omega, \mathscr{A}, \mathbb{P})$ to be the joint probability space associated with the construction of the random geometric graph \mathcal{G}_r and the independent assignment of the random passage times $\{\tau_e\}_{e \in E}$. The joint probability space can be constructed as a product space and we will write $\mathbb{E} = \mathbb{E}_{\mathbb{P}}$ for short.

Given any path $\gamma = (x, x_1, \dots, x_m, y) \in \mathscr{P}(x, y)$ for $x, y \in V(\mathcal{H})$, we write $e \in \gamma$ for an edge $e \in E(\mathcal{H})$ between a pair of consecutive vertices of γ . We denote the passage time of the path γ by

$$T(\gamma) := \sum_{e \in \gamma} \tau_e.$$

The *passage time* between $x, y \in \mathbb{R}^d$ is then defined by the random variable

$$T(x,y) := \inf \left\{ T(\gamma) : \gamma \in \mathscr{P}(q(x),q(y)) \right\}$$

In fact, we can see later that T(x, y) is a random pseudo-metric when associated with a group action. To avoid cumbersome notation, we set T(x) := T(o, x) for all $x \in \mathbb{R}^d$.

Using these definitions, we have

$$H_t := \{ x \in \mathbb{R}^d : T(x) \le t \}$$

for the set of Voronoi cells induced by \mathcal{H} reached up to time t with the FPP starting in q(o).

Next, consider $\vartheta : \mathbb{R}^d \curvearrowright \Omega$ to be an extension of the group action introduced in Section 2 such that ϑ_z will induce $\tau_{\{x,y\}} \mapsto \tau_{\{x-z,y-z\}}$ independently in the product space. It is easily seen that ϑ inherits the ergodic property of the previously defined group action and that Remark 2.1 still holds for actions on Ω associated with isometries by extending them in the same fashion. Then, we observe that $T(x, x + y) = T(y) \circ \vartheta_x$ and that the subadditivity

$$T(x+y) \le T(x) + T(y) \circ \vartheta_x \tag{3.1}$$

for all $x, y \in \mathbb{R}^d$ is straightforward.

We have the following lemma.

Lemma 3.1. Let $d \ge 2$, $r > r_c$, and $\mathbb{P}(\tau = 0) < 1/(\upsilon_d r^d)$. Then, there exists a > 0 depending on r such that, for all $x \in \mathbb{R}^d$,

$$a\|x\| \le \mathbb{E}[T(x)].$$

Proof. It suffices to prove this statement for large ||x|| as due to subadditivity (3.1) and stationarity of T(nx, (n+1)x) for all $n \in \mathbb{N}$ we have

$$\mathbb{E}[T(mx)]/\|mx\| \le \sum_{i=1}^{m} \mathbb{E}[T((i-1)x, ix)]/\|mx\| = \mathbb{E}[T(x)]/\|x\|.$$

Define the event

$$A_x^1 := \left\{ \max\{ \|q(o)\|, \|q(x) - x\| \} \le \|x\|/4 \right\}.$$

In order to simplify the notation, let us abbreviate $m_x = \lceil ||x||/(2r)\rceil$. Note that on A_x^1 , every path $\gamma \in \mathscr{P}(q(o), q(x))$ has $|\gamma| \ge m_x$ and therefore includes a sub-path of length at least m_x . Consequently, for any t > 0,

$$\mathbb{P}\left(\{T(x) \le t\} \cap A_x^1\right) = \mathbb{P}\left(\left\{\exists \gamma \in \mathscr{P}(q(o), q(x)) : T(\gamma) \le t\right\} \cap A_x^1\right)$$
$$\le \mathbb{P}\left(\exists \gamma \in W_{m_x}^{q(o)} : T(\gamma) \le t\right),$$

In order to proceed, first observe that, using Chernoff's bound for the binomial distribution with $X \sim \text{Binomial}(n, p)$, we have

$$\mathbb{P}(X \le cn) \le \exp\left(-n\left(c\log\frac{c}{p} + (1-c)\log\frac{1-c}{1-p}\right)\right) \\ = \left(p^{-c}(1-p)^{(1-c)} \cdot c^{c}(1-c)^{(1-c)}\right)^{-n},$$

where for $c \to 0$ the base converges to $(1-p)^{-1}$. Also, because of the right continuity of the cumulative distribution function associated to τ , there exist $\kappa > 1$ and $\delta > 0$ such that

$$\mathbb{P}(\tau \le \delta) < 1/(\kappa \boldsymbol{\upsilon}_d \, r^d).$$

Further, consider a random variable $X' \sim \text{Binomial}(n, \mathbb{P}(\tau > \delta))$ with respect to \mathbb{P} . Note that $\tau \geq \delta \mathbb{1}\{\tau > \delta\}$ and therefore on any (self-avoiding path) of length $|\gamma| = n$ the sum of n i.i.d. copies of τ , stochastically dominates $\delta X'$. Therefore, there exists c > 0 and $\kappa' > 1$ such that for all $n \in \mathbb{N}$ and $|\gamma| = n$

$$\mathbb{P}(T(\gamma) \le cn) \le \mathbb{P}(X' > cn/\delta) \le \left(\kappa' \mathfrak{v}_d r^d\right)^{-n}.$$

Fix $1 < \kappa'' < \kappa'$ and define

$$A_x^2 := \{ |W_{m_x}^{q(o)}| \le (\kappa'' \upsilon_d r^d)^{m_x} \} \text{ and } A_x := A_x^1 \cap A_x^2.$$

Then, we have

$$\mathbb{P}\left(T(x) \le cm_x\right) \le \mathbb{P}\left((A_x)^c\right) + \mathbb{P}\left(A_x \cap \left\{\exists \gamma \in W_{m_x}^{q(o)} \colon T(\gamma) \le cm_x\right\}\right)$$
$$\le \mathbb{P}\left((A_x)^c\right) + \mathbb{E}\left[\mathbbm{1}_{A_x} \cdot \sum_{\gamma \in W_{m_x}^{q(o)}} \mathbbm{1}\left\{T(\gamma) \le cm_x\right\}\right]$$
$$\le \mathbb{P}\left((A_x)^c\right) + (\kappa'' \upsilon_d r^d)^{m_x} \cdot (\kappa' \upsilon_d r^d)^{-m_x},$$

where $\mathbb{P}((A_x)^c)$ can be made arbitrarily small by choosing ||x|| large enough via Lemma 2.3 and Lemma 2.6. As $\kappa'' < \kappa'$, the exponent in the second summand is negative and dominates the polynomial term for large x. Therefore, there is a k such that for all $x \in \mathbb{R}^d$ with $||x|| \ge k$,

$$\mathbb{P}\left(T(x) \le cm_x\right) \le 1/2$$

Setting a = c/4r we arrive at the statement $a||x|| \leq \mathbb{E}[T(x)]$ when ||x|| > k.

Remark 3.1. Observe that $\mathbb{E}[T(x)] \leq \mathbb{E}[D(o, x)] \mathbb{E}[\tau]$ due to the subadditivity and Fubini's Theorem. Moreover, condition (A_2) implies that $\mathbb{E}[\tau] < +\infty$. One can easily see from Proposition 2.4 and L^1 convergence given by Kingman's subadditive ergodic theorem [6] applied to the \mathcal{H} -distance, that, for all $x \in \mathbb{R}^d$,

$$b := \rho_r \mathbb{E}[\tau] \ge \limsup_{n \uparrow \infty} \mathbb{E}[T(nx)] / \|nx\|.$$

Denote by \mathbb{P}_{ξ} the quenched probability of the propagation model given a realization $\xi \in \Xi'$. The lemma below ensures the at-least linear growth of the passage times.

Lemma 3.2. Let $d \ge 2$, $r > r_c$, and assume that (A_2) holds. Then, there exist deterministic $\beta > 0$ and $\kappa > 1$ such that, for every $x, y \in \mathbb{R}^d$, and for each $\xi \in \Xi'$,

$$\mathbb{P}_{\xi}\left(T(x,y) \ge t\right) \le t^{-(d+\kappa)}$$

for all $t \ge \beta D(x, y)$.

Proof. Let $\gamma \in \mathscr{P}(x, y)(\xi)$ be a geodesic given by the \mathcal{H} -distance. Then, one has by Markov's inequality that, for $t > \mathbb{E}[\tau] D(x, y)$ and η from (A_2),

$$\mathbb{P}_{\xi}\left(T(x,y) \ge t\right) \le \mathbb{P}_{\xi}\left(T(\gamma) \ge t\right) \le \frac{\mathbb{E}_{\xi}\left[\left(\sum_{e \in \gamma} \left(\tau_e - \mathbb{E}[\tau]\right)\right)^{\eta}\right]}{\left(t - \mathbb{E}[\tau] \operatorname{D}(x,y)\right)^{\eta}}.$$
(3.2)

Rosenthal's inequality [13] states that, if Y_1, \ldots, Y_n are independent random variables with mean zero and finite moment of order p, with p > 2, then

$$\mathbb{E}\left[\left(\sum_{i=1}^{n} Y_{i}\right)^{p}\right] \leq C_{p} \cdot \max\left\{\sum_{i=1}^{n} \mathbb{E}[|Y_{i}|^{p}]; \left(\sum_{i=1}^{n} \mathbb{E}[(Y_{i})^{2}]\right)^{p/2}\right\},\$$

where $C_p > 0$ is a constant that depends only on p. Since (A_2) holds and $(\tau_e - \mathbb{E}[\tau_e])$ are identically distributed for all $e \in \gamma$, it yields

$$\mathbb{E}_{\xi}\left[\left(\sum_{e\in\gamma}(\tau_e - \mathbb{E}[\tau])\right)^{\eta}\right] \le C \operatorname{D}(x, y)^{\eta/2},\tag{3.3}$$

where C is now a constant that depends both on η and on the distribution of the random variables.

In case $t \ge 2 \mathbb{E}[\tau] D(x, y)$ we have $t - \mathbb{E}[\tau] D(x, y) \ge t/2$. Using this and (3.3), the right-hand side of (3.2) is smaller than

$$2^{\eta}C \operatorname{D}(x,y)^{\eta/2} t^{-\eta}.$$

If we also have that $t \ge 4C^{2/\eta} D(x, y)$, this is smaller than $t^{-\eta/2}$. We have thus proved that

$$\mathbb{P}_{\xi}\left(T(x,y) \geq t\right) \leq t^{-(d+\kappa)} \quad \text{for all } t \geq \beta \operatorname{D}(x,y),$$

with $\beta := \max\left\{2 \mathbb{E}[\tau], 4C^{2/\eta}\right\}$ and $\kappa := \eta/2 - d > 1$.

Before proving our first main theorem, we state and prove the following result.

Lemma 3.3. Let $d \ge 2$, $r > r_c$. Consider the i.i.d. FPP on the RGG satisfying (A_2). Then, there exist constants δ , C > 0, and $\kappa > 1$ such that for all t > 0 and all $x \in \mathbb{R}^d$, one has

$$\mathbb{P}\left(\sup_{y\in B_{\delta t}(x)}T(x,y)\geq t\right)\leq Ct^{-\kappa}.$$

DOI 10.20347/WIAS.PREPRINT.2877

Berlin 2021

Proof. Due to the translation invariance it suffices to prove the lemma for x = 0. Let $\delta = (\beta'\beta)^{-1}$ with β' and β from Lemmas 2.5 and 3.2. Set $c_d > 0$ to be such that $B_{2\delta}(o) \subseteq [-c_d/2, c_d/2]^d$ and write $C_d := 2\theta_r c_d$ with $\theta_r > 0$ from Proposition 2.2. Let us now define the following events:

$$G_{1} := \left\{ q(o) \in B_{\delta t}(o) \right\} \cap \left\{ |B_{\delta 2t}(o) \cap V(\mathcal{H})| \le C_{d} \cdot t^{d} \right\}$$
$$G_{2} := \left\{ \sup_{\|y\| \le \delta t} \mathcal{D}(o, y) \le t/\beta \right\} \cap G_{1}$$
$$G_{3} := \left\{ \sup_{\|y\| \le \delta t} T(o, y) \ge t \right\} \cap G_{1} \cap G_{2}$$

Now,

$$\mathbb{P}\left(\sup_{y\in B_{\delta t}(o)}T(o,y)\geq t\right)\leq \mathbb{P}(G_3)+\mathbb{P}(G_1^c)+\mathbb{P}(G_2^c),\tag{3.4}$$

where the last two summands decrease exponentially in *t* due to Proposition 2.2, and Lemmas 2.3 and 2.5. By Lemma 3.2, there exists $\kappa > 1$ such that

$$\mathbb{P}(G_3) \le \mathbb{E}\left[\mathbb{1}\left\{\xi \in G_1 \cap G_2\right\} \mathbb{P}_{\xi}\left(\sup_{\|y\| < \delta t} T(y) \ge t\right)\right] \le C_d t^d / t^{d+\kappa}.$$
(3.5)

Combining (3.5) with (3.4), the desired bound is obtained by choosing a suitable C > 0. \Box

After this preparatory work, we now proceed to prove Theorem 1.1.

Proof of Theorem 1.1. We begin by verifying properties of T(nx). Note that for every $x \in \mathbb{R}^d$ one has that $\mathbb{E}[T(x)] < +\infty$ by Lemmas 2.5 and 3.2. Recall that the process is mixing on $(\Omega, \mathscr{A}, \mathbb{P}, \vartheta)$ by Lemma 2.1. Then, by the subadditivity (3.1), we apply Kingman's subadditive ergodic theorem to obtain that, \mathbb{P} -*a.s.*, for all $x \in \mathbb{R}^d$,

$$\lim_{n\uparrow+\infty}\frac{T(nx)}{n} = \phi(x), \tag{3.6}$$

where $\phi:\mathbb{R}^d\rightarrow [0,+\infty)$ is a homogeneous and subadditive function given by

$$\phi(x) = \inf_{n \ge 1} \frac{\mathbb{E}[T(nx)]}{n} = \lim_{n \uparrow +\infty} \frac{\mathbb{E}[T(nx)]}{n}$$

Since the process is rotation invariant, there exists a constant φ (the time constant) such that $\phi(x) = \varphi^{-1} ||x||$ for all $x \in \mathbb{R}^d$. In fact, one has from Lemma 3.1 and Remark 3.1 that

$$0 < a \le \varphi^{-1} \le b = \rho_r \mathbb{E}[\tau] < \infty.$$

Let us now prove the \mathbb{P} -*a.s.* asymptotic equivalence

$$\lim_{\|y\|\uparrow+\infty} \frac{T(y)}{\|y\|} = \frac{1}{\varphi}.$$
(3.7)

For the lower bound of the theorem we prove the equivalent statement that for every $\epsilon \in (0,1),$

$$\limsup_{s\uparrow\infty} \quad \sup_{\|y\| \le (1-\epsilon)s} \frac{T(y)}{s} = \limsup_{m \in \mathbb{N}, m\uparrow\infty} \quad \sup_{\|y\| \le (1-\epsilon)m} \frac{T(y)}{m} < \frac{1}{\varphi} \quad \mathbb{P}-a.s.,$$

DOI 10.20347/WIAS.PREPRINT.2877

Berlin 2021

where the first equation holds as $\lfloor s \rfloor / s$ converges to 1. Fix $\epsilon \in (0,1)$ and let δ be given by Lemma 3.3. Due to compactness, there exists a finite cover of open balls with centers $(y_i)_{i \in \{1,...,n\}} \subseteq \mathbb{R}^d$ with $\|y_i\| \leq 1 - \epsilon$ such that

$$\overline{B_{1-\epsilon}(o)} \subseteq \bigcup_{i \in \{1,\dots,n\}} B_{\delta\epsilon/(2\varphi)}(yi).$$

Furthermore $B_{m(1-\epsilon)}(o) \subseteq \bigcup_{i \in \{1,...,n\}} B_{m\delta\epsilon/(2\varphi)}(my_i)$ for every $m \in \mathbb{N}$. Applying Lemma 3.3 we obtain

$$\sum_{m \in \mathbb{N}} \mathbb{P}\left(\sup_{\|y - my_i\| \le m\delta\epsilon/(2\varphi)} T(my_i, y) > m\epsilon/(2\varphi)\right) < \infty.$$

Therefore, by the Borel-Cantelli lemma,

$$\limsup_{m \in \mathbb{N}, m \uparrow \infty} \quad \sup_{\|my_i - y\| \le m\delta\epsilon/(2\varphi)} \frac{T(my_i, y)}{m} < \frac{\epsilon}{2\varphi} \quad \mathbb{P} - a.s.$$

Applying (3.6) and subadditivity, we obtain

$$\begin{split} \limsup_{m\uparrow\infty; \|y\|\leq (1-\epsilon)m} \frac{T(y)}{m} &\leq \limsup_{m\uparrow\infty} \left(\max_{i\in\{1,\dots,n\}} \frac{T(o,my_i)}{m} + \sup_{\|my_i-y\|\leq m\delta\epsilon/(2\varphi)} \frac{T(my_i,y)}{m} \right) \\ &\leq \max_{i\in\{1,\dots,n\}} \|y_i\|/\varphi + \epsilon/(2\varphi) < 1/\varphi \quad \mathbb{P}-a.s., \end{split}$$

where we used that $||y_i|| < 1 - \epsilon$.

For the upper bound of the theorem define $A_t := B_{t(1+2\epsilon)}(o) \setminus B_{t(1+\epsilon)}(o)$ and observe that it suffices to prove

$$\liminf_{m \in \mathbb{N}, m \uparrow \infty} \quad \inf_{y \in A_t} \frac{T(y)}{m} > \frac{1}{\varphi} \quad \mathbb{P}-a.s.$$

for arbitrary but fixed $\epsilon > 0$, as for (for $t > \epsilon$) any x with $||x|| > t(1 + 2\epsilon)$ there exists an $\tilde{x} \in A_t$ with $T(\tilde{x}) \leq T(x)$.

Similar as in the lower bound of the theorem, fix $\epsilon > 0$ and $\delta > 0$ small enough that Lemma 3.3 holds. There exists a set of centers $(y_i)_{i \in \{1,...,n\}} \subseteq \mathbb{R}^d$ with $||y_i|| \ge 1 + \epsilon$ such that

$$A_t \subseteq \bigcup_{i \in \{1, \dots, n\}} B_{\delta \epsilon/(2\varphi)}(y_i)$$

and hence

$$\liminf_{m \in \mathbb{N}, m \uparrow \infty} \quad \inf_{y \in A_m} \frac{T(y)}{m} \ge \liminf_{m \in \mathbb{N}, m \uparrow \infty} \left(\min_{i \in \{1, \dots, n\}} \frac{T(o, my_i)}{m} - \sup_{\|my_i - y\| \le m\delta\epsilon/(2\varphi)} \frac{T(my_i, y)}{m} \right)$$
$$\ge \min_{i \in \{1, \dots, n\}} \|y_i\| / \varphi - \epsilon / (2\varphi) > 1/\varphi,$$

which concludes the proof of the asymptotic equivalence (3.7). The proof of the theorem is now complete by standard arguments of the \mathbb{P} -*a.s.* uniform convergence given by (3.7).

4 **Proof of Theorem 1.2**

Throughout this section, we fix $r, \lambda, \lambda_{\mathrm{I}}$. For the process $(\mathcal{H}_t^{\alpha\lambda,\lambda_{\mathrm{I}}/\alpha})_{t\geq 0}$, we let $T'_{\alpha,0} = 0$ and let $T'_{\alpha,1}, T'_{\alpha,2}, \ldots$ denote the jump times in increasing order. We let $Z_{\alpha,0} := q(o)$ and for $i \in \mathbb{N}$ we let $Z_{\alpha,i}$ denote the point of \mathbb{R}^d that joins the process at time $T'_{\alpha,i}$, so that

$$\mathcal{H}_t^{\alpha\lambda,\lambda_{\mathrm{I}}/\alpha} = \{Z_{\alpha,0}, Z_{\alpha,1}, \dots, Z_{\alpha,i}\} \quad \text{for } t \in [T'_{\alpha,i}, T'_{\alpha,i+1}), \ i \in \mathbb{N}_0$$

We also let

$$T_{\alpha,0} := 0, \qquad T_{\alpha,i} := T'_{\alpha,i} - T'_{\alpha,i-1}, \ i \in \mathbb{N}.$$

For the process $(\mathcal{T}_t^{\lambda,\lambda_1})_{t\geq 0}$, we define sequences Z_0, Z_1, Z_2, \ldots and T_0, T_1, T_2, \ldots in the same way (except that $Z_0 := o$ in this case). With this notation, we can state:

Proposition 4.1. Let $k \in \mathbb{N}$, $f_0, \ldots, f_k : \mathbb{R}^d \to \mathbb{R}$ and $g_1, \ldots, g_k : (0, \infty) \to \mathbb{R}$; assume all these functions are continuous with compact support. We then have

$$\mathbb{E}\left[f_0(Z_{\alpha,0})\prod_{i=1}^k f_i(Z_{\alpha,i})g_i(T_{\alpha,i})\right] \xrightarrow{\alpha \to \infty} f_0(o) \cdot \mathbb{E}\left[\prod_{i=1}^k f_i(Z_i)g_i(T_i)\right]$$
(4.1)

We will prove this proposition later; for now let us show how it implies Theorem 1.2.

Proof of Theorem 1.2. Let \mathcal{M} denote the space of finite point measures on \mathbb{R}^d endowed with the topology of vague convergence, and let $D([0,\infty),\mathcal{M})$ denote the space of càdlàg functions $\gamma : [0,\infty) \to \mathcal{M}$; this space of functions is endowed with the Skorokhod topology.

We now want to define the space to which the sequences of jump times and growth locations defined above belong, and endow this space with a topology. We let

$$\Theta := \left\{ (z_0, z_1, t_1, z_2, t_2, \ldots) \in \mathbb{R}^d \times (\mathbb{R}^d \times (0, \infty))^{\mathbb{N}} : \sum_{i=1}^\infty t_i = \infty \right\}.$$

The topology we take in Θ is the infinite product topology, where each copy of \mathbb{R}^d and each copy of $(0, \infty)$ is endowed with the corresponding Euclidean topology.

For $\mathbf{z} = (z_0, z_1, t_1, z_2, t_2, ...) \in \Theta$, define $\Gamma \mathbf{z} \in D([0, \infty), \mathcal{M})$ as follows (with the convention $t_0 = 0$):

$$(\Gamma \mathbf{z})(t) := \sum_{k=0}^{\infty} \delta_{\{z_k\}} \cdot \mathbb{1}\{t \ge t_k\}, \quad t \ge 0.$$

In words, Γz is the càdlàg function constructed from z by starting from $\delta_{\{z_0\}}$ at time zero, and prescribing that at each time of the form $\sum_{i=0}^{k} t_i$, a pointmass at z_k is added to the process (and the trajectory is kept constant between jumps). We then have

$$(\mathcal{H}_t^{\alpha\lambda,\lambda_I/\alpha})_{t\geq 0} = \Gamma(Z_{\alpha,0}, Z_{\alpha,1}, T_{\alpha,1}, Z_{\alpha,2}, T_{\alpha,2}, \ldots),$$

$$(\mathcal{T}_t^{\lambda,\lambda_I})_{t\geq 0} = \Gamma(Z_0, Z_1, T_1, Z_2, T_2, \ldots).$$

Using the definition of the Skorokhod topology, it is standard to prove that Γ is a homeomorphism between Θ and a measurable subset of $D([0,\infty), \mathcal{M})$. Hence, in order to prove the convergence stated in the theorem, it suffices to prove that

$$(Z_{\alpha,0}, Z_{\alpha,1}, T_{\alpha,1}, Z_{\alpha,2}, T_{\alpha,2}, \ldots) \xrightarrow{\alpha \to \infty} (Z_0, Z_1, T_1, Z_2, T_2, \ldots),$$

in distribution. Convergence in the infinite product topology is implied by convergence of all finite-dimensional distributions, so it is sufficient to prove that for all k we have

$$(Z_{\alpha,0}, Z_{\alpha,1}, T_{\alpha,1}, \dots, Z_{\alpha,k}, T_{\alpha,k}) \xrightarrow{\alpha \to \infty} (Z_0, Z_1, T_1, \dots, Z_k, T_k)$$

in distribution. In order to check this, by approximation arguments it is sufficient to check convergence of the integral of functions as taken in (4.1), so the result readily follows from Proposition 4.1. \Box

In order to prove Proposition 4.1, it will be important to have a more explicit description of the expectations that appear in (4.1). To this end, let us give some definitions. Recall that $\mathcal{P}_{\alpha\lambda}$ denotes a PPP on \mathbb{R}^d with intensity $\alpha\lambda$; we assume that this is the point process that gives rise to the infinite cluster in which the growth process $(\mathcal{H}_t^{\alpha\lambda,\lambda_{I}/\alpha})_{t\geq 0}$ is defined. Given a realization of $\mathcal{P}_{\alpha\lambda}$ and a finite set $S \subseteq \mathcal{P}_{\alpha\lambda}$, define

$$\mathcal{N}_{\alpha}(S) := \sum_{x \in S} \left| (\mathcal{P}_{\alpha\lambda} \cap B_r(x)) \backslash S \right| = \sum_{y \in \mathcal{P}_{\alpha\lambda} \backslash S} \left| \{ x \in S : \|x - y\| \le r \} \right|.$$

We now introduce probability kernels for each value of α ; these encode the jump rates of the dynamics of the growth process, conditioned on the realization of $\mathcal{P}_{\alpha\lambda}$. We start with the temporal kernel

$$\mathcal{L}_{\alpha}(S, \mathrm{d}t) := \frac{\mathcal{N}_{\alpha}(S)\lambda_{\mathrm{I}}}{\alpha} \cdot t \cdot \exp\left(-\frac{\mathcal{N}_{\alpha}(S)\lambda_{\mathrm{I}}}{\alpha} \cdot t\right) \mathrm{d}t,$$

where S is any finite subset of $\mathcal{P}_{\alpha\lambda}$ and $\mathcal{L}_{\alpha}(S, \cdot)$ gives a measure on the Borel sets of $(0, \infty)$, described above in terms of its density with respect to Lebesgue measure. Next, define the spatial kernel

$$\mathcal{K}_{\alpha}(S,A) := \frac{1}{\mathcal{N}_{\alpha}(S)} \cdot \sum_{y \in \mathcal{P}_{\alpha\lambda} \setminus S} \left| \{x \in S : \|x - y\| \le r\} \right| \cdot \delta_{\{y\}}(A).$$

where S is any finite subset of $\mathcal{P}_{\alpha\lambda}$ and A is a Borel subset of \mathbb{R}^d , so that $\mathcal{K}_{\alpha}(S, \cdot)$ gives a probability measure on \mathbb{R}^d . Finally, define the kernel

$$K_{\alpha}(S, \mathrm{d}(x, t)) := \mathcal{K}_{\alpha}(S, \mathrm{d}x) \otimes \mathcal{L}_{\alpha}(S, \mathrm{d}t),$$

that is, $K_{\alpha}(S, \cdot)$ is the measure on $\mathbb{R}^d \times (0, \infty)$ that satisfies, for any functions $f : \mathbb{R}^d \to \mathbb{R}$ and $g : (0, \infty) \to \mathbb{R}$ both continuous with compact support:

$$\int_{\mathbb{R}^d \times (0,\infty)} f(x)g(t) \ K_{\alpha}(S, \mathbf{d}(x, t)) = \int_{\mathbb{R}^d} f(x)\mathcal{K}_{\alpha}(S, \mathbf{d}x) \times \int_{(0,\infty)} g(t) \ \mathcal{L}_{\alpha}(S, \mathbf{d}t).$$

Now, using the strong Markov property, we can write

$$\mathbb{E}\left[f_{0}(Z_{\alpha,0})\prod_{i=1}^{k}f_{i}(Z_{\alpha,i})g_{i}(T_{\alpha,i})\right]$$

$$=\mathbb{E}\left[f_{0}(q(o))\int K_{\alpha}(\{q(o)\}, d(z_{1}, t_{1}))f_{1}(z_{1})g_{1}(t_{1}) \int K_{\alpha}(\{q(o), z_{1}\}, d(z_{2}, t_{2}))f_{2}(z_{2})g_{2}(t_{2})\cdots \int K_{\alpha}(\{q(o), z_{1}, \dots, z_{k-1}\}, d(z_{k}, t_{k}))f_{k}(z_{k})g_{k}(t_{k})\right].$$
(4.2)

DOI 10.20347/WIAS.PREPRINT.2877

Berlin 2021

We now define the analogous kernels for the limiting growth process. The temporal kernel is given by

$$\mathcal{L}(S, \mathrm{d}t) := |S| \mathbf{v}_d r^d \lambda \lambda_{\mathrm{I}} \cdot t \cdot \exp\left(-|S| \mathbf{v}_d r^d \lambda \lambda_{\mathrm{I}} \cdot t\right) \, \mathrm{d}t$$

where $S \subseteq \mathbb{R}^d$ is finite, and again we obtain a measure on $(0,\infty)$. Next, the spatial kernel is given by

$$\mathcal{K}(S, \mathrm{d}y) := \left(\frac{1}{|S|\boldsymbol{v}_d r^d} \sum_{x \in S} \mathbb{1}\{y \in B_r(x)\}\right) \mathrm{d}y,$$

for $S \subseteq \mathbb{R}^d$ finite. So $\mathcal{K}(S, \cdot)$ is the probability measure on \mathbb{R}^d obtained from first choosing $x \in S$ uniformly at random, and then choosing a point $y \in B_r(x)$ uniformly at random. We again let $K(S, d(x, t)) := \mathcal{K}(S, dx) \otimes \mathcal{L}(S, dt)$. Again by the Markov property, the equality in (4.2) holds for the limiting process with respect to this kernel (that is, the same equality with q(o) replaced by o and all α 's omitted).

The following will be the essential ingredient in the proof of Proposition 4.1.

Lemma 4.2. Let $f : \mathbb{R}^d \to \mathbb{R}$ and $g : (0, \infty) \to \mathbb{R}$ be both continuous with compact support, $k \in \mathbb{N}$ and $\varepsilon > 0$. There exists $\alpha_0 > 0$ such that for any $\alpha \ge \alpha_0$, we have that with probability larger than $1 - \varepsilon$, $\mathcal{P}_{\alpha\lambda}$ satisfies the following. For any set $S \subseteq B_{kr}(o) \cap \mathcal{P}_{\alpha\lambda}$ with $|S| \le k$, we have

$$\left| \int_{\mathbb{R}^d} f(x)g(t) \ K_{\alpha}(S, \mathbf{d}(x, t)) - \int_{\mathbb{R}^d} f(x)g(t) \ K(S, \mathbf{d}(x, t)) \right| < \varepsilon$$

We postpone the proof of this lemma; it will immediately follow from Lemma 4.4 and Lemma 4.2 below, which separately treat the spatial and temporal kernels. For now, let us show how Lemma 4.2 implies Proposition 4.1.

Proof of Proposition 4.1. We will abbreviate

$$[K_{\alpha}fg](S) := \int_{\mathbb{R}^d \times (0,\infty)} f(x)g(t)K_{\alpha}(S, \mathbf{d}(x, t)),$$

and also

$$S_{\alpha,k} := \{ Z_{\alpha,0}, Z_{\alpha,1}, \dots, Z_{\alpha,k} \}, \quad k \in \mathbb{N}_0,$$

and similarly when α is absent.

We proceed by induction on $k \in \mathbb{N}_0$. We interpret the case k = 0 to mean that

$$\mathbb{E}[f_0(Z_{\alpha,0})] \xrightarrow{\alpha \to \infty} f_0(0),$$

which holds because $Z_{\alpha,0} = q(o)$ converges in probability to o as $\alpha \to \infty$, as is easily seen. Now assume that the statement has been proved for $k - 1 \ge 0$, and take functions $f_1, \ldots, f_k, g_1, \ldots, g_k$ as in the statement. It will be convenient to add and subtract as follows (for k = 1, we interpret ' $\prod_{i=1}^{0}$ ' as being equal to one):

$$\mathbb{E}\left[f_0(Z_{\alpha,0})\prod_{i=1}^{k-1}f_i(Z_{\alpha,i})g_i(T_{\alpha,i})\right]$$
$$=\mathbb{E}\left[f_0(Z_{\alpha,0})\prod_{i=1}^{k-1}f_i(Z_{\alpha,i})g_i(T_{\alpha,i})\cdot[K_{\alpha}f_kg_k](S_{\alpha,k})\right]$$
$$=\mathbb{E}\left[\prod_{i=1}^{k-1}f_i(Z_{\alpha,i})g_i(T_{\alpha,i})\cdot\left([K_{\alpha}f_kg_k](S_{\alpha,k})\pm[Kf_kg_k](S_{\alpha,k})\right)\right]$$

Noting that the function that maps $(z_0, z_1, t_1, \ldots, z_{k-1}, t_{k-1})$ into

$$f_0(z_0) \prod_{i=1}^{k-1} f_i(z_i) g_i(t_i) \cdot [K f_k g_k](\{z_0, z_1, \dots, z_{k-1}\})$$

is continuous, the induction hypothesis and the definition of weak convergence give

$$\mathbb{E}\left[f_0(Z_{\alpha,0})\prod_{i=1}^{k-1}f_i(Z_{\alpha,i})g_i(T_{\alpha,i})[Kf_kg_k](S_{\alpha,k})\right]$$

$$\xrightarrow{\alpha\to\infty} \mathbb{E}\left[f_0(Z_0)\prod_{i=1}^{k-1}f_i(Z_i)g_i(T_i)[Kf_kg_k](S_k)\right] = \mathbb{E}\left[f_0(Z_0)\prod_{i=1}^kf_i(Z_i)g_i(T_i)\right].$$

Next, we bound

$$\mathbb{E}\left[f_0(Z_{\alpha,0})\prod_{i=1}^{k-1}f_i(Z_{\alpha,i})g_i(T_{\alpha,i})\cdot|[K_{\alpha}f_kg_k](S_{\alpha,k})-[Kf_kg_k](S_{\alpha,k})|\right]$$

$$\leq \left(\max_{i\leq k-1}(\|f_i\|_{\infty}\vee\|g_i\|_{\infty})\right)^k\cdot\mathbb{E}\left[|[K_{\alpha}f_kg_k](S_{\alpha,k})-[Kf_kg_k](S_{\alpha,k})|\right].$$

By Lemma 4.2, the right-hand side converges to zero as $\alpha \to \infty$. This completes the proof. \Box

It remains to prove Lemma 4.2. To do so, let us now introduce some more notation. For each $\delta > 0$, we define the collection of cubes

$$\mathcal{C}_{\delta} := \left\{ \delta z + \left[-\frac{\delta}{2}, \frac{\delta}{2} \right]^d : \ z \in \mathbb{Z}^d \right\}.$$

Additionally, given $\alpha > 0$, $\ell \in \mathbb{N}$ and $\delta > 0$, we define the event (involving the set $\mathcal{P}_{\alpha\lambda}$, but not the passage times):

$$\operatorname{REG}_{\alpha}(\ell,\delta) := \left\{ \left| \frac{|\mathcal{P}_{\alpha\lambda} \cap Q|}{\alpha\lambda\delta^d} - 1 \right| < \delta \text{ for all } Q \in \mathcal{C}_{\delta} \text{ with } Q \subseteq B_{\ell}(o) \right\}.$$

By the law of large numbers we have

$$\lim_{\alpha \to \infty} \mathbb{P}\left(\operatorname{REG}_{\alpha}(\ell, \delta)\right) = 1.$$
(4.3)

Lemma 4.3. For any $\varepsilon > 0$ and $k \in \mathbb{N}$ there exists $\delta_1 = \delta_1(\varepsilon, k)$ such that the following holds for $\delta \in (0, \delta_1]$. For any $\ell \ge r$, if α is large enough and the event $\operatorname{REG}_{\alpha}(\ell+1, \delta)$ occurs, then for any $S \subseteq \mathcal{P}_{\alpha\lambda} \cap B_{\ell-r}(o)$ with $|S| \le k$ we have

$$\sum_{x \in S} \sum_{Q \in \mathcal{C}_{\delta}} \left| \frac{\left| (\mathcal{P}_{\alpha\lambda} \cap Q \cap B_{r}(x)) \setminus S \right|}{\alpha \lambda} - \int_{Q} \mathbb{1}\{ \|y - x\| \le r \} \, \mathrm{d}y \right| < \varepsilon.$$
(4.4)

Proof. It is not hard to see that we can choose $\delta' > 0$ so that for any $\delta < \delta'$ we have

$$\delta^{d} \sup_{x \in \mathbb{R}^{d}} \sum_{Q \in \mathcal{C}_{\delta}} \mathbb{1}\{Q \cap \partial B_{r}(x) \neq \emptyset\} < \frac{\varepsilon}{3k},\tag{4.5}$$

DOI 10.20347/WIAS.PREPRINT.2877

Berlin 2021

where
$$\partial B_r(x) := \{y \in \mathbb{R}^d : ||x - y|| = r\}$$
. Next, we let $\delta_1 := \min\left(\delta', \frac{\varepsilon}{2\nu_d r^d}\right)$

Now, assume that $\delta < \delta_1$ and that $\operatorname{REG}_{\alpha}(\ell+1, \delta)$ occurs. Fix $S \subseteq \mathcal{P}_{\alpha\lambda} \cap B_{\ell-r}(o)$ with $|S| \leq k$, and also fix $x \in S$. For each $Q \in \mathcal{C}_{\delta}$ define

$$\mathscr{E}_x(Q) := \left| \frac{|(\mathcal{P}_{\alpha\lambda} \cap Q \cap B_r(x)) \setminus S|}{\alpha\lambda} - \int_Q \mathbb{1}\{||y - x|| \le r\} \, \mathrm{d}y \right|.$$

If $Q \cap \partial B_r(x) \neq \emptyset$ we bound

$$\mathscr{E}_x(Q) \le \frac{|\mathcal{P}_{\alpha\lambda} \cap Q|}{\alpha\lambda} + \delta^d \le (1+\delta)\delta^d + \delta^d = (2+\delta)\delta^d,$$

by the triangle inequality and the definition of $\operatorname{REG}_{\alpha}(\ell, \delta)$. If $Q \subseteq B_r(x)$ with $Q \cap \partial B_r(x) = \emptyset$ we have

$$\frac{|\mathcal{P}_{\alpha\lambda} \cap Q \cap B_r(x)|}{\alpha\lambda} = \frac{|\mathcal{P}_{\alpha\lambda} \cap Q|}{\alpha\lambda} \in ((1-\delta)\delta^d, (1+\delta)\delta^d),$$

so we bound

$$\mathscr{E}_x(Q) \le \frac{k}{\alpha\lambda} + \delta \cdot \delta^d \le 2\delta^{d+1}$$

(the factor $\frac{k}{\alpha\lambda}$ is there to account for the possibility that Q contains some points of S; the second inequality holds if α is large enough that $k/(\alpha\lambda) < \delta^{d+1}$). Now, also using (4.5), we have that the left-hand side of (4.4) is at most

$$\begin{split} \sum_{x \in S} \sum_{Q \in \mathcal{C}_{\delta}} \mathscr{E}_{x}(Q) &\leq \sum_{x \in S} \left((2+\delta)\delta^{d} \sum_{Q \in \mathcal{C}_{\delta}} \mathbb{1}\{Q \cap \partial B_{r}(x) \neq \varnothing\} + 2\delta^{d+1} \frac{\mathbf{v}_{d}r^{d}}{\delta^{d}} \right) \\ &\leq k \cdot \left(\frac{(2+\delta)\varepsilon}{3k} + 2\delta^{d+1} \frac{\mathbf{v}_{d}r^{d}}{\delta^{d}} \right). \end{split}$$

If δ is small enough, the right-hand side is smaller than ε , completing the proof.

Lemma 4.4. Let $f : \mathbb{R}^d \to \mathbb{R}$ be continuous with compact support, $k \in \mathbb{N}$ and $\varepsilon > 0$. There exists $\alpha_0 > 0$ such that for any $\alpha \ge \alpha_0$, we have that with probability larger than $1 - \varepsilon$, $\mathcal{P}_{\alpha\lambda}$ satisfies the following. For any set $S \subseteq B_{kr}(o) \cap \mathcal{P}_{\alpha\lambda}$ with $|S| \le k$, we have

$$\left|\int_{\mathbb{R}^d} f(x) \, \mathcal{K}_{\alpha}(S, \mathrm{d}x) - \int_{\mathbb{R}^d} f(x) \, \mathcal{K}(S, \mathrm{d}x)\right| < \varepsilon$$

Proof. Fix f, k and ε as in the statement of the lemma. We choose constants as follows:

since *f* is continuous with compact support, it is easy to see that we can choose δ_0 small enough that for any $\delta \in (0, \delta_0)$ we have

$$\sup_{\mu',\mu''} \left| \int_{\mathbb{R}^d} f \mathrm{d}\mu' - \int_{\mathbb{R}^d} f \mathrm{d}\mu'' \right| < \varepsilon,$$

where the supremum is taken over all pairs of probability measures μ',μ'' on Borel sets of \mathbb{R}^d with

$$\sum_{Q \in \mathcal{C}_{\delta}} |\mu'(Q) - \mu''(Q)| < \delta.$$
(4.6)

Berlin 2021

• next, let $\varepsilon' := \frac{\delta_0 v_d r^d}{2}$, and choose $\delta_1 = \delta_1(\varepsilon', k)$ as in Lemma 4.3.

Letting ℓ be large enough that the support of f is contained in $B_{\ell}(o)$, assume that the event $\operatorname{REG}_{\alpha}(\ell + rk + 1, \delta_1)$ occurs, and let $S \subseteq B_{rk}(o) \cap \mathcal{P}_{\alpha\lambda}$ be a set with at most k points. Using the triangle inequality and Lemma 4.3, we bound

$$\sum_{Q \in \mathcal{C}_{\delta_1}} \left| \frac{\mathcal{N}_{\alpha}(S)}{\alpha \lambda} \cdot \mathcal{K}_{\alpha}(S,Q) - |S| \mathbf{v}_d r^d \cdot \mathcal{K}(S,Q) \right|$$

$$\leq \sum_{x \in S} \sum_{Q \in \mathcal{C}_{\delta_1}} \left| \frac{|(\mathcal{P}_{\alpha \lambda} \cap Q \cap B_r(x)) \setminus S|}{\alpha \lambda} - \int_Q \mathbb{1}\{ \|y - x\| \le r \} \, \mathrm{d}y \right| \le \varepsilon'.$$
(4.7)

Using the fact that $\sum_Q \mathcal{K}_{\alpha}(S,Q) = \sum_Q \mathcal{K}(S,Q) = 1$, this readily gives

$$\left|\frac{\mathcal{N}_{\alpha}(S)}{\alpha\lambda} - |S|\boldsymbol{v}_{d}r^{d}\right| = \left|\frac{\mathcal{N}_{\alpha}(S)}{\alpha\lambda}\sum_{Q\in\mathcal{C}_{\delta}}\mathcal{K}_{\alpha}(S,Q) - |S|\boldsymbol{v}_{d}r^{d}\sum_{Q\in\mathcal{C}_{\delta}}\mathcal{K}(S,Q)\right|$$

$$\leq \sum_{Q\in\mathcal{C}_{\delta_{1}}}\left|\frac{\mathcal{N}_{\alpha}(S)}{\alpha\lambda}\cdot\mathcal{K}_{\alpha}(S,Q) - |S|\boldsymbol{v}_{d}r^{d}\cdot\mathcal{K}(S,Q)\right| \leq \varepsilon'.$$
(4.8)

Next, the triangle inequality gives, for any $Q \in C_{\delta}$,

$$\begin{aligned} |\mathcal{K}_{\alpha}(S,Q) - \mathcal{K}(S,Q)| &\leq \frac{1}{|S|\boldsymbol{v}_{d}r^{d}} \left(\left| |S|\boldsymbol{v}_{d}r^{d} - \frac{\mathcal{N}_{\alpha}(S)}{\alpha\lambda} \right| \cdot \mathcal{K}_{\alpha}(S,Q) \right. \\ &+ \left| \frac{\mathcal{N}_{\alpha}(S)}{\alpha\lambda} \cdot \mathcal{K}_{\alpha}(S,Q) - |S|\boldsymbol{v}_{d}r^{d} \cdot \mathcal{K}(S,Q) \right| \right). \end{aligned}$$

Combining this with (4.7) and (4.8), we obtain

$$\sum_{Q \in \mathcal{C}_{\delta_1}} |\mathcal{K}_{\alpha}(S,Q) - \mathcal{K}(S,Q)| \le \frac{\varepsilon'}{|S| \boldsymbol{v}_d r^d} \left(\sum_{Q \in \mathcal{C}_{\delta_1}} \mathcal{K}_{\alpha}(S,Q) + 1 \right) \le \frac{2\varepsilon'}{\boldsymbol{v}_d r^d} = \delta_0.$$

This shows that $\mathcal{K}_{\alpha}(S, \cdot)$ and $\mathcal{K}(S, \cdot)$ are close enough in the sense that (4.6) is satisfied. The proof is now completed using (4.3).

The following is proved in a similar manner as Lemma 4.4, only simpler, so we omit the details.

Lemma 4.5. Let $g: (0, \infty) \to \mathbb{R}$ be continuous with compact support, $k \in \mathbb{N}$ and $\varepsilon > 0$. There exists $\alpha_0 > 0$ such that for any $\alpha \ge \alpha_0$, we have that with probability larger than $1 - \varepsilon$, $\mathcal{P}_{\alpha\lambda}$ satisfies the following. For any set $S \subseteq B_{kr}(o) \cap \mathcal{P}_{\alpha\lambda}$ with $|S| \le k$, we have

$$\left|\int_{(0,\infty)} g(t) \mathcal{L}_{\alpha}(S, \mathrm{d}t) - \int_{(0,\infty)} g(t) \mathcal{L}(S, \mathrm{d}t)\right| < \varepsilon.$$

References

 [1] A. Auffinger, M. Damron, and J. Hanson. 50 years of first-passage percolation, volume 68 of University Lecture Series. American Mathematical Society, Providence, RI, 2017. ISBN 978-1-4704-4183-8. doi: 10.1090/ulect/068. URL https://doi.org/10.1090/ulect/068.

- M. Deijfen. Asymptotic shape in a continuum growth model. *Adv. in Appl. Probab.*, 35(2):303–318, 2003. ISSN 0001-8678. doi: 10.1239/aap/1051201647. URL https://doi.org/10.1239/aap/1051201647.
- [3] J. Hammersley and D. Welsh. First-passage percolation, subadditive processes, stochastic networks, and generalized renewal theory. In *Proc. Internat. Res. Semin., Statist. Lab., Univ. California, Berkeley, Calif*, pages 61–110. Springer-Verlag, New York, 1965.
- [4] C. Hirsch, D. Neuhäuser, C. Gloaguen, and V. Schmidt. First passage percolation on random geometric graphs and an application to shortest-path trees. *Adv. in Appl. Probab.*, 47(2):328–354, 2015. ISSN 0001-8678. doi: 10.1239/aap/1435236978. URL https://doi.org/10.1239/aap/1435236978.
- [5] B. Jahnel and W. König. Probabilistic Methods In Telecommunications. Compact Textbooks in Mathematics. Birkhäuser Basel, 2020. ISBN 978-3-030-36090-0. doi: 10.1007/978-3-030-36090-0. URL https://doi.org/10.1007/978-3-030-36090-0.
- [6] J. Kingman. The ergodic theory of subadditive stochastic processes. J. Roy. Statist. Soc. Ser. B, 30:499-510, 1968. ISSN 0035-9246. URL http://links.jstor.org/sici?sici= 0035-9246(1968)30:3<499:TETOSS>2.0.CO;2-E&origin=MSN.
- [7] R. Meester and R. Roy. Continuum percolation, volume 119 of Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 1996. ISBN 0-521-47504-X. doi: 10.1017/ CBO9780511895357. URL https://doi.org/10.1017/CBO9780511895357.
- [8] L. Ménard and A. Singh. Percolation by cumulative merging and phase transition for the contact process on random graphs. Ann. Sci. Éc. Norm. Supér. (4), 49(5):1189–1238, 2016. ISSN 0012-9593. doi: 10.24033/asens.2307. URL https://doi.org/10.24033/asens.2307.
- [9] M. Penrose. Random geometric graphs, volume 5 of Oxford Studies in Probability. Oxford University Press, Oxford, 2003. ISBN 0-19-850626-0. doi: 10.1093/acprof:oso/9780198506263.001.0001. URL https://doi.org/10.1093/acprof:oso/9780198506263.001.0001.
- [10] M. Penrose and A. Pisztora. Large deviations for discrete and continuous percolation. Adv. in Appl. Probab., 28(1):29-52, 1996. ISSN 0001-8678. doi: 10.2307/1427912. URL https://doi.org/10.2307/1427912.
- [11] T. Riblet. Le processus de contact sur le graphe Booléen. PhD thesis, Université de Lorraine, 2019.
- [12] D. Richardson. Random growth in a tessellation. *Proc. Cambridge Philos. Soc.*, 74:515–528, 1973.
 ISSN 0008-1981. doi: 10.1017/s0305004100077288. URL https://doi.org/10.1017/s0305004100077288.
- [13] H. Rosenthal. On the subspaces of L^p (p > 2) spanned by sequences of independent random variables. *Israel J. Math.*, 8:273–303, 1970. ISSN 0021-2172. doi: 10.1007/BF02771562. URL https://doi.org/10.1007/BF02771562.
- [14] S. Torquato and Y. Jiao. Effect of dimensionality on the continuum percolation of overlapping hyperspheres and hypercubes. ii. simulation results and analyses. *The Journal of Chemical Physics*, 137(7):074106, 2012. doi: 10.1063/1.4742750. URL https://doi.org/10.1063/ 1.4742750.
- [15] C.-L. Yao, G. Chen, and T.-D. Guo. Large deviations for the graph distance in supercritical continuum percolation. J. Appl. Probab., 48(1):154–172, 2011. ISSN 0021-9002. doi: 10.1239/jap/ 1300198142. URL https://doi.org/10.1239/jap/1300198142.