

**Periodic L_p estimates by \mathcal{R} -boundedness:
Applications to the Navier–Stokes equations**

Thomas Eiter¹, Mads Kyed², Yoshihiro Shibata³

submitted: April 26, 2022

¹ Weierstrass Institute
Mohrenstr. 39
10117 Berlin
Germany
E-Mail: thomas.eiter@wias-berlin.de

² Hochschule Flensburg
Kanzleistraße 91-93
24943 Flensburg
Germany
E-Mail: mads.kyed@hs-flensburg.de

³ Department of Mathematics
Waseda University
Ohkubo 3-4-1, Shinjuku-ku
Tokyo 169-8555
Japan
E-Mail: yshibata325@gmail.com

No. 2931
Berlin 2022



2020 *Mathematics Subject Classification.* 47J35, 35K90, 35B10, 35B45.

Key words and phrases. Evolution equations, time-periodic solutions, L_p estimates, Navier–Stokes equations, inhomogeneous boundary data.

M. Kyed: Joint Appointment Professor of Waseda University, Tokyo, Japan

Y. Shibata: Adjunct faculty member in the Department of Mechanical Engineering and Materials Science, University of Pittsburgh, USA; Partially supported by Top Global University Project, JSPS Grant-in-aid for Scientific Research (A) 17H0109, and Toyota Central Research Institute Joint Research Fund.

Edited by
Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS)
Leibniz-Institut im Forschungsverbund Berlin e. V.
Mohrenstraße 39
10117 Berlin
Germany

Fax: +49 30 20372-303
E-Mail: preprint@wias-berlin.de
World Wide Web: <http://www.wias-berlin.de/>

Periodic L_p estimates by \mathcal{R} -boundedness: Applications to the Navier–Stokes equations

Thomas Eiter, Mads Kyed, Yoshihiro Shibata

Abstract

General evolution equations in Banach spaces are investigated. Based on an operator-valued version of de Leeuw’s transference principle, time-periodic L_p estimates of maximal regularity type are established from \mathcal{R} -bounds of the family of solution operators (\mathcal{R} -solvers) to the corresponding resolvent problems. With this method, existence of time-periodic solutions to the Navier–Stokes equations is shown for two configurations: in a periodically moving bounded domain and in an exterior domain, subject to prescribed time-periodic forcing and boundary data.

1 Introduction

The study of time-periodic solutions to evolution equations is the study of oscillations. In this article we investigate time-periodic solutions corresponding to time-periodic data, that is, systems of forced oscillation. A number of different methods, further described below, are traditionally used to carry out a mathematical investigation of such solutions. In the following we introduce a new technique to establish *a priori* estimates of maximal L_p regularity type for linearized equations. Such estimates are essential in the study of nonlinear problems, which we will demonstrate on some examples. For notational simplicity, we consider only 2π -periodic problems. By a simple scaling argument, however, all our results extend to \mathcal{T} -periodic problems for any $\mathcal{T} > 0$.

The study of 2π -time-periodic solutions to evolution equations can be carried out in a framework where the time axis is replaced with a torus $\mathbb{T} := \mathbb{R}/2\pi\mathbb{Z}$. Consider for example an abstract evolution equation

$$\partial_t u + Au = f \text{ in } \mathbb{T} \tag{1.1}$$

in a Banach space X , where A is a linear operator on X . Since the time domain is a torus, a solution to (1.1) is intrinsically time-periodic. We refer to estimates of the solution in $L_p(\mathbb{T}; X)$ norms as *periodic L_p estimates*. Estimates that include all highest-order norms of the solution are said to be of maximal regularity, which in the case (1.1) above means an estimate of type

$$\|\partial_t u\|_{L_p(\mathbb{T}, X)} + \|Au\|_{L_p(\mathbb{T}, X)} + \|u\|_{L_p(\mathbb{T}, X)} \leq C\|f\|_{L_p(\mathbb{T}, X)}. \tag{1.2}$$

Such estimates lead to a characterization of $\partial_t + A$ as a homeomorphism in an $L_p(\mathbb{T}, X)$ setting, which is critical to the analysis of non-linear problems. In the following, we show how to establish periodic L_p estimates of maximal regularity type for a large class of abstract evolution equations based on their \mathcal{R} -solvers, that is, solution operators of the associated resolvent problems that satisfy specific \mathcal{R} -bounds. In particular, we include cases where 0 lies in the spectrum of A , which constitutes a particular challenge and where traditional methods have shortcomings. As we explain below in more detail, in this case the classical maximal regularity estimate (1.2) is not available and $\partial_t + A$ can only be realized as a homeomorphism in an adapted framework of function spaces.

Typically, L_p estimates are established via Fourier multipliers, often via the multiplier theorem of Mikhlin, which was extended to operator-valued multipliers by WEIS [67]. He showed that Mikhlin's theorem remains valid in the operator-valued case if boundedness is replaced with \mathcal{R} -boundedness in the assumptions. On the strength of this result, it is possible to establish maximal regularity in $L_p((0, T), X)$ norms for initial-value problems such as

$$\partial_t u + Au = f \quad \text{in } (0, T), \quad u(0) = u_0 \quad (1.3)$$

by establishing \mathcal{R} -boundedness on the resolvent family

$$\{\lambda R(\lambda, A) \mid \lambda \in \Sigma_{\theta, \gamma_0}\}, \quad (1.4)$$

where $R(\lambda, A) = (\lambda I - A)^{-1}$ denotes the resolvent operator of A and

$$\Sigma_{\theta, \gamma_0} := \{\lambda \in \mathbb{C} \mid \arg(\lambda) \leq \pi - \theta, \quad |\lambda| \geq \lambda_0\} \quad (0 \leq \theta < \frac{\pi}{2})$$

is a sector, which excludes a ball around the origin if $\lambda_0 > 0$. Note that for many problems, in particular in unbounded domains, the inclusion of the origin is not possible since 0 does not belong to the resolvent set. However, for the derivation of L_p estimates for the initial-value problem (1.3), the origin can be excluded from the sector unless $T = \infty$ is required. Since the appearance of [67], \mathcal{R} -bounds for resolvent families of the form (1.4) have been established for a substantial number of boundary-value problems, which lead to L_p maximal regularity estimates for the associated initial-value problem.

In this article, we develop a technique to obtain periodic L_p estimates of maximal regularity type from these \mathcal{R} -bounds. In particular, we focus on problems where the origin belongs to the spectrum $\sigma(A)$ of A , so that \mathcal{R} -bounds can at best be established in $\Sigma_{\theta, \gamma_0}$ for some $\gamma_0 > 0$. In the case that 0 is included in the resolvent set as well as in the \mathcal{R} -bounds, that is, the operator family (1.4) is \mathcal{R} -bounded for $\gamma_0 = 0$, classical maximal L_p regularity can be established, which was shown by ARENDT and BU [2]. If this is not the case, classical periodic L_p estimates of maximal regularity type such as (1.2) cannot be established because invertibility of the linear time-periodic problem would require invertibility of the associated steady-state problem, that is, that 0 belongs to the resolvent set $\rho(A)$. Therefore, we introduce an alternative functional setting in order to characterize the parabolic operator $\partial_t + A$ as a homeomorphism with respect to data in L_p spaces.

The technique developed in the following is based on the transference principle introduced by DE LEEUW in [19] for scalar-valued multipliers and generalized to the operator-valued case by HYTÖNEN, VAN NEERVEN, VERAAR, and WEIS [28]. It states that L_p boundedness of a continuous Fourier multiplier on $L_p(\mathbb{R})$ is retained when the multiplier is restricted to \mathbb{Z} and thus becomes a multiplier in the torus $\mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$ setting. Despite the transference principle seeing little usage outside the field of harmonic analysis, we believe it to be an effective tool in the analysis of periodic solutions to partial differential equations. The promotion of this viewpoint is one of the main purposes of this article since it provides us with an extremely useful tool to derive periodic L_p estimates from existing \mathcal{R} -bounds on resolvent families. If for example the resolvent family (1.4) is \mathcal{R} -bounded in a sector containing the full imaginary axis $i\mathbb{R}$, that is, for $\gamma_0 = 0$, then the operator-valued version of the transference principle combined with the operator-valued version of Mikhlin's multiplier theorem by WEIS [67] immediately yields the periodic L_p estimates (1.2). If, however, \mathcal{R} -bounds are only available in a sector excluding the origin, that is, $\gamma_0 > 0$, a decomposition technique has to be introduced. Expanding (1.1) into a Fourier series, we introduce the projection of u into a lower frequency part $u_\ell := \sum_{|k| \leq \gamma_0} u_k e^{ikt}$ corresponding to the finite number of modes $k \in \mathbb{Z}$ with $|k| \leq \gamma_0$, and a complementary higher frequency part $u_h = u - u_\ell$. Based on the \mathcal{R} -bounds of (1.4), we can combine the transference

principle with Mikhlin's multiplier theorem to establish periodic L_p estimates for the higher frequency part. Provided that a (possibly different) framework of Banach spaces can be identified that ensures periodic maximal L_p regularity for the lower frequency part, we can combine the two parts to establish periodic maximal L_p regularity for the full problem. Due to the bespoke framework introduced for the lower frequency part, the resulting type of L_p estimates are not classical in the sense of (1.2). They are, however, effective in the investigation of time-periodic solutions to corresponding nonlinear problems, which we demonstrate by specific examples.

Whereas the study of time-periodic solutions to ordinary differential equations goes back to the nineteenth century, one of the first investigations of time-periodic partial differential equations is due to PRODI [47], who examined the (parabolic) 1D heat equation. Although the work of PRODI is predated by a few other articles [3, 61, 31, 69], it seems that [47] is the first rigorous, by contemporary standards, investigation into the matter. Around the same time, articles also appeared on time-periodic solutions to the (hyperbolic) wave equation [48, 21]. In the following years, the foundation was laid for the methods that have nowadays become standard in the study of time-periodic partial differential equations. We shall give a brief overview of the main ideas. Consider for this purpose a time-periodic abstract evolution equation

$$\partial_t u + Au = F(t, u) \text{ in } \mathbb{R}, \quad u(t + 2\pi) = u(t) \quad (1.5)$$

in the classical setting with the whole of \mathbb{R} as time axis, but still considered as equation in a Banach space X for some operator A and for 2π -time-periodic (nonlinear) data F .

By far the most popular method that emerged is based on the identification of solutions to (1.5) as fixed points of the so-called Poincaré operator¹. The basic idea goes back to the pioneering work of Poincaré [45, 46] on dynamical systems. The Poincaré operator, sometimes also referred to as *translation operator along trajectories*, is the mapping $\Phi : X \rightarrow X$ that maps an initial value u_0 to the value $u(2\pi)$ of the solution u to the associated initial-value problem

$$\partial_t u + Au = F(t, u) \text{ in } (0, \infty), \quad u(0) = u_0. \quad (1.6)$$

In other words, if $t \mapsto S(t, u_0)$ is the solution operator to (1.5) for the initial value u_0 , the Poincaré operator is given by $\Phi(u_0) := S(2\pi, u_0)$. It is obvious that a fixed point $w_0 = \Phi(w_0)$ of Φ is the initial value of a 2π -periodic solution. In this sense a fixed point of the Poincaré operator induces a solution to (1.5). The main challenge in the application of this method is to construct a setting of Banach spaces such that the Poincaré operator is well defined and admits a fixed point. In some cases, this can be carried out directly for the nonlinear problem, but often the method is first employed to obtain suitable *a priori* estimates of maximal regularity type for the linearization of (1.5), which are subsequently used to investigate the nonlinear problem with classical nonlinear functional analysis. In the context of time-periodic partial differential equations, BROWDER introduced the Poincaré operator approach in [9], and around the same time KRASNOSEL'SKIĬ [32] and his student KOLESOV [35, 36, 37] advanced the method. The investigation of time-periodic solutions as fixed points of the Poincaré operator depends heavily on the framework in which a solution operator to the initial-value problem (1.6) can be realized. To illustrate this issue, assume that A generates a sufficiently regular semi-group and consider the linear case $F(t, u) = F(t)$. The solution operator S then takes the form

$$S(t, u_0) := e^{-tA} u_0 + \int_0^t e^{-(t-\tau)A} F(\tau) \, d\tau,$$

¹Not to be confused with the *Poincaré mapping*, which is a related but different notion from the theory of dynamical systems.

and a fixed point w_0 of the Poincaré operator is therefore given by

$$w_0 = S(2\pi, w_0) \iff w_0 = (I - e^{-2\pi A})^{-1} \int_0^{2\pi} e^{-(2\pi-\tau)A} F(\tau) d\tau$$

provided $0 \in \rho(A)$, so that $1 \in \rho(e^{-2\pi A})$ and $I - e^{-2\pi A}$ is thereby invertible. In this case, *a priori* estimates for the time-periodic solution $u(t) := S(t, w_0)$ can be established in the setting of Banach spaces in which S is realized. If, on the other hand, $0 \in \sigma(A)$, the representation formula for w_0 above is not valid and it becomes much more difficult to establish *a priori* estimates for the corresponding time-periodic solution. For this reason, the Poincaré operator approach is seemingly always carried out in a setting where $0 \in \rho(A)$. General applications of the method can be found in articles going back to [1, 51, 62, 66] for example, but also in more recent work such as [40, 4, 13, 34, 44]. More examples can be found in articles devoted to specific equations; so many that an exhaustive list is beyond the scope of our exposition here. We shall mention only the work of GEISSERT, HIEBER and NGUYEN [26] in which the restriction $0 \in \rho(A)$ is circumvented by introducing interpolation spaces.

Provided one is able to establish suitable energy estimates for the problem under consideration, time-periodic solutions can also be obtained via a Galerkin approximation scheme. The existence of a time-periodic solution then has to be accomplished in a finite-dimensional setting and is thus reduced to finding periodic solutions to an ordinary differential equation. In the finite-dimensional setting, the Poincaré operator is compact, and it is therefore much less critical to establish existence of a fixed point. The time-periodic incompressible Navier–Stokes problem is a good example of a system that can be treated with energy methods; see for example [49, 24]. Also the time-periodic wave equation with suitable damping can be solved in this way [50]. Moreover, time-periodic solutions to the compressible Navier–Stokes equations can be established, as was first shown for the one-dimensional case by MATSUMURA and NISHIDA [42] and then extended by several authors, *cf.* [30, 64] and the references therein. Since energy estimates typically lead to *a priori* estimates in Hilbert space settings, this method is not always suited to establish optimal *a priori* estimates for linear parabolic problems though. Nevertheless, it gives a strong tool in the case of hyperbolic or hyperbolic-parabolic mixed systems.

We also want to mention a very different method, which is due to SEIDMAN [54], who intentionally avoids using the Poincaré operator and shows existence of weak time-periodic solutions in $L_p(\mathbb{T} \times \Omega)$ spaces to a nonlinear evolution equation based on the theory of monotone operators.

A different approach is based on a representation formula that arises from the principle that a solution to the initial-value problem (1.6) (at least in the linear case $F(t, u) = F(t)$) tends to a periodic orbit as $t \rightarrow \infty$ regardless of the initial value. Equivalently formulated, a solution to the initial-value problem with time-periodic right-hand side

$$\partial_t u + Au = F(t) \text{ in } (R, \infty), \quad u(R) = u_0,$$

tends to a periodic orbit as $R \rightarrow -\infty$. Assuming again that A generates a sufficiently regular semi-group, this principle leads to the formula

$$u(t) = \int_{-\infty}^t e^{-(t-\tau)A} F(\tau) d\tau \tag{1.7}$$

for the time-periodic solution. It is easy to verify that this integral expression indeed leads to a periodic solution of the same period as F . As with the Poincaré operator approach, the challenge with the method based on (1.7) is to construct a framework of Banach spaces such that the integral expression is well defined. Since F is time-periodic and therefore non-decaying, this clearly requires suitable

decay properties of the semi-group, which again leads to $0 \notin \sigma(A)$ as a critical assumption. Under this assumption, however, the representation (1.7) can be very useful, which was demonstrated already in the paper [47] by PRODI. A similar idea was used by FIFE [22] and in subsequent papers [63, 5, 43, 23] as well as a number of articles on specific equations such as the Navier–Stokes equations [38, 68].

The principle described above gives rise to yet another approach. If namely the solution $u(t)$ to the initial-value problem (1.6) tends to a periodic orbit as $t \rightarrow \infty$, then the sequence $u_n(t) := u(t+n2\pi)$, $n \in \mathbb{N}$, will tend to a periodic solution to (1.5) as $n \rightarrow \infty$. This idea was employed in the context of partial differential equations by FICKEN and FLEISHMAN [21] as early as 1957, and later used to investigate time-periodic solutions to the Navier–Stokes equations in the incompressible case [55, 41] and the compressible case [65].

Finally, we mention the perhaps most natural approach to time-periodic partial differential equations, namely the decomposition of data and solution into a Fourier series with respect to time. In the linear case, the investigation then reduces to an analysis of the individual Fourier coefficients, each of which satisfying a resolvent problem, that is, a time-independent problem; see for example [48, 12, 52, 53, 27, 11]. This technique, however, has some limitations since it is difficult to obtain satisfactory estimates of the Fourier series based only on estimates of the individual coefficients. Typically, the method only leads to suitable *a priori* estimates when working in a framework of absolutely convergent Fourier series, see the recent articles [18, 14, 15] for examples from fluid dynamics, or when Parseval’s identity can be invoked, which requires a Hilbert space setting, that is, an investigation in $L_2((0, 2\pi), H)$ for some Hilbert space H . Our following analysis based on Fourier multipliers on the torus \mathbb{T} offers one way to overcome these limitations and to establish *a priori* estimates in a general $L_p((0, 2\pi), X)$ Banach space setting.

In order to illustrate another significant novelty of our approach, we return to the requirement $0 \notin \sigma(A)$ that is needed in both the Poincaré operator approach and in the method based on the representation formula (1.7). The root cause of this restriction is the necessity in both techniques that the investigation of (1.5) is carried out in the framework of function spaces of the corresponding initial-value problem (1.6). Specifically, in both methods the time-periodic solution is characterized as a special solution to the initial-value problem, and can therefore only be estimated in the framework in which the initial-value problem is rendered well posed. However, this framework is not suitable for *a priori* estimates of solutions to the corresponding stationary problem $Au = F$ when $0 \in \sigma(A)$. Since a stationary solution is trivially also time periodic, it is clear why the restriction $0 \notin \sigma(A)$ is imposed. In our approach, based solely on Fourier multipliers, both the Poincaré operator and the representation formula (1.7) are avoided, and we are able to construct a bespoke setting of Banach spaces that enables us to also treat cases where $0 \in \sigma(A)$.

The article is divided into a more theoretical first part (Section 3), where an abstract linear time-periodic problem is investigated, and a second part devoted to applications (Sections 4 and 5). The theoretical part focuses on an abstract time-periodic boundary-value problem, and we show statements in general terms of the periodic maximal L_p regularity. It is based on a combination of the decomposition technique described above with the existence of suitable \mathcal{R} -solvers; see Theorems 3.1 and 3.2. As examples of the effectiveness of this approach, we subsequently investigate time-periodic solutions to the the N -dimensional Navier–Stokes equations in a periodically moving bounded domain in Section 4, and to the three-dimensional Navier–Stokes equations in an exterior domain (at rest) in Section 5. Existence of time-periodic solutions to the first problem can be shown in a framework of Sobolev spaces, and in the final results the described decomposition technique is not visible, which is the case since 0 is not in the spectrum of the underlying linear operator. However, for the second problem the situation is different, and the zero-order mode, which is the time mean of the periodic function, has

to be treated in a separate functional framework. Moreover, in order to handle the nonlinear terms, we consider spaces of functions with additional pointwise spatial decay, and a large part of Section 5 is concerned with the asymptotic properties of solutions. Since this analysis of the exterior-domain problem is already quite extensive, the even more involved case of the Navier–Stokes flow inside a periodically moving exterior domain is postponed to a future work.

2 Notation and preliminaries

2.1 General notation

Let \mathbb{N} , \mathbb{Z} , \mathbb{R} and \mathbb{C} denote the set of all natural numbers, integers, and real and complex numbers, respectively. To denote generic constants, we use the symbol C , and $C_{a,b,\dots}$ indicates the dependency of the constant on the quantities a, b, \dots . Here, the constants C and $C_{a,b,\dots}$ may change from line to line.

For any domain $D \subset \mathbb{R}^N$, $N \in \mathbb{N}$, we denote Lebesgue spaces, Sobolev spaces, and Besov spaces on D by $L_q(D)$, $H_q^m(D)$ and $B_{q,p}^s(D)$, respectively, while $\|\cdot\|_{L_q(D)}$, $\|\cdot\|_{H_q^m(D)}$, and $\|\cdot\|_{B_{q,p}^s(D)}$ denote their norms. For partial derivatives, we write $\partial_t = \partial/\partial t$ and $\partial_j = \partial/\partial x_j$. Let $\nabla f = (\partial_1 f, \partial_2 f, \partial_3 f)$ and $\nabla^2 f = (\partial_i \partial_j f \mid i, j = 1, 2, 3)$. Let $\hat{H}_q^1(D)$ be the homogeneous space defined by

$$\hat{H}_q^1(D) = \{\varphi \in L_{q,\text{loc}}(D) \mid \nabla \varphi \in L_q(D)^N\}.$$

For a topological vector space V , we let V' denote its dual space. In the following, X and Y will always denote Banach spaces, and $\mathcal{L}(X, Y)$ denotes the space of bounded linear operators from X to Y , and we simply write $\mathcal{L}(X) = \mathcal{L}(X, X)$. Sometimes, we do not distinguish between a space X and its vector-valued analog X^N , and we simply write $\|\cdot\|_X$ for the norm of X^N . The set of all X -valued holomorphic functions defined on $U \subset \mathbb{C}$ is denoted by $\text{Hol}(U, X)$. For $\varepsilon \in (0, \pi)$ and $\lambda_0 > 0$ we define the sectors

$$\Sigma_\varepsilon = \{\lambda \in \mathbb{C} \setminus \{0\} \mid |\arg \lambda| \leq \pi - \varepsilon\}, \quad \Sigma_{\varepsilon, \lambda_0} = \{\lambda \in \Sigma_\varepsilon \mid |\lambda| \geq \lambda_0\}.$$

2.2 Time-periodic framework

The study of partial differential equations in a setting where both the data and the corresponding solutions are time periodic can conveniently be carried out in a framework where the time axis is replaced with a torus group. In the following, we consider only the torus

$$\mathbb{T} := \mathbb{R}/2\pi\mathbb{Z},$$

which provides us with a framework to study 2π -periodic solutions. We endow \mathbb{T} with the quotient topology inherited from \mathbb{R} via the quotient mapping

$$\pi_Q : \mathbb{R} \rightarrow \mathbb{T}, \quad \pi_Q(t) := [t] = \{t + 2n\pi \mid n \in \mathbb{Z}\}.$$

Additionally, the quotient mapping induces a differentiable structure on the torus, and we can therefore investigate equations such as (1.1) as a differential equation on the smooth manifold \mathbb{T} . A solution u in this setting corresponds to a classical time-periodic solution $u \circ \pi_Q$ in the Euclidean setting and vice versa. Usually, we tacitly identify functions u on \mathbb{T} with their time-periodic analogue $u \circ \pi_Q$ on \mathbb{R} .

The topology on \mathbb{T} turns it into a compact group with a (normalized) Haar measure $d\tau$ such that

$$\forall u \in C(\mathbb{T}) : \int_{\mathbb{T}} u(\tau) d\tau = \frac{1}{2\pi} \int_0^{2\pi} u \circ \pi_Q(t) dt,$$

where $C(\mathbb{T})$ is the class of all continuous functions on \mathbb{T} . Bochner–Lebesgue spaces $L_p(\mathbb{T}, X)$ for $p \in [1, \infty]$ are then defined in the usual manner.

The differentiable structure gives rise to the space

$$C^\infty(\mathbb{T}, X) := \{u : \mathbb{T} \rightarrow X \mid u \circ \pi_Q \in C^\infty(\mathbb{R}, X)\}$$

of vector-valued smooth functions on \mathbb{T} for any Banach space X . The simple structure of \mathbb{T} implies that the set of Schwartz–Bruhat functions $\mathcal{S}(\mathbb{T}, X)$ (see [10, 16]) coincides with the set of smooth functions, that is,

$$\mathcal{S}(\mathbb{T}, X) = C^\infty(\mathbb{T}, X),$$

which is endowed with the semi-norm topology induced by the family $\rho_\ell(u) := \sup_{\tau \in \mathbb{T}} \|\partial_t^\ell u(\tau)\|_X$, $\ell \in \mathbb{N}_0$. We refer to the space

$$\mathcal{S}'(\mathbb{T}, X) = \mathcal{L}(\mathcal{S}(\mathbb{T}, X))$$

as the space of X -valued tempered distributions on \mathbb{T} . One may observe that the notion of classical distributions on \mathbb{T} (also known as periodic distributions) coincides with the notion of tempered distributions on \mathbb{T} . Derivatives of distributions $u \in \mathcal{S}'(\mathbb{T}, X)$ are defined as distributions $\partial_t^\ell u \in \mathcal{S}'(\mathbb{T}, X)$ by duality in the usual way.

As a (locally) compact abelian group, the torus \mathbb{T} has a Fourier transform $\mathcal{F}_{\mathbb{T}}$ associated to it. Obviously, this Fourier transform corresponds to the classic expansion of a function on \mathbb{T} into a Fourier series. In this paper, however, it is essential to treat it as Fourier transform in the same framework as the Fourier transform $\mathcal{F}_{\mathbb{R}}$ on the real line, which is defined by

$$\begin{aligned} \mathcal{F}_{\mathbb{R}} : \mathcal{S}(\mathbb{R}, X) &\rightarrow \mathcal{S}(\mathbb{R}, X), & \mathcal{F}_{\mathbb{R}}[u](\xi) &:= \frac{1}{2\pi} \int_{\mathbb{R}} u(x) e^{-i\xi x} dx, \\ \mathcal{F}_{\mathbb{R}}^{-1} : \mathcal{S}(\mathbb{R}, X) &\rightarrow \mathcal{S}(\mathbb{R}, X), & \mathcal{F}_{\mathbb{R}}^{-1}[v](x) &:= \int_{\mathbb{R}} v(\xi) e^{i\xi x} d\xi \end{aligned}$$

and extended to mappings $\mathcal{F}_{\mathbb{R}}, \mathcal{F}_{\mathbb{R}}^{-1} : \mathcal{S}'(\mathbb{R}, X') \rightarrow \mathcal{S}'(\mathbb{R}, X')$ by duality. To this end, we recall that \mathbb{Z} , endowed with discrete topology and counting measure, can be viewed as the dual group of \mathbb{T} . The X -valued Schwartz–Bruhat space on \mathbb{Z} is given by

$$\mathcal{S}(\mathbb{Z}, X) = \{\psi : \mathbb{Z} \rightarrow X \mid \forall \ell \in \mathbb{N}_0 : \sup_{k \in \mathbb{Z}} |k|^\ell \|\psi(k)\|_X < \infty\}$$

and is equipped with the locally convex topology induced by the family of semi-norms $\hat{\rho}_\ell$, $\ell \in \mathbb{N}_0$, where $\hat{\rho}_\ell(\psi) := \sup_{k \in \mathbb{Z}} |k|^\ell \|\psi(k)\|_X$.

In the setting of vector-valued Schwartz–Bruhat spaces, the Fourier transform $\mathcal{F}_{\mathbb{T}}$ is the homeomorphism given by

$$\mathcal{F}_{\mathbb{T}} : \mathcal{S}(\mathbb{T}, X) \rightarrow \mathcal{S}(\mathbb{Z}, X), \quad \mathcal{F}_{\mathbb{T}}[u](k) := \int_{\mathbb{T}} u(t) e^{-ikt} dt,$$

with inverse mapping

$$\mathcal{F}_{\mathbb{T}}^{-1} : \mathcal{S}(\mathbb{Z}, X) \rightarrow \mathcal{S}(\mathbb{T}, X), \quad \mathcal{F}_{\mathbb{T}}^{-1}[w](t) := \sum_{k \in \mathbb{Z}} w(k) e^{ikt}.$$

As above, by a duality argument, $\mathcal{F}_{\mathbb{T}}$ extends to a homeomorphism on the space of tempered distributions $\mathcal{F}_{\mathbb{T}} : \mathcal{S}'(\mathbb{T}, X) \rightarrow \mathcal{S}'(\mathbb{Z}, X)$ in the usual way.

A standard verification shows that Lebesgue spaces $L_p(\mathbb{T}, X)$ are embedded into $\mathcal{S}'(\mathbb{T}, X)$, which enables us to define vector-valued Sobolev spaces as

$$\begin{aligned} H_p^m(\mathbb{T}, X) &:= \{u \in L_p(\mathbb{T}, X) \mid \partial_t^\ell u \in L_p(\mathbb{T}, X) \text{ for } \ell = 0, \dots, m\}, \\ \|u\|_{H_p^m(\mathbb{T}, X)} &:= \left(\sum_{\ell=0}^m \|\partial_t^\ell u\|_{L_p(\mathbb{T}, X)}^p \right)^{\frac{1}{p}} \end{aligned}$$

for $m \in \mathbb{N}$ and $p \in [1, \infty)$. A standard mollification argument shows that $\mathcal{S}(\mathbb{T}, X) = C^\infty(\mathbb{T}, X)$ lies dense in $L_p(\mathbb{T}, X)$ and $H_p^m(\mathbb{T}, X)$. For $\theta \in (0, 1)$ we further define the fractional Sobolev space $H_p^\theta(\mathbb{T}, X)$ via Fourier transform by

$$\begin{aligned} H_p^\theta(\mathbb{T}, X) &:= \{u \in L_p(\mathbb{T}, X) \mid \mathcal{F}_{\mathbb{T}}^{-1}[|k|^\theta \mathcal{F}_{\mathbb{T}}[u](k)] \in L_p(\mathbb{T}, X)\}, \\ \|u\|_{H_p^\theta(\mathbb{T}, X)} &:= \left\| \mathcal{F}_{\mathbb{T}}^{-1}[(1 + |k|)^\theta \mathcal{F}_{\mathbb{T}}[u](k)] \right\|_{L_p(\mathbb{T}, X)}. \end{aligned}$$

2.3 \mathcal{R} -boundedness and operator-valued Fourier multipliers

A family of operators $\mathcal{T} \subset \mathcal{L}(X, Y)$ is called \mathcal{R} -bounded in $\mathcal{L}(X, Y)$ if there exists some $C > 0$ such that for all $n \in \mathbb{N}$, $\{T_j\}_{j=1}^n \in \mathcal{T}^n$, and $\{f_j\}_{j=1}^n \in X^n$, we have

$$\left\| \sum_{k=1}^n r_k T_k f_k \right\|_{L_1((0,1), Y)} \leq C \left\| \sum_{k=1}^n r_k f_k \right\|_{L_1((0,1), X)}, \quad (2.1)$$

where $r_k, k \in \mathbb{N}$, denote the Rademacher functions given by $r_k : [0, 1] \rightarrow \{-1, 1\}$, $t \mapsto \text{sign}(\sin 2^k \pi t)$. The smallest constant C such that (2.1) holds is called the \mathcal{R} -bound of \mathcal{T} and denoted by $\mathcal{R}_{\mathcal{L}(X, Y)} \mathcal{T}$. If $\mathcal{S} \subset \mathcal{L}(X, Y)$ and $\mathcal{U} \subset \mathcal{L}(Y, Z)$ are further operator families, we have

$$\begin{aligned} \mathcal{R}_{\mathcal{L}(X, Y)} \{S + T \mid S \in \mathcal{S}, T \in \mathcal{T}\} &\leq \mathcal{R}_{\mathcal{L}(X, Y)} \mathcal{S} + \mathcal{R}_{\mathcal{L}(X, Y)} \mathcal{T}, \\ \mathcal{R}_{\mathcal{L}(X, Z)} \{UT \mid U \in \mathcal{U}, T \in \mathcal{T}\} &\leq \mathcal{R}_{\mathcal{L}(Y, Z)} \mathcal{U} \cdot \mathcal{R}_{\mathcal{L}(X, Y)} \mathcal{T}; \end{aligned} \quad (2.2)$$

see [28, Remark 5.3.14] for example. Due to Kahane's inequality, one may further replace the spaces $L_1((0, 1), X)$ and $L_1((0, 1), Y)$ in (2.1) with $L_p((0, 1), X)$ and $L_p((0, 1), Y)$, respectively, for any $p \in [1, \infty)$. In what follows, this choice of p makes no difference.

We recall the notion of operator-valued Fourier multipliers on \mathbb{R} and \mathbb{T} . For $M \in L_\infty(\mathbb{R}, \mathcal{L}(X, Y))$ we define the operator

$$\text{op}_{\mathbb{R}}[M] : \mathcal{S}(\mathbb{R}, X) \rightarrow \mathcal{S}'(\mathbb{R}, Y), \quad \text{op}_{\mathbb{R}}[M]f := \mathcal{F}_{\mathbb{R}}^{-1}[M \mathcal{F}_{\mathbb{R}}[f]].$$

For $m \in L_\infty(\mathbb{Z}, \mathcal{L}(X, Y))$ we define the operator

$$\text{op}_{\mathbb{T}}[m] : \mathcal{S}(\mathbb{T}, X) \rightarrow \mathcal{S}'(\mathbb{T}, Y), \quad \text{op}_{\mathbb{T}}[m]f := \mathcal{F}_{\mathbb{T}}^{-1}[m \mathcal{F}_{\mathbb{T}}[f]].$$

If there exists a continuous extension of $\text{op}_{\mathbb{T}}[m]$ to a bounded operator

$$\text{op}_{\mathbb{T}}[m] : L_p(\mathbb{T}, X) \rightarrow L_p(\mathbb{T}, Y),$$

we call m an $L_p(\mathbb{T})$ -multiplier. To identify such $L_p(\mathbb{T})$ -multipliers, we shall make use of an operator-valued transference principle, which relates $L_p(\mathbb{T})$ -multipliers to $L_p(\mathbb{R})$ -multipliers, and combine it

with an operator-valued multiplier theorem due to WEIS [67]. For its formulation, we need the notion of \mathcal{R} -boundedness of operator families introduced above as well as the notion of UMD spaces. Recall that a Banach space X is called a UMD space (or a space of class \mathcal{HT}) if the Hilbert transform H defined by

$$Hf(t) := \frac{1}{\pi} \lim_{\varepsilon \rightarrow 0} \int_{|x| \geq \varepsilon} \frac{f(t-s)}{s} ds, \quad f \in \mathcal{S}(\mathbb{R}, X),$$

extends to a bounded linear operator $\mathcal{L}(L_p(\mathbb{R}, X))$ for $p \in (1, \infty)$.

Now we can state the multiplier theorem due to WEIS [67, Theorem 3.4], which is an operator-valued version of the classical Mihlin theorem.

Theorem 2.1 (WEIS). *Let X and Y be UMD-spaces and $p \in (1, \infty)$. Let $M \in L_\infty(\mathbb{R}, \mathcal{L}(X, Y))$ be differentiable in $\mathbb{R} \setminus \{0\}$ and such that*

$$\mathcal{R}_{\mathcal{L}(X,Y)}\{M(t) \mid t \in \mathbb{R} \setminus \{0\}\} \leq r_0, \quad \mathcal{R}_{\mathcal{L}(X,Y)}\{tM'(t) \mid t \in \mathbb{R} \setminus \{0\}\} \leq r_0, \quad (2.3)$$

for some $r_0 > 0$. Then $\text{op}_{\mathbb{R}}[M]$ extends to a bounded operator $\text{op}_{\mathbb{R}}[M] : L_p(\mathbb{R}, X) \rightarrow L_p(\mathbb{R}, Y)$, that is, M is an $L_p(\mathbb{R})$ -multiplier, and

$$\|\text{op}_{\mathbb{R}}[M]\|_{\mathcal{L}(L_p(\mathbb{R}, X), L_p(\mathbb{R}, Y))} \leq C_p r_0 \quad (2.4)$$

for some constant $C_p > 0$ depending only on p but independent of r_0 .

In order to investigate L_p -boundedness of operators associated with Fourier multipliers on \mathbb{T} , we combine this theorem with the following result that allows an investigation of L_p -boundedness of a Fourier multiplier $m \in L_\infty(\mathbb{Z}, \mathcal{L}(X, Y))$ in the torus setting by extending it to a multiplier $M \in L_\infty(\mathbb{R}; \mathcal{L}(X, Y))$ in the Euclidean setting.

Theorem 2.2 (Operator-valued transference principle). *Let X, Y be Banach spaces and $p \in (1, \infty)$. If*

$$M \in L_\infty(\mathbb{R}, \mathcal{L}(X, Y)) \cap C(\mathbb{R}, \mathcal{L}(X, Y))$$

is an $L_p(\mathbb{R})$ -multiplier, that is, $\text{op}_{\mathbb{T}}[M] \in \mathcal{L}(L_p(\mathbb{R}, X), L_p(\mathbb{R}, Y))$, then $M|_{\mathbb{Z}}$ is an $L_p(\mathbb{T})$ -multiplier, that is, $\text{op}_{\mathbb{T}}[M|_{\mathbb{Z}}] \in \mathcal{L}(L_p(\mathbb{T}, X), L_p(\mathbb{T}, Y))$, and

$$\|\text{op}_{\mathbb{T}}[M|_{\mathbb{Z}}]\|_{\mathcal{L}(L_p(\mathbb{T}, X), L_p(\mathbb{T}, Y))} \leq \|\text{op}_{\mathbb{R}}[M]\|_{\mathcal{L}(L_p(\mathbb{R}, X), L_p(\mathbb{T}, Y))}. \quad (2.5)$$

Proof. This is a special version of [28, Prop.5.7.1], which is a generalization of the scalar-valued case originally due to DE LEEUW [19]. \square

Combining the operator-valued transference principle with the Weis multiplier theorem, we directly obtain the following result, which we employ when studying Fourier multipliers on the torus \mathbb{T} .

Corollary 2.3. *Let X and Y be UMD spaces, and let*

$$M \in L_\infty(\mathbb{R}, \mathcal{L}(X, Y)) \cap C(\mathbb{R}, \mathcal{L}(X, Y)) \cap C^1(\mathbb{R} \setminus \{0\}, \mathcal{L}(X, Y))$$

satisfy (2.3) for some $r_0 > 0$. Then $M|_{\mathbb{Z}}$ is an $L_p(\mathbb{T})$ -multiplier, and

$$\|\text{op}_{\mathbb{T}}[M|_{\mathbb{Z}}]\|_{\mathcal{L}(L_p(\mathbb{T}, X), L_p(\mathbb{T}, Y))} \leq C_p r_0 \quad (2.6)$$

for some constant $C_p > 0$ only depending on p .

Proof. By Theorem 2.1, M is an $L_p(\mathbb{R})$ -multiplier, and Theorem 2.2 implies that $M|_{\mathbb{Z}}$ is an $L_p(\mathbb{T})$ -multiplier. Finally, estimate (2.6) follows from (2.5) and (2.4). \square

3 Maximal L_p regularity for periodic evolution equations

We next show how the operator-valued transference principle (Theorem 2.2) can be utilized to establish periodic L_p estimates of the maximal regularity type for a large class of abstract linear evolution equations based on their \mathcal{R} -solvers, that is, the \mathcal{R} -bounded family of solution operators to the associated resolvent problem. We consider the abstract time-periodic boundary-value problem

$$\begin{cases} \partial_t u + Au = F & \text{in } \mathbb{T}, \\ Bu = TG & \text{in } \mathbb{T}. \end{cases} \quad (3.1)$$

Here, A is an abstract (differential) operator, B is a boundary (differential) operator and T plays the role of a trace operator. Using this notion, we avoid the introduction of suitable trace classes, whose identification strongly relies on the underlying function spaces. We show that the existence of a unique solution to (3.1) can be derived from suitable properties of the associated generalized resolvent problem

$$\begin{cases} i\sigma w + Aw = f, \\ Bw = Tg, \end{cases} \quad (3.2)$$

for $\sigma \in \mathbb{R}$. More precisely, while for small k the existence of *a priori* bounds in a suitable functional framework is sufficient, for large k we require the existence of \mathcal{R} -solvers, that is, a functional framework such that the solution operator satisfies suitable \mathcal{R} -bounds. As we shall also see in the examples of the subsequent sections, this lack of \mathcal{R} -bounds for the whole line often appears in applications. Moreover, the presented approach allows to use different function spaces for different modes, which can be a useful and necessary modification; see also Theorem 5.2 below, where the zero-order mode requires separate treatment.

Theorem 3.1. *Let X, Y, Z and W be UMD spaces such that X and W are continuously embedded into Y . Assume that $A \in \mathcal{L}(X, Y)$, $B \in \mathcal{L}(X, Z)$ and $T \in \mathcal{L}(W, Z)$. Let $\gamma_0 \in \mathbb{R}$, $\beta \in [0, 1]$, and let*

$$A \in C^1(\mathbb{R} \setminus (-\gamma_0, \gamma_0), \mathcal{L}(Y \times Y \times W, X))$$

be an operator such that for all $\sigma \in \mathbb{R} \setminus (-\gamma_0, \gamma_0)$ and all $(f, g) \in Y \times W$, the function $w = \mathcal{A}(\sigma)(f, \sigma^\beta g, g)$ is the unique solution to the generalized resolvent problem (3.2). Assume the validity of the \mathcal{R} -bounds

$$\mathcal{R}_{\mathcal{L}(Y \times Y \times W, X)}(\{(\sigma \frac{d}{d\sigma})^\ell \mathcal{A}(\sigma) \mid \sigma \in \mathbb{R} \setminus (-\gamma_0, \gamma_0)\}) \leq r_0, \quad (3.3)$$

$$\mathcal{R}_{\mathcal{L}(Y \times Y \times W, Y)}(\{(\sigma \frac{d}{d\sigma})^\ell (i\sigma \mathcal{A}(\sigma)) \mid \sigma \in \mathbb{R} \setminus (-\gamma_0, \gamma_0)\}) \leq r_0 \quad (3.4)$$

for $\ell = 0, 1$ and some constant $r_0 > 0$.

Moreover, for $k \in \mathbb{Z}$ with $|k| \leq k_0 := \max\{k \in \mathbb{Z} \mid k \leq \gamma_0\}$, let $\mathcal{X}_k, \mathcal{Y}_k, \mathcal{Z}_k$ and \mathcal{W}_k be Banach spaces such that $(ikI + A) \in \mathcal{L}(\mathcal{X}_k, \mathcal{Y}_k)$, $B \in \mathcal{L}(\mathcal{X}_k, \mathcal{Z}_k)$ and $T \in \mathcal{L}(\mathcal{W}_k, \mathcal{Z}_k)$, and such that for all $(f, g) \in \mathcal{Y}_k \times \mathcal{W}_k$ there exists a unique solution $w \in \mathcal{X}_k$ to (3.2) with $\sigma = k$ such that

$$\|w\|_{\mathcal{X}_k} \leq C_k(\|f\|_{\mathcal{Y}_k} + \|g\|_{\mathcal{W}_k}) \quad (3.5)$$

for some constant $C_k > 0$.

Then for any $p \in (1, \infty)$ and (F, G) defined by

$$F(t) = \sum_{k=-k_0}^{k_0} F_k e^{ikt} + F_h(t), \quad G(t) = \sum_{k=-k_0}^{k_0} G_k e^{ikt} + G_h(t), \quad (3.6)$$

with $(F_k, G_k) \in \mathcal{Y}_k \times \mathcal{W}_k$ for $k \in \mathbb{Z}$, $|k| \leq k_0$, and $(F_h, G_h) \in L_p(\mathbb{T}; Y) \times (L_p(\mathbb{T}, W) \cap H_p^\beta(\mathbb{T}, Y))$ such that $(\mathcal{F}_\mathbb{T}[F_h](k), \mathcal{F}_\mathbb{T}[G_h](k)) = 0$ for all $|k| \leq k_0$, there exists a unique element

$$(u_{-k_0}, \dots, u_{k_0}, u_h) \in \mathcal{X}_{-k_0} \times \dots \times \mathcal{X}_{k_0} \times (L_p(\mathbb{T}, X) \cap H_p^1(\mathbb{T}, Y)) \quad (3.7)$$

with $\mathcal{F}_\mathbb{T}[u_h](k) = 0$ for $|k| \leq k_0$, such that

$$u(t) := \sum_{k=-k_0}^{k_0} u_k e^{ikt} + u_h(t) \quad (3.8)$$

is the unique solution to the time-periodic problem (3.1), and

$$\|u_k\|_{\mathcal{X}_k} \leq C_k (\|F_k\|_{\mathcal{Y}_k} + \|G_k\|_{\mathcal{W}_k}) \quad (3.9)$$

$$\|u_h\|_{L_p(\mathbb{T}, X) \cap H_p^1(\mathbb{T}, Y)} \leq C r_0 (\|F_h\|_{L_p(\mathbb{T}, Y)} + \|G_h\|_{H_p^\beta(\mathbb{T}, Y) \cap L_p(\mathbb{T}, W)}), \quad (3.10)$$

for some constant $C > 0$. In particular,

$$\begin{aligned} \|u\| &:= \sum_{k=-k_0}^{k_0} \|u_k\|_{\mathcal{X}_k} + \|u_h\|_{L_p(\mathbb{T}, X) \cap H_p^1(\mathbb{T}, Y)} \\ &\leq C r_0 (\|F_h\|_{L_p(\mathbb{T}, Y)} + \|G_h\|_{H_p^\beta(\mathbb{T}, Y) \cap L_p(\mathbb{T}, W)}) + \sum_{k=-k_0}^{k_0} C_k \|(F_k, G_k)\|_{\mathcal{Y}_k \times \mathcal{W}_k}. \end{aligned} \quad (3.11)$$

Proof. Let $\varphi = \varphi(\sigma)$ be a $C^\infty(\mathbb{R})$ function which equals 1 for $|\sigma| \geq (\gamma_0 + k_0 + 1)/2$, and equals 0 for $|\sigma| \leq \gamma_0$. We define u as in (3.8), where u_k is the unique solution to (3.2) with $\sigma = k$ and $(f, g) = (F_k, G_k)$ for $|k| \leq k_0$, and

$$u_h = \mathcal{F}_\mathbb{T}^{-1}[\varphi(k)\mathcal{A}(k)(\mathcal{F}_\mathbb{T}[F](k), k^\beta \mathcal{F}_\mathbb{T}[G](k), \mathcal{F}_\mathbb{T}[G](k))].$$

One readily verifies that u formally satisfies the time-periodic problem (3.1). Moreover, for $|k| \leq k_0$ we directly conclude $u_k \in \mathcal{X}_k$ and estimate (3.9) since $(F_k, G_k) \in \mathcal{Y}_k \times \mathcal{W}_k$. To show that u_h also belongs to the claimed function space, first observe that $\varphi(\sigma)\mathcal{A}(\sigma) = 0$ for $|\sigma| \leq \gamma_0$, so that $\varphi\mathcal{A} \in C^1(\mathbb{R}, \mathcal{L}(Y \times Y \times W, X))$. From (3.3) and (3.4) it follows that

$$\begin{aligned} \mathcal{R}_{\mathcal{L}(Y \times Y \times W, X)}(\{(\sigma \frac{d}{d\sigma})^\ell(\varphi(\sigma)\mathcal{A}(\sigma)) \mid \sigma \in \mathbb{R}\}) &\leq \|\varphi\|_{H_\infty^1(\mathbb{R})} r_0, \\ \mathcal{R}_{\mathcal{L}(Y \times Y \times W, Y)}(\{(\sigma \frac{d}{d\sigma})^\ell(i\sigma\varphi(\sigma)\mathcal{A}(\sigma)) \mid \sigma \in \mathbb{R}\}) &\leq \|\varphi\|_{H_\infty^1(\mathbb{R})} r_0. \end{aligned}$$

We can thus apply Corollary 2.3 to conclude that $k \mapsto \varphi(k)\mathcal{A}(k) \in \mathcal{L}(Y \times Y \times W, X)$ and $k \mapsto ik\varphi(k)\mathcal{A}(k) \in \mathcal{L}(Y \times Y \times W, Y)$ are $L_p(\mathbb{T})$ -multipliers. We thus deduce $u_h \in L_p(\mathbb{T}, X) \cap H_p^1(\mathbb{T}, Y)$, and (3.10) follows from the above \mathcal{R} -bounds together with (2.6). In total, we have shown existence of a solution in the asserted function class.

To prove the uniqueness statement, let u be of the form (3.8) and satisfy the homogeneous equations, that is, (3.1) with $(f, g) = 0$. For $k \in \mathbb{Z}$ with $|k| > k_0$, we set $u_k = \mathcal{F}_\mathbb{T}[u_h](k)$. Then an application of the Fourier transform to (3.1) gives

$$\begin{cases} ik u_k + A u_k = 0, \\ B u_k = 0. \end{cases}$$

We have $u_k \in X$ for $|k| > k_0$ and $u_k \in \mathcal{X}_k$ for $|k| \leq k_0$, so that $u_k = 0$ follows from the uniqueness assumption for the respective resolvent problem. We thus conclude $u = 0$, which completes the proof. \square

The important case of homogeneous boundary conditions can be incorporated in the above setting in several different manners. Of course, the simplest way is to take boundary data $g = 0$, but one may also consider $T = 0$ as an (abstract) trace operator. Another very common way is to simply set $B = 0$ and $T = 0$, that is, to drop the abstract boundary condition in (3.1), and to incorporate the boundary condition in the function space X . Then (3.1) reduces to the time-periodic X -valued ordinary differential equation

$$\partial_t u + Au = F \quad \text{in } \mathbb{T}, \quad (3.12)$$

and (3.2) becomes a proper resolvent problem

$$i\sigma w + Aw = f. \quad (3.13)$$

In this case, the statement of Theorem 3.1 also simplifies significantly, and it can be formulated using the notion of closed linear operators.

Theorem 3.2. *Let Y be a UMD space, and let $A : D(A) \rightarrow Y$ be a closed operator with domain $D(A) \subset Y$ and resolvent set $\rho(A)$. Let $\gamma_0 \in \mathbb{R}$ such that $\{i\sigma \mid \sigma \in \mathbb{R} \setminus (-\gamma_0, \gamma_0)\} \subset \rho(A)$, and assume the validity of the \mathcal{R} -bounds*

$$\mathcal{R}_{\mathcal{L}(Y)}(\{\lambda(\lambda I - A)^{-1} \mid \lambda = i\sigma, \sigma \in \mathbb{R} \setminus (-\gamma_0, \gamma_0)\}) \leq r_0 \quad (3.14)$$

for some constant $r_0 > 0$.

Moreover, for $k \in \mathbb{Z}$ with $|k| \leq k_0 := \max\{k \in \mathbb{Z} \mid k \leq \gamma_0\}$, let \mathcal{X}_k and \mathcal{Y}_k be Banach spaces such that $(ikI + A) \in \mathcal{L}(\mathcal{X}_k, \mathcal{Y}_k)$ is a linear homeomorphism.

Then for any $p \in (1, \infty)$ and F defined as in (3.6), the function u defined in (3.8) is the unique solution to the time-periodic problem (3.12), where $u_k = (ikI + A)^{-1}F_k$, and

$$\|u_h\|_{L_p(\mathbb{T}, X) \cap H_p^1(\mathbb{T}, Y)} \leq C r_0 \|F_h\|_{L_p(\mathbb{T}, Y)} \quad (3.15)$$

for some constant $C > 0$.

Proof. We set $X = D(A) \subset Y$, equipped with the usual graph norm, which is a UMD space since $(i\gamma_0 I - A) : X \rightarrow Y$ is a homeomorphism. We further set $\mathcal{A}(\sigma) = (i\sigma I + A)^{-1}$ for $|\sigma| \geq \gamma_0$. Then $\mathcal{A}(\sigma)$ is a solution operator for the resolvent problem (3.13) for $|\sigma| \geq \gamma_0$, and we have $\mathcal{A} \in C^\infty(\mathbb{R} \setminus (-\gamma_0, \gamma_0), \mathcal{L}(Y, X))$ by analyticity of the resolvent mapping. Moreover, due to the identities

$$i\sigma \mathcal{A}(\sigma) = i\sigma(i\sigma I + A)^{-1}, \quad \sigma \frac{d}{d\sigma}(i\sigma \mathcal{A}(\sigma)) = i\sigma(i\sigma I + A)^{-1} + \sigma^2(i\sigma I + A)^{-2},$$

and the formulas from (2.2), the assumed \mathcal{R} -bound (3.14) directly implies (3.4). Since

$$\mathcal{A}(\sigma) = \mathcal{A}(\gamma_0) \mathcal{A}(\gamma_0)^{-1} \mathcal{A}(\sigma) = \frac{\gamma_0 - \sigma}{\sigma} \mathcal{A}(\gamma_0) i\sigma(i\sigma I + A)^{-1} + I, \quad \sigma \frac{d}{d\sigma} \mathcal{A}(\sigma) = i\sigma(i\sigma I + A)^{-2},$$

and since $\mathcal{A}(\gamma_0)$ is a homeomorphism in $\mathcal{L}(Y, X)$, we infer (3.3) from (3.14) in the same way. Now the statement directly follows from 3.1 with $W = Z = \mathcal{W}_k = \mathcal{Z}_k = \{0\}$ and $B = T = 0$. \square

4 The Navier–Stokes equations in a bounded periodically moving domain

Let Ω be a bounded domain in \mathbb{R}^N , $N \geq 2$, whose boundary, Γ , is a compact C^2 hypersurface. We assume that for each $t \in \mathbb{R}$ there exists an injective map $\phi(\cdot, t) : \Omega \rightarrow \mathbb{R}^N$ such that $\phi(y, 0) = 0$ and $\phi(y, t + 2\pi) = \phi(y, t)$ for $t \in \mathbb{R}$ and $y \in \Omega$; possessing the regularity

$$\phi \in C^0(\mathbb{T}; C^3(\Omega)^N) \cap C^1(\mathbb{T}; C^1(\Omega)^N). \quad (4.1)$$

Again, we identify 2π -periodic on \mathbb{R} with functions on \mathbb{T} . Let Ω_t be a domain in \mathbb{R}^N given by setting

$$\Omega_t = \{x = y + \phi(y, t) \mid y \in \Omega\} \quad (t \in \mathbb{R}), \quad (4.2)$$

that is, Ω_t is the image of the transformation $\Phi_t : \Omega \rightarrow \mathbb{R}^N$, $\Phi_t(y) = y + \phi(y, t)$, for $t \in \mathbb{R}$. Notice that Ω_t is a given bounded periodically moving domain in \mathbb{R}^N such that $\Omega_{t+2\pi} = \Omega_t$. Let Γ_t be the boundary of Ω_t , which is given by $\Gamma_t = \{x = y + \phi(y, t) \mid y \in \Gamma\}$. We consider the Navier–Stokes equations in Ω_t :

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} - \Delta \mathbf{u} + \nabla \mathbf{p} = \mathbf{F}, \quad \operatorname{div} \mathbf{u} = 0 \quad \text{in } \Omega_t, \quad \mathbf{u}|_{\Gamma_t} = \mathbf{h}|_{\Gamma_t} \quad (4.3)$$

for $t \in (0, 2\pi)$. Here, $\mathbf{u} = (u_1, \dots, u_N)^\top$ is an unknown velocity field, M^\top being the transposed M , \mathbf{p} an unknown pressure field, $\mathbf{F} = (F_1, \dots, F_N)^\top$ a prescribed external force, and $\mathbf{h} = (h_1, \dots, h_N)^\top$ a velocity field that prescribes the boundary velocity. Assume that $\mathbf{F}(t + 2\pi) = \mathbf{F}(t)$ and $\mathbf{h}(t + 2\pi) = \mathbf{h}(t)$ for any $t \in \mathbb{R}$. Then system (4.3) describes the flow of an incompressible viscous fluid around a periodically moving body, subject to a time-periodic external force and with prescribed time-periodic boundary conditions. Note that a natural choice for \mathbf{h} would be the flow velocity associated with the transformation Φ_t , which means that the fluid particles adhere to the boundary. This choice corresponds to a no-slip condition, which we further discuss in Remark 4.5 below.

We transform (4.3) into a system in Ω by using the change of variables induced by Φ_t , namely $x = y + \phi(y, t)$. For this purpose, we assume that

$$\|\phi\|_{L_\infty(\mathbb{T}, H_\infty^3(\Omega))} + \|\partial_t \phi\|_{L_\infty(\mathbb{T}, H_\infty^1(\Omega))} < \varepsilon_0 \quad (4.4)$$

with some small number $\varepsilon_0 > 0$. By this smallness assumption, we may assume the existence of the inverse transformation: $y = x + \psi(x, t)$. The associated Jacobi matrix $\partial(t, y)/\partial(t, x)$ is given by the formulas:

$$\frac{\partial t}{\partial t} = 1, \quad \frac{\partial t}{\partial x_j} = 0, \quad \frac{\partial y_\ell}{\partial t} = \frac{\partial \psi_\ell}{\partial t}, \quad \frac{\partial y_\ell}{\partial x_j} = \delta_{\ell j} + \frac{\partial \psi_\ell}{\partial x_j}$$

for $j, \ell = 1, \dots, N$. Set $a_{\ell 0}(y, t) = (\partial \psi_\ell / \partial t)(y + \phi(y, t), t)$, and $a_{\ell j}(y, t) = (\partial \psi_\ell / \partial x_j)(y + \phi(y, t), t)$. Then partial derivatives transform as

$$\frac{\partial f}{\partial t} = \frac{\partial g}{\partial t} + \sum_{\ell=1}^N a_{\ell 0}(y, t) \frac{\partial g}{\partial y_\ell}, \quad \frac{\partial f}{\partial x_j} = \frac{\partial g}{\partial y_j} + \sum_{\ell=1}^N a_{\ell j}(y, t) \frac{\partial g}{\partial y_\ell} \quad (4.5)$$

for $f(x, t) = g(y, t)$. Set $J = \det(\partial x / \partial y) = 1 + J_0(y, t)$, which is the Jacobian of Φ_t . By the L_∞ -bounds in (4.4) we have

$$\begin{aligned} & \sup_{t \in \mathbb{R}} \|a_{\ell j}(\cdot, t)\|_{H_\infty^2(\Omega)} + \sup_{t \in \mathbb{R}} \|\partial_t a_{\ell j}(\cdot, t)\|_{L_\infty(\Omega)} + \sup_{t \in \mathbb{R}} \|a_{0j}(\cdot, t)\|_{L_\infty(\Omega)} \\ & + \sup_{t \in \mathbb{R}} \|J_0(\cdot, t)\|_{H_\infty^2(\Omega)} + \sup_{t \in \mathbb{R}} \|\partial_t J_0(\cdot, t)\|_{L_\infty(\Omega)} \leq C\varepsilon_0 \end{aligned} \quad (4.6)$$

with some constant $C > 0$ for $j, \ell = 1, \dots, N$. For simplicity, we set $\mathbf{v}(y, t) = (v_1, \dots, v_N)^\top = \mathbf{u}(x, t)$, and $\mathbf{q}(y, t) = \mathbf{p}(x, t)$. By (4.5) we have

$$\begin{aligned} \partial_t \mathbf{u} &= \partial_t \mathbf{v} + \sum_{\ell=1}^N a_{\ell 0} \frac{\partial \mathbf{v}}{\partial y_\ell}, \quad \mathbf{u} \cdot \nabla \mathbf{u} = \mathbf{v} \cdot (\mathbf{I} + \mathbf{A}) \nabla \mathbf{v}, \\ \Delta \mathbf{u} &= \Delta \mathbf{v} + \sum_{\ell=1}^N (a_{\ell j} + a_{j\ell}) \frac{\partial^2 \mathbf{v}}{\partial y_\ell \partial y_j} + \sum_{j, \ell, m=1}^N a_{\ell j} a_{mj} \frac{\partial^2 \mathbf{v}}{\partial y_\ell \partial y_m} \\ &\quad + \sum_{\ell, m=1}^N \left(\frac{\partial a_{m\ell}}{\partial y_\ell} + \sum_{j=1}^N a_{\ell j} \frac{\partial a_{mj}}{\partial y_\ell} \right) \frac{\partial \mathbf{v}}{\partial y_m}, \\ \operatorname{div} \mathbf{u} &= \mathbf{J}^{-1} \{ \operatorname{div} \mathbf{v} + \operatorname{div} (\mathbf{J}_0 \mathbf{v}) + \sum_{j, \ell=1}^N \frac{\partial}{\partial y_\ell} (a_{\ell j} \mathbf{J} v_j) \}. \quad \nabla \mathbf{p} = (\mathbf{I} + \mathbf{A}) \nabla \mathbf{q}, \end{aligned} \quad (4.7)$$

where \mathbf{A} is an $(N \times N)$ -matrix whose (j, k) -th component is a_{jk} . Setting $w_\ell = v_\ell + \mathbf{J}_0 v_\ell + \sum_{j=1}^N a_{\ell j} \mathbf{J} v_j$, we have $\mathbf{J} \operatorname{div} \mathbf{u} = \operatorname{div} \mathbf{w}$ with $\mathbf{w} = (w_1, \dots, w_N)^\top$. Notice that $\mathbf{w} = (\mathbf{I} + \mathbf{J}_0 \mathbf{I} + \mathbf{A}^\top \mathbf{J}) \mathbf{v}$. In view of (4.6), choosing $\varepsilon > 0$, we see that there exists an $(N \times N)$ -matrix \mathbf{B}_{-1} such that $(\mathbf{I} + \mathbf{J}_0 \mathbf{I} + \mathbf{A}^\top \mathbf{J})^{-1} = \mathbf{I} + \mathbf{B}_{-1}$ and

$$\sup_{t \in \mathbb{R}} \|\mathbf{B}_{-1}(\cdot, t)\|_{\mathbf{H}_\infty^2(\Omega)} \leq C\varepsilon_0, \quad \sup_{t \in \mathbb{R}} \|\partial_t \mathbf{B}_{-1}(\cdot, t)\|_{\mathbf{L}_\infty(\Omega)} \leq C\varepsilon_0. \quad (4.8)$$

Summing up, we see that (4.3) is transformed to the following equations:

$$\partial_t \mathbf{w} - \Delta \mathbf{w} + \nabla \mathbf{q} = \mathbf{G} + \mathcal{L}(\mathbf{w}, \mathbf{q}) + \mathcal{N}(\mathbf{w}), \quad \operatorname{div} \mathbf{w} = 0 \quad \text{in } \Omega \times \mathbb{T}, \quad \mathbf{w}|_\Gamma = \mathbf{H}|_\Gamma, \quad (4.9)$$

where \mathbf{G} and \mathbf{H} are prescribed data and

$$\begin{aligned} \mathcal{L}(\mathbf{w}, \mathbf{q}) &= -\partial_t (\mathbf{B}_{-1} \mathbf{w}) - \sum_{\ell=1}^N a_{\ell 0} \frac{\partial}{\partial y_\ell} ((\mathbf{I} + \mathbf{B}_{-1}) \mathbf{w}) + \Delta (\mathbf{B}_{-1} \mathbf{w}) \\ &\quad + \sum_{\ell=1}^N (a_{\ell j} + a_{j\ell}) \frac{\partial^2}{\partial y_\ell \partial y_j} ((\mathbf{I} + \mathbf{B}_{-1}) \mathbf{w}) + \sum_{j, \ell, m=1}^N a_{\ell j} a_{mj} \frac{\partial^2}{\partial y_\ell \partial y_m} ((\mathbf{I} + \mathbf{B}_{-1}) \mathbf{w}) \\ &\quad + \sum_{\ell, m=1}^N \left(\frac{\partial a_{m\ell}}{\partial y_\ell} + \sum_{j=1}^N a_{\ell j} \frac{\partial a_{mj}}{\partial y_\ell} \right) \frac{\partial}{\partial y_m} ((\mathbf{I} + \mathbf{B}_{-1}) \mathbf{w}) - \mathbf{A} \nabla \mathbf{q} \\ \mathcal{N}(\mathbf{v}) &= ((\mathbf{I} + \mathbf{B}_{-1}) \mathbf{w}) \cdot (\mathbf{I} + \mathbf{A}) \nabla ((\mathbf{I} + \mathbf{B}_{-1}) \mathbf{w}). \end{aligned} \quad (4.10)$$

Observe that, due to $\operatorname{div} \mathbf{w} = 0$ and the boundedness of Ω , for the existence of solutions to (4.9) it is necessary that the boundary data satisfy

$$\int_\Gamma \mathbf{H} \cdot \mathbf{n} \, d\sigma = 0 \quad (4.11)$$

for all $t \in \mathbb{R}$, where \mathbf{n} denotes the unit outer normal vector at Γ . The following theorem is our main result in this section.

Theorem 4.1. *Let $2 < p < \infty$ and $N < q < \infty$. Then, there exists numbers $\varepsilon, \varepsilon_0 > 0$ such that if the condition (4.4) is valid, and if the prescribed terms $\mathbf{G} \in \mathbf{L}_p(\mathbb{T}, \mathbf{L}_q(\Omega)^N)$ and $\mathbf{H} \in$*

$H_p^1(\mathbb{T}, L_q(\Omega)^N) \cap L_p(\mathbb{T}, H_q^2(\Omega)^N)$ satisfy the compatibility condition (4.11) as well as the smallness condition

$$\|\mathbf{G}\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{H}\|_{H_p^1(\mathbb{T}, L_q(\Omega))} + \|\mathbf{H}\|_{L_p(\mathbb{T}, H_q^2(\Omega))} \leq \varepsilon^2,$$

then problem (4.9) admits a unique² solution (\mathbf{w}, \mathbf{q}) with

$$\mathbf{w} \in H_p^1(\mathbb{T}, L_q(\Omega)^N) \cap L_p(\mathbb{T}, H_q^2(\Omega)^N), \quad \mathbf{q} \in L_p(\mathbb{T}, \hat{H}_q^1(\Omega)),$$

possessing the estimate

$$\|\partial_t \mathbf{w}\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{w}\|_{L_p(\mathbb{T}, H_q^2(\Omega))} + \|\nabla \mathbf{q}\|_{L_p(\mathbb{T}, L_q(\Omega))} \leq \varepsilon.$$

To prove Theorem 4.1, we consider the following linearization of equations (4.9):

$$\partial_t \mathbf{u} - \Delta \mathbf{u} + \nabla \mathbf{p} = \mathbf{F}, \quad \operatorname{div} \mathbf{u} = 0 \quad \text{in } \Omega \times \mathbb{T}, \quad \mathbf{u}|_\Gamma = \mathbf{H}|_\Gamma. \quad (4.12)$$

For a maximal-regularity theorem for problem (4.12), we consider the associated resolvent problem

$$\lambda \mathbf{w} - \Delta \mathbf{w} + \nabla \mathbf{p} = \mathbf{f}, \quad \operatorname{div} \mathbf{w} = 0 \quad \text{in } \Omega, \quad \mathbf{w}|_\Gamma = \mathbf{h}|_\Gamma. \quad (4.13)$$

At first, we consider the case of homogeneous boundary conditions $\mathbf{h} = 0$, for which we have the following theorem, which holds for bounded and exterior domains simultaneously.

Theorem 4.2. *Let $1 < q < \infty$ and $0 < \varepsilon < \pi/2$. Let Ω be a bounded domain or an exterior domain in \mathbb{R}^N ($N \geq 2$) with C^2 boundary. There exist operator families $(\mathcal{S}(\lambda)) \subset \mathcal{L}(L_q(\Omega)^N, H_q^2(\Omega)^N)$ and $(\mathcal{P}(\lambda)) \subset \mathcal{L}(L_q(\Omega)^N, \hat{H}_q^1(\Omega))$ such that for every $\lambda \in \Sigma_\varepsilon \setminus \{0\}$ and every $\mathbf{f} \in L_q(\Omega)^N$ the pair $(\mathbf{w}, \mathbf{p}) = (\mathcal{S}(\lambda)\mathbf{f}, \mathcal{P}(\lambda)\mathbf{f})$ is the unique solution to (4.13) with $\mathbf{h} = 0$ and satisfies the estimate*

$$|\lambda| \|\mathcal{S}(\lambda)\mathbf{f}\|_{L_q(\Omega)} + \|\nabla^2 \mathcal{S}(\lambda)\mathbf{f}\|_{L_q(\Omega)} + \|\nabla \mathcal{P}(\lambda)\mathbf{f}\|_{L_q(\Omega)} \leq C \|\mathbf{f}\|_{L_q(\Omega)} \quad (4.14)$$

with some constant $C > 0$ depending on Ω , q and ε . Moreover, there exist constants $\lambda_0, r_0 > 0$, depending on Ω , q and ε , such that

$$\mathcal{S} \in \operatorname{Hol}(\Sigma_{\varepsilon, \lambda_0}, \mathcal{L}(L_q(\Omega)^N, H_q^2(\Omega)^N)), \quad \mathcal{P} \in \operatorname{Hol}(\Sigma_{\varepsilon, \lambda_0}, \mathcal{L}(L_q(\Omega)^N, \hat{H}_q^1(\Omega))),$$

and

$$\begin{aligned} \mathcal{R}_{\mathcal{L}(L_q(\Omega)^N, H_q^{2-j}(\Omega)^N)}(\{(\lambda \partial_\lambda)^\ell (\lambda^{j/2} \mathcal{S}(\lambda)) \mid \lambda \in \Sigma_{\varepsilon, \lambda_0}\}) &\leq r_0, \\ \mathcal{R}_{\mathcal{L}(L_q(\Omega)^N, L_q(\Omega)^N)}(\{(\lambda \partial_\lambda)^\ell (\nabla \mathcal{P}(\lambda)) \mid \lambda \in \Sigma_{\varepsilon, \lambda_0}\}) &\leq r_0 \end{aligned}$$

for $\ell = 0, 1, j = 0, 1, 2$. Additionally, if Ω is bounded, there exist $\mathcal{S}(0) \in \mathcal{L}(L_q(\Omega)^N, H_q^2(\Omega)^N)$ and $\mathcal{P}(0) \in \mathcal{L}(L_q(\Omega)^N, \hat{H}_q^1(\Omega))$ such that $(\mathbf{w}, \mathbf{p}) = (\mathcal{S}(0)\mathbf{f}, \mathcal{P}(0)\mathbf{f})$ is the unique solution to (4.13) for $\lambda = 0$ and satisfies

$$\|\mathcal{S}(0)\mathbf{f}\|_{H_q^2(\Omega)} + \|\nabla \mathcal{P}(0)\mathbf{f}\|_{L_q(\Omega)} \leq C \|\mathbf{f}\|_{L_q(\Omega)}. \quad (4.15)$$

Proof. Existence of unique solutions to (4.13) satisfying (4.14) with a uniform constant C for $\lambda \in \Sigma \setminus \{0\}$ was shown in [8]. The analyticity of the associated family of solution operators in $\Sigma_{\varepsilon, \lambda_0}$ for some $\lambda_0 > 0$ together with the asserted \mathcal{R} -bounds was established in [57, Theorem 1.6] and [58, Theorem 9.1.4]. Finally, the statement for $\lambda = 0$ follows from Fredholm's alternative principle since the embedding $H_q^2(\Omega) \hookrightarrow L_q(\Omega)$ is compact if Ω is bounded, and the solution to (4.13) is unique. \square

²Note that here and in what follows, the uniqueness statement refers to uniqueness of the velocity field, but the pressure term is usually merely unique up to a constant.

As in the results from Section 3, in the case of non-zero boundary values, the \mathcal{R} -solvers for the resolvent problem (4.13) are more involved than in the situation of Theorem 4.2. To state the result, we introduce the space

$$\mathcal{X}_q = \{(F_1, F_2, F_3, F_4) \mid F_1, F_2 \in L_q(\Omega)^N, F_3 \in H_q^1(\Omega)^N, F_4 \in H_q^2(\Omega)^N\}.$$

Then we have the following result.

Theorem 4.3. *Let $1 < q < \infty$ and $0 < \varepsilon < \pi/2$. Let Ω be a bounded domain or an exterior domain in \mathbb{R}^N ($N \geq 2$) with C^2 boundary. There exist constants $\lambda_0, r_0 > 0$ and operator families*

$$\mathcal{S} \in \text{Hol}(\Sigma_{\varepsilon, \lambda_0}, \mathcal{L}(\mathcal{X}_q(\Omega), H_q^2(\Omega)^N)), \quad \mathcal{P} \in \text{Hol}(\Sigma_{\varepsilon, \lambda_0}, \mathcal{L}(\mathcal{X}_q(\Omega), \hat{H}_q^1(\Omega)))$$

such that for any $\mathbf{f} \in L_q(\Omega)^N$ and $\mathbf{h} \in H_q^2(\Omega)^N$, satisfying $\int_{\Gamma} \mathbf{h} \cdot \mathbf{n} \, d\sigma = 0$ if Ω is bounded, the pair (\mathbf{w}, \mathbf{p}) defined by $\mathbf{w} = \mathcal{S}(\lambda)(\mathbf{f}, \lambda \mathbf{h}, \lambda^{1/2} \mathbf{h}, \mathbf{h})$ and $\mathbf{p} = \mathcal{P}(\lambda)(\mathbf{f}, \lambda \mathbf{h}, \lambda^{1/2} \mathbf{h}, \mathbf{h})$ is the unique solution to (4.13), and

$$\begin{aligned} \mathcal{R}_{\mathcal{L}(\mathcal{X}_q(\Omega), H_q^{2-j}(\Omega)^N)}(\{(\lambda \frac{d}{d\lambda})^\ell (\lambda^{j/2} \mathcal{S}(\lambda)) \mid \lambda \in \Sigma_{\varepsilon, \lambda_0}\}) &\leq r_0, \\ \mathcal{R}_{\mathcal{L}(\mathcal{X}_q(\Omega), L_q(\Omega)^N)}(\{(\lambda \frac{d}{d\lambda})^\ell (\nabla \mathcal{P}(\lambda)) \mid \lambda \in \Sigma_{\varepsilon, \lambda_0}\}) &\leq r_0 \end{aligned}$$

for $\ell = 0, 1, j = 0, 1, 2$.

Proof. See [57, Theorem 1.6]. □

Now we can prove the following theorem on the time-periodic linear problem (4.12).

Theorem 4.4. *Let $1 < p, q < \infty$, and let $\Omega \subset \mathbb{R}^N$ be a bounded domain with C^2 boundary. Then, for any $\mathbf{F} \in L_p(\mathbb{T}, L_q(\Omega)^N)$ and $\mathbf{H} \in H_p^1(\mathbb{T}, L_q(\Omega)^N) \cap L_p(\mathbb{T}, H_q^2(\Omega)^N)$ satisfying (4.11), problem (4.12) admits a unique solution (\mathbf{u}, \mathbf{p}) with*

$$\mathbf{u} \in H_p^1(\mathbb{T}, L_q(\Omega)^N) \cap L_p(\mathbb{T}, H_q^2(\Omega)^N), \quad \mathbf{p} \in L_p(\mathbb{T}, \hat{H}_q^1(\Omega))$$

possessing the estimate

$$\begin{aligned} &\|\partial_t \mathbf{u}\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{u}\|_{L_p(\mathbb{T}, H_q^2(\Omega))} + \|\nabla \mathbf{p}\|_{L_p(\mathbb{T}, L_q(\Omega))} \\ &\leq C(\|\mathbf{F}\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\partial_t \mathbf{H}\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{H}\|_{L_p(\mathbb{T}, H_q^2(\Omega))}). \end{aligned} \quad (4.16)$$

Proof. We first consider the case $\mathbf{H} = 0$, for which proceed analogously to the proof of Theorem 3.1. Let $\varphi = \varphi(\sigma)$ be a $C^\infty(\mathbb{R})$ function that equals 1 for $|\sigma| \geq \lambda_0 + 1/2$ and 0 for $|\sigma| \leq \lambda_0 + 1/4$. Set

$$\mathbf{u}_h = \mathcal{F}_{\mathbb{T}}^{-1}[\mathcal{L}(ik)\varphi(k)\mathcal{F}_{\mathbb{T}}[\mathbf{F}](k)], \quad \mathbf{p}_h = \mathcal{F}_{\mathbb{T}}^{-1}[\mathcal{P}(ik)\varphi(k)\mathcal{F}_{\mathbb{T}}[\mathbf{F}](k)].$$

Then \mathbf{u}_h and \mathbf{p}_h satisfy the equations

$$\partial_t \mathbf{u}_h - \mu \Delta \mathbf{u}_h + \nabla \mathbf{p}_h = \mathbf{F}_h, \quad \text{div } \mathbf{u}_h = 0 \quad \text{in } \Omega \times \mathbb{T}, \quad \mathbf{u}_h|_{\Gamma} = 0,$$

where we have set $\mathbf{F}_h = \mathcal{F}_{\mathbb{T}}^{-1}[\varphi(k)\mathcal{F}_{\mathbb{T}}[\mathbf{F}](k)]$. Moreover, arguing as for the proof of Theorem 3.1, we can use the \mathcal{R} -bounds from Theorem 4.2 and invoke Corollary 2.3 to deduce

$$\|\partial_t \mathbf{u}_h\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{u}_h\|_{L_p(\mathbb{T}, H_q^2(\Omega))} + \|\nabla \mathbf{p}_h\|_{L_p(\mathbb{T}, L_q(\Omega))} \leq C\|\mathbf{F}_h\|_{L_p(\mathbb{T}, L_q(\Omega))} \leq C\|\mathbf{F}\|_{L_p(\mathbb{T}, L_q(\Omega))}. \quad (4.17)$$

Now, in view of Theorem 4.2, we set

$$\mathbf{u}(t) = \mathbf{u}_h(t) + \sum_{|k| \leq \lambda_0} e^{ikt} \mathcal{S}(ik) \mathcal{F}_{\mathbb{T}}[\mathbf{f}](k), \quad \mathbf{p}(t) = \mathbf{p}_h(t) + \sum_{|k| \leq \lambda_0} e^{ikt} \mathcal{P}(ik) \mathcal{F}_{\mathbb{T}}[\mathbf{f}](k). \quad (4.18)$$

Then, \mathbf{u} and \mathbf{p} satisfy (4.12) with $\mathbf{H} = 0$, and from (4.14), (4.15) and (4.17) we conclude the estimate

$$\|\partial_t \mathbf{u}\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{u}\|_{L_p(\mathbb{T}, H_q^2(\Omega))} + \|\nabla \mathbf{p}\|_{L_p(\mathbb{T}, L_q(\Omega))} \leq C \|\mathbf{F}\|_{L_p(\mathbb{T}, L_q(\Omega))}. \quad (4.19)$$

Thus, we have shown existence for $\mathbf{H} = 0$.

Now consider arbitrary $\mathbf{H} \in H_p^1(\mathbb{T}, L_q(\Omega)^N) \cap L_p(\mathbb{T}, H_q^2(\Omega)^N)$ satisfying (4.11). Fix $\lambda_1 > \lambda_0$ with λ_0 from Theorem 4.3. We use the \mathcal{R} -bounded solution operators \mathcal{S} and \mathcal{P} from Theorem 4.3 to define

$$\begin{aligned} \mathbf{u}_1 &= \mathcal{F}_{\mathbb{T}}^{-1}[\mathcal{S}(ik + \lambda_1)(0, (ik + \lambda_1)\tilde{\mathbf{H}}_k, (ik + \lambda_1)^{1/2}\tilde{\mathbf{H}}_k, \tilde{\mathbf{H}}_k)], \\ \mathbf{p}_1 &= \mathcal{F}_{\mathbb{T}}^{-1}[\mathcal{P}(ik + \lambda_1)(0, (ik + \lambda_1)\tilde{\mathbf{H}}_k, (ik + \lambda_1)^{1/2}\tilde{\mathbf{H}}_k, \tilde{\mathbf{H}}_k)], \end{aligned}$$

where $\tilde{\mathbf{H}}_k = \mathcal{F}[\mathbf{H}](k)$. Then $(\mathbf{u}_1, \mathbf{p}_1)$ is a solution to the auxiliary problem

$$\partial_t \mathbf{u}_1 + \lambda_1 \mathbf{u}_1 - \mu \Delta \mathbf{u}_1 + \nabla \mathbf{p}_1 = 0, \quad \operatorname{div} \mathbf{u}_1 = 0 \quad \text{in } \Omega \times \mathbb{T}, \quad \mathbf{u}_1|_{\Gamma} = \mathbf{H},$$

and by the multiplier theorem from Corollary 2.3, we have

$$\mathbf{u}_1 \in H_p^1(\mathbb{T}, L_q(\Omega)^N) \cap L_p(\mathbb{T}, H_q^2(\Omega)^N), \quad \mathbf{p}_1 \in L_p(\mathbb{T}, \hat{H}_q^1(\Omega))$$

and the estimate

$$\|\partial_t \mathbf{u}_1\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{u}_1\|_{L_p(\mathbb{T}, H_q^2(\Omega))} + \|\nabla \mathbf{p}_1\|_{L_p(\mathbb{T}, L_q(\Omega))} \leq C(\|\partial_t \mathbf{H}\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{H}\|_{L_p(\mathbb{T}, H_q^2(\Omega))}).$$

Here, we have used the interpolation inequality

$$\|\mathbf{H}\|_{H_p^{1/2}(\mathbb{T}, H_q^1(\Omega))} \leq C(\|\partial_t \mathbf{H}\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{H}\|_{L_p(\mathbb{T}, H_q^2(\Omega))}).$$

as well as the trivial estimate

$$\begin{aligned} \|\mathcal{F}_{\mathbb{T}}^{-1}[(ik + \lambda_1)\tilde{\mathbf{H}}_k]\|_{L_p(\mathbb{T}, L_q(\Omega))} &\leq \|\mathcal{F}_{\mathbb{T}}^{-1}[ik\tilde{\mathbf{H}}_k]\|_{L_p(\mathbb{T}, L_q(\Omega))} + \lambda_1 \|\mathcal{F}_{\mathbb{T}}^{-1}[\tilde{\mathbf{H}}_k]\|_{L_p(\mathbb{T}, L_q(\Omega))} \\ &= \|\partial_t \mathbf{H}\|_{L_p(\mathbb{T}, L_q(\Omega))} + \lambda_1 \|\mathbf{H}\|_{L_p(\mathbb{T}, L_q(\Omega))}. \end{aligned}$$

Now consider the problem

$$\partial_t \mathbf{u}_2 - \mu \Delta \mathbf{u}_2 + \nabla \mathbf{p}_2 = \mathbf{F} + \lambda_1 \mathbf{u}_1, \quad \operatorname{div} \mathbf{u}_2 = 0 \quad \text{in } \Omega \times \mathbb{T}, \quad \mathbf{u}_2|_{\Gamma} = 0.$$

As shown in the first part of the proof, there exists a solution $(\mathbf{u}_2, \mathbf{q}_2)$ in the claimed function class and satisfying (4.19) with \mathbf{F} replaced with $\mathbf{F} + \lambda_1 \mathbf{u}_1$. Then $(\mathbf{u}, \mathbf{q}) = (\mathbf{u}_1 + \mathbf{u}_2, \mathbf{q}_1 + \mathbf{q}_2)$ is a solution to (4.12), belongs to the correct function class, and satisfies estimate (4.16).

For the uniqueness statement, let (\mathbf{u}, \mathbf{p}) be in the considered function class and satisfy (4.12) with $\mathbf{F} = 0$ and $\mathbf{H} = 0$. Then, for each $k \in \mathbb{Z}$, setting $\tilde{\mathbf{u}}_k = \mathcal{F}[\mathbf{u}](k)$ and $\tilde{\mathbf{p}}_k = \mathcal{F}_{\mathbb{T}}[\mathbf{p}](k)$, we see that $\tilde{\mathbf{u}}_k \in H_q^2(\Omega)^3$ and $\tilde{\mathbf{p}}_k \in \hat{H}_q^1(\Omega)$ satisfy the homogeneous equations

$$ik\tilde{\mathbf{u}}_k - \mu \Delta \tilde{\mathbf{u}}_k + \nabla \tilde{\mathbf{p}}_k = 0, \quad \operatorname{div} \tilde{\mathbf{u}}_k = 0 \quad \text{in } \Omega, \quad \tilde{\mathbf{u}}_k|_{\Gamma} = 0.$$

Thus, the uniqueness statement from Theorem 4.2 yields that $\tilde{\mathbf{u}}_k = \nabla \tilde{\mathbf{p}}_k = 0$, which shows that $\mathbf{u} = \nabla \mathbf{p} = 0$. This completes the proof of Theorem 5.5. \square

Proof of Theorem 4.1. We conclude the proof using the contraction mapping principle in the underlying space \mathcal{I}_ε defined by

$$\begin{aligned} \mathcal{I}_\varepsilon = \{(\mathbf{u}, \mathbf{p}) \mid & \mathbf{u} \in H_p^1(\mathbb{T}, L_q(\Omega)) \cap L_p(\mathbb{T}, H_q^2(\Omega)^N), \mathbf{p} \in L_p(\mathbb{T}, \hat{H}_q^1(\Omega)), \\ & E(\mathbf{u}, \mathbf{p}) := \|\partial_t \mathbf{u}\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{u}\|_{L_p(\mathbb{T}, H_q^2(\Omega))} + \|\nabla \mathbf{p}\|_{L_p(\mathbb{T}, L_q(\Omega))} \leq \varepsilon\}. \end{aligned}$$

Given $(\mathbf{u}, \mathbf{p}) \in \mathcal{I}_\varepsilon$, let \mathbf{v} and \mathbf{q} satisfy

$$\partial_t \mathbf{v} - \Delta \mathbf{v} + \nabla \mathbf{q} = \mathbf{G} + \mathcal{L}(\mathbf{u}, \mathbf{p}) + \mathcal{N}(\mathbf{u}), \quad \operatorname{div} \mathbf{v} = 0 \quad \text{in } \Omega \times \mathbb{T}, \quad \mathbf{v}|_\Gamma = \mathbf{H}|_\Gamma. \quad (4.20)$$

The existence of (\mathbf{v}, \mathbf{q}) follows from Theorem 4.4 if we can show that the forcing term in (4.20) belongs to $L_p(\mathbb{T}, L_q(\Omega)^N)$. Firstly, by (4.4), (4.6) and (4.8) we have

$$\|\mathcal{L}(\mathbf{u}, \mathbf{p})\|_{L_p(\mathbb{T}, L_q(\Omega))} \leq C\varepsilon E(\mathbf{u}, \mathbf{p}). \quad (4.21)$$

In a similar way, noting that $N < q < \infty$, by Sobolev's imbedding theorem we have

$$\|\mathcal{N}(\mathbf{u}(\cdot, t))\|_{L_q(\Omega)} \leq C\|\mathbf{u}(\cdot, t)\|_{L_q(\Omega)}\|\mathbf{u}(\cdot, t)\|_{H_q^2(\Omega)}. \quad (4.22)$$

By real interpolation theorem, we know that

$$\begin{aligned} H_p^1(\mathbb{T}, L_q(\Omega)) \cap L_p(\mathbb{T}, H_q^2(\Omega)) & \hookrightarrow C^0(\mathbb{T}, B_{q,p}^{2(1-1/p)}(\Omega)), \\ \sup_{t \in \mathbb{T}} \|f(\cdot, t)\|_{B_{q,p}^{2(1-1/p)}(\Omega)} & \leq C(\|f\|_{H_p^1(\mathbb{T}, L_q(\Omega))} + \|f\|_{L_p(\mathbb{T}, H_q^2(\Omega))}) \end{aligned} \quad (4.23)$$

since we have $p > 2$, and we obtain

$$\|f\|_{L_\infty(\mathbb{T}, L_q(\Omega))} \leq C\|f\|_{L_\infty(\mathbb{T}, B_{q,p}^{2(1-1/p)}(\Omega))} \leq C(\|f\|_{H_p^1(\mathbb{T}, L_q(\Omega))} + \|f\|_{L_p(\mathbb{T}, H_q^2(\Omega))}). \quad (4.24)$$

Therefore,

$$\|\mathcal{N}(\mathbf{u})\|_{L_p(\mathbb{T}, L_q(\Omega))} \leq CE(\mathbf{u}, \mathbf{p})^2. \quad (4.25)$$

Combining estimates (4.21) and (4.25) with the smallness assumption on \mathbf{G} and \mathbf{H} , choosing $\varepsilon_0 \leq C\varepsilon$ and applying Theorem 4.4 gives the unique existence of a solution (\mathbf{v}, \mathbf{q}) of (4.20) with

$$\mathbf{v} \in H_p^1(\mathbb{T}, L_q(\Omega)^N) \cap L_p(\mathbb{T}, H_q^2(\Omega)^N), \quad \mathbf{q} \in L_p(\mathbb{T}, \hat{H}_q^1(\Omega)),$$

possessing the estimate

$$E(\mathbf{v}, \mathbf{q}) \leq C(\varepsilon^2 + \varepsilon E(\mathbf{u}, \mathbf{p}) + E(\mathbf{u}, \mathbf{p})^2). \quad (4.26)$$

Since we assume that $E(\mathbf{u}, \mathbf{p}) \leq \varepsilon$, by (4.26) we have $E(\mathbf{v}, \mathbf{q}) \leq 3C\varepsilon^2$. Therefore, choosing $\varepsilon > 0$ so small that $3C\varepsilon \leq 1$, we have $E(\mathbf{v}, \mathbf{q}) \leq \varepsilon$, so that $(\mathbf{v}, \mathbf{q}) \in \mathcal{I}_\varepsilon$. Thus, the map Ξ acting on $(\mathbf{u}, \mathbf{p}) \in \mathcal{I}_\varepsilon$ by setting $\Xi(\mathbf{u}, \mathbf{p}) = (\mathbf{v}, \mathbf{q})$ is a map from \mathcal{I}_ε into \mathcal{I}_ε .

We next prove that Ξ is a contraction map. Let $(\mathbf{u}_i, \mathbf{p}_i)$ ($i = 1, 2$) be any two elements of \mathcal{I}_ε and set $(\mathbf{v}_i, \mathbf{q}_i) = \Xi(\mathbf{u}_i, \mathbf{p}_i)$. Since

$$\begin{aligned} \mathcal{N}(\mathbf{u}_1) - \mathcal{N}(\mathbf{u}_2) & = ((\mathbf{I} + \mathbf{B}_{-1})(\mathbf{u}_1 - \mathbf{u}_2)) \cdot (\mathbf{I} + \mathbf{A})\nabla((\mathbf{I} + \mathbf{B}_{-1})\mathbf{u}_1) \\ & \quad + ((\mathbf{I} + \mathbf{B}_{-1})\mathbf{u}_2) \cdot (\mathbf{I} + \mathbf{A})\nabla((\mathbf{I} + \mathbf{B}_{-1})(\mathbf{u}_1 - \mathbf{u}_2)), \end{aligned}$$

by Sobolev's inequality and the assumption $N < q < \infty$, we have

$$\|\mathcal{N}(\mathbf{u}_1) - \mathcal{N}(\mathbf{u}_2)\|_{L_q(\Omega)} \leq C(\|\mathbf{u}_1 - \mathbf{u}_2\|_{L_q(\Omega)}\|\mathbf{u}_1\|_{H_q^2(\Omega)} + \|\mathbf{u}_2\|_{L_q(\Omega)}\|\mathbf{u}_1 - \mathbf{u}_2\|_{H_q^2(\Omega)}).$$

Thus, from (4.24) we deduce

$$\begin{aligned} \|\mathcal{N}(\mathbf{u}_1) - \mathcal{N}(\mathbf{u}_2)\|_{L_p(\mathbb{T}, L_q(\Omega))} &\leq C(\|(\mathbf{u}_1, \mathbf{u}_2)\|_{H_p^1(\mathbb{T}, L_q(\Omega))} + \|(\mathbf{u}_1, \mathbf{u}_2)\|_{L_p(\mathbb{T}, H_q^2(\Omega))}) \\ &\quad \times (\|\mathbf{u}_1 - \mathbf{u}_2\|_{H_p^1(\mathbb{T}, L_q(\Omega))} + \|\mathbf{u}_1 - \mathbf{u}_2\|_{L_p(\mathbb{T}, H_q^2(\Omega))}) \\ &\leq C\varepsilon E(\mathbf{u}_1 - \mathbf{u}_2, \mathbf{p}_1 - \mathbf{p}_2). \end{aligned} \quad (4.27)$$

Since \mathcal{L} is a linear operation, by (4.21), we have

$$\|\mathcal{L}(\mathbf{u}_1 - \mathbf{u}_2, \mathbf{p}_1 - \mathbf{p}_2)\|_{L_p(\mathbb{T}, L_q(\Omega))} \leq C\varepsilon E(\mathbf{u}_1 - \mathbf{u}_2, \mathbf{p}_1 - \mathbf{p}_2). \quad (4.28)$$

Moreover, $\mathbf{v} = \mathbf{v}_1 - \mathbf{v}_2$ and $\mathbf{q} = \mathbf{q}_1 - \mathbf{q}_2$ satisfy the equations

$$\partial_t \mathbf{v} - \Delta \mathbf{v} + \nabla \mathbf{q} = \mathcal{L}(\mathbf{u}_1 - \mathbf{u}_2, \mathbf{p}_1 - \mathbf{p}_2) + (\mathcal{N}(\mathbf{u}_1) - \mathcal{N}(\mathbf{u}_2)), \quad \operatorname{div} \mathbf{v} = 0 \quad \text{in } \Omega \times \mathbb{T}, \quad \mathbf{v}|_\Gamma = 0.$$

Applying Theorem 4.4 and using (4.27) and (4.28) now gives that

$$E(\mathbf{v}_1 - \mathbf{v}_2, \mathbf{q}_1 - \mathbf{q}_2) \leq C\varepsilon E(\mathbf{u}_1 - \mathbf{u}_2, \mathbf{p}_1 - \mathbf{p}_2).$$

Choosing $\varepsilon > 0$ smaller if necessary, we may assume that $C\varepsilon < 1$, and so Ξ is a contraction map on \mathcal{I}_ε , which yields the unique existence of $(\mathbf{u}, \mathbf{p}) \in \mathcal{I}_\varepsilon$ such that $(\mathbf{u}, \mathbf{p}) = \Xi(\mathbf{u}, \mathbf{p})$. Obviously, (\mathbf{u}, \mathbf{p}) is the required solution of (4.9). This completes the proof of Theorem 4.1. \square

Remark 4.5. In the case of no-slip boundary conditions, the fluid particles at the boundary are attached to the body, so that the fluid velocity coincides with the velocity of boundary particles. Then, the boundary data \mathbf{h} in (4.3) are given by

$$\mathbf{h}(x, t) = \partial_t \Phi_t(y) = \partial_t \phi(t, y),$$

where $x = \Phi_t(y) = y + \phi(y, t)$, and in the formulation (4.9) on a time-independent domain, this corresponds to boundary data

$$\mathbf{H} = (\mathbf{I} + \mathbf{J}_0 \mathbf{I} + \mathbf{A}^\top \mathbf{J}) \partial_t \phi.$$

Therefore, the assumptions on \mathbf{H} in Theorem 4.1 are additional regularity and smallness assumptions on ϕ in this case. Moreover, the compatibility condition (4.11) is satisfied if and only if Φ_t preserves the volume of Ω .

5 On periodic Navier–Stokes flow around a body at rest

5.1 Problem and main results

Let Ω be an exterior domain in \mathbb{R}^3 , that is, a domain that is the complement of a compact set. We assume that its boundary Γ is a C^2 hypersurface. Let $b > 0$ be a sufficiently large number such that $\Omega^c \subset B_b$, where $\Omega^c = \mathbb{R}^3 \setminus \Omega$ and $B_b = \{x \in \mathbb{R}^N \mid |x| < b\}$. We further set $S_b = \{x \in \mathbb{R}^N \mid |x| = b\}$. We consider the Navier–Stokes equations in Ω :

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} - \mu \Delta \mathbf{u} + \nabla \mathbf{p} = \mathbf{F}, \quad \operatorname{div} \mathbf{u} = 0 \quad \text{in } \Omega, \quad \mathbf{u}|_\Gamma = \mathbf{h}|_\Gamma. \quad (5.1)$$

Here, $\partial_j = \partial/\partial x_j$, $x = (x_1, x_2, x_3) \in \mathbb{R}^3$, $\mathbf{u} = (u_1, u_2, u_3)^\top$ is an unknown velocity field, \mathbf{p} an unknown pressure field, $\mathbf{F} = (F_1, \dots, F_N)^\top$ a prescribed external force, and $\mathbf{h} = (h_1, \dots, h_N)^\top$ are prescribed boundary data. Assume that $\mathbf{F}(t + 2\pi) = \mathbf{F}(t)$ and $\mathbf{h}(t + 2\pi) = \mathbf{h}(t)$ for any $t \in \mathbb{R}$.

Then (5.1) describes the fluid flow around a body, subject to time-periodic external forcing \mathbf{F} and with boundary data \mathbf{h} .

Given any time-periodic function $f = f(x, t)$, with period 2π , we write

$$f_S(x) = \int_{\mathbb{T}} f(x, t) dt = \frac{1}{2\pi} \int_0^{2\pi} f(x, t) dt, \quad f_{\perp}(x, t) = f(x, t) - f_S(x), \quad (5.2)$$

and f_S and f_{\perp} are called the stationary part of f and the oscillatory part of f , respectively. By means of this decomposition, we divide the data and the solution into two parts, which have different asymptotic properties as $|x| \rightarrow \infty$. To quantify this spatial decay, we set

$$\langle f_S \rangle_{\ell} = \sup_{x \in \Omega} |f_S(x)| (1 + |x|)^{\ell}, \quad \langle f_{\perp} \rangle_{p, \ell} = \sup_{x \in \Omega} \|f_{\perp}(x, \cdot)\|_{L_p(\mathbb{T})} (1 + |x|)^{\ell}$$

for $\ell \in \mathbb{R}$ and $p \in (1, \infty)$. We shall prove the following theorem.

Theorem 5.1. *Let $2 < p < \infty$ and $3 < q < \infty$. Assume that $\mathbf{F} = \mathbf{F}_S + \mathbf{F}_{\perp}$ with $\mathbf{F}_S = \operatorname{div} \mathbf{G}_S$ and $\mathbf{F}_{\perp} = \operatorname{div} \mathbf{G}_{\perp}$. Then there exists a small constant $\varepsilon > 0$ depending on p and q such that if \mathbf{F} and $\mathbf{h} \in H_p^1(\mathbb{T}, L_q(\Omega)^3) \cap L_p(\mathbb{T}, H_q^2(\Omega)^3)$ satisfy the smallness condition*

$$\langle \mathbf{F}_S \rangle_3 + \langle \mathbf{G}_S \rangle_2 + \langle \mathbf{F}_{\perp} \rangle_{p, 2} + \langle \mathbf{G}_{\perp} \rangle_{p, 1} + \|\mathbf{h}\|_{H_p^1(\mathbb{T}, L_q(\Omega))} + \|\mathbf{h}\|_{L_p(\mathbb{T}, H_q^2(\Omega))} < \varepsilon^2, \quad (5.3)$$

then problem (5.1) admits a unique solution (\mathbf{u}, \mathbf{p}) such that $\mathbf{u} = \mathbf{u}_S + \mathbf{u}_{\perp}$ and $\mathbf{p} = \mathbf{p}_S + \mathbf{p}_{\perp}$ with

$$\mathbf{u}_S \in H_q^2(\Omega)^3, \quad \mathbf{u}_{\perp} \in H_p^1(\mathbb{T}, L_q(\Omega)) \cap L_p(\mathbb{T}, H_q^2(\Omega)), \quad \mathbf{p}_S \in H_q^1(\Omega), \quad \mathbf{p}_{\perp} \in L_p(\mathbb{T}, \hat{H}_q^1(\Omega))$$

satisfying the estimate

$$\begin{aligned} & \langle \mathbf{u}_S \rangle_1 + \langle \nabla \mathbf{u}_S \rangle_2 + \|\mathbf{u}_S\|_{H_q^2(\Omega)} + \|\mathbf{p}_S\|_{H_q^1(\Omega)} \\ & + \langle \mathbf{u}_{\perp} \rangle_{p, 1} + \langle \nabla \mathbf{u}_{\perp} \rangle_{p, 2} + \|\mathbf{u}_{\perp}\|_{L_p(\mathbb{T}, H_q^2(\Omega))} + \|\partial_t \mathbf{u}_{\perp}\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\nabla \mathbf{p}_{\perp}\|_{L_p(\mathbb{T}, L_q(\Omega))} \leq \varepsilon. \end{aligned}$$

Our proof of Theorem 5.1 is based on the study of the associated linearized system, the time-periodic Stokes problem

$$\partial_t \mathbf{v} - \mu \Delta \mathbf{v} + \nabla \mathbf{p} = \mathbf{f}, \quad \operatorname{div} \mathbf{v} = 0 \quad \text{in } \Omega \times \mathbb{T}, \quad \mathbf{v}|_{\Gamma} = \mathbf{h}. \quad (5.4)$$

We shall derive the following theorem, which ensures existence of a unique solution to (5.4) in a framework of spatially weighted spaces. For shorter notation, we set $L_{q, 3b}(\Omega) = \{f \in L_q(\Omega) \mid \operatorname{supp} f \subset B_{3b}\}$.

Theorem 5.2. *Let $1 < p < \infty$, $3 < q < \infty$ and $\ell \in (0, 3]$. For all $\mathbf{f} = \mathbf{f}_S + \mathbf{f}_{\perp} \in L_p(\mathbb{T}, L_q(\Omega)^3)$ such that $\mathbf{f}_S = \operatorname{div} \mathbf{G}_S + \mathbf{g}_S$ and $\mathbf{f}_{\perp} = \operatorname{div} \mathbf{G}_{\perp} + \mathbf{g}_{\perp}$ with $\mathbf{g} = \mathbf{g}_S + \mathbf{g}_{\perp} \in L_p(\mathbb{T}, L_{q, 3b}(\Omega)^3)$ and*

$$\langle \mathbf{G}_S \rangle_2 + \langle \operatorname{div} \mathbf{G}_S \rangle_3 + \langle \mathbf{G}_{\perp} \rangle_{p, \ell} + \langle \operatorname{div} \mathbf{G}_{\perp} \rangle_{p, \ell+1} < \infty,$$

problem (5.4) admits a unique solution (\mathbf{v}, \mathbf{p}) with

$$\mathbf{v} \in H_p^1(\mathbb{T}, L_q(\Omega)^3) \cap L_p(\mathbb{T}, H_q^2(\Omega)^3), \quad \mathbf{p} \in L_p(\mathbb{T}, \hat{H}_q^1(\Omega)),$$

possessing the estimate

$$\begin{aligned} & \|\mathbf{v}_S\|_{H_q^2(\Omega)} + \langle \mathbf{v}_S \rangle_1 + \langle \nabla \mathbf{v}_S \rangle_2 + \|\mathbf{p}_S\|_{H_q^1(\Omega)} + \langle \mathbf{p}_S \rangle_2 \\ & + \|\partial_t \mathbf{v}_{\perp}\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{v}_{\perp}\|_{L_p(\mathbb{T}, H_q^2(\Omega))} + \langle \mathbf{v}_{\perp} \rangle_{p, \ell} + \langle \nabla \mathbf{v}_{\perp} \rangle_{p, \ell+1} + \|\nabla \mathbf{p}_{\perp}\|_{L_p(\mathbb{T}, L_q(\Omega))} \\ & \leq C(\langle \operatorname{div} \mathbf{G}_S \rangle_3 + \langle \mathbf{G}_S \rangle_2 + \langle \operatorname{div} \mathbf{G}_{\perp} \rangle_{p, 1+\ell} + \langle \mathbf{G}_{\perp} \rangle_{p, \ell} \\ & \quad + \|\mathbf{g}\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{h}\|_{H_p^1(\mathbb{T}, L_q(\Omega))} + \|\mathbf{h}\|_{L_p(\mathbb{T}, H_q^2(\Omega))}). \end{aligned} \quad (5.5)$$

For the proof of this theorem, we split (5.4) into two separate problems by means of the decomposition (5.2). In the following subsections these problems are analyzed independently of each other in the case of vanishing boundary data. In Subsection 5.4 we return to the original linear and nonlinear problems (5.4) and (5.1) and complete the proofs of Theorems 5.1 and 5.2.

5.2 Stationary solutions to the Stokes problem

Here we examine time-independent solutions to (5.4) with vanishing boundary data $\mathbf{h} = 0$, that is, solutions (\mathbf{u}, \mathbf{p}) to the stationary problem

$$-\mu\Delta\mathbf{u} + \nabla\mathbf{p} = \operatorname{div}\mathbf{F} + \mathbf{g}, \quad \operatorname{div}\mathbf{u} = 0 \quad \text{in } \Omega, \quad \mathbf{u}|_{\Gamma} = 0. \quad (5.6)$$

We shall derive the following theorem.

Theorem 5.3. *Let $3 < q < \infty$. If \mathbf{F} satisfies the condition $\langle \operatorname{div}\mathbf{F} \rangle_3 + \langle \mathbf{F} \rangle_2 < \infty$ and $\mathbf{g} \in L_{q,3b}(\Omega)^3$, then problem (5.6) admits a unique solution $(\mathbf{u}, \mathbf{p}) \in H_q^2(\Omega)^3 \times H_q^1(\Omega)$ possessing the estimate:*

$$\|\mathbf{u}\|_{H_q^2(\Omega)} + \langle \mathbf{u} \rangle_1 + \langle \nabla\mathbf{u} \rangle_2 + \|\mathbf{p}\|_{H_q^1(\Omega)} + \langle \mathbf{p} \rangle_2 \leq C(\langle \operatorname{div}\mathbf{F} \rangle_3 + \langle \mathbf{F} \rangle_2 + \|\mathbf{g}\|_{L_q(\Omega)}) \quad (5.7)$$

with some constant $C > 0$.

For the proof, we first consider the Stokes equations in \mathbb{R}^3 :

$$-\mu\Delta\mathbf{u} + \nabla\mathbf{p} = \mathbf{f}, \quad \operatorname{div}\mathbf{u} = 0 \quad \text{in } \mathbb{R}^3. \quad (5.8)$$

As is well-known (cf. Galdi [25, pp.239-240]), there exist fundamental solutions $\mathbf{U} = (U_{ij}(x))$ and $\mathbf{q} = (q_1(x), q_2(x), q_3(x))^T$ of equations (5.8) with

$$U_{ij}(x) = -\frac{1}{8\pi\mu} \left(\frac{\delta_{ij}}{|x|} + \frac{x_i x_j}{|x|^3} \right), \quad q_j(x) = \frac{1}{4\pi} \frac{x_j}{|x|^3}. \quad (5.9)$$

If we set

$$\mathbf{u}(x) = \mathbf{U} * \mathbf{f}(x) := \int_{\mathbb{R}^3} \mathbf{U}(y) \mathbf{f}(x-y) dy, \quad \mathbf{p}(x) = \mathbf{q} * \mathbf{f} := \int_{\mathbb{R}^3} \mathbf{q}(y) \cdot \mathbf{f}(x-y) dy, \quad (5.10)$$

then, \mathbf{u} and \mathbf{p} formally satisfy equations (5.8). We prove the following lemma.

Lemma 5.4. *Let $3 < q < \infty$.*

(1) *Let \mathbf{F} be a function satisfying $\langle \operatorname{div}\mathbf{F} \rangle_3 + \langle \mathbf{F} \rangle_2 < \infty$ and set $\mathbf{u} = \mathbf{U} * (\operatorname{div}\mathbf{F})$ and $\mathbf{p} = \mathbf{q} * (\operatorname{div}\mathbf{F})$. Then, we have*

$$\|\mathbf{u}\|_{H_q^2(\mathbb{R}^3)} + \|\mathbf{p}\|_{H_q^1(\mathbb{R}^3)} + \langle \mathbf{u} \rangle_1 + \langle \nabla\mathbf{u} \rangle_2 + \langle \mathbf{p} \rangle_2 \leq C(\langle \mathbf{F} \rangle_2 + \langle \operatorname{div}\mathbf{F} \rangle_3)$$

with some constant $C > 0$.

(2) *Let $\mathbf{g} \in L_q(\mathbb{R}^3)$ such that \mathbf{g} vanishes for $|x| > b$ with some constant $b > 0$. Let $\mathbf{v} = \mathbf{U} * \mathbf{g}$ and $\mathbf{q} = \mathbf{q} * \mathbf{g}$. Then, we have*

$$\|\mathbf{v}\|_{H_q^2(\mathbb{R}^3)} + \|\mathbf{q}\|_{H_q^1(\mathbb{R}^3)} + \langle \mathbf{v} \rangle_1 + \langle \nabla\mathbf{v} \rangle_2 + \langle \mathbf{q} \rangle_2 \leq C\|\mathbf{g}\|_{L_q(\mathbb{R}^3)}$$

for some constant C .

Proof. (1) The theory of singular integrals yields that

$$\begin{aligned} \|\nabla^2 \mathbf{u}\|_{L_q(\mathbb{R}^3)} + \|\nabla \mathbf{p}\|_{L_q(\mathbb{R}^3)} &\leq C \|\operatorname{div} \mathbf{F}\|_{L_q(\mathbb{R}^3)} \leq C_q \langle \operatorname{div} \mathbf{F} \rangle_3; \\ \|\nabla \mathbf{u}\|_{L_q(\mathbb{R}^3)} + \|\mathbf{p}\|_{L_q(\mathbb{R}^3)} &\leq C \|\mathbf{F}\|_{L_q(\mathbb{R}^3)} \leq C_q \langle \mathbf{F} \rangle_2. \end{aligned}$$

For notational simplicity, set $\gamma = \langle \operatorname{div} \mathbf{F} \rangle_3 + \langle \mathbf{F} \rangle_2$. By the Gaussian divergence theorem, we write

$$\begin{aligned} \mathbf{u}(x) &= \int_{|y| \leq |x|/2} \mathbf{U}(y) (\operatorname{div} \mathbf{F})(x-y) dy - \int_{|y|=|x|/2} \mathbf{U}(y) \frac{y}{|y|} \cdot \mathbf{F}(x-y) d\omega \\ &\quad + \int_{|x|/2 \leq |y| \leq 2|x|} \nabla \mathbf{U}(y) \mathbf{F}(x-y) dy + \int_{|y| \geq 2|x|} \nabla \mathbf{U}(y) \mathbf{F}(x-y) dy. \end{aligned}$$

Noting that $|x-y| \geq |x|/2$ for $|y| \leq |x|/2$, $|x-y| \leq 3|x|$ for $|x|/2 \leq |y| \leq 2|x|$, and $|x-y| \geq |y|/2$ for $|y| \geq 2|x|$, by (5.9) we have

$$\begin{aligned} |\mathbf{u}(x)| &\leq C\gamma \left\{ (1+|x|)^{-3} \int_{|y| \leq |x|/2} |y|^{-1} dy + |x|^{-1} (1+|x|)^{-2} \int_{|y|=|x|/2} d\omega \right. \\ &\quad \left. + |x|^{-2} \int_{|z| \leq 3|x|} |z|^{-2} dz + \int_{|y| \geq 2|x|} |y|^{-4} dy \right\} \leq C\gamma |x|^{-1} \end{aligned}$$

for $x \neq 0$. When $|x| \leq 1$, noting that $|(\operatorname{div} \mathbf{F})(x-y)| \leq \gamma$ for $|y| \leq 2$ and $|(\operatorname{div} \mathbf{F})(x-y)| \leq C\gamma |y|^{-3}$ for $|y| \geq 2$, we have

$$\begin{aligned} |\mathbf{u}(x)| &\leq \int_{|y| \leq 2} |\mathbf{U}(y) (\operatorname{div} \mathbf{F})(x-y)| dy + \int_{|y| \geq 2} |\mathbf{U}(y) (\operatorname{div} \mathbf{F})(x-y)| dy \\ &\leq C\gamma \left\{ \int_{|y| \leq 2} |y|^{-1} dy + \int_{|y| \geq 2} |y|^{-4} dy \right\} \leq C\gamma. \end{aligned}$$

In total, we thus have $\langle \mathbf{u} \rangle_1 \leq C\gamma$. In particular, noting that $3 < q < \infty$, we obtain

$$\|\mathbf{u}\|_{L_q(\mathbb{R}^3)} \leq C_q \langle \mathbf{u} \rangle_1 \leq C_q \gamma.$$

Similarly to before, we proceed with the estimate of $\nabla \mathbf{u}$ and write

$$\begin{aligned} \nabla \mathbf{u}(x) &= \int_{|y| \leq |x|/2} \nabla \mathbf{U}(y) (\operatorname{div} \mathbf{F})(x-y) dy - \int_{|y|=|x|/2} \nabla \mathbf{U}(y) \frac{y}{|y|} \cdot \mathbf{F}(x-y) d\omega \\ &\quad + \int_{|x|/2 \leq |y| \leq 2|x|} \nabla^2 \mathbf{U}(y) \mathbf{F}(x-y) dy + \int_{|y| \geq 2|x|} \nabla^2 \mathbf{U}(y) \mathbf{F}(x-y) dy. \end{aligned}$$

Then, we have

$$\begin{aligned} |\nabla \mathbf{u}(x)| &\leq C\gamma \left\{ (1+|x|)^{-3} \int_{|y| \leq |x|/2} |y|^{-2} dy + |x|^{-2} (1+|x|)^{-2} \int_{|y|=|x|/2} d\omega \right. \\ &\quad \left. + |x|^{-3} \int_{|z| \leq 3|x|} |z|^{-2} dz + \int_{|y| \geq 2|x|} |y|^{-5} dy \right\} \leq C\gamma |x|^{-2} \end{aligned}$$

for $x \neq 0$. When $|x| \leq 1$, arguing as above, we have

$$\begin{aligned} |\nabla \mathbf{u}(x)| &\leq \int_{|y| \leq 2} |\nabla \mathbf{U}(y) (\operatorname{div} \mathbf{F})(x-y)| dy + \int_{|y| \geq 2} |\nabla \mathbf{U}(y) (\operatorname{div} \mathbf{F})(x-y)| dy \\ &\leq C\gamma \left\{ \int_{|y| \leq 2} |y|^{-2} dy + \int_{|y| \geq 2} |y|^{-5} dy \right\} \leq C\gamma. \end{aligned}$$

Summing up, we have $\langle \nabla \mathbf{u} \rangle_2 \leq C\gamma$.

In the very same way, we can use $|\mathbf{q}(y)| \leq C|y|^{-2}$ and $|\nabla \mathbf{q}(y)| \leq C|y|^{-3}$ for $y \neq 0$ to deduce $\langle \mathbf{p} \rangle_2 \leq C\gamma$. In particular, $\|\mathbf{p}\|_{L_q(\mathbb{R}^3)} \leq C_q \langle \mathbf{p} \rangle_2 \leq C_q\gamma$, because $3 < q < \infty$. This completes the proof of the first part of Lemma 5.4.

(2) As before, the theory of singular integral operators yields that

$$\|\nabla^2 \mathbf{v}\|_{L_q(\mathbb{R}^3)} + \|\nabla \mathbf{q}\|_{L_q(\mathbb{R}^3)} \leq C\|\mathbf{g}\|_{L_q(\mathbb{R}^3)}. \quad (5.11)$$

By estimates for weak singular integral operators (cf. [25, II.11]), we further have

$$\|\nabla \mathbf{v}\|_{L_q(B_{2b})} + \|\mathbf{v}\|_{L_q(B_{2b})} + \|\mathbf{q}\|_{L_q(B_{2b})} \leq C_{b,q}\|\mathbf{g}\|_{L_q(\mathbb{R}^3)}. \quad (5.12)$$

Since $|\mathbf{U}(x)| \leq C|x|^{-1}$, $|\nabla \mathbf{U}(x)| \leq C|x|^{-2}$ and $|\mathbf{q}(x)| \leq C|x|^{-2}$, noting that $\mathbf{g}(y) = 0$ for $|y| > b$, for $|x| > 2b$ we have

$$\begin{aligned} |\mathbf{v}(x)| &\leq C \int_{|y|<b} \frac{|\mathbf{g}(y)|}{|x-y|} dy \leq C_b|x|^{-1} \int_{|y|<b} |\mathbf{g}(y)| dy \leq C_b|x|^{-1}\|\mathbf{g}\|_{L_q(\mathbb{R}^3)}; \\ |\nabla \mathbf{v}(x)| &\leq C \int_{|y|<b} \frac{|\mathbf{g}(y)|}{|x-y|^2} dy \leq C_b|x|^{-2} \int_{|y|<b} |\mathbf{g}(y)| dy \leq C_b|x|^{-2}\|\mathbf{g}\|_{L_q(\mathbb{R}^3)}; \\ |\mathbf{q}(x)| &\leq C \int_{|y|<b} \frac{|\mathbf{g}(y)|}{|x-y|^2} dy \leq C_b|x|^{-2} \int_{|y|<b} |\mathbf{g}(y)| dy \leq C_b|x|^{-2}\|\mathbf{g}\|_{L_q(\mathbb{R}^3)}. \end{aligned} \quad (5.13)$$

In particular, setting $(B_{2b})^c = \mathbb{R}^3 \setminus B_{2b}$, we conclude

$$\|\mathbf{v}\|_{L_q((B_{2b})^c)} + \|\nabla \mathbf{v}\|_{L_q((B_{2b})^c)} + \|\mathbf{q}\|_{L_q((B_{2b})^c)} \leq C_{b,q}\|\mathbf{g}\|_{L_q(\mathbb{R}^3)},$$

because $3 < q < \infty$, which, combined with (5.11) and (5.12), yields that

$$\|\mathbf{v}\|_{H_q^2(\mathbb{R}^3)} + \|\mathbf{q}\|_{H_q^1(\mathbb{R}^3)} \leq C_{q,b}\|\mathbf{g}\|_{L_q(\mathbb{R}^3)}.$$

By Sobolev's inequality, we have

$$\begin{aligned} \sup_{|x| \leq 2b} (1+|x|)|\mathbf{v}(x)| + \sup_{|x| \leq 2b} (1+|x|)^2|\nabla \mathbf{v}(x)| + \sup_{|x| \leq 2b} (1+|x|)^2|\mathbf{q}(x)| \\ \leq C(\|\mathbf{v}\|_{H_q^2(\mathbb{R}^3)} + \|\mathbf{q}\|_{H_q^1(\mathbb{R}^3)}) \leq C_{q,b}\|\mathbf{g}\|_{L_q(\mathbb{R}^3)}, \end{aligned}$$

which, combined with (5.13), yields that

$$\langle \mathbf{v} \rangle_1 + \langle \nabla \mathbf{v} \rangle_2 + \langle \mathbf{q} \rangle_2 \leq C_{q,b}\|\mathbf{g}\|_{L_q(\mathbb{R}^3)}.$$

This completes the proof of the second part of Lemma 5.4. \square

Proof of Theorem 5.3. To construct a solution operator for problem (5.6), we first consider the case where $\mathbf{f} \in L_{q,3b}(\Omega)^3$. Firstly, let \mathbf{f}_0 be the zero extension of \mathbf{f} to the complement of Ω , that is, we set $\mathbf{f}_0(x) = \mathbf{f}(x)$ for $x \in \Omega$ and $\mathbf{f}_0(x) = 0$ for $x \notin \Omega$. Let $\mathcal{T}_0 \mathbf{f}_0 = \mathbf{U} * \mathbf{f}_0$ and $P_0 \mathbf{f}_0 = \mathbf{q} * \mathbf{f}_0$. Secondly, let \mathbf{f}_b be the restriction of $\mathbf{f} \in L_{q,3b}(\Omega)^3$ to Ω_{4b} , and let \mathcal{A}_0 and \mathcal{B}_0 be the operators acting on $\mathbf{f}_b \in L_q(\Omega_{4b})^3$ such that $\mathcal{A}_0 \mathbf{f}_b \in H_q^2(\Omega_{4b})^3$, $\mathcal{B}_0 \mathbf{f}_b \in \hat{H}_q^1(\Omega_{4b})$ satisfy the equations

$$-\mu \Delta \mathcal{A}_0 \mathbf{f}_b + \nabla \mathcal{B}_0 \mathbf{f}_b = \mathbf{f}_b, \quad \operatorname{div} \mathcal{A}_0 \mathbf{f}_b = 0 \quad \text{in } \Omega_{4b}, \quad \mathcal{A}_0 \mathbf{f}_b|_{\Gamma} = \mathcal{A}_0 \mathbf{f}_b|_{S_{4b}} = 0, \quad (5.14)$$

and possess the estimate

$$\|\mathcal{A}_0 \mathbf{f}_b\|_{\mathbb{H}_q^2(\Omega_{4b})} + \|\mathcal{B}_0 \mathbf{f}_b\|_{\dot{\mathbb{H}}_q^1(\Omega_{4b})} \leq C \|\mathbf{f}_b\|_{L_q(\Omega_{4b})}. \quad (5.15)$$

Since $\mathcal{B}_0 \mathbf{f}_b$ is only defined up to a constant, choosing a constant suitably, we may assume that

$$\int_{\Omega_{4b}} (P_0 \mathbf{f}_0 - \mathcal{B}_0 \mathbf{f}_b) dx = 0. \quad (5.16)$$

In what follows, let φ be a function in $C^\infty(\mathbb{R}^3)$ that equals 1 for $x \in B_{2b}$ and 0 for $x \notin B_{3b}$, and let \mathbb{B} be the Bogovskiĭ operator. For $\mathbf{f} \in L_{q,3b}(\Omega)^3$, let

$$\mathcal{V}_0 \mathbf{f} = (1 - \varphi) \mathcal{T}_0 \mathbf{f}_0 + \varphi \mathcal{A}_0 \mathbf{f}_b + \mathbb{B}[(\nabla \varphi)(\mathcal{T}_0 \mathbf{f}_0 - \mathcal{A}_0 \mathbf{f}_b)], \quad \mathcal{W}_0 \mathbf{f} = (1 - \varphi) P_0 \mathbf{f}_0 + \varphi \mathcal{B}_0 \mathbf{f}_b. \quad (5.17)$$

Inserting these formulas into equations (5.6), we have

$$-\mu \Delta \mathcal{V}_0 \mathbf{f} + \nabla \mathcal{W}_0 \mathbf{f} = \mathbf{f} + \mathcal{R}_1 \mathbf{f}, \quad \operatorname{div} \mathcal{V}_0 \mathbf{f} = 0 \quad \text{in } \Omega, \quad \mathcal{V}_0 \mathbf{f}|_\Gamma = 0, \quad (5.18)$$

where

$$\begin{aligned} \mathcal{R}_1 \mathbf{f} &= 2\mu(\nabla \varphi) \cdot (\nabla \mathcal{T}_0 \mathbf{f}_0 - \nabla \mathcal{A}_0 \mathbf{f}_b) + \mu(\Delta \varphi)(\mathcal{T}_0 \mathbf{f}_0 - \mathcal{A}_0 \mathbf{f}_b) \\ &\quad - \mu \Delta \mathbb{B}[(\nabla \varphi) \cdot (\mathcal{T}_0 \mathbf{f}_0 - \mathcal{A}_0 \mathbf{f}_b)] - (\nabla \varphi)(P_0 \mathbf{f}_0 - \mathcal{B}_0 \mathbf{f}_b). \end{aligned}$$

Employing the same arguments as in [29] and [33], we shall show that the inverse operator $(I + \mathcal{R}_1)^{-1} \in \mathcal{L}(L_{q,3b}(\Omega)^3)$ exists and

$$\|(I + \mathcal{R}_1)^{-1} \mathbf{f}\|_{L_q(\Omega)} \leq C \|\mathbf{f}\|_{L_q(\Omega)} \quad (5.19)$$

for any $\mathbf{f} \in L_{q,3b}(\Omega)^3$. Postponing proving (5.19), we continue the proof of Theorem 5.3. Setting

$$\begin{aligned} \mathcal{U}_0 \mathbf{f} &:= \mathcal{V}_0 (I + \mathcal{R}_1)^{-1} \mathbf{f} \\ &= (1 - \varphi) \mathcal{T}_0 ((I + \mathcal{R}_1)^{-1} \mathbf{f})_0 + \varphi \mathcal{A}_0 ((I + \mathcal{R}_1)^{-1} \mathbf{f})_b \\ &\quad + \mathbb{B}[(\nabla \varphi)(\mathcal{T}_0 ((I + \mathcal{R}_1)^{-1} \mathbf{f})_0 - \mathcal{A}_0 ((I + \mathcal{R}_1)^{-1} \mathbf{f})_b)], \\ \mathcal{Q}_0 \mathbf{f} &:= \mathcal{W}_0 (I + \mathcal{R}_1)^{-1} \mathbf{f} = (1 - \varphi) P_0 ((I + \mathcal{R}_1)^{-1} \mathbf{f})_0 + \varphi \mathcal{B}_0 ((I + \mathcal{R}_1)^{-1} \mathbf{f})_b, \end{aligned}$$

we see that $(\mathbf{u}, \mathbf{p}) = (\mathcal{U}_0 \mathbf{f}, \mathcal{Q}_0 \mathbf{f})$ is a solution to problem (5.6). Moreover, by (5.19), Lemma 5.4 (2) and (5.15), we have

$$\begin{aligned} \sup_{x \in \Omega} (1 + |x|) |\mathcal{U}_0 \mathbf{f}(x)| &\leq C \|\mathbf{f}\|_{L_q(\Omega)}, & \sup_{x \in \Omega} (1 + |x|)^2 |\nabla \mathcal{U}_0 \mathbf{f}(x)| &\leq C \|\mathbf{f}\|_{L_q(\Omega)}, \\ \sup_{x \in \Omega} (1 + |x|)^2 |\mathcal{Q}_0 \mathbf{f}(x)| &\leq C \|\mathbf{f}\|_{L_q(\Omega)}, & \|\mathcal{U}_0 \mathbf{f}\|_{\mathbb{H}_q^2(\Omega)} + \|\mathcal{Q}_0 \mathbf{f}\|_{\mathbb{H}_q^1(\Omega)} &\leq C \|\mathbf{f}\|_{L_q(\Omega)}. \end{aligned} \quad (5.20)$$

We now consider the case where $\mathbf{f} = \operatorname{div} \mathbf{F} + \mathbf{g}$ with $\langle \operatorname{div} \mathbf{F} \rangle_3 + \langle \mathbf{F} \rangle_2 < \infty$ and $\mathbf{g} \in L_{q,3b}(\Omega)$. We write $\mathbf{f} = \operatorname{div}((1 - \varphi)\mathbf{F}) + \mathbf{h}$ with $\mathbf{h} = \varphi \operatorname{div} \mathbf{F} + (\nabla \varphi) \cdot \mathbf{F} + \mathbf{g}$. Let

$$\mathbf{u} = (1 - \varphi) \mathcal{T}_0 \mathbf{f}_0 + \mathbb{B}[(\nabla \varphi) \mathcal{T}_0 \mathbf{f}_0], \quad \mathbf{p} = (1 - \varphi) P_0 \mathbf{f}_0.$$

Notice that $\mathbf{f}_0 = \operatorname{div}((1 - \varphi)\mathbf{F}) + \mathbf{h}_0$. We see that \mathbf{u} and \mathbf{p} satisfy the equations:

$$-\mu \Delta \mathbf{u} + \nabla \mathbf{p} = (1 - \varphi) \mathbf{f}_0 + \mathcal{R}_2 \mathbf{f}, \quad \operatorname{div} \mathbf{u} = 0 \quad \text{in } \Omega, \quad \mathbf{u}|_\Gamma = 0,$$

where we have set

$$\mathcal{R}_2 \mathbf{f} = 2\mu(\nabla \varphi) \cdot \nabla \mathcal{T}_0 \mathbf{f}_0 + \mu(\Delta \varphi) \mathcal{T}_0 \mathbf{f}_0 - \mu \Delta \mathbb{B}[(\nabla \varphi) \cdot \mathcal{T}_0 \mathbf{f}_0] - (\nabla \varphi) P_0 \mathbf{f}_0.$$

Moreover, by Lemma 5.4, we have

$$\begin{aligned} & \sup_{x \in \Omega} (1 + |x|) |\mathbf{u}(x)| + \sup_{x \in \Omega} (1 + |x|)^2 |\nabla \mathbf{u}(x)| + \sup_{x \in \Omega} (1 + |x|)^2 |\mathbf{p}(x)| + \|\mathbf{u}\|_{H_q^2(\Omega)} + \|\mathbf{p}\|_{H_q^1(\Omega)} \\ & \leq C(\langle \operatorname{div} \mathbf{F} \rangle_3 + \langle \mathbf{F} \rangle_2 + \|\mathbf{g}\|_{L_q(\Omega)}). \end{aligned}$$

Notice that

$$\|\mathcal{R}_2 \mathbf{f}\|_{L_q(\Omega)} \leq C(\langle \operatorname{div} \mathbf{F} \rangle_3 + \langle \mathbf{F} \rangle_2 + \|\mathbf{g}\|_{L_q(\Omega)})$$

and $\operatorname{supp} \mathcal{R}_2 \mathbf{f} \subset B_{3b}$. Thus, if we define operators \mathcal{S}_0 and \mathcal{P}_0 acting on \mathbf{f} by setting

$$\begin{aligned} \mathcal{S}_0 \mathbf{f} &= (1 - \varphi) \mathcal{T}_0 \mathbf{f}_0 + \mathbb{B}[(\nabla \varphi) \cdot \mathcal{T}_0 \mathbf{f}_0] + \mathcal{U}_0(\varphi \mathbf{f} + \mathcal{R}_2 \mathbf{f}), \\ \mathcal{P}_0 \mathbf{f} &= (1 - \varphi) \mathcal{P}_0 \mathbf{f}_0 + \mathcal{Q}_0(\varphi \mathbf{f} + \mathcal{R}_2 \mathbf{f}), \end{aligned}$$

then, $\mathbf{u} = \mathcal{S}_0 \mathbf{f}$ and $\mathbf{p} = \mathcal{P}_0 \mathbf{f}$ satisfy (5.6), and combining the estimates above gives that

$$\begin{aligned} & \sup_{x \in \Omega} (1 + |x|) |(\mathcal{S}_0 \mathbf{f})(x)| + \sup_{x \in \Omega} (1 + |x|)^2 |(\nabla \mathcal{S}_0 \mathbf{f})(x)| + \sup_{x \in \Omega} (1 + |x|)^2 |(\mathcal{P}_0 \mathbf{f})(x)| \\ & + \|\mathcal{S}_0 \mathbf{f}\|_{H_q^2(\Omega)} + \|\mathcal{P}_0 \mathbf{f}\|_{H_q^1(\Omega)} \leq C(\langle \operatorname{div} \mathbf{F} \rangle_3 + \langle \mathbf{F} \rangle_2 + \|\mathbf{g}\|_{L_q(\Omega)}), \end{aligned}$$

which completes the proof of Theorem 5.3. □

Proof of existence of $(I + \mathcal{R}_1)^{-1}$. In what follows, we shall prove (5.19). In view of (5.16) and since $\nabla(P_0 \mathbf{f}_0 - \mathcal{B}_0 \mathbf{f}_b) \in L_q(\Omega_{4b})^3$, we have $(P_0 \mathbf{f}_0 - \mathcal{B}_0 \mathbf{f}_b) \in H_q^1(\Omega_{4b})$. Thus, $\mathcal{R}_1 \mathbf{f} \in H_q^1(\Omega)^3$ and $\operatorname{supp} \mathcal{R}_1 \mathbf{f} \subset D_{2b,3b}$, where $D_{2b,3b} = \{x \in \mathbb{R}^3 \mid 2b \leq |x| \leq 3b\}$. Thus, by Rellich's compactness theorem, \mathcal{R}_1 is a compact operator on $L_{q,3b}(\Omega)^3$. Let $\operatorname{Ker}(I + \mathcal{R}_1) = \{\mathbf{f} \in L_{q,3b}(\Omega)^3 \mid (I + \mathcal{R}_1) \mathbf{f} = 0\}$. By Fredholm's alternative principle, if $\operatorname{Ker}(I + \mathcal{R}_1) = \{0\}$, then $I + \mathcal{R}_1$ is invertible, and therefore we have (5.19). To verify this, we choose $\mathbf{f} \in \operatorname{Ker}(I + \mathcal{R}_1)$ arbitrarily, and we shall show that $\mathbf{f} = 0$. Since $(I + \mathcal{R}_1) \mathbf{f} = 0$, we have $\mathbf{f} = -\mathcal{R}_1 \mathbf{f} \in H_q^1(\Omega)$ and $\operatorname{supp} \mathbf{f} \subset D_{2b,3b}$. Let $\mathbf{u} = \mathcal{V}_0 \mathbf{f}$ and $\mathbf{p} = \mathcal{W}_0 \mathbf{f}$. Then by (5.18) we see that

$$-\mu \Delta \mathbf{u} + \nabla \mathbf{p} = 0, \quad \operatorname{div} \mathbf{u} = 0 \quad \text{in } \Omega, \quad \mathbf{u}|_{\Gamma} = 0. \tag{5.21}$$

Since $3 < q < \infty$ and $\mathbf{u} \in H_q^2(\Omega)^3$ and $\mathbf{p} \in \hat{H}_q^1(\Omega)$, we have $\mathbf{u} \in H_{2,loc}^2(\Omega)$ and $\mathbf{p} \in H_{2,loc}^1(\Omega)$. Let ψ be a $C^\infty(\mathbb{R}^3)$ function which equals 1 for $|x| < 1$ and 0 for $|x| > 2$ and set $\psi_R(x) = \psi(x/R)$ for $R > 4b$. From (5.21) it follows that

$$0 = (-\mu \Delta \mathbf{u} + \nabla \mathbf{p}, \psi_R \mathbf{u}) = \mu(\nabla \mathbf{u}, \psi_R \nabla \mathbf{u}) + \mu(\nabla \mathbf{u}, (\nabla \psi_R) \mathbf{u}) - (\mathbf{p}, (\nabla \psi_R) \cdot \mathbf{u}). \tag{5.22}$$

Using Lemma 5.4 (2), we obtain

$$|\mathbf{u}(x)| \leq C|x|^{-1}, \quad |\nabla \mathbf{u}(x)| \leq C|x|^{-2}, \quad |\mathbf{p}| \leq C|x|^{-2}$$

for $|x| > 4b$, and so we have

$$\begin{aligned} |(\nabla \mathbf{u}, (\nabla \psi_R) \mathbf{u})| &\leq C \|\nabla \psi\|_{L^\infty(\mathbb{R}^3)} R^{-1} \int_{2R \leq |x| \leq 3R} |x|^{-3} dx = O(R^{-1}) \rightarrow 0, \\ |(\mathbf{p}, (\nabla \psi_R) \mathbf{u})| &\leq C \|\nabla \psi\|_{L^\infty(\mathbb{R}^3)} R^{-1} \int_{2R \leq |x| \leq 3R} |x|^{-3} dx = O(R^{-1}) \rightarrow 0 \end{aligned}$$

as $R \rightarrow \infty$, and so taking $R \rightarrow \infty$ in (5.22), we have $\|\nabla \mathbf{u}\|_{L_2(\Omega)} = 0$, which implies that \mathbf{u} is a constant vector. But, $\mathbf{u}|_{\Gamma} = 0$, and so $\mathbf{u} = 0$. Thus, by the first equation of (5.21), $\nabla \mathbf{p} = 0$, which

shows that \mathbf{p} is a constant. But, $\mathbf{p}(x) = O(|x|^{-2})$ as $|x| \rightarrow \infty$, and so $\mathbf{p} = 0$. Therefore, by (5.17) we have

$$\begin{aligned} (1 - \varphi)\mathcal{T}_0\mathbf{f}_0 + \varphi\mathcal{A}_0\mathbf{f}_b + \mathbb{B}[(\nabla\varphi) \cdot (\mathcal{T}_0\mathbf{f}_0 - \mathcal{A}_0\mathbf{f}_b)] &= 0, \\ (1 - \varphi)P_0\mathbf{f}_0 + \varphi\mathcal{B}_0\mathbf{f}_b &= 0 \end{aligned} \quad (5.23)$$

in Ω_{4b} . Since $\mathbb{B}[(\nabla\varphi) \cdot (\mathcal{T}_0\mathbf{f}_0 - \mathcal{A}_0\mathbf{f}_b)]$ vanishes for $x \notin D_{2b,3b}$ and $\varphi(x) = 0$ for $|x| > 3b$ and $1 - \varphi(x) = 0$ for $|x| < 2b$, we have

$$\mathcal{A}_0\mathbf{f}_b = 0, \quad \mathcal{B}_0\mathbf{f}_b = 0 \quad \text{for } |x| < 2b, \quad \mathcal{T}_0\mathbf{f}_0 = 0, \quad P_0\mathbf{f}_0 = 0 \quad \text{for } |x| > 3b. \quad (5.24)$$

Let

$$\mathbf{w}(x) = \begin{cases} (\mathcal{A}_0\mathbf{f}_b)(x) & \text{for } x \in \Omega_{4b}, \\ 0 & \text{for } x \notin \Omega, \end{cases} \quad \mathbf{q}(x) = \begin{cases} (\mathcal{B}_0\mathbf{f}_b)(x) & \text{for } x \in \Omega_{4b}, \\ 0 & \text{for } x \notin \Omega, \end{cases}$$

and then, by (5.15) and (5.24) $\mathbf{w} \in H_q^2(B_{4b})^3$ and $\mathbf{q} \in H_q^1(B_{4b})$, and \mathbf{w} and \mathbf{q} satisfy equations:

$$-\mu\Delta\mathbf{w} + \nabla\mathbf{q} = \mathbf{f}_0, \quad \operatorname{div}\mathbf{w} = 0 \quad \text{in } B_{4b}, \quad \mathbf{w}|_{S_{4b}} = 0. \quad (5.25)$$

On the other hand, by (5.24), we know that $\mathcal{T}_0\mathbf{f}_0$ and $P_0\mathbf{f}_0$ also satisfy equations (5.25), and so the uniqueness of solutions yields that $\mathbf{w} = \mathcal{T}_0\mathbf{f}_0$ and $\nabla(\mathbf{q} - P_0\mathbf{f}_0) = 0$ in B_{4b} . Noting that $\mathbf{q} = \mathcal{B}_0\mathbf{f}_b$, by (5.16) we have $\mathbf{q} = P_0\mathbf{f}_0$. In particular, $(\nabla\varphi) \cdot (\mathcal{T}_0\mathbf{f}_0 - \mathcal{A}_0\mathbf{f}_b) = 0$. Thus, from (5.23) we even have

$$\begin{aligned} 0 &= \mathcal{T}_0\mathbf{f}_0 - \varphi(\mathcal{T}_0\mathbf{f}_0 - \mathcal{A}_0\mathbf{f}_b) = \mathcal{T}_0\mathbf{f}_0, \\ 0 &= P_0\mathbf{f}_0 - \varphi(P_0\mathbf{f}_0 - \mathcal{B}_0\mathbf{f}_b) = P_0\mathbf{f}_0, \end{aligned}$$

which gives that $\mathbf{f} = -\mu\Delta\mathcal{T}_0\mathbf{f}_0 + \nabla P_0\mathbf{f}_0 = 0$ in Ω_{4b} . Thus, we have $\mathbf{f} = 0$. This completes the proof of existence of $(I + \mathcal{R}_1)^{-1}$. \square

5.3 Purely oscillatory solutions to the Stokes problem

In this section we consider the oscillatory part of the linearization (5.4) for $\mathbf{h} = 0$, which is given by

$$\partial_t\mathbf{v}_\perp - \mu\Delta\mathbf{v}_\perp + \nabla\mathbf{p}_\perp = \mathbf{f}_\perp, \quad \operatorname{div}\mathbf{v}_\perp = 0 \quad \text{in } \Omega \times \mathbb{T}, \quad \mathbf{v}_\perp|_\Gamma = 0, \quad (5.26)$$

where the subscript \perp indicates that all functions have vanishing time mean. We shall prove the following theorem.

Theorem 5.5. *Let $1 < p, q < \infty$. Then, for any $\mathbf{f}_\perp \in L_p(\mathbb{T}, L_q(\Omega))$ with $\int_{\mathbb{T}} \mathbf{f}_\perp(\cdot, s) ds = 0$, problem (5.26) admits a unique solution $(\mathbf{v}_\perp, \mathbf{p}_\perp)$ with*

$$\mathbf{v}_\perp \in H_p^1(\mathbb{T}, L_q(\Omega)^3) \cap L_p(\mathbb{T}, H_q^2(\Omega)^3), \quad \mathbf{p}_\perp \in L_p(\mathbb{T}, \hat{H}_q^1(\Omega)), \quad \int_{\mathbb{T}} \mathbf{v}_\perp ds = 0, \quad \int_{\mathbb{T}} \mathbf{p}_\perp ds = 0,$$

possessing the estimate

$$\|\partial_t\mathbf{v}_\perp\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{v}_\perp\|_{L_p(\mathbb{T}, H_q^2(\Omega))} + \|\nabla\mathbf{p}_\perp\|_{L_p(\mathbb{T}, L_q(\Omega))} \leq C\|\mathbf{f}_\perp\|_{L_p(\mathbb{T}, L_q(\Omega))}. \quad (5.27)$$

In order to prove Theorem 5.5, we first consider the corresponding resolvent problem

$$\lambda\mathbf{w} - \mu\Delta\mathbf{w} + \nabla\mathbf{t} = \mathbf{f}, \quad \operatorname{div}\mathbf{w} = 0 \quad \text{in } \Omega, \quad \mathbf{w}|_\Gamma = 0, \quad (5.28)$$

for which we have the results from Theorem 4.2. Observe that, with regard to the time-periodic problem (5.4), we are mainly interested in the resolvent problem (5.28) with $\lambda = ik$ for $k \in \mathbb{Z}$, and Theorem 4.2 gives a framework where this problem is uniquely solvable, but merely for $k \in \mathbb{Z} \setminus \{0\}$. This is the main reason why we only consider the purely oscillatory problem (5.26) in Theorem 5.5. Apart from this, Theorem 5.5 can be proved in the same way as Theorem 4.4.

Proof of Theorem 5.5. Let λ_0 be as in Theorem 4.2, and let $\varphi = \varphi(\sigma)$ be a $C^\infty(\mathbb{R})$ function that equals 1 for $|\sigma| \geq \lambda_0 + 1/2$ and 0 for $|\sigma| \leq \lambda_0 + 1/4$. Using the operator families \mathcal{S} and \mathcal{P} from Theorem 4.2, we set

$$\mathbf{v}_h = \mathcal{F}_\mathbb{T}^{-1}[\mathcal{S}(ik)\varphi(k)\mathcal{F}_\mathbb{T}[\mathbf{H}_\perp](k)], \quad \mathbf{p}_h = \mathcal{F}_\mathbb{T}^{-1}[\mathcal{P}(ik)\varphi(k)\mathcal{F}_\mathbb{T}[\mathbf{H}_\perp](k)].$$

Then \mathbf{v}_h and \mathbf{p}_h satisfy the equations

$$\partial_t \mathbf{v}_h - \mu \Delta \mathbf{v}_h + \nabla \mathbf{p}_h = \mathbf{H}_h, \quad \operatorname{div} \mathbf{v}_h = 0 \quad \text{in } \Omega \times \mathbb{T}, \quad \mathbf{v}_h|_\Gamma = 0,$$

where we have set $\mathbf{H}_h = \mathcal{F}_\mathbb{T}^{-1}[\varphi(k)\mathcal{F}_\mathbb{T}[\mathbf{H}_\perp](k)]$. Moreover, arguing as in the proof of Theorem 3.1, we can use the \mathcal{R} -bounds from Theorem 4.2 and employ Corollary 2.3 to deduce

$$\begin{aligned} \|\partial_t \mathbf{v}_h\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{v}_h\|_{L_p(\mathbb{T}, H_q^2(\Omega))} + \|\nabla \mathbf{p}_h\|_{L_p(\mathbb{T}, L_q(\Omega))} \\ \leq C \|\mathbf{H}_h\|_{L_p(\mathbb{T}, L_q(\Omega))} \leq C \|\mathbf{H}_\perp\|_{L_p(\mathbb{T}, L_q(\Omega))}. \end{aligned} \quad (5.29)$$

Now, in view of Theorem 4.2, we set

$$\mathbf{v}_\perp(t) = \mathbf{v}_h(t) + \sum_{0 < |k| \leq \lambda_0} e^{ikt} \mathcal{S}(ik) \mathcal{F}_\mathbb{T}[\mathbf{H}_\perp](k), \quad \mathbf{p}_\perp(t) = \mathbf{p}_h(t) + \sum_{0 < |k| \leq \lambda_0} e^{ikt} \mathcal{P}(ik) \mathcal{F}_\mathbb{T}[\mathbf{H}_\perp](k).$$

Then, \mathbf{v}_\perp and \mathbf{p}_\perp satisfy equations (5.26), and from (4.14) and (5.29) we conclude estimate (5.27). Thus, we have shown the existence part of Theorem 5.5. The uniqueness statement follows exactly as in the proof of Theorem 4.4 noting that $\mathcal{F}_\mathbb{T}[\mathbf{v}_\perp](0) = 0$ and $\mathcal{F}_\mathbb{T}[\mathbf{p}_\perp](0) = 0$ by assumption. \square

Next we examine the pointwise decay of the solution $(\mathbf{v}_\perp, \mathbf{p}_\perp)$. More precisely, we show decay properties of $\|\mathbf{v}_\perp(x, \cdot)\|_{L_p(\mathbb{T})}$ with respect to the x -variable, as stated in the following theorem.

Theorem 5.6. *In the situation of Theorem 5.5, let $3 < q < \infty$ and $\ell \in (0, 3]$ such that $\mathbf{f}_\perp = \operatorname{div} \mathbf{F}_\perp + \mathbf{g}_\perp$ with*

$$\begin{aligned} \int_{\mathbb{T}} \mathbf{F}_\perp(x, t) dt = 0, \quad \langle \mathbf{F}_\perp \rangle_{p, \ell} + \langle \operatorname{div} \mathbf{F}_\perp \rangle_{p, \ell+1} < \infty, \\ \int_{\mathbb{T}} \mathbf{g}_\perp(x, t) dt = 0, \quad \mathbf{g}_\perp \in L_p(\mathbb{T}, L_{q, 3b}(\Omega)). \end{aligned} \quad (5.30)$$

Then, \mathbf{v}_\perp has the following asymptotics:

$$\langle \mathbf{v}_\perp \rangle_{p, \ell} + \langle \nabla \mathbf{v}_\perp \rangle_{p, \ell+1} \leq C (\langle \operatorname{div} \mathbf{F}_\perp \rangle_{p, \ell+1} + \langle \mathbf{F}_\perp \rangle_{p, \ell} + \|\mathbf{g}_\perp\|_{L_p(\mathbb{T}, L_q(\Omega))}) \quad (5.31)$$

with some constant $C > 0$.

Remark 5.7. Since $3 < q < \infty$, we have $\|\operatorname{div} \mathbf{F}_\perp\|_{L_p(\mathbb{T}, L_q(\Omega))} \leq C_q \langle \operatorname{div} \mathbf{F}_\perp \rangle_{p, \ell+1}$ and so

$$\|\mathbf{f}_\perp\|_{L_p(\mathbb{T}, L_q(\Omega))} \leq C_q (\langle \operatorname{div} \mathbf{F}_\perp \rangle_{p, \ell+1} + \|\mathbf{g}_\perp\|_{L_p(\mathbb{T}, L_q(\Omega))}). \quad (5.32)$$

Therefore, Theorem 5.5 really shows existence for \mathbf{f}_\perp as in Theorem 5.6.

To prove (5.31), we use the following theorem due to EITER and KYED [17], which collects properties of the velocity fundamental solution Γ_\perp to (5.26), which is a tensor field Γ_\perp such that $\mathbf{v}_\perp := \Gamma_\perp * \mathbf{H}_\perp$ is formally a solution to (5.26) for $\Omega = \mathbb{R}^3$.

Theorem 5.8. *Let*

$$\Gamma_{\perp} = \mathcal{F}_{\mathbb{R}^3 \times \mathbb{T}}^{-1} \left[\frac{1 - \delta_{\mathbb{Z}}}{|\mu|\xi|^2 + ik} \left(\mathbf{I} - \frac{\xi \otimes \xi}{|\xi|^2} \right) \right]. \quad (5.33)$$

Then, it holds $\Gamma_{\perp} \in L_q(\mathbb{R}^3 \times \mathbb{T})^{3 \times 3}$ for $q \in (1, 5/3)$, and $\partial_j \Gamma_{\perp} \in L_q(\mathbb{R}^3 \times \mathbb{T})$ for $q \in (1, 5/4)$, $j = 1, 2, 3$. Moreover, for any $\alpha \in \mathbb{N}_0^3$, $\delta > 0$ and $r \in [1, \infty)$, there exists a constant $C_{\alpha, \delta} > 0$ such that

$$\forall |x| \geq \delta : \quad \|\partial_x^{\alpha} \Gamma_{\perp}(x, \cdot)\|_{L_r(\mathbb{T})} \leq \frac{C_{\alpha, \delta}}{|x|^{3+|\alpha|}}.$$

Remark 5.9. This theorem holds for any dimension $N \geq 2$ replacing $5/3$, $5/4$ and $3 + |\alpha|$ with $(N + 2)/N$, $(N + 2)/(N + 1)$ and $N + |\alpha|$, respectively.

Proof of Theorem 5.6. Since we assume that $3 < q < \infty$, by Sobolev's inequality, we have

$$\sup_{|x| \leq 4b} \|\mathbf{v}_{\perp}(\cdot, x)\|_{L_p(\mathbb{T})} + \sup_{|x| \leq 4b} \|(\nabla \mathbf{v}_{\perp})(\cdot, x)\|_{L_p(\mathbb{T})} \leq C \|\mathbf{v}_{\perp}\|_{L_p(\mathbb{T}, H_q^2(\Omega))} \leq C \|\mathbf{H}_{\perp}\|_{L_p(\mathbb{T}, L_q(\Omega))}.$$

It thus remains to estimate \mathbf{v}_{\perp} for $|x| > 4b$. To this end, recall the operator families \mathcal{S} and \mathcal{P} given in Theorem 4.2. As seen in the proof of Theorem 5.5, we have $\mathbf{v}_{\perp} = \mathcal{F}_{\mathbb{T}}^{-1}[\mathcal{S}(ik)\mathcal{F}_{\mathbb{T}}[\mathbf{H}_{\perp}]](k)$ and $\mathbf{p}_{\perp} = \mathcal{F}_{\mathbb{T}}^{-1}[\mathcal{P}(ik)\mathcal{F}_{\mathbb{T}}[\mathbf{H}_{\perp}]](k)$. We shall first give a representation formula of $\mathcal{S}(ik)$ for $k \in \mathbb{Z} \setminus \{0\}$ for $|x| > 3b$, which will be used to investigate the asymptotic behavior of \mathbf{v}_{\perp} for $|x| > 3b$. Notice that $\mathcal{S}(ik) \in \mathcal{L}(L_q(\Omega)^3, H_q^2(\Omega)^3)$ and $\mathcal{P}(ik) \in \mathcal{L}(L_q(\Omega)^3, \hat{H}_q^1(\Omega))$ satisfy the estimate

$$\|\mathcal{S}(ik)\mathbf{f}\|_{H_q^2(\Omega)} + \|\nabla \mathcal{P}(ik)\mathbf{f}\|_{L_q(\Omega)} \leq C \|\mathbf{f}\|_{L_q(\Omega)} \quad (5.34)$$

for $\mathbf{f} \in L_q(\Omega)^3$, where C depends solely on q and Ω . Moreover, the functions $\mathbf{u} = \mathcal{F}_{\mathbb{T}}[\mathbf{v}_{\perp}](k) = \mathcal{S}(ik)\mathcal{F}_{\mathbb{T}}[\mathbf{H}_{\perp}](k)$ and $\mathbf{q} = \mathcal{F}_{\mathbb{T}}[\mathbf{p}_{\perp}](k) = \mathcal{P}(ik)\mathcal{F}_{\mathbb{T}}[\mathbf{H}_{\perp}](k)$ satisfy the equations

$$ik\mathbf{u} - \mu\Delta\mathbf{u} + \nabla\mathbf{q} = \mathbf{f}_k, \quad \operatorname{div}\mathbf{u} = 0 \quad \text{in } \Omega, \quad \mathbf{u}|_{\Gamma} = 0, \quad (5.35)$$

where $\mathbf{f}_k = \mathcal{F}_{\mathbb{T}}[\mathbf{f}_{\perp}](k)$. Let φ be a function in $C_0^{\infty}(\mathbb{R}^3)$ that equals 1 for $|x| < 2b$ and 0 for $|x| > 3b$. Let

$$\mathbf{w} = (1 - \varphi)\mathcal{S}(ik)\mathbf{f}_k + \mathbb{B}[(\nabla\varphi) \cdot \mathcal{S}(ik)\mathbf{f}_k], \quad \mathbf{r} = (1 - \varphi)\mathcal{P}(ik)\mathbf{f}_k. \quad (5.36)$$

Then $\mathbf{w} \in H_q^2(\mathbb{R}^3)^3$ and $\mathbf{r} \in \hat{H}_q^1(\mathbb{R}^3)$. Moreover, by (5.35) \mathbf{w} and \mathbf{r} satisfy the equations

$$ik\mathbf{w} - \mu\Delta\mathbf{w} + \nabla\mathbf{r} = (1 - \varphi)\mathbf{f}_k + \mathcal{R}_3(ik)\mathbf{f}_k, \quad \operatorname{div}\mathbf{w} = 0 \quad \text{in } \mathbb{R}^3,$$

where we have set

$$\begin{aligned} \mathcal{R}_3(\lambda)\mathbf{f} &= 2\mu(\nabla\varphi) \cdot \nabla\mathcal{S}(\lambda)\mathbf{f} + \mu(\Delta\varphi)\mathcal{S}(\lambda)\mathbf{f} \\ &\quad - (\nabla\varphi)\mathcal{P}(\lambda)\mathbf{f} + (\lambda - \mu\Delta)\mathbb{B}[(\nabla\varphi) \cdot \mathcal{S}(\lambda)\mathbf{f}]. \end{aligned} \quad (5.37)$$

By the uniqueness of solutions to the Stokes resolvent problem in \mathbb{R}^3 , we have $\mathbf{w} = \mathcal{T}(ik)((1 - \varphi)\mathbf{f}_k + \mathcal{R}_3(ik)\mathbf{f}_k)$, where

$$\mathcal{T}(\lambda)\mathbf{f} = \mathcal{F}_{\mathbb{R}^3}^{-1} \left[\frac{1}{|\mu|\xi|^2 + \lambda} \left(\mathbf{I} - \frac{\xi \otimes \xi}{|\xi|^2} \right) \mathcal{F}_{\mathbb{R}^3}[\mathbf{f}] \right]. \quad (5.38)$$

Since $1 - \varphi = 1$ and $\mathbb{B}[(\nabla\varphi) \cdot \mathcal{S}(ik)\mathbf{f}_k] = 0$ for $|x| > 4b$, by (5.36) we thus have

$$\mathcal{S}(ik)\mathbf{f}_k = \mathcal{T}(ik)((1 - \varphi)\mathbf{f}_k) + \mathcal{T}(ik)(\mathcal{R}_3(ik)\mathbf{f}_k) \quad (|x| > 4b) \quad (5.39)$$

for any $k \in \mathbb{Z} \setminus \{0\}$. Thus, we have

$$\begin{aligned} \mathbf{v}_\perp &= \mathcal{F}_\mathbb{T}^{-1}[(1 - \delta_\mathbb{Z}(k))\mathcal{S}(ik)\mathcal{F}_\mathbb{T}[\mathbf{H}_\perp](k)] \\ &= \mathcal{F}_\mathbb{T}^{-1}[(1 - \delta_\mathbb{Z}(k))\mathcal{T}(ik)\mathcal{F}_\mathbb{T}[(1 - \varphi)\mathbf{H}_\perp](k)] \\ &\quad + \mathcal{F}_\mathbb{T}^{-1}[(1 - \delta_\mathbb{Z}(k))\mathcal{T}(ik)(\mathcal{R}_3(ik)\mathcal{F}_\mathbb{T}[\mathbf{H}_\perp](k))] \end{aligned} \quad (5.40)$$

for $|x| > 4b$. Moreover, from Theorem 4.2 we conclude

$$\mathcal{R}_{\mathcal{L}(\mathbb{L}_q(\Omega)^3, \mathbb{H}_q^1(\mathbb{R}^3)^3)}(\{(\lambda\partial_\lambda)^\ell \mathcal{R}_3(\lambda) \mid \lambda \in \mathbb{R} \setminus [-\lambda_0, \lambda_0]\}) \leq r_0 \quad (\ell = 0, 1), \quad (5.41)$$

$$\|\mathcal{R}_3(ik)\mathbf{f}_k\|_{\mathbb{H}_q^1(\mathbb{R}^3)} \leq r_0 \|\mathbf{f}_k\|_{\mathbb{L}_q(\Omega)} \quad (5.42)$$

for any $k \in \mathbb{Z} \setminus \{0\}$ with some constant r_0 . In particular, we define $\mathcal{R}_4\mathbf{H}_\perp$ by setting $\mathcal{R}_4\mathbf{H}_\perp = \mathcal{F}_\mathbb{T}^{-1}[(1 - \delta_\mathbb{Z}(k))\mathcal{R}_3(ik)\mathbf{f}_k]$. Then, employing Corollary 2.3 in the same way as in the proof of Theorem 5.5, we see that

$$\begin{aligned} \text{supp } \mathcal{R}_4\mathbf{H}_\perp &\subset D_{2b,3b} := \{(x, t) \in \mathbb{R}^3 \times \mathbb{R} \mid 2b < |x| < 3b\}, \\ \|\mathcal{R}_4\mathbf{H}_\perp\|_{\mathbb{L}_p(\mathbb{T}, \mathbb{L}_q(\Omega))} &\leq C \|\mathbf{H}_\perp\|_{\mathbb{L}_p(\mathbb{T}, \mathbb{L}_q(\Omega))}. \end{aligned} \quad (5.43)$$

Recalling that $\mathbf{H}_\perp = \text{div } \mathbf{F}_\perp + \mathbf{g}_\perp$, we set $\mathbf{G} = (1 - \varphi)\mathbf{F}_\perp$ and $\mathbf{h} = (\nabla\varphi)\mathbf{F}_\perp + (1 - \varphi)\mathbf{g}_\perp + \mathcal{R}_4\mathbf{H}_\perp$. In virtue of (5.33), (5.38) and (5.40), we then have

$$\begin{aligned} \mathbf{v}_\perp &= \Gamma_\perp * (\text{div } \mathbf{G}) + \Gamma_\perp * \mathbf{h} \\ &= \int_{\mathbb{T}} \int_{\mathbb{R}^3} \Gamma_\perp(y, s) (\text{div } \mathbf{G})(x - y, t - s) dy ds + \int_{\mathbb{T}} \int_{\mathbb{R}^3} \Gamma_\perp(y, s) \mathbf{h}(x - y, t - s) dy ds. \end{aligned} \quad (5.44)$$

for $|x| > 4b$. Set $\mathbf{v}_1 = \Gamma_\perp * (\text{div } \mathbf{G})$ and $\mathbf{v}_2 = \Gamma_\perp * \mathbf{h}$. By the divergence theorem of Gauß, we write

$$\begin{aligned} \mathbf{v}_1(x, t) &= \nabla \Gamma_\perp * \mathbf{G}(x, t) \\ &= \int_{\mathbb{T}} \int_{|y| \leq 1} \nabla \Gamma_\perp(y, s) \mathbf{G}(x - y, t - s) dy ds + \int_{\mathbb{T}} \int_{1 \leq |y| \leq |x|/2} \nabla \Gamma_\perp(y, s) \mathbf{G}(x - y, t - s) dy ds \\ &\quad + \int_{\mathbb{T}} \int_{|x|/2 \leq |y| \leq 2|x|} \nabla \Gamma_\perp(y, s) \mathbf{G}(x - y, t - s) dy ds \\ &\quad + \int_{\mathbb{T}} \int_{|y| \geq 2|x|} \nabla \Gamma_\perp(y, s) \mathbf{G}(x - y, t - s) dy ds. \end{aligned}$$

Let r_0 and r_1 be exponents such that $p < r_0 < \infty$, $r_1 \in (1, 5/4)$ and $1 + 1/r_0 = 1/r_1 + 1/p$. Then, we have Young's inequality

$$\|f * g\|_{\mathbb{L}_{r_0}(\mathbb{T})} \leq \|f\|_{\mathbb{L}_{r_1}(\mathbb{T})} \|g\|_{\mathbb{L}_p(\mathbb{T})} \quad (5.45)$$

for $f * g(t) = \int_{\mathbb{T}} f(s)g(t - s) ds$. Setting $\gamma_\ell = \langle \mathbf{G} \rangle_{p, \ell}$, from Theorem 5.8 we thus conclude

$$\begin{aligned} \|\mathbf{v}_1(x, \cdot)\|_{\mathbb{L}_{r_0}(\mathbb{T})} &\leq \gamma_\ell \|\nabla \Gamma_\perp\|_{\mathbb{L}_{r_1}(B_1 \times \mathbb{T})} (1 + |x|)^{-\ell} + C_1 \gamma_\ell \int_{1 \leq |y| \leq |x|/2} |y|^{-4} dy (1 + |x|)^{-\ell} \\ &\quad + C_1 \gamma_\ell (|x|/2)^{-4} \int_{|z| \leq 3|x|} (1 + |z|)^{-\ell} dz + C_1 \gamma_\ell \int_{|y| \geq 2|x|} |y|^{-4-\ell} dy. \end{aligned}$$

Noting that $p \leq r_0$ and $\gamma_\ell \leq \langle \mathbf{F}_\perp \rangle_{p, \ell}$, we infer

$$\|\mathbf{v}_1(x, \cdot)\|_{\mathbb{L}_p(\mathbb{T})} \leq C_b |x|^{-\min\{\ell, 4\}} \langle \mathbf{F}_\perp \rangle_{p, \ell} \quad \text{for } |x| \geq 4b.$$

Analogously, we write

$$\begin{aligned}\nabla \mathbf{v}_1(x, t) &= \int_{\mathbb{T}} \int_{|y| \leq 1} \nabla \Gamma_{\perp}(y, s) (\operatorname{div} \mathbf{G})(x - y, t - s) dy ds \\ &+ \int_{\mathbb{T}} \int_{1 \leq |y| \leq |x|/2} \nabla \Gamma_{\perp}(y, s) (\operatorname{div} \mathbf{G})(x - y, t - s) dy ds \\ &+ \int_{\mathbb{T}} \int_{|x|/2 \leq |y| \leq 2|x|} \nabla \Gamma_{\perp}(y, s) (\operatorname{div} \mathbf{G})(x - y, t - s) dy ds \\ &+ \int_{\mathbb{T}} \int_{|y| \geq 2|x|} \nabla^{\ell} \Gamma_{\perp}(y, s) (\operatorname{div} \mathbf{G})(x - y, t - s) dy ds.\end{aligned}$$

Setting $\gamma_{\ell+1} = \langle \operatorname{div} \mathbf{G} \rangle_{p, \ell+1}$, by Theorem 5.8 and (5.45) we have

$$\begin{aligned}\|\nabla \mathbf{v}_1(x, \cdot)\|_{L_{r_0}(\mathbb{T})} &\leq \gamma_{\ell+1} \|\nabla \Gamma_{\ell}\|_{L_{r_1}(B_1 \times \mathbb{T})} (1 + |x|)^{-\ell-1} + C_1 \gamma_{\ell+1} \int_{1 \leq |y| \leq |x|/2} |y|^{-4} dy (1 + |x|)^{-\ell-1} \\ &+ C_1 \gamma_{\ell+1} (|x|/2)^{-4} \int_{|z| \leq 3|x|} (1 + |z|)^{-\ell-1} dz + C_1 \gamma_{\ell+1} \int_{|y| \geq 2|x|} |y|^{-5-\ell} dy.\end{aligned}$$

Since we have

$$\begin{aligned}\langle \operatorname{div} \mathbf{G} \rangle_{p, \ell+1} &\leq \langle \operatorname{div} \mathbf{F}_{\perp} \rangle_{p, \ell+1} + \langle (\nabla \varphi) \mathbf{F}_{\perp} \rangle_{p, \ell+1} \\ &\leq \langle \operatorname{div} \mathbf{F} \rangle_{p, \ell+1} + \|\nabla \varphi\|_{L_{\infty}(\mathbb{R}^3)} 3b \langle \mathbf{F} \rangle_{p, \ell}\end{aligned}$$

and $p \leq r_0$, we thus obtain

$$\|\nabla \mathbf{v}_1(x, \cdot)\|_{L_p(\mathbb{T})} \leq C_b |x|^{-\min\{\ell+1, 4\}} (\langle \operatorname{div} \mathbf{F}_{\perp} \rangle_{p, \ell+1} + \langle \mathbf{F}_{\perp} \rangle_{p, \ell}) \quad \text{for } |x| \geq 4b.$$

Finally, we use that $\mathbf{h}(y, s)$ vanishes for $|y| \geq 3b$. For $m = 0, 1$ we thus have

$$\nabla^m \mathbf{v}_2(x, t) = \int_{\mathbb{T}} \int_{|x-y| \leq 3b} \nabla^m \Gamma_{\perp}(y, s) \mathbf{h}(x - y, t - s) dy ds$$

Since $|x| \geq 4b$ and $|x - y| \leq 3b$ implies $|y| \geq |x|/4 \geq b$, by Theorem 5.8 and (5.45), we deduce

$$\begin{aligned}\|\nabla^m \mathbf{v}_2(x, \cdot)\|_{L_p(\mathbb{T})} &\leq \int_{|x-y| \leq 3b} \|\nabla^m \Gamma_{\perp}(y, \cdot)\|_{L_p(\mathbb{T})} \|\mathbf{h}(x - y, \cdot)\|_{L_1(\mathbb{T})} dy \\ &\leq C_m |x|^{-3-m} \|\mathbf{h}\|_{L_1(B_{3b} \times \mathbb{T})}.\end{aligned}$$

Noting (5.43), we can estimate the last term as

$$\begin{aligned}\|\mathbf{h}\|_{L_1(B_{3b} \times \mathbb{T})} &\leq C \|\mathbf{h}\|_{L_p(\mathbb{T}, L_q(B_{3b}))} \leq C (\langle \mathbf{F}_{\perp} \rangle_{p, \ell} + \|\mathbf{g}_{\perp}\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathcal{R}_4 \mathbf{H}_{\perp}\|_{L_p(\mathbb{T}, L_q(\Omega))}) \\ &\leq C (\langle \mathbf{F}_{\perp} \rangle_{p, \ell} + \|\mathbf{g}_{\perp}\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{H}_{\perp}\|_{L_p(\mathbb{T}, L_q(\Omega))}),\end{aligned}$$

For $|x| \geq 4b$ we now conclude

$$\|\nabla^m \mathbf{v}_2(x, \cdot)\|_{L_p(\mathbb{T})} \leq C |x|^{-3-m} (\langle \mathbf{F}_{\perp} \rangle_{p, \ell} + \|\mathbf{g}_{\perp}\|_{L_p(\mathbb{T}, L_q(\Omega))} + \langle \operatorname{div} \mathbf{F}_{\perp} \rangle_{p, \ell+1})$$

in virtue of estimate (5.32). Since $\mathbf{v} = \mathbf{v}_1 + \mathbf{v}_2$ for $|x| \geq 4b$, this completes the proof of Theorem 5.6. \square

5.4 Existence of periodic solutions

The linear theory from Theorem 5.2 is now a direct consequence of Theorem 5.3 and Theorem 5.6 if $\mathbf{h} = 0$. For the case of non-zero boundary data \mathbf{h} we proceed similarly to the proof of Theorem 4.4.

Proof of Theorem 5.2. At first consider the case $\mathbf{h} = 0$. Let $(\mathbf{v}_S, \mathbf{p}_S) = (\mathbf{u}, \mathbf{p})$ be the unique solution to (5.6) with $\mathbf{F} = \mathbf{G}_S$ and $\mathbf{g} = \mathbf{g}_S$, which exists due to Theorem 5.3, and let $(\mathbf{v}_\perp, \mathbf{p}_\perp)$ be the unique solution to (5.26), which exists due to Theorem 5.5 and Theorem 5.6. Then $\mathbf{v} = \mathbf{v}_S + \mathbf{v}_\perp$ and $\mathbf{p} = \mathbf{p}_S + \mathbf{p}_\perp$ defines a solution (\mathbf{v}, \mathbf{p}) to (5.4) with $\mathbf{h} = 0$, and (5.5) follows from (5.7), (5.27), (5.31) and (5.32).

To show existence for arbitrary $\mathbf{h} \in H_p^1(\mathbb{T}, L_q(\Omega)^N) \cap L_p(\mathbb{T}, H_q^2(\Omega)^N)$, we fix $\lambda_1 > \lambda_0$ with λ_0 from Theorem 4.3 and define

$$\begin{aligned} \mathbf{v}_1 &= \mathcal{F}_\mathbb{T}^{-1}[\mathcal{S}(ik + \lambda_1)(0, (ik + \lambda_1)\tilde{\mathbf{h}}_k, (ik + \lambda_1)^{1/2}\tilde{\mathbf{h}}_k, \tilde{\mathbf{h}}_k)], \\ \mathbf{p}_1 &= \mathcal{F}_\mathbb{T}^{-1}[\mathcal{P}(ik + \lambda_1)(0, (ik + \lambda_1)\tilde{\mathbf{h}}_k, (ik + \lambda_1)^{1/2}\tilde{\mathbf{h}}_k, \tilde{\mathbf{h}}_k)], \end{aligned}$$

where \mathcal{S} and \mathcal{P} are the \mathcal{R} -bounded solution operators from Theorem 4.3, and $\tilde{\mathbf{h}}_k = \mathcal{F}[\mathbf{h}](k)$. Then $(\mathbf{v}_1, \mathbf{p}_1)$ is a solution to the auxiliary problem

$$\partial_t \mathbf{v}_1 + \lambda_1 \mathbf{v}_1 - \mu \Delta \mathbf{v}_1 + \nabla \mathbf{p}_1 = 0, \quad \operatorname{div} \mathbf{v}_1 = 0 \quad \text{in } \Omega \times \mathbb{T}, \quad \mathbf{v}_1|_\Gamma = \mathbf{h}|_\Gamma.$$

Following the proof of Theorem 4.4 and invoking Corollary 2.3, we further conclude

$$\mathbf{v}_1 \in H_p^1(\mathbb{T}, L_q(\Omega)^N) \cap L_p(\mathbb{T}, H_q^2(\Omega)^N), \quad \mathbf{p}_1 \in L_p(\mathbb{T}, \hat{H}_q^1(\Omega))$$

and the estimate

$$\|\partial_t \mathbf{v}_1\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{v}_1\|_{L_p(\mathbb{T}, H_q^2(\Omega))} + \|\nabla \mathbf{p}_1\|_{L_p(\mathbb{T}, L_q(\Omega))} \leq C(\|\partial_t \mathbf{h}\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{h}\|_{L_p(\mathbb{T}, H_q^2(\Omega))}).$$

Now let $\varphi \in C_0^\infty(\Omega)$ such that $\varphi \equiv 1$ in B_{2b} and $\varphi \equiv 0$ in $\mathbb{R}^3 \setminus B_{3b}$. Let $D_{2b,3b} = \{x \in \mathbb{R}^N \mid 2b < |x| < 3b\}$ and

$$H_{q,0,a}^2(D_{2b,3b}) = \{f \in H_q^2(D_{2b,3b}) \mid \partial_x^\alpha f|_{S_L} = 0 \text{ for } L = 2b, 3b \text{ and } |\alpha| \leq 1, \int_{D_{2b,3b}} f(x) dx = 0\}.$$

According to [59, Lemma 5], we know that $(\nabla \varphi) \cdot \mathbf{v}_1(t) \in H_{q,0,a}^2(D_{2b,3b})$ for a.a. $t \in \mathbb{R}$, and setting $\mathbf{w}_1 = \varphi \mathbf{v}_1 - \mathbb{B}[(\nabla \varphi) \cdot \mathbf{v}_1]$, we see that

$$\begin{aligned} \mathbf{w}_1 &\in H_p^1(\mathbb{T}, L_q(\Omega)^3) \cap L_p(\mathbb{T}, H_q^2(\Omega)^3), \quad \operatorname{supp} \mathbf{w}_1 \subset B_{3b} \cap \Omega, \quad \operatorname{div} \mathbf{w}_1 = 0, \quad \mathbf{w}_1|_\Gamma = \mathbf{h}, \\ \|\partial_t \mathbf{w}_1\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{w}_1\|_{L_p(\mathbb{T}, H_q^2(\Omega))} &\leq C(\|\partial_t \mathbf{h}\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{h}\|_{L_p(\mathbb{T}, H_q^2(\Omega))}). \end{aligned} \tag{5.46}$$

Now let $(\mathbf{w}_2, \mathbf{q}_2)$ be the unique solution to

$$\partial_t \mathbf{w}_2 - \Delta \mathbf{w}_2 + \nabla \mathbf{q}_2 = \mathbf{f} - \partial_t \mathbf{w}_1 - \Delta \mathbf{w}_1, \quad \operatorname{div} \mathbf{w}_2 = 0 \quad \text{in } \Omega \times \mathbb{T}, \quad \mathbf{w}_2|_\Gamma = 0,$$

which exists due to the first part of the proof. Note that \mathbf{w}_1 vanishes in $\mathbb{R}^3 \setminus B_{3b}$, so that $(\mathbf{v}, \mathbf{p}) = (\mathbf{w}_1 + \mathbf{w}_2, \mathbf{q}_2)$ is a solution to (5.4), and estimate (5.5) follows from the corresponding estimate for \mathbf{w}_2 and the properties listed in (5.46).

The uniqueness assertion follows by decomposing a solution (\mathbf{v}, \mathbf{p}) into a stationary and an oscillatory part by means of (5.2) and employing the uniqueness statements from Theorem 5.3 and Theorem 5.5. \square

Proof of Theorem 5.1. We employ Banach's contraction mapping principle. Define

$$\mathcal{I}_\varepsilon = \{(\mathbf{v}, \mathbf{q}) \mid \mathbf{v} = \mathbf{v}_\perp + \mathbf{v}_S, \mathbf{q} = \mathbf{q}_\perp + \mathbf{q}_S, \mathbf{v}_\perp \in \mathbf{H}_p^1(\mathbb{T}, L_q(\Omega)^3) \cap L_p(\mathbb{T}, \mathbf{H}_q^2(\Omega)^3), \\ \mathbf{v}_S \in \mathbf{H}_q^2(\Omega)^3, \operatorname{div} \mathbf{v} = 0, \mathbf{q}_\perp \in L_p(\mathbb{T}, \hat{\mathbf{H}}_q^1(\Omega)), \mathbf{q}_S \in \mathbf{H}_q^1(\Omega), \|(\mathbf{v}, \mathbf{q})\|_{\mathcal{I}_\varepsilon} \leq \varepsilon\},$$

where we set

$$\|(\mathbf{v}, \mathbf{q})\|_{\mathcal{I}_\varepsilon} = \|\partial_t \mathbf{v}_\perp\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{v}_\perp\|_{L_p(\mathbb{T}, \mathbf{H}_q^2(\Omega))} + \|\mathbf{v}_S\|_{\mathbf{H}_q^2(\Omega)} + \|\nabla \mathbf{q}_\perp\|_{L_p(\mathbb{T}, L_q(\Omega))} \\ + \|\mathbf{q}_S\|_{\mathbf{H}_q^1(\Omega)} + \langle \mathbf{v}_\perp \rangle_{p,1} + \langle \nabla \mathbf{v}_\perp \rangle_{p,2} + \langle \mathbf{v}_S \rangle_1 + \langle \nabla \mathbf{v}_S \rangle_2.$$

For $(\mathbf{v}, \mathbf{q}) \in \mathcal{I}_\varepsilon$, let (\mathbf{u}, \mathbf{p}) be the solution of the linear system of equations

$$\partial_t \mathbf{u} - \mu \Delta \mathbf{u} + \nabla \mathbf{p} = \mathbf{F} + \mathcal{N}(\mathbf{v}), \quad \operatorname{div} \mathbf{u} = 0 \quad \text{in } \Omega \times \mathbb{T}, \quad \mathbf{u}|_\Gamma = \mathbf{h}|_\Gamma, \quad (5.47)$$

where $\mathcal{N}(\mathbf{v}) = \mathbf{v} \cdot \nabla \mathbf{v}$. Theorem 5.2 yields that

$$\|(\mathbf{u}, \mathbf{p})\|_{\mathcal{I}_\varepsilon} \leq C(\langle \mathbf{G}_S \rangle_3 + \langle \mathbf{H}_S \rangle_2 + \langle \mathcal{N}(\mathbf{v})_S \rangle_3 + \langle \tilde{\mathcal{N}}(\mathbf{v})_S \rangle_2 \\ + \langle \mathbf{G}_\perp \rangle_{p,2} + \langle \mathbf{H}_\perp \rangle_{p,1} + \langle \mathcal{N}(\mathbf{v})_\perp \rangle_{p,2} \\ + \langle \tilde{\mathcal{N}}(\mathbf{v})_\perp \rangle_{p,1} + \|\mathbf{h}\|_{\mathbf{H}_p^1(\mathbb{T}, L_q(\Omega))} + \|\mathbf{h}\|_{L_p(\mathbb{T}, \mathbf{H}_q^2(\Omega))}) \quad (5.48)$$

provided that the right-hand side of (5.48) is finite. Here, we write $\tilde{\mathcal{N}}(\mathbf{v}) = \mathbf{v} \otimes \mathbf{v}$, so that $\operatorname{div} \tilde{\mathcal{N}}(\mathbf{v}) = \mathcal{N}(\mathbf{v})$ since $\operatorname{div} \mathbf{v} = 0$. We further have

$$\mathcal{N}(\mathbf{v})_S = \mathbf{v}_S \cdot \nabla \mathbf{v}_S + \int_{\mathbb{T}} \mathbf{v}_\perp \cdot \nabla \mathbf{v}_\perp dt \\ \tilde{\mathcal{N}}(\mathbf{v})_S = \mathbf{v}_S \otimes \mathbf{v}_S + \int_{\mathbb{T}} \mathbf{v}_\perp \otimes \mathbf{v}_\perp dt; \\ \mathcal{N}(\mathbf{v})_\perp = \mathbf{v}_S \cdot \nabla \mathbf{v}_\perp + \mathbf{v}_\perp \cdot \nabla \mathbf{v}_S + \mathbf{v}_\perp \cdot \nabla \mathbf{v}_\perp - \int_{\mathbb{T}} \mathbf{v}_\perp \cdot \nabla \mathbf{v}_\perp dt \\ \tilde{\mathcal{N}}(\mathbf{v})_\perp = \mathbf{v}_S \otimes \mathbf{v}_\perp + \mathbf{v}_\perp \otimes \mathbf{v}_S + \mathbf{v}_\perp \otimes \mathbf{v}_\perp - \int_{\mathbb{T}} \mathbf{v}_\perp \otimes \mathbf{v}_\perp dt. \quad (5.49)$$

Notice that $\operatorname{div} \tilde{\mathcal{N}}(\mathbf{v})_S = \mathcal{N}(\mathbf{v})_S$ and $\operatorname{div} \tilde{\mathcal{N}}(\mathbf{v})_\perp = \mathcal{N}(\mathbf{v})_\perp$. To estimate these nonlinear terms, we choose $\sigma > 0$ so small that $\sigma + 3/q < 2(1 - 1/p)$, which is possible since $2/p + 3/q < 2$ by assumption. By Sobolev inequality and real interpolation, we then have

$$\|\mathbf{v}_\perp\|_{L_\infty(\mathbb{T}, L_\infty(\Omega))} \leq C\|\mathbf{v}_\perp\|_{L_\infty(\mathbb{T}, W_q^{\sigma+3/q}(\Omega))} \leq C\|\mathbf{v}_\perp\|_{L_\infty(\mathbb{T}, B_{q,p}^{2(1-1/p)}(\Omega))} \\ \leq C(\|\partial_t \mathbf{v}_\perp\|_{L_p(\mathbb{T}, L_q(\Omega))} + \|\mathbf{v}_\perp\|_{L_p(\mathbb{T}, \mathbf{H}_q^2(\Omega))}), \quad (5.50)$$

Using Hölder's inequality with $p > 2$, we further obtain

$$\langle \mathcal{N}(\mathbf{v})_S \rangle_3 \leq C(\langle \mathbf{v}_S \rangle_1 \langle \nabla \mathbf{v}_S \rangle_2 + \langle \mathbf{v}_\perp \rangle_{p,1} \langle \nabla \mathbf{v}_\perp \rangle_{p,2}); \\ \langle \tilde{\mathcal{N}}(\mathbf{v})_S \rangle_2 \leq C(\langle \mathbf{v}_S \rangle_1^2 + \langle \mathbf{v}_\perp \rangle_{p,1}^2); \\ \langle \mathcal{N}(\mathbf{v})_\perp \rangle_{p,2} \leq C(\langle \mathbf{v}_S \rangle_1 \langle \nabla \mathbf{v}_\perp \rangle_{p,2} + \langle \mathbf{v}_\perp \rangle_{p,1} \langle \nabla \mathbf{v}_S \rangle_2 \\ + \|\mathbf{v}_\perp\|_{L_\infty(\mathbb{T}, L_\infty(\Omega))} \langle \nabla \mathbf{v}_\perp \rangle_{p,2} + \langle \mathbf{v}_\perp \rangle_{p,1} \langle \nabla \mathbf{v}_\perp \rangle_{p,2}); \\ \langle \tilde{\mathcal{N}}(\mathbf{v})_\perp \rangle_{p,1} \leq C(\langle \mathbf{v}_S \rangle_1 \langle \mathbf{v}_\perp \rangle_{p,1} + \|\mathbf{v}_\perp\|_{L_\infty(\mathbb{T}, L_\infty(\Omega))} \langle \mathbf{v}_\perp \rangle_{p,1} + \langle \mathbf{v}_\perp \rangle_{p,1}^2). \quad (5.51)$$

Combining (5.48) with (5.51) and (5.50) yields that

$$\|(\mathbf{u}, \mathbf{p})\|_{\mathcal{I}_\varepsilon} \leq C(\langle \mathbf{G}_S \rangle_3 + \langle \mathbf{H}_S \rangle_2 + \langle \mathbf{G}_\perp \rangle_{p,2} + \langle \mathbf{H}_\perp \rangle_{p,1} \\ + \|\mathbf{h}\|_{\mathbf{H}_p^1(\mathbb{T}, L_q(\Omega))} + \|\mathbf{h}\|_{L_p(\mathbb{T}, \mathbf{H}_q^2(\Omega))} + \|(\mathbf{v}, \mathbf{q})\|_{\mathcal{I}_\varepsilon}^2).$$

Recalling the smallness assumption (5.3) and that $(\mathbf{v}, \mathbf{q}) \in \mathcal{I}_\varepsilon$, we have $\|(\mathbf{u}, \mathbf{p})\|_{\mathcal{I}_\varepsilon} \leq C_0\varepsilon^2$ for some constant C_0 . Thus, choosing $\varepsilon > 0$ so small that $C_0\varepsilon \leq 1$, we have $\|(\mathbf{u}, \mathbf{p})\|_{\mathcal{I}_\varepsilon} \leq \varepsilon$, which implies that $(\mathbf{u}, \mathbf{p}) \in \mathcal{I}_\varepsilon$. Therefore, if we define a map Ξ acting on $(\mathbf{v}, \mathbf{q}) \in \mathcal{I}_\varepsilon$ by setting $\Xi(\mathbf{v}, \mathbf{q}) = (\mathbf{u}, \mathbf{p})$, then Ξ is a map from \mathcal{I}_ε into itself.

In an analogous way, we see that for any $(\mathbf{v}_i, \mathbf{q}_i) \in \mathcal{I}_\varepsilon$ ($i = 1, 2$),

$$\|\Xi(\mathbf{v}_1, \mathbf{q}_1) - \Xi(\mathbf{v}_2, \mathbf{q}_2)\|_{\mathcal{I}_\varepsilon} \leq C_1\varepsilon\|(\mathbf{v}_1, \mathbf{q}_1) - (\mathbf{v}_2, \mathbf{q}_2)\|_{\mathcal{I}_\varepsilon}$$

for some constant C_1 . Thus, choosing $\varepsilon > 0$ smaller if necessary, we have $C_1\varepsilon < 1$, which shows that Ξ is a contraction map on \mathcal{I}_ε . Therefore, there exists a unique $(\mathbf{u}, \mathbf{p}) \in \mathcal{I}_\varepsilon$ such that $\Xi(\mathbf{u}, \mathbf{p}) = (\mathbf{u}, \mathbf{p})$, which is the required unique solution to (4.9). This completes the proof of Theorem 5.1. \square

References

- [1] H. Amann, *Periodic solutions of semilinear parabolic equations*, In *Nonlinear Analysis*, pages 1–29. Elsevier, 1978.
- [2] W. Arendt, S. Bu, *The operator-valued Marcinkiewicz multiplier theorem and maximal regularity*, *Math. Z.*, **240**(2) (2002), 311–343.
- [3] N. Artemiev, *Über die periodischen Lösungen der nichtlinearen partiellen Differentialgleichungen*, *Izv. Akad. Nauk SSSR, Ser. Mat.*, **1937** (1937), 15–50
- [4] R. Bader, W. Kryszewski, *On the solution sets of differential inclusions and the periodic problem in Banach spaces*, *Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods*, **54**(4) (2003), 707–754.
- [5] D. W. Bange, *Periodic solutions of a quasilinear parabolic differential equation*, *J. Diff. Eqns.*, **17**(1)(1975), 61–72.
- [6] M. E. Bogovskiĭ, *Solution of the first boundary value problem for the equation of continuity of an incompressible medium*, *Dokl. Acad. Nauk SSSR* **248** (5), 1037–1040; English Transl: *Soviet Math Dokl.*, **20** (1979), 1094–1098.
- [7] M. E. Bogovskiĭ, *Solution of some vector analysis problems connected with operators Div and Grad*, *Trudy Seminar S.L.Sobolev #1*, 80, Akademia Nauk SSSR, Sibirskoe Otdelnie Matematiki, Nowosibirsk, 5-40 (in Russian).
- [8] W. Borchers, H. Sohr, *On the semigroup of the Stokes operator for exterior domains in L^q -spaces*, *Mathematische Zeitschrift* **196** (3) (1987), 415–425.
- [9] F. E. Browder, *Existence of periodic solutions for nonlinear equations of evolution*, *Proceedings of the National Academy of Sciences* **53** (5) (1965), 1100–1103.
- [10] F. Bruhat, *Distributions sur un groupe localement compact et applications à l'étude des représentations des groupes p -adique*, *Bull. Soc. Math. Fr.*, **89** (1961), 43–75
- [11] H. Brézis, L. Nirenberg, *Forced vibrations for a nonlinear wave equation*, *Commun. Pure, Appl. Math.*, **31**(1) (1978), 1–30.

- [12] L. Cesari, *Existence in the large of periodic solutions of hyperbolic partial differential equations*, Arch. Rational Mech. Anal., **20**(3) (1965), 170–190.
- [13] A. Ówiszewski, *Positive periodic solutions of parabolic evolution problems: a translation along trajectories approach*, Cent. Eur. J. Math., **9**(2) (2011), 244–268.
- [14] T. Eiter, *On the Stokes-type resolvent problem associated with time-periodic flow around a rotating obstacle*, J. Math. Fluid Mech. **24**, 52 (2022).
- [15] T. Eiter, *On the Oseen-type resolvent problem associated with time-periodic flow around a rotating body*, arXiv:2111.00984, 2021.
- [16] T. Eiter, M. Kyed, *Time-periodic linearized Navier–Stokes equations: An approach based on Fourier multipliers*. In *Particles in flows*, Adv. Math. Fluid Mech., pp. 77–137. Birkhäuser/Springer, Cham, 2017.
- [17] T. Eiter, M. Kyed, *Estimates of time-periodic fundamental solutions to the linearized Navier-Stokes equations*, J. Math. Fluid Mech., 20(2):517–529, 2018.
- [18] T. Eiter, M. Kyed, *Viscous Flow Around a Rigid Body Performing a Time-periodic Motion*, J. Math. Fluid Mech., 23(1):28, 2021.
- [19] K. de Leeuw, *On L_p multipliers*, Ann. Math. **81** (1965), 364–379.
- [20] R. Farwig, H. Kozono, K. Tsuda, D. Wegmann, *The time periodic problem for the Navier-Stokes equations in a bounded domain with moving boundary*, Preprint in 2020.
- [21] F. A. Ficken, B. A. Fleishman, *Initial value problems and time-periodic solutions for a nonlinear wave equation*, Commun Pure Appl. Math., **10**(3) (1957), 331–356.
- [22] P. Fife, *Solutions of parabolic boundary problems existing for all time*, Arch. Rational Mech. Anal., **16**(3) (1964), 155–186.
- [23] R. Gaines, W. Walter, *Periodic solutions to nonlinear parabolic differential equations*, Rocky Mountain Journal of Mathematics, **7**(2) (1977), 297–312.
- [24] G. P. Galdi, H. Sohr, *Existence and uniqueness of time-periodic physically reasonable Navier-Stokes flow past a body*, Arch. Rational. Mech. Anal., **172**(3) (2004), 363–406.
- [25] G. P. Galdi, *An Introduction to the Mathematical Theory of the Navier-Stokes Equations, Steady-State Problems*, Second ed. 2011 Springer.
- [26] M. Geissert, M. Hieber, T. H. Nguyen, *A general approach to time periodic incompressible viscous fluid flow problems*, Arch. Ration. Mech. Anal., **220**(3) (2016), 1095–1118.
- [27] W. S. Hall, *Periodic solutions of a class of weakly nonlinear evolution equations*, Arch. Rational Mech. Anal., **39** (1970), 294–322.
- [28] T. Hytönen, J. van Neerven, M. Veraar, L. Weis, *Analysis in Banach spaces. Vol. I. Martingales and Littlewood-Paley theory*, 2016 Springer, Cham.
- [29] H. Iwashita $L_q - L_r$ estimates for solutions of the nonstationary Stokes equations in an exterior domain and the Navier–Stokes initial value problems in L_q spaces, Math. Ann., **285** (1989), 265–288.

- [30] C. Jin, T. Yang *Time periodic solution for a 3-D compressible Navier-Stokes system with an external force in \mathbb{R}^3* , J. Diff. Eqns **259** (2015), 2576–2601.
- [31] D. H. Karimov, *Über die periodischen Lösungen der nichtlinearen differentialen Gleichungen des parabolischen Typus*, C.R.(Dokl.) Acad. Sci. URSS, Ser. **28** (1940), 403–406.
- [32] M. Krasnosel'skiĭ, *The operator of translation along the trajectories of differential equations.*, Translations of Mathematical Monographs, 1968.
- [33] T. Kobayashi, Y. Shibata, *On the Oseen equation in exterior domains*, Math. Ann., **310** (1) (1998), 1–45.
- [34] P. Kokocki, *Krasnosel'skii type formula and translation along trajectories method on the scale of fractoinal spaces*, Commun. Pure Appl. Anal **14**(6) (2015), 2315–2334.
- [35] Yu. S. Kolesov, *Vertain tests for the existence of stable periodic solutions of quasi-linear parabolic equations*, Sov. Math., Dokl. **5** (1964), 1118–1120
- [36] Yu. S. Kolesov, *A test for the existence of periodic solutions to parabolic equations*, Sov. Math., Dokl., **7** (1966), 1318–1320.
- [37] Yu. S. Kolesov, *Periodische Lösungen quasilinear parabolischer Gleichungen zweiter Ordnung*, Tr. Mosk. Mat. O.-va, **21** (1970), 103–134.
- [38] H. Kozono, M. Nakao, *Periodic solutions of the Navier-Stokes equations in unbounded domains*, Tohoku Math. J. (2), **48**(1) (1996), 33–50.
- [39] M. Kyed, *Time-Periodic Solutions to the Navier-Stokes Equations*, Habilitationsschrift, Technische Universität Darmstadt, 2012.
- [40] G. M. Liebermann, *Time-periodic solutions of linear parabolic differential equations*, Commun. Parital Differ. Equations, **24** (3-4) (1999), 631–663.
- [41] P. Maremonti, *Existence and stability of time-periodic solutions to the Navier-Stokes equations in the whole space*, Nonlinearity, **4**(2)(1991), 503–529.
- [42] A. Matsumura, T. Nishida, *Periodic solutions of a viscous gas equation*, in: Recent Topics in Nonlinear PDE, Vol IV, Eds. M. Mimura and T. Nishida, North-Holland Math. Studies, **160** (1989), 49–82.
- [43] M. Nakao, *On boundedness, operiodicity, and almost periodicity of solutions of some nonlinear parabolic equations*, J. Diff. Eqns., **19**(2)(1975), 371–385.
- [44] T. H. Nguyen, T. K. O. Tran, *Periodicity of inhomogeneous trajectories and applications*, J. Math. Anal. Appl., **468**(1) (2018), 161–168.
- [45] H. Poincaré, *Sur le problème des trois corps et les équations de la dynamique*, Acta mathematica, F. & G. Beijer, 1890.
- [46] H. Poincaré, *The three-body problem and the equations of dynamics*, Poincaré's foundational work on dynamical systems theory. Translated from the French by Bruce D. Popp. Originally published by Institute Mittag-Leffler, Sweden, 1890., volume 443. Cham: Springer, originally published by institut mittag-leffler, weden, 1890 edition, 2017.

- [47] G. Prodi, *Soluzioni periodiche di equazioni alle derivate parziali di tipo parabolico e non lineari*, Riv. Mat. Univ. Parma, **3** (1952), 265–290.
- [48] G. Prodi, *Soluzioni periodiche di equazioni a derivate parziali di tipo iperbolico non lineari*, Annali di Matematica Pura ed Applicata, Series 4, **42**(1) (1956), 25–49.
- [49] G. Prodi, *Qualche risultato riguardo alle equazioni di Navier-Stokes nel caso bidimensionale*, Rend. Sem. Mat. Univ. Padova, **30** (1960), 1–15.
- [50] G. Prouse, *Soluzioni periodiche dell'equazione delle onde non omogenea con termine dissipativo quadratico*, Ric. Mat., **13** (1964), 261–280.
- [51] J. Prüß, *Periodic solutions of semilinear evolution equations*, Nonlinear Anal., Theory Methods Appl., **3** (1979), 601–612.
- [52] P. H. Rabinowitz, *Periodic solutions of nonlinear hyperbolic partial differential equations*, Commun. Pure Appl. Math., **20** (1967), 145–205
- [53] P. H. Rabinowitz, *Periodic solutions of nonlinear hyperbolic partial differential equations, II*, Commun. Pure Appl. Math., **22** (1969), 15–39
- [54] T. I. Seidman, *Periodic solutions of a non-linear parabolic equation*, J. Diff. Eqns., **19**(2) (1975), 242–257.
- [55] J. Serrin, *A note on the existence of periodic solutions of the Navier-Stokes equations*, Arch. Rational Mech. Anal., **3** (1959), 120–122.
- [56] Y. Shibata, *On an exterior initial-boundary value problem for Navier-Stokes equations*, Quart. Appl. Math., **57**(1) (1999), 117–155.
- [57] Y. Shibata, *On the \mathcal{R} -boundedness of solution operators for the Stokes equations with free boundary condition*, Differ. Int. Eqns., **27** (3–4), (2014), 313–368.
- [58] Y. Shibata, *On the \mathcal{R} -bounded solution operators in the study of free boundary problem for the Navier-Stokes equations*, In Springer Proceedings in Mathematics & Statistics, ed. by Y. Shibata, Y. Suzuki, vol. 183 (Mathematical Fluid Dynamics, Present and Future, Tokyo, 2016), pp.203–285
- [59] Y. Shibata, *On the L_p - L_q decay estimate for the Stokes equations with free boundary conditions in an exterior domain*, Asymptotic Anal. **107** (2018), 33–72.
- [60] Y. Shibata, *\mathcal{R} Boundedness, Maximal Regularity and Free Boundary Problems for the Navier Stokes equations*, Centrarò, Italy 2017, Lecture Notes in Mathematics 2254, CIME Foundation Subseries, Springer 2020, pp.193–462.
- [61] P. Solovieff, *Quelques remarques sur les solutions des équations nonlinéaires du type hyperbolique*, Izv. Acad. Nauk SSSR, Ser. Mat., **3** (1939), 149–164.
- [62] I. Straskraba, O. Vejvoda, *Periodic solutions to a singular abstract differential equations*, Cech. Math. J., **24** (1974), 528–540.
- [63] C. T. Taam, *Stability, periodicity, and almost periodicity of solutions of nonlinear differential equations in Banach spaces*, J. Math. Mech., **15** (1966), 849–876.

- [64] K. Tsuda, *On the existence and stability of time periodic solution to the compressible Navier-Stokes equation on the whole space*, Arch. Rational Mech. Anal. **219** (2016) 637–678.
- [65] A. Valli, *Periodic and stationary solutions for compressible Navier-Stokes equations via a stability method*, Ann. Sc. Norm. Super. Pisa C1. Sci. (4) 10 (1983), 607–647.
- [66] O. Vejvoda, *Parital differential equations: time-periodic solutions*. The Hague-Boston-London: Martinus Nijhoff Publishers; Prague: SNTL, Publishers of Technical Literature, 1982.
- [67] L. Weis, *Operator-valued Fourier multiplier theorems and maximal L_p -regularity*. Math. Ann. **319** (2001), 735–758.
- [68] M. Yamazaki, *The Navier-Stokes equations in the weak- L^n space with time-dependent external force*, Math. Ann., **317**(4) (2000), 635–675.
- [69] M. E. Zabolinskij *Über periodische Lösungen nichtlinearer partieller Differentialgleichungen*, Dokl. Acad. Nauk SSSR, Ser. **56** (1947), 469–472.