

Weierstraß–Institut für Angewandte Analysis und Stochastik

im Forschungsverbund Berlin e.V.

Preprint

ISSN 0946 – 8633

Singularly perturbed partly dissipative reaction-diffusion systems in case of exchange of stabilities

Valentin F. Butuzov,¹ Nikolai N. Nefedov,¹ Klaus R. Schneider²


submitted: 6th April 2000

¹ Moscow State University
Faculty of Physics
Department of Mathematics
Vorob'jovi Gori
119899 Moscow, Russia
E-Mail: butuzov@mt384.phys.msu.su
E-Mail: nefedov@mt384.phys.msu.su

² Weierstrass Institute for Applied
Analysis and Stochastics
Mohrenstraße 39
D – 10117 Berlin
Germany
E-Mail: schneider@wias-berlin.de

Preprint No. 572

Berlin 2000



1991 *Mathematics Subject Classification.* 35B25, 35K57.

Key words and phrases. Initial boundary value problem, singularly perturbed partly dissipative reaction-diffusion system, exchange of stabilities, asymptotic lower and upper solutions.

Edited by
Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS)
Mohrenstraße 39
D — 10117 Berlin
Germany

Fax: + 49 30 2044975
E-Mail (X.400): c=de;a=d400-gw;p=WIAS-BERLIN;s=preprint
E-Mail (Internet): preprint@wias-berlin.de
World Wide Web: <http://www.wias-berlin.de/>

Abstract

We consider the singularly perturbed partly dissipative reaction-diffusion system $\varepsilon^2 \left(\frac{\partial u}{\partial t} - \frac{\partial^2 u}{\partial x^2} \right) = g(u, v, x, t, \varepsilon)$, $\frac{\partial v}{\partial t} = f(u, v, x, t, \varepsilon)$ under the condition that the degenerate equation $g(u, v, t, 0) = 0$ has two solutions $u = \varphi_i(v, x, t)$, $i = 1, 2$, that intersect (exchange of stabilities). Our main result concerns existence and asymptotic behavior in ε of the solution of the initial boundary value problem under consideration. The proof is based on the method of asymptotic lower and upper solutions.

1 Introduction.

We consider the singularly perturbed partly dissipative reaction-diffusion system

$$\begin{aligned} \varepsilon^2 \left(\frac{\partial u}{\partial t} - \frac{\partial^2 u}{\partial x^2} \right) &= g(u, v, x, t, \varepsilon), \\ \frac{\partial v}{\partial t} &= f(u, v, x, t, \varepsilon), \end{aligned} \tag{1.1}$$

where u and v are scalar functions, ε is a small positive parameter. Partly dissipative systems are often used to model reaction-diffusion processes in different fields (chemical kinetics, biology, astrophysics) when the effect of diffusion of one of the species is negligible (see, for example, [10, 11, 6, 7, 5, 12, 4]).

If the degenerate equation

$$g(u, v, x, t, 0) = 0$$

has an isolated solution with respect to u then the standard theory (see [14]) can be applied to derive asymptotic properties for the solution of initial boundary value problems to system (1.1).

In the present paper we consider system (1.1) under the assumption that the degenerate equation has two intersecting solutions. This assumption implies an exchange of stabilities for families of equilibria of the associated differential equation to (1.1). The main result of this paper concerns the existence and asymptotic behavior in ε of the solution of some initial boundary value problem related to system (1.1). The proof of our results is based on the method of asymptotic lower and upper solutions. The results of this paper are extensions of corresponding results in [9, 8, 2, 3], they can be used to study differential systems modelling bimolecular reactions with fast reaction rates and to explain the jumping behavior of the reaction rate in systems of this type.

2 Formulation of the problem. Assumptions.

We study the singularly perturbed nonlinear initial boundary value problem

$$\begin{aligned}
\varepsilon^2 \left(\frac{\partial u}{\partial t} - \frac{\partial^2 u}{\partial x^2} \right) &= g(u, v, x, t, \varepsilon), \\
\frac{\partial v}{\partial t} &= f(u, v, x, t, \varepsilon), \\
(x, t) \in Q &:= \{(x, t) \in \mathbb{R}^2 : 0 < x < 1, 0 < t \leq T\}, \\
\varepsilon \in I_{\varepsilon_0} &:= \{\varepsilon \in \mathbb{R} : 0 < \varepsilon \leq \varepsilon_0\}, 0 < \varepsilon_0 \ll 1, \\
\frac{\partial u}{\partial x}(0, t, \varepsilon) &= \frac{\partial u}{\partial x}(1, t, \varepsilon) = 0 \quad \text{for } 0 < t \leq T, \\
u(x, 0, \varepsilon) &= u^0(x), \quad v(x, 0, \varepsilon) = v^0(x) \quad \text{for } 0 \leq x \leq 1
\end{aligned} \tag{2.1}$$

under the following assumptions:

(A₀). $f, g \in C^2(D, \mathbb{R})$, where $D := \mathbb{R} \times \mathbb{R} \times [0, 1] \times [0, T] \times \overline{I}_{\varepsilon_0}$, $u^0, v^0 \in C^2([0, 1], \mathbb{R})$.

If we set $\varepsilon = 0$ in (2.1) then we get the degenerate system

$$\begin{aligned}
0 &= g(u, v, x, t, 0), \\
\frac{dv}{dt} &= f(u, v, x, t, 0).
\end{aligned} \tag{2.2}$$

Concerning the solution set of the equation

$$g(u, v, x, t, 0) = 0 \tag{2.3}$$

we assume

(A₁). Equation (2.3) has exactly two solutions $u = \varphi_1(v, x, t)$ and $u = \varphi_2(v, x, t)$ defined for $(v, x, t) \in \overline{I}_v \times \overline{Q}$, where I_v is some open interval, and that have the same smoothness properties as g .

From assumption (A₁) we obtain for $(v, x, t) \in \overline{I}_v \times \overline{Q}$, $i = 1, 2$:

$$\begin{aligned}
g(\varphi_i(v, x, t), v, x, t, 0) &\equiv 0, \\
g_u(\varphi_i(v, x, t), v, x, t, 0) \frac{\partial \varphi_i}{\partial v}(v, x, t) + g_v(\varphi_i(v, x, t), v, x, t, 0) &\equiv 0.
\end{aligned} \tag{2.4}$$

Different from the standard case (see [13],[14]), assumption (A₁) does not require that the solutions φ_1 and φ_2 are isolated. The following assumption expresses the

property that the surfaces $u = \varphi_1(v, x, t)$ and $u = \varphi_2(v, x, t)$ intersect in a smooth surface whose projection into the (v, x, t) -space can be described by $v = s(x, t)$.

(A₂). *There exists a smooth function $s : \overline{Q} \rightarrow R$ such that*

$$\begin{aligned}\varphi_1(v, x, t) &= \varphi_2(v, x, t) & \text{for } v &= s(x, t), \\ \varphi_1(v, x, t) &> \varphi_2(v, x, t) & \text{for } v &< s(x, t), \\ \varphi_1(v, x, t) &< \varphi_2(v, x, t) & \text{for } v &> s(x, t).\end{aligned}$$

The differential equation

$$\frac{du}{d\tau} = g(u, v, x, t, 0), \quad (2.5)$$

where v, x, t are considered as parameters, is said to be the associated equation to (2.1). It follows from hypothesis (A₁) that $u = \varphi_i(v, x, t), i = 1, 2$, are families of equilibria of (2.5). The families φ_i are stable (unstable) if $g_u(\varphi_i, v, x, t, 0)$ is negative (positive). Concerning the stability of these families we assume

(A₃). *For $(x, t) \in \overline{Q}$ it holds*

$$\begin{aligned}g_u(\varphi_1(v, x, t), v, x, t, 0) &< 0, & g_u(\varphi_2(v, x, t), v, x, t, 0) &> 0 & \text{for } v < s(x, t), \\ g_u(\varphi_1(v, x, t), v, x, t, 0) &> 0, & g_u(\varphi_2(v, x, t), v, x, t, 0) &< 0 & \text{for } v > s(x, t).\end{aligned}$$

From assumption (A₃) we get that $g_u(u, v, x, t, 0)$ changes its sign on the surface $v = s(x, t)$ where $u = \varphi_1(v, x, t)$ and $u = \varphi_2(v, x, t)$ intersect. This sign change of g_u expresses an exchange of stabilities of the families of equilibria of the associated equation (2.5).

From (A₃) we get for $(x, t) \in \overline{Q}$

$$g_u(\varphi_1(s(x, t), x, t), s(x, t), x, t, 0) \equiv g_u(\varphi_2(s(x, t), x, t), s(x, t), x, t, 0) \equiv 0.$$

In what follows we will construct the so-called composed stable solution to the degenerate system (2.2). To this purpose we have to assume $v^0(x) \neq s(x, 0)$ for all x . First we consider the case

$$v^0(x) < s(x, 0) \quad \text{for } 0 \leq x \leq 1. \quad (2.6)$$

(A₄). *For $x \in [0, 1]$, the initial value problem*

$$\begin{aligned}\frac{dv}{dt} &= f(\varphi_1(v, x, t), v, x, t, 0), \\ v(x, 0) &= v^0(x)\end{aligned} \quad (2.7)$$

where $v^0(x)$ satisfies (2.6) has a unique solution $v = v_1(x, t)$ defined on \overline{Q} . There exists a smooth curve \mathcal{C} defined by $\mathcal{C} := \{(x, t) \in \overline{Q} : t = t_c(x), 0 \leq x \leq 1\}$ where $t_c \in C^2[0, 1]$ and such that

$$\begin{aligned}
 0 < t_c(x) < T & \quad \text{for} & \quad 0 \leq x \leq 1, \\
 v_1(x, t) < s(x, t) & \quad \text{for} & \quad 0 \leq t < t_c(x), 0 \leq x \leq 1, \\
 v_1(x, t) = s(x, t) & \quad \text{for} & \quad t = t_c(x), 0 \leq x \leq 1, \\
 v_1(x, t) > s(x, t) & \quad \text{for} & \quad t_c(x) < t \leq T, 0 \leq x \leq 1.
 \end{aligned} \tag{2.8}$$

Assumption (A₄) says that the surfaces $v = v_1(x, t)$ and $v = s(x, t)$ intersect in a curve whose projection into \overline{Q} can be described by $t = t_c(x)$. We denote by Q_1 and Q_2 all points (x, t) of \overline{Q} satisfying $t < t_c(x)$ and $t > t_c(x)$ respectively (see Fig 1.).




Fig. 1: Decomposition of \overline{Q} into Q_1 and Q_2 by the curve \mathcal{C} .

(A₅). For $x \in [0, 1]$, the initial value problem

$$\begin{aligned}
 \frac{dv}{dt} &= f(\varphi_2(v, x, t), v, x, t, 0), \\
 v(x, t_c(x)) &= s(x, t_c(x))
 \end{aligned} \tag{2.9}$$

has a unique solution $v = v_2(x, t)$ defined on \overline{Q} such that

$$\begin{aligned}
 v_2(x, t) > s(x, t) & \quad \text{for} & \quad (x, t) \in Q_2, \\
 v_2(x, t) < s(x, t) & \quad \text{for} & \quad (x, t) \in Q_1.
 \end{aligned} \tag{2.10}$$

Let

$$\begin{aligned}\psi_1(x, t) &:= \varphi_1(v_1(x, t), x, t) & \text{for } (x, t) \in \overline{Q}, \\ \psi_2(x, t) &:= \varphi_2(v_2(x, t), x, t) & \text{for } (x, t) \in \overline{Q}.\end{aligned}\tag{2.11}$$

From assumption (A₂) and from (2.11) we obtain

$$\psi_1(x, t) \equiv \psi_2(x, t) \quad \text{on } \mathcal{C}.\tag{2.12}$$

Now, we introduce the functions $\hat{u}(x, t)$ and $\hat{v}(x, t)$ by

$$\begin{aligned}\hat{u}(x, t) &:= \begin{cases} \psi_1(x, t) & \text{for } (x, t) \in \overline{Q}_1, \\ \psi_2(x, t) & \text{for } (x, t) \in \overline{Q}_2, \end{cases} \\ \hat{v}(x, t) &:= \begin{cases} v_1(x, t) & \text{for } (x, t) \in \overline{Q}_1, \\ v_2(x, t) & \text{for } (x, t) \in \overline{Q}_2. \end{cases}\end{aligned}\tag{2.13}$$

The pair of functions $(\hat{u}(x, t), \hat{v}(x, t))$ is referred to as the *composed stable solution* of the degenerate system (2.2).

The function $\hat{v}(x, t)$ is obviously continuously differentiable with respect to t . But $\hat{u}(x, t)$ is in general not smooth on the curve \mathcal{C} , since we get from (2.8), (2.10) and (2.11)

$$\frac{\partial \psi_1}{\partial t} \leq \frac{\partial \psi_2}{\partial t} \quad \text{on } \mathcal{C}.$$

For the sequel it is convenient to introduce the following notation: the symbol $\hat{\cdot}$ over g and f or some derivative of g and f denotes that we have to consider the arguments (u, v, ε) at $(\hat{u}(x, t), \hat{v}(x, t), 0)$.

It follows from assumption (A₁) that

$$\hat{g}(x, t) := g(\hat{u}(x, t), \hat{v}(x, t), x, t, 0) \equiv 0 \quad \text{in } \overline{Q},\tag{2.14}$$

by assumption (A₃) we have

$$\hat{g}_u(x, t) < 0 \quad \text{in } \overline{Q} \setminus \mathcal{C},\tag{2.15}$$

$$\hat{g}_u(x, t) \equiv 0 \quad \text{on } \mathcal{C}.\tag{2.16}$$

Remark 2.1 *The case $v^0(x) > s(x, 0)$ can be treated analogously. In that case we have to use the function $\varphi_2(v, x, t)$ to construct $v_1(x, t)$ (see assumption (A₄)) and the function $\varphi_1(v, x, t)$ to construct $v_2(x, t)$ (see assumption (A₅)). The case when $v^0(x) = s(x, 0)$ for some x requires a special treatment.*

In what follows we prove that under the hypotheses (A₁) – (A₅) and under some additional assumptions (see (A₆) – (A₈) below) problem (2.1) has a solution $(u(x, t, \varepsilon), v(x, t, \varepsilon))$ satisfying

$$\begin{aligned}\lim_{\varepsilon \rightarrow 0} u(x, t, \varepsilon) &= \hat{u}(x, t) \quad \text{in } \overline{Q} \setminus \{t = 0, 0 \leq x \leq 1\}, \\ \lim_{\varepsilon \rightarrow 0} v(x, t, \varepsilon) &= \hat{v}(x, t) \quad \text{in } \overline{Q}.\end{aligned}\tag{2.17}$$

Concerning the initial condition $u^0(x)$ for $u(x, t, \varepsilon)$ we assume

(A₆). For $x \in [0, 1]$, $u^0(x)$ lies in the basin of attraction of the equilibrium point $\varphi_1(v^0(x), x, 0)$ of the associated equation (2.5) for $v = v^0(x)$, $t = 0$.

Assumption (A₆) implies that for $v = v^0(x)$, $t = 0$ equation (2.5) with the initial condition

$$u(x, 0) = u^0(x)$$

has a unique solution $u = \bar{u}(x, \tau)$ defined for $\tau \geq 0$, and such that $\lim_{\tau \rightarrow \infty} \bar{u}(x, \tau) = \varphi_1(v^0(x), x, 0)$. Finally, we assume

(A₇). $\hat{g}_{uu}(x, t) := g_{uu}(\hat{u}(x, t), \hat{v}(x, t), x, t, 0) < 0$ on \mathcal{C} .

(A₈). $\hat{g}_\varepsilon(x, t) > 0$ on \mathcal{C} .

Concerning assumption (A₈) we would like to mention that the sign of $\hat{g}_\varepsilon(x, t)$ on \mathcal{C} plays an important role (see [1] - [3]).

Our approach to prove the asymptotic behavior of the solution of problem (2.1) is based on the concept of ordered lower and upper solutions. Let us recall its definition [10].

Definition 2.1 Let the vector-functions $\alpha(x, t, \varepsilon) = (\alpha^u(x, t, \varepsilon), \alpha^v(x, t, \varepsilon))$ and $\beta(x, t, \varepsilon) = (\beta^u(x, t, \varepsilon), \beta^v(x, t, \varepsilon))$ be defined for $(x, t, \varepsilon) \in \overline{Q} \times \overline{I}_{\varepsilon_1}$, $\varepsilon_1 \leq \varepsilon_0$, and satisfy the smoothness conditions $\alpha^u, \beta^u \in C_{x,t,\varepsilon}^{2,1,0}(Q \times I_{\varepsilon_1}) \cap C_{x,t,\varepsilon}^{1,0,0}(\overline{Q} \times \overline{I}_{\varepsilon_1})$, $\alpha^v, \beta^v \in C_{x,t,\varepsilon}^{0,1,0}(Q \times I_{\varepsilon_1}) \cap C_{x,t,\varepsilon}^{0,0,0}(\overline{Q} \times \overline{I}_{\varepsilon_1})$. Furthermore, we assume $\alpha^u(x, t, \varepsilon) \leq \beta^u(x, t, \varepsilon)$, $\alpha^v(x, t, \varepsilon) \leq \beta^v(x, t, \varepsilon)$ for $(x, t, \varepsilon) \in \overline{Q} \times I_{\varepsilon_1}$. Let the operators L_v and M_u be defined by

$$(L_v w)(x, t, \varepsilon) := \varepsilon^2 \left(\frac{\partial w}{\partial t} - \frac{\partial^2 w}{\partial x^2} \right) - g(w, v, x, t, \varepsilon),\tag{2.18}$$

$$(M_u w)(x, t, \varepsilon) := \frac{\partial w}{\partial t} - f(u, w, x, t, \varepsilon).\tag{2.19}$$

Then, $\alpha(x, t, \varepsilon)$ and $\beta(x, t, \varepsilon)$ are called ordered lower and upper solutions of problem (2.1) respectively, if they satisfy the following inequalities

$$(L_v \alpha^u)(x, t, \varepsilon) \leq 0 \leq (L_v \beta^u)(x, t, \varepsilon) \text{ for } (x, t, \varepsilon) \in Q \times I_{\varepsilon_1}, \quad \alpha^v \leq v \leq \beta^v,\tag{2.20}$$

$$(M_u \alpha^v)(x, t, \varepsilon) \leq 0 \leq (M_u \beta^v)(x, t, \varepsilon) \text{ for } (x, t, \varepsilon) \in Q \times I_{\varepsilon_1}, \quad \alpha^u \leq u \leq \beta^u,\tag{2.21}$$

$$\frac{\partial \alpha^u}{\partial x}(0, t, \varepsilon) \geq 0 \geq \frac{\partial \beta^u}{\partial x}(0, t, \varepsilon), \quad \frac{\partial \alpha^u}{\partial x}(1, t, \varepsilon) \leq 0 \leq \frac{\partial \beta^u}{\partial x}(1, t, \varepsilon), \quad (2.22)$$

$$\alpha^u(x, 0, \varepsilon) \leq u^0(x) \leq \beta^u(x, 0, \varepsilon), \quad \alpha^v(x, 0, \varepsilon) \leq v^0(x) \leq \beta^v(x, 0, \varepsilon) \quad (2.23)$$

for $(x, t, \varepsilon) \in \overline{Q} \times I_{\varepsilon_1}$.

This definition can be obviously adapted to any subdomain of \overline{Q} . It is known (see, for example, [10]) that the existence of ordered lower and upper solutions of (2.1) implies the existence of a unique solution $(u(x, t, \varepsilon), v(x, t, \varepsilon))$ of (2.1) satisfying for $(x, t, \varepsilon) \in \overline{Q} \times I_{\varepsilon_1}$

$$\begin{aligned} \alpha^u(x, t, \varepsilon) &\leq u(x, t, \varepsilon) \leq \beta^u(x, t, \varepsilon), \\ \alpha^v(x, t, \varepsilon) &\leq v(x, t, \varepsilon) \leq \beta^v(x, t, \varepsilon). \end{aligned}$$

The goal of the following investigations is to characterize the asymptotic behavior of the solution of (2.1), in particular, we prove the limit behavior (2.17) by constructing lower and upper solutions to the initial boundary value problem (2.1).

3 Existence and asymptotic behavior of the solution.

In this section we will prove that the initial boundary value problem (2.1) has a unique solution. Taking into account an initial layer correction we can show that for small ε the solution of (2.1) is close to the composed stable solution $(\hat{u}(x, t), \hat{v}(x, t))$. In order to be able to formulate our main result we decompose the domain \overline{Q} into two disjoint sub-domains Q_1 and Q_2 and introduce a function which represents an approximation of the initial layer correction.

First we decompose \overline{Q} . Let t_{min} be the minimum of the function $t_c(x)$ in $[0, 1]$, let ν be any small positive number such that $t_1 := t_{min} - \nu$ is positive. Let Q_c be the domain defined by $Q_c := \{(x, t) \in R^2 : 0 < x < 1, t_1 < t \leq T\}$, (see Fig. 2).

Next we introduce an initial layer correction. According to [14] we define the zeroth order initial layer function $\Pi_0(x, \tau)$ ($\tau = t/\varepsilon^2$) as the solution of the initial value problem ($x \in [0, 1]$ has to be considered as a parameter)

$$\begin{aligned} \frac{d\Pi_0}{d\tau} &= g(\psi_1(x, 0) + \Pi_0, v^0(x), x, 0, 0), \quad \tau > 0, \\ \Pi_0(x, 0) &= u^0(x) - \psi_1(x, 0). \end{aligned} \quad (3.1)$$




Fig. 2: Decomposition of \bar{Q} .

By (2.11) we have $\psi_1(x, 0) = \varphi_1(v^0(x), x, 0)$. Thus, from assumptions (A₃) and (A₆) it follows that the initial value problem (3.1) has a solution which satisfies the estimate $|\Pi_0(x, \tau)| < c \exp(-\kappa\tau)$, $\tau \geq 0$ for some positive constants c and κ .

Theorem 3.1 *Assume hypotheses (A₀)–(A₈) to be valid. Then, for sufficiently small ε , the initial boundary value problem (2.1) has a unique solution $(u(x, t, \varepsilon), v(x, t, \varepsilon))$ satisfying*

$$u(x, t, \varepsilon) = \begin{cases} \hat{u}(x, t) + \Pi_0(x, \tau) + O(\varepsilon) & \text{for } (x, t) \in \bar{Q} \setminus Q_c, \\ \hat{u}(x, t) + O(\sqrt{\varepsilon}) & \text{for } (x, t) \in Q_c, \end{cases} \quad (3.2)$$

$$v(x, t, \varepsilon) = \begin{cases} \hat{v}(x, t) + O(\varepsilon) & \text{for } (x, t) \in \bar{Q} \setminus Q_c, \\ \hat{v}(x, t) + O(\sqrt{\varepsilon}) & \text{for } (x, t) \in Q_c. \end{cases} \quad (3.3)$$

Corollary 3.1 *From (3.2), (3.3) it is obvious that the relations (2.17) hold.*

Proof. The proof Theorem 3.1 consists of two steps. In the first step we consider the initial boundary value problem (2.1) in the sub-domain $\bar{Q} \setminus Q_c$. From our assumptions it follows that the exchange of stabilities takes place in Q_c . Therefore, we can apply the standard theory [14] to solve the initial boundary value problem in $\bar{Q} \setminus Q_c$. We get the following result.

Lemma 3.1 *Assume hypotheses (A₀) – (A₆) to be valid. Then, for sufficiently small ε , the initial boundary value problem (2.1) has a unique solution $(u(x, t, \varepsilon), v(x, t, \varepsilon))$ in $\bar{Q} \setminus Q_c$ satisfying*

$$\begin{aligned} u(x, t, \varepsilon) &= \hat{u}(x, t) + \Pi_0(x, \tau) + O(\varepsilon), \\ v(x, t, \varepsilon) &= \hat{v}(x, t) + O(\varepsilon). \end{aligned} \quad (3.4)$$

Let $u^1(x, \varepsilon) := u(x, t_1, \varepsilon)$, $v^1(x, \varepsilon) := v(x, t_1, \varepsilon)$. Now we consider the initial boundary value problem (2.1) in Q_c with the initial conditions $u(x, t_1, \varepsilon) = u^1(x, \varepsilon)$, $v(x, t_1, \varepsilon) = v^1(x, \varepsilon)$ for $0 \leq x \leq 1$.

Our approach to study this problem is based on the method of ordered lower and upper solutions. We construct these solutions for (2.1) by means of the composed stable solution $(\hat{u}(x, t), \hat{v}(x, t))$.

As we noticed above, $\hat{u}(x, t)$ in general is not smooth on the curve \mathcal{C} . In order to be able to use $\hat{u}(x, t)$ for the construction of lower and upper solutions we have to smooth $\hat{u}(x, t)$ in some neighbourhood Q_ν of \mathcal{C} . Let Q_ν be defined by $Q_\nu := \{(x, t) \in \bar{Q} : |t - t_c(x)| < \nu, 0 \leq x \leq 1\}$, where ν is any sufficiently small positive number such that Q_ν has no common point with $t = T$ (see Fig. 2).

Using the function

$$\omega(\xi) := \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\xi} \exp(-s^2) ds,$$

where

$$\xi := (t - t_c(x))/\varepsilon^a, \quad a \in (1/2, 1)$$

we introduce the function \tilde{u} by

$$\tilde{u}(x, t, \varepsilon) := \psi_1(x, t)\omega(-\xi) + \psi_2(x, t)\omega(\xi). \quad (3.5)$$

It is easy to show that \tilde{u} is smooth and satisfies

$$\tilde{u}(x, t, \varepsilon) = \hat{u}(x, t) + \eta(x, t, \varepsilon), \quad (3.6)$$

where

$$\eta(x, t, \varepsilon) = \begin{cases} O(\varepsilon^a) & \text{for } (x, t) \in Q_\nu, \\ O(\exp -(\nu/\varepsilon)) & \text{for } (x, t) \in Q \setminus Q_\nu \end{cases} \quad (3.7)$$

(see [1]).

Now we construct lower and upper solutions for (2.1) in Q_c by using the smooth function \tilde{u} as follows

$$\begin{aligned} \beta^u(x, t, \varepsilon) &:= \tilde{u}(x, t, \varepsilon) + \sqrt{\varepsilon}\gamma h(x, t) + \varepsilon^a z(x, \varepsilon), \\ \alpha^u(x, t, \varepsilon) &:= \tilde{u}(x, t, \varepsilon) - \sqrt{\varepsilon}\sigma h(x, t) - \varepsilon^a z(x, \varepsilon), \\ \beta^v(x, t, \varepsilon) &:= \hat{v}(x, t) + \sqrt{\varepsilon}\sigma^2 h(x, t), \\ \alpha^v(x, t, \varepsilon) &:= \hat{v}(x, t) - \sqrt{\varepsilon}\sigma^2 h(x, t), \end{aligned} \quad (3.8)$$

where

$$\begin{aligned} h(x, t) &:= \exp(\lambda(t - t_c(x))), \\ z(x, \varepsilon) &:= \exp(-kx/\varepsilon^a) + \exp(-k(1-x)/\varepsilon^a) \end{aligned} \quad (3.9)$$

are positive functions in $\overline{Q} \times I_{\varepsilon_1}$, $\gamma, \sigma, \lambda, k$ are positive numbers. We will determine these numbers in such a way that α and β will be ordered lower and upper solutions, i.e. they will satisfy all conditions of Definition 2.1 in \overline{Q}_c .

It is obvious that for any choice of γ, σ, λ and k we have

$$\alpha^u(x, t, \varepsilon) \leq \beta^u(x, t, \varepsilon), \quad \alpha^v(x, t, \varepsilon) \leq \beta^v(x, t, \varepsilon) \quad \text{in } \overline{Q}_c,$$

hence, if $\alpha(x, t, \varepsilon)$ and $\beta(x, t, \varepsilon)$ are lower and upper solutions for (2.1) then they are ordered.

Taking into account the exponential decay of $\Pi_0(x, \tau)$ we get from (3.8), (3.4) for sufficiently small ε

$$\begin{aligned} \alpha^u(x, t_1, \varepsilon) &\leq u(x, t_1, \varepsilon) = u^1(x, \varepsilon) \leq \beta^u(x, t_1, \varepsilon), \\ \alpha^v(x, t_1, \varepsilon) &\leq v(x, t_1, \varepsilon) = v^1(x, \varepsilon) \leq \beta^v(x, t_1, \varepsilon). \end{aligned}$$

Consequently, the inequalities (2.23) for the initial data hold.

Now we check that $\alpha^u(x, t, \varepsilon)$ and $\beta^u(x, t, \varepsilon)$ satisfy the inequalities (2.20) in Q_ν for sufficiently small ε .

From (2.11) and (2.12) we obtain

$$\psi_2(x, t) - \psi_1(x, t) = O(|t - t_c(x)|).$$

Using this relation it can be shown (see [1, 3]) that

$$\varepsilon^2 \left(\frac{\partial \tilde{u}}{\partial t} - \frac{\partial^2 \tilde{u}}{\partial x^2} \right) = \begin{cases} O(\varepsilon^{2-a}) & \text{in } Q_\nu, \\ O(\varepsilon^2) & \text{in } Q_c \setminus Q_\nu. \end{cases} \quad (3.10)$$

From (3.9) we get

$$\begin{aligned} \varepsilon^2 \varepsilon^{1/2} \left(\frac{\partial h}{\partial t} - \frac{\partial^2 h}{\partial x^2} \right) &= O(\varepsilon^{5/2}) \quad \text{in } Q_c, \\ \varepsilon^2 \varepsilon^a \left(\frac{\partial z}{\partial t} - \frac{\partial^2 z}{\partial x^2} \right) &= O(\varepsilon^{2-a}) \quad \text{in } Q_c. \end{aligned} \quad (3.11)$$

Thus, because of $1/2 < a < 1$, we obtain from (3.8) - (3.11)

$$\varepsilon^2 \left(\frac{\partial \beta^u}{\partial t} - \frac{\partial^2 \beta^u}{\partial x^2} \right) = O(\varepsilon^{2-a}) = o(\varepsilon) \quad \text{in } Q_c, \quad (3.12)$$

$$\varepsilon^2 \left(\frac{\partial \alpha^u}{\partial t} - \frac{\partial^2 \alpha^u}{\partial x^2} \right) = O(\varepsilon^{2-a}) = o(\varepsilon) \quad \text{in } Q_c. \quad (3.13)$$

To treat the expression $g(\beta^u(x, t, \varepsilon), v, x, t, \varepsilon)$ in $L_\nu \beta^u$ we use the relations $\tilde{u}(x, t, \varepsilon) = \hat{u}(x, t) + O(\varepsilon^a)$ which follows from (3.6) and (3.7) and $\varepsilon^a z(x, \varepsilon) = O(\varepsilon^a)$ in Q_ν due to

(3.9). Moreover, we note that the set of all v satisfying $\alpha^v(x, t, \varepsilon) \leq v \leq \beta^v(x, t, \varepsilon)$ can be represented in the form

$$v = \hat{v}(x, t) + \sqrt{\varepsilon}\sigma^2 h(x, t)\theta, \quad -1 \leq \theta \leq 1.$$

Thus, we have

$$\begin{aligned} g(\beta^u(x, t, \varepsilon), v, x, t, \varepsilon) &= g(\hat{u}(x, t) + \sqrt{\varepsilon}\gamma h(x, t) + O(\varepsilon^a), \hat{v} + \sqrt{\varepsilon}\sigma^2 h(x, t)\theta, x, t, \varepsilon) \\ &= \hat{g}(x, t) + \sqrt{\varepsilon}[\hat{g}_u(x, t)(\gamma + O(\varepsilon^{a-1/2})) + \hat{g}_v(x, t)\sigma^2\theta]h(x, t) + \\ &+ \frac{1}{2}\varepsilon[\hat{g}_{uu}(x, t)\gamma^2 + 2\hat{g}_{uv}(x, t)\gamma\sigma^2\theta + \hat{g}_{vv}(x, t)\sigma^4\theta^2]h^2(x, t) + \varepsilon\hat{g}_\varepsilon(x, t) + o(\varepsilon). \end{aligned} \quad (3.14)$$

Our goal is to prove $g(\beta^u(x, t, \varepsilon), v, x, t, \varepsilon) = -c\varepsilon + o(\varepsilon)$ for $(x, t) \in Q_\nu$ and some positive constant c .

From (2.4) we get

$$\hat{g}_v(x, t) = -\hat{g}_u(x, t)\hat{\varphi}_v(x, t), \quad (3.15)$$

where

$$\hat{\varphi}_v(x, t) = \begin{cases} \varphi_{1v}(v_1(x, t), x, t) & \text{in } Q_1, \\ \varphi_{2v}(v_2(x, t), x, t) & \text{in } Q_2. \end{cases}$$

Since $\hat{\varphi}_v(x, t)$ is uniformly bounded in \overline{Q} , $|\theta| \leq 1$, we have by (3.15) and (2.15), (2.16) for any fixed σ and for sufficiently large γ

$$\begin{aligned} \hat{g}_u(x, t)(\gamma + O(\varepsilon^{a-1/2})) + \hat{g}_v(x, t)\sigma^2\theta &= \\ = \hat{g}_u(x, t)[\gamma + O(\varepsilon^{a-1/2}) - \hat{\varphi}_v(x, t)\sigma^2\theta] &\leq 0. \end{aligned} \quad (3.16)$$

According to assumption (A₇) there is a positive constant c_ν such that for sufficiently small ν

$$\hat{g}_{uu}(x, t) \leq -c_\nu < 0 \quad \text{in } Q_\nu. \quad (3.17)$$

Hence, for sufficiently large γ , we have for $(x, t) \in Q_\nu$

$$\gamma[\hat{g}_{uu}(x, t)\gamma + 2\hat{g}_{uv}(x, t)\sigma^2\theta + \gamma^{-1}\hat{g}_{vv}(x, t)\sigma^4\theta^2] < -2\gamma\bar{c}, \quad (3.18)$$

where \bar{c} is some positive constant.

Now we set $\lambda = 1/\nu$. Then, by (3.9), it holds

$$e^{-1} \leq h(x, t) \leq e \quad \text{for } (x, t) \in \overline{Q}_\nu. \quad (3.19)$$

Under our smoothness assumption there is a positive constant c_g such that

$$|\hat{g}_\varepsilon(x, t)| \leq c_g \quad \text{for } (x, t) \in \overline{Q}_\nu. \quad (3.20)$$

By (2.14), (3.15) - (3.20) we get from (3.14)

$$g(\beta^u(x, t, \varepsilon), v, x, t, \varepsilon) < -(\gamma\bar{c}e^{-2} - c_g)\varepsilon + o(\varepsilon). \quad (3.21)$$

Taking into account (3.12) and (3.21) we have for sufficiently small ν and ε and for sufficiently large γ

$$\begin{aligned} (L_\nu \beta^u)(x, t, \varepsilon) &\equiv \varepsilon^2 \left(\frac{\partial \beta^u}{\partial t} - \frac{\partial^2 \beta^u}{\partial x^2} \right) - g(\beta^u(x, t, \varepsilon), v, x, t, \varepsilon) \\ &> (\gamma \bar{c} e^{-2} - c_g) \varepsilon + o(\varepsilon) \geq 0, \end{aligned}$$

i.e. the inequality (2.20) holds for β^u in Q_ν .

Now we verify the inequality (2.20) for α^u in Q_ν . Using (3.8), (3.13), and a representation for $g(\alpha^u(x, t, \varepsilon), v, x, t, \varepsilon)$ similar to (3.14) we get

$$\begin{aligned} L_\nu \alpha^u(x, t, \varepsilon) &\equiv \varepsilon^2 \left(\frac{\partial \alpha^u}{\partial t} - \frac{\partial^2 \alpha^u}{\partial x^2} \right) - g(\alpha^u(x, t, \varepsilon), v, x, t, \varepsilon) = \\ &\sqrt{\varepsilon} \hat{g}_u(x, t) \left[\sigma + O(\varepsilon^{a-1/2}) + \hat{\varphi}_v(x, t) \sigma^2 \theta \right] h(x, t) \\ &- \frac{1}{2} \sigma^2 \varepsilon \left[\hat{g}_{uu}(x, t) - 2 \hat{g}_{uv}(x, t) \sigma \theta + \hat{g}_{vv}(x, t) \sigma^2 \theta^2 \right] h^2(x, t) - \varepsilon \hat{g}_\varepsilon(x, t) + o(\varepsilon). \end{aligned} \quad (3.22)$$

There is a sufficiently small σ_0 such that for $0 < \sigma \leq \sigma_0$

$$1 + \sigma \hat{\varphi}_v(x, t) \theta \geq 1/2 \quad \text{for } (x, t) \in Q_\nu, \quad |\theta| \leq 1.$$

Thus, because of $a - 1/2 > 0$ and taking into account (2.15), (2.16) and (3.9), we have for sufficiently small ε

$$\hat{g}_u(x, t) \left[\sigma + O(\varepsilon^{a-1/2}) + \hat{\varphi}_v(x, t) \sigma^2 \theta \right] h(x, t) \leq 0. \quad (3.23)$$

By assumption (A₈) there is a positive constant k_g such that for sufficiently small ν

$$-\hat{g}_\varepsilon(x, t) \leq -k_g < 0 \quad \text{for } (x, t) \in Q_\nu.$$

Now we choose σ_0 so small that for $0 < \sigma \leq \sigma_0$

$$\frac{1}{2} \sigma^2 \left| \hat{g}_{uu}(x, t) - 2 \hat{g}_{uv}(x, t) \sigma \theta + \hat{g}_{vv}(x, t) \sigma^2 \theta^2 \right| h^2(x, t) \leq k_g / 2 \quad \text{for } (x, t) \in Q_\nu. \quad (3.24)$$

Therefore, for $0 < \sigma \leq \sigma_0$, and for sufficiently small ε we get from (3.22), (3.23), and (3.24)

$$(L_\nu \alpha^u)(x, t, \varepsilon) \leq 0,$$

i.e. inequality (2.20) is satisfied for α^u in Q_ν .

Now we will prove that α^u and β^u will satisfy the inequalities (2.20), (2.21) in $Q_c \setminus Q_\nu$. From (3.14) we get

$$g(\beta^u(x, t, \varepsilon), v, x, t, \varepsilon) = \sqrt{\varepsilon} \left[\hat{g}_u(x, t) \gamma + \hat{g}_v(x, t) \sigma^2 \theta \right] h(x, t) + o(\sqrt{\varepsilon}). \quad (3.25)$$

It follows from (2.15) that there is a positive constant c_1 such that for sufficiently large γ

$$\hat{g}_u(x, t)\gamma + \hat{g}_v(x, t)\sigma^2\theta \leq -c_1 \quad \text{in } Q_c \setminus Q_\nu. \quad (3.26)$$

Therefore, by (2.18), (3.12), (3.25), and (3.26) we have for γ sufficiently large and ε sufficiently small

$$(L_\nu \beta^u)(x, t, \varepsilon) \geq 0 \quad \text{in } Q_c \setminus Q_\nu.$$

Analogously, we get from (3.22) for σ and ε sufficiently small

$$(L_\nu \alpha^u)(x, t, \varepsilon) = \sqrt{\varepsilon} \hat{g}_u(x, t)(\sigma + \hat{\varphi}_v(x, t)\sigma^2\theta)h(x, t) + o(\sqrt{\varepsilon}) \leq 0 \quad \text{for } (x, t) \in Q_c \setminus Q_\nu.$$

Thus, the inequalities (2.20) for α^u, β^u hold in $Q_c \setminus Q_\nu$.

Now we verify the inequality (2.21) in Q_c . For u we use the representation

$$u = \hat{u}(x, t) + \sqrt{\varepsilon} \kappa h(x, t) + O(\varepsilon^a), \quad -\sigma \leq \kappa \leq \gamma.$$

By (2.19) and (3.8) we have

$$\begin{aligned} (M_u \beta^v)(x, t, \varepsilon) &\equiv \frac{\partial \beta^v}{\partial t} - f(u, \beta^v(x, t, \varepsilon), x, t, \varepsilon) = \frac{\partial \hat{v}}{\partial t} + \sqrt{\varepsilon} \frac{\sigma^2}{\nu} h(x, t) - \\ &\quad - f(\hat{u}(x, t) + \sqrt{\varepsilon} \kappa h(x, t) + O(\varepsilon^a), \hat{v} + \sqrt{\varepsilon} \sigma^2 h(x, t), x, t, \varepsilon). \end{aligned} \quad (3.27)$$

Using the representation

$$\begin{aligned} f(\hat{u}(x, t) + \sqrt{\varepsilon} \kappa h(x, t) + O(\varepsilon^a), \hat{v} + \sqrt{\varepsilon} \sigma^2 h(x, t), x, t, \varepsilon) &= f(\hat{u}, \hat{v}, x, t, 0) + \\ &\quad \sqrt{\varepsilon} [\hat{f}_u(x, t)\kappa + \hat{f}_v(x, t)\sigma^2] h(x, t) + o(\sqrt{\varepsilon}) \end{aligned}$$

and taking into account

$$\frac{\partial \hat{v}}{\partial t} - f(\hat{u}, \hat{v}, x, t, 0) \equiv 0$$

we get from (3.27)

$$(M_u \beta^v)(x, t, \varepsilon) = \sqrt{\varepsilon} \left[\frac{\sigma^2}{\nu} - \hat{f}_u(x, t)\kappa - \hat{f}_v(x, t)\sigma^2 \right] h(x, t) + o(\sqrt{\varepsilon}). \quad (3.28)$$

To given $\sigma > 0$ we choose ν so small such that

$$\left[\frac{\sigma^2}{\nu} - \hat{f}_u(x, t)\kappa - \hat{f}_v(x, t)\sigma^2 \right] h(x, t) \geq c_2 \quad \text{for } (x, t) \in Q_c,$$

where c_2 is some positive number. Thus, for sufficiently small ε , we have

$$(M_u \beta^v)(x, t, \varepsilon) \geq 0 \quad \text{for } (x, t) \in Q_c.$$

Similarly we can verify the inequality (2.21) for α^v .

Finally we verify the inequalities (2.22). If we differentiate β^u with respect to x at $x = 0$ and $x = 1$ respectively we get from (3.8)

$$\begin{aligned}\frac{\partial \beta^u}{\partial x}(0, t, \varepsilon) &= \frac{\partial \tilde{u}}{\partial x}(0, t, \varepsilon) - k + O(\sqrt{\varepsilon}), \\ \frac{\partial \beta^u}{\partial x}(1, t, \varepsilon) &= \frac{\partial \tilde{u}}{\partial x}(1, t, \varepsilon) + k + O(\sqrt{\varepsilon}).\end{aligned}$$

Using (3.5) it can be shown that there exists a positive constant c_3 such that

$$\left| \frac{\partial \tilde{u}}{\partial x}(x, t, \varepsilon) \right| \leq c_3 \quad \text{for } (x, t) \in \overline{Q}.$$

Consequently, the inequalities (2.22) for β^u from Definition 2.1 are satisfied if k is chosen sufficiently large and ε is sufficiently small. The inequalities (2.22) for α^u can be verified in a similar way.

From our considerations above it follows that the functions $\alpha(x, t, \varepsilon), \beta(x, t, \varepsilon)$ fulfil all conditions of Definition 2.1, and we can conclude that for sufficiently small ε there exists a unique solution $(u(x, t, \varepsilon), v(x, t, \varepsilon))$ of problem (2.1) satisfying for $(x, t) \in \overline{Q}_c$

$$\begin{aligned}\alpha^u(x, t, \varepsilon) &\leq u(x, t, \varepsilon) \leq \beta^u(x, t, \varepsilon), \\ \alpha^v(x, t, \varepsilon) &\leq v(x, t, \varepsilon) \leq \beta^v(x, t, \varepsilon).\end{aligned}$$

From these inequalities and from (3.8) it follows that the representations (3.2) and (3.3) for $u(x, t, \varepsilon)$ and $v(x, t, \varepsilon)$ in \overline{Q}_c are valid. This completes the proof of Theorem 3.1.

□

4 Example.

Consider the initial boundary value problem

$$\begin{aligned}\varepsilon^2 \left(\frac{\partial u}{\partial t} - \frac{\partial^2 u}{\partial x^2} \right) &= g(u, v, x, t, \varepsilon) \equiv -u(u - v + x + t + 2) + \varepsilon I(x, t), \\ \frac{\partial v}{\partial t} &= f(u, v, x, t, \varepsilon) \equiv u + 2 \\ (x, t) \in Q &:= \{(x, t) \in R^2 : 0 < x < 1, 0 < t \leq T\}, \quad T > 2, \\ \frac{\partial u}{\partial x}(0, t, \varepsilon) &= \frac{\partial u}{\partial x}(1, t, \varepsilon) = 0 \quad \text{for } 0 < t \leq T, \\ u(x, 0, \varepsilon) = u^0(x) &> 0, \quad v(x, 0, \varepsilon) = v^0(x) \equiv 1 \quad \text{for } 0 \leq x \leq 1,\end{aligned}\tag{4.1}$$

where $I : \overline{Q} \rightarrow R$ is smooth and positive, u^0 is a smooth function on $0 \leq x \leq 1$. The degenerate equation

$$-u(u - v + x + t + 2) = 0$$

has two solutions

$$u = \varphi_1(v, x, t) \equiv 0 \quad \text{and} \quad u = \varphi_2(v, x, t) \equiv v - x - t - 2 \quad (4.2)$$

intersecting in a smooth surface with the representation

$$v = s(x, t) \equiv x + t + 2.$$

The inequality $\varphi_1(v, x, t) > (<) \varphi_2(v, x, t)$ holds for $v < (>) s(x, t)$, $(x, t) \in \overline{Q}$, i.e. the assumptions (A₁) and (A₂) are fulfilled.

From (4.1) and (4.2) we get

$$g_u(\varphi_1(v, x, t), x, t, 0) \equiv v - x - t - 2 \equiv -g_u(\varphi_2(v, x, t), x, t, 0).$$

Obviously we have for $(x, t) \in \overline{Q}$

$$\begin{aligned} g_u(\varphi_1(v, x, t), x, t, 0) &< 0, \quad g_u(\varphi_2(v, x, t), x, t, 0) > 0 \quad \text{for } v < s(x, t), \\ g_u(\varphi_1(v, x, t), x, t, 0) &> 0, \quad g_u(\varphi_2(v, x, t), x, t, 0) < 0 \quad \text{for } v > s(x, t), \end{aligned}$$

i.e. assumption (A₃) holds.

Note that $1 \equiv v^0(x) < s(x, 0) = x + 2$ for $x \in [0, 1]$ and $f(\varphi_1(v, x, t), v, x, t, 0) \equiv 2$. Therefore, the initial value problem for $v_1(x, t)$ reads

$$\frac{dv_1}{dt} = 2, \quad 0 < t \leq T; \quad v_1(x, 0) = 1.$$

It has the solution

$$v_1(x, t) = 2t + 1.$$

The equation

$$v_1(x, t) = s(x, t), \quad \text{i.e. } 2t + 1 = x + t + 2$$

defines the curve \mathcal{C} :

$$t = t_c(x) = x + 1.$$

It is obvious that

$$v_1(x, t) < s(x, t) \quad \text{for } 0 \leq t < t_c(x)$$

and

$$v_1(x, t) > s(x, t) \quad \text{for } t_c(x) < t \leq T,$$

i.e. assumption (A₄) is fulfilled.

From $f(\varphi_2(v, x, t), v, x, t, 0) \equiv v - x - t$ and $v_1(x, t_c(x)) = 2x + 3$ it follows that the initial value problem for $v_2(x, t)$ reads

$$\frac{dv_2}{dt} = v_2 - x - t, \quad v_2(x, t_c(x)) = 2x + 3.$$

Its solution is

$$v_2(x, t) = \exp(t - x - 1) + x + t + 1.$$

It is easy to check that

$$v_2(x, t) > s(x, t) \quad \text{for } t_c(x) < t \leq T \quad (\text{i.e. in } Q_2),$$

and

$$v_2(x, t) < s(x, t) \quad \text{for } 0 \leq t < t_c(x) \quad (\text{i.e. in } Q_1).$$

Therefore, assumption (A₅) holds.

In our example the composed stable solution has the form

$$\hat{u}(x, t) = \begin{cases} \psi_1(x, t) \equiv 0 & \text{in } \overline{Q}_1, \\ \psi_2(x, t) \equiv \exp(t - x - 1) - 1 & \text{in } \overline{Q}_2, \end{cases} \quad (4.3)$$

$$\hat{v}(x, t) = \begin{cases} v_1(x, t) \equiv 2t + 1 & \text{in } \overline{Q}_1, \\ v_2(x, t) \equiv \exp(t - x - 1) + x + t + 1 & \text{in } \overline{Q}_2. \end{cases} \quad (4.4)$$

Now we verify the hypotheses (A₆) - (A₈). The associated equation (2.5) for $v = v^0(x) \equiv 1, t = 0$ reads

$$\frac{du}{d\tau} = -u(u + x + 1), \quad \tau > 0.$$

It is easy to see that the solution $\bar{u}(x, \tau)$ of this equation with the initial condition

$$\bar{u}(x, 0) = u^0(x) > 0$$

exists for $\tau > 0$ and tends to $\varphi_1(v^0(x), x, 0) = 0$ as $\tau \rightarrow \infty$. Hence, assumption (A₆) is fulfilled.

Moreover, the solution $\Pi_0(x, \tau)$ of problem (3.1) which reads in our case

$$\frac{d\Pi_0}{d\tau} = -\Pi_0(\Pi_0 + x + 1), \quad \tau > 0; \quad \Pi_0(x, 0) = u^0(x)$$

can be found in the explicit form

$$\Pi_0(x, \tau) = u^0(x)(x + 1) \left[u^0(x)(1 - \exp(-(x + 1)\tau) + x + 1 \right]^{-1} \exp(-(x + 1)\tau). \quad (4.5)$$

Assumptions (A₇) and (A₈) are obviously satisfied since

$$g_{uu} \equiv -2 < 0 \quad \text{in } \overline{Q} \quad \text{and} \quad g_\varepsilon \equiv I(x, t) > 0 \quad \text{in } \overline{Q}.$$

Thus, all assumptions (A₁)-(A₈) of Theorem 3.1 are fulfilled. Therefore, the initial boundary value problem (2.1) for our example has a unique solution $(u(x, t, \varepsilon), v(x, t, \varepsilon))$ satisfying (3.2), (3.3) where $\hat{u}(x, t), \hat{v}(x, t)$, and $\Pi_0(x, \tau)$ are defined in (4.3), (4.4) and (4.5) respectively.

5 Acknowledgment

The authors acknowledge the financial support by a RFFI-DFG grant.

References

- [1] V.F. BUTUZOV AND N.N. NEFEDOV, *Singularly perturbed boundary value problems for a second order equation in case of exchange of stability* (in Russian), *Mat. Zamet*, **63** (1998), pp. 354-362.
- [2] V.F. BUTUZOV, N.N. NEFEDOV AND K. R. SCHNEIDER, *Singularly perturbed boundary value problems in case of exchange of stabilities*, *J. Math. Anal. Appl.*, **229** (1999), pp. 543-562.
- [3] V.F. BUTUZOV AND I. SMUROV, *Initial boundary value problem for singularly perturbed parabolic equation in case of exchange of stability*, *J. Math. Anal. Appl.*, **234** (1999), pp. 183-192.
- [4] P. FABRIE, C. GALUSINSKI, *Exponential attractors for a partially dissipative reaction system*, *Asymptotic Anal.*, **12** (1996), pp. 329-354.
- [5] S.L. HOLLIS, J.J. MORGAN, *Partly dissipative reaction-diffusion systems and a model of phosphorus diffusion in silicon*, *Nonlinear Anal. Theory Methods Appl.*, **19** (1992), pp. 427-440.
- [6] M. MARION, *Inertial manifolds associated to partly dissipative reaction-diffusion systems*, *J. Math. Anal. Appl.*, **143** (1989), pp. 295-326.
- [7] M. MARION, *Finite-dimensional attractors associated with partly dissipative reaction-diffusion systems*, *SIAM J. Math. Anal.*, **20** (1989), pp. 816-844.
- [8] N.N. NEFEDOV, K.R. SCHNEIDER, *Immediate exchange of stabilities in singularly perturbed systems*, *Differential and Integral Equations*, **12** (1999), pp. 583-599.
- [9] N.N. NEFEDOV, K.R. SCHNEIDER, A. SCHUPPERT, *Jumping behavior in singularly perturbed systems modelling bimolecular reactions*, Weierstraß-Institut für Angewandte Analysis und Stochastik Berlin, Preprint No. 137, Berlin 1994.
- [10] C.V. PAO, *Nonlinear Parabolic and Elliptic Equations*, Plenum Press, New York and London, 1992.
- [11] K.R. SCHNEIDER, *On the existence of wave trains in partly dissipative systems*, *Proc. Int. Conf. Differential Equations*. Ed. by C. Perello, C. Simo, J. Sola-Morales. World Scientific Publ. Singapore 1993, vol. 2, pp. 893-898.
- [12] Z. SHAO, *Existence of inertial manifolds for partly dissipative reaction diffusion systems in higher space dimensions*, *J. Differ. Equations*, **144** (1998), pp. 1-43.
- [13] A.N. TIKHONOV, *Systems of differential equations containing small parameters* (in Russian), *Mat. Sb.*, **73** (1952), 575-586.
- [14] A.B. VASIL'EVA, V.F. BUTUZOV AND L.V. KALACHEV, *The boundary function method for singular perturbation problems*, SIAM, Philadelphia, 1995.