

Modeling boundary conditions for solving stationary Schrödinger equations

X. ANTOINE¹, C. BESSE², M. EHRHARDT³, P. KLEIN¹

¹Institut Elie Cartan Nancy, Nancy-Universität, CNRS UMR 7502, INRIA CORIDA Team, Boulevard des Aiguillettes B.P. 239, 54506 Vandoeuvre-lès-Nancy, France,
{Xavier.Antoine, Pauline.Klein}@iecn.u-nancy.fr

²Equipe Projet Simpaf - Inria CR Lille Nord Europe, Laboratoire Paul Painlevé, Unité Mixte de Recherche CNRS (UMR 8524), UFR de Mathématiques Pures et Appliquées, Université des Sciences et Technologies de Lille, Cité Scientifique, 59655 Villeneuve d'Ascq Cedex, France,
Christophe.Besse@math.univ-lille1.fr

³Lehrstuhl für Angewandte Mathematik und Numerische Analysis, Fachbereich C – Mathematik und Naturwissenschaften, Bergische Universität Wuppertal, Gaußstr. 20, 42119 Wuppertal, Germany,
ehrhardt@math.uni-wuppertal.de

Abstract

In this talk we present some novel absorbing boundary conditions (ABCs) for modeling the solution of linear and nonlinear variable potentials one-dimensional stationary Schrödinger equations. Using pseudodifferential calculus and factorization theorems we construct a hierarchy of novel ABCs and generalize the well-known quantum transmitting boundary condition of Kirk and Lentner to the case of space-dependent potential. Moreover, we propose a rapidly converging iterative method based on finite elements suitable for computing scattering solutions and bound states.

The accuracy of our new absorbing boundary conditions is investigated numerically for two different situations. The first problem is related to the computation of linear scattering problems. The second application concerns the computation of energies and ground-states for linear and nonlinear Schrödinger equations. It turns out that these absorbing boundary conditions and their variants lead to a higher accuracy than the usual Dirichlet boundary condition. Finally, our approach also offers the possibility to construct ABCs for higher dimensional problems.

References

- [1] X. Antoine, A. Arnold, C. Besse, M. Ehrhardt, A. Schädle, *A review of transparent and artificial boundary conditions techniques for linear and nonlinear Schrödinger equations*, Commun. Comput. Phys. **4** (2008), 729–796. (open-access article)
- [2] X. Antoine, C. Besse and P. Klein, *Absorbing boundary conditions for the one-dimensional Schrödinger equation with an exterior repulsive potential*, J. Comput. Phys. **228** (2009), 312–335.
- [3] X. Antoine, C. Besse, M. Ehrhardt and P. Klein, *Modeling boundary conditions for solving stationary Schrödinger equations*, Preprint 10/04 of the chairs of Applied Mathematics & Numerical Analysis and Optimization & Approximation, University of Wuppertal, February 2010.
- [4] P. Klein, X. Antoine, C. Besse and M. Ehrhardt, *Absorbing boundary conditions for solving stationary Schrödinger equations*, Proceedings of the 16th European Conference on Mathematics for Industry, July 26-30, 2010, Wuppertal, Germany, Springer, Berlin, Heidelberg, 2011.
- [5] P. Klein, X. Antoine, C. Besse and M. Ehrhardt, *Absorbing boundary conditions for solving N-dimensional stationary Schrödinger equations with unbounded potentials and nonlinearities*, Commun. Comput. Phys. **10** (2011), 1280–1304.