

Department for Mathematics and Computer Science
Freie Universität Berlin
Prof. Dr. V. John, john@wias-berlin.de
Marwa Zainelabdeen, marwazinabdeen@gmail.com

Berlin, 07.11.2022

Numerical Mathematics II

Exercise Problems 04

Attention: The approach for getting a solution has to be clearly presented. All statements have to be proved, auxiliary calculations have to be written down. Statements given in the lectures can be used without proof.

1. *Embedded Runge–Kutta scheme with two stages.* Derive the embedded explicit Runge–Kutta scheme $p(q) = 1(2)$ with two stages and the condition $a_{21} = b_1$, i.e., write the Butcher tableau in terms of c_2 . Which schemes are obtained in the special case $c_2 = 1$? **3 points**
2. *Convergence of damped Jacobi method.* Proof the following statement: If the Jacobi method converges for each initial iterate, then also the damped Jacobi method with $0 < \omega \leq 1$ converges for each initial iterate. **3 points**
3. *Optimal relaxation parameter for SOR method.* Continue Problem 3, Exercise sheet 03. One can show that the optimal relaxation parameter for the SOR method is

$$\omega_{\text{opt}} = \frac{2}{1 + \sqrt{1 - \rho^2}},$$

where ρ is the spectral radius of the iteration matrix of the Jacobi method. In the considered example, one finds that

$$\omega_{\text{opt}} = \frac{2}{1 + \sin(\pi h)}.$$

Use this relaxation parameter in the code from Problem 3, Exercise sheet 03. How does the optimal parameter behave if h decreases? Compare the number of iterations with the numbers obtained for the other relaxation parameters from the solution of Problem 3, Exercise sheet 03. What can be observed?

4 points

The exercise problems should be solved in groups of four to five students. The solutions have to be submitted until **Monday, Nov. 14, 2022, 10:00 a.m.** via the whiteboard.