Fachbereich Mathematik und Informatik/ Institut für Mathematik Prof. Dr. Volker John, john@wias-berlin.de Joshua Röber, roebej98@zedat.fu-berlin.de

Berlin, 20.06.2022

Numerik I

English translation of Übungsserie 09

Attention: Only solutions which provide a comprehensible reasoning will be graded. Every statement has to be argued. You can use results from the lecture. Statments without reasoning won't get any points.

1. Absolute condition for the computation of roots. Let $f \in C^n([a, b])$ and $x_0 \in (a, b)$ a root of

$$f(x) - \alpha = 0, \quad \alpha \in \mathbb{R},$$

with multiplicity n but not (n+1).

(a) Consider a small perturbation ε of the data α and prove the following estimate for the absolute condition K

$$K \approx \left| \frac{n!}{f^{(n)}(x_0)} \right| \varepsilon^{1/n-1}.$$

What does this imply for arbitrary small perturbations?

(b) The computation of roots of polynomials is a special case of the above mentioned assignment. This is for example necessary if one want to compute the Eigenvalues of a matrix via the characteristic polynomial. Let f(x) a polynomial of *n*-th degree and root x_0 with multiplicity *n*. How can you improve the statement of the first part.

Hint: Use Taylor expansion of f(x) at x_0 and approximate the term including the unknown intermediate point by the same term at x_0 . 4 points

2. Gershgorin circles. Estimate the Eigenvalues of

,

$$A = \begin{pmatrix} 2 & -1/2 & 0 & -1/2 \\ 0 & 4 & 0 & 2 \\ -1/2 & 0 & 6 & 1/2 \\ 0 & 0 & 1 & 9 \end{pmatrix}$$

by the means of the Gershgorin circles.

3. *Programming exercise: Power iteration*. Implement the Power iteration and approximate the Eigenvalue with largest absolute value of the following matrix

$$A(\alpha) = \begin{pmatrix} \alpha & 2 & 3 & 13\\ 5 & 11 & 10 & 8\\ 9 & 7 & 6 & 12\\ 4 & 14 & 15 & 1 \end{pmatrix}$$

for $\alpha \in \{-30, 30\}$. As initial guess one is supposed to use the unit vector e_1 . The iteration should be stopped if one obtains the following bound on the relative decay of the iterate

$$\frac{|\lambda^{(k+1)} - \lambda^{(k)}|}{|\lambda^{(k+1)}|} < 10^{-10}.$$

2 points

How many iterations are necessary? Which statement from the lectures can explain the difference in the number of iterations. 4 points

The exercises should be solved in groups of three or four students. They have to be submitted until Sie Monday, 27.06.2022, 12:00, either in the box of the tutor or electronically.