

# EFFICIENT AND ACCURATE LOG-LÉVY APPROXIMATIONS TO LÉVY DRIVEN LIBOR MODELS

ANTONIS PAPAPANTOLEON, JOHN SCHOENMAKERS,  
AND DAVID SKOVMAND

ABSTRACT. The LIBOR market model is very popular for pricing interest rate derivatives, but is known to have several pitfalls. In addition, if the model is driven by a jump process, then the complexity of the drift term is growing exponentially fast (as a function of the tenor length). In this work, we consider a Lévy-driven LIBOR model and aim at developing accurate and efficient log-Lévy approximations for the dynamics of the rates. The approximations are based on truncation of the drift term and Picard approximation of suitable processes. Numerical experiments for FRAs, caps and swaptions show that the approximations perform very well. In addition, we also consider the log-Lévy approximation of annuities, which offers good approximations for high volatility regimes.

## 1. INTRODUCTION

The LIBOR market model (LMM) has become a standard model for the pricing of interest rate derivatives in recent years, because the evolution of discretely compounded, market-observable forward rates is modeled directly and not deduced from the evolution of unobservable factors, as is the case in short rate and forward rate (HJM) models. See Miltersen et al. (1997), Brace et al. (1997) and Jamshidian (1997) for the seminal papers in LIBOR modeling. In addition, the lognormal LIBOR model provides a theoretical justification to the market practice of pricing caps according to Black's formula (cf. Black 1976). However, despite its apparent popularity, the LIBOR market model has certain well-known pitfalls.

An interest rate model is typically calibrated to the implied volatility surface from the cap market and the correlation structure of at-the-money swaptions. The implied volatility from caplets has a "smile" shape as a function of strike, while its term structure is typically decreasing. The standard lognormal LMM cannot be calibrated adequately to the observed market data. Therefore, several extensions of the LMM have been proposed in the literature using jump-diffusions, Lévy processes or general semimartingales as the driving motion (cf. e.g. Glasserman and Kou 2003, Eberlein and Özkan 2005, Jamshidian 1999), or incorporating stochastic volatility effects (cf. e.g. Andersen and Brotherton-Ratcliffe 2005, Wu and Zhang 2006, Belomestny, Mathew, and Schoenmakers 2009).

---

2000 *Mathematics Subject Classification.* 91G30, 91G60, 60G51.

*Key words and phrases.* LIBOR market model, Lévy processes, drift term, Picard approximation, option pricing, caps, swaptions, annuities.

J. S. acknowledges the financial support by the DFG Research Center MATHEON "Mathematics for Key Technologies" in Berlin.

The dynamics of LIBOR models are typically not tractable under different forward measures, due to the random terms that enter the dynamics of LIBOR rates. In particular, LIBOR rates are tractable under their corresponding forward measure only in the lognormal setting or simple perturbations thereof, for example displaced diffusion, CEV or extended with an uncorrelated volatility process. If the driving process is a general diffusion process or semimartingale, then the dynamics of LIBOR rates are usually not tractable even under their own forward measures. Consequently, even caplets cannot be priced exactly in “closed form” (meaning, e.g. by Fourier methods), let alone swaptions and other multi-LIBOR products. In order to calibrate the model, closed form solutions are necessary, and these are typically involving approximations.

The standard approximation is the so-called “frozen drift” approximation; it was first proposed by Brace et al. (1997) for the pricing of swaptions and has been used by several authors ever since. The frozen drift approximation typically leads to closed-form solutions for caplet pricing in realistic LIBOR models, see Eberlein and Özkan (2005) and Belomestny et al. (2009). Although some authors (Brace et al. 2001, Dun et al. 2001 and Schlögl 2002) argue that freezing the drift is justified in the lognormal LMM, it is well-known that it does not yield acceptable results in more advanced LIBOR model extensions, especially for exotic derivatives and longer time horizons. See also the numerical experiments in section 5.

Several alternative approximations have been developed in the literature. In one line of research, Kurbanmuradov, Sabelfeld, and Schoenmakers (2002) and Daniluk and Gałtarek (2005) have derived lognormal approximations to the forward LIBOR dynamics (for deterministic volatility structures). Other authors have been using linear interpolations and predictor-corrector Monte Carlo methods to get a more accurate discretization of the drift term (cf. e.g. Hunter et al. 2001 and Glasserman and Zhao 2000). We refer the reader to Joshi and Stacey (2008) and Gałtarek, Bachert, and Maksymiuk (2006, Ch. 10) for a detailed overview of that literature, some new approximation schemes and numerical experiments. Although most of this literature focuses on the lognormal LIBOR market model, Glasserman and Merener (2003b, 2003a) have developed approximation schemes for the pricing of caps and swaptions in jump-diffusion LIBOR market models, based on freezing the drift.

In this article, we consider a LIBOR market model driven by a Lévy process and aim at deriving efficient and accurate log-Lévy approximations. As a main result, we develop accurate log-Lévy LIBOR approximations which may be represented as a *deterministic* drift term plus a stochastic integral of a deterministic function with respect to a Lévy process. In particular, in the context of Monte Carlo simulation the drift term can be computed outside the Monte Carlo loop, while the stochastic integrals can be computed efficiently for each trajectory. In contrast, standard Euler stepping of the original LIBOR SDE involves, for each LIBOR trajectory, an accurate computation of a complex-structured random drift term at each Euler step and

is therefore significantly more time-consuming<sup>1</sup>. Theoretical investigations as well as numerical experiments show that the log-Lévy approximations are both fast and accurate when the LIBOR volatilities are not too high, and thus provide an effective alternative to simulation methods based on standard Euler discretizations. Finally, as a generalization of Gatarek et al. (2006), we derive log-Lévy approximations for annuity terms, which allow for pricing options in high volatility regimes.

The article is structured as follows: in section 2 we review the Lévy-driven LIBOR model, in section 3 we construct the log-Lévy approximations to the model and in section 4 we provide some error estimates. Section 5 demonstrates numerically the effect of the approximations, while section 6 deals with approximation of annuities. The final section provides some recommendations on the construction of multi-dimensional Lévy LIBOR models, while the appendices collect various calculations.

## 2. LÉVY LIBOR FRAMEWORK

Let  $0 = T_0 < T_1 < \dots < T_N < T_{N+1} = T_*$  denote a discrete tenor structure where  $\delta_i = T_{i+1} - T_i$ ,  $i = 0, 1, \dots, N$ , are the so called day-count fractions. For this tenor structure we consider an arbitrage free system of zero coupon bond processes  $B_i$ ,  $i = 1, \dots, N + 1$ , on a filtered probability space  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{0 \leq t \leq T_*}, \mathbb{P}_*)$ , where  $\mathbb{P}_* := \mathbb{P}_{N+1}$  is a numeraire measure connected with the terminal bond  $B_{N+1}$ . From this bond system we may deduce a forward rate system, also called LIBOR rate system, defined by

$$L_i(t) := \frac{1}{\delta_i} \left( \frac{B_i(t)}{B_{i+1}(t)} - 1 \right), \quad 0 \leq t \leq T_i, \quad 1 \leq i \leq N. \quad (2.1)$$

$L_i$  is the annualized effective forward rate contracted at date  $t \leq T_i$  for the period  $[T_i, T_{i+1}]$ . Jamshidian (1999) derived a general representation for the LIBOR dynamics in a semimartingale framework. In this article we consider a Lévy LIBOR framework as constructed by Eberlein and Özkan (2005); see also Glasserman and Kou (2003) and Belomestny and Schoenmakers (2011) for jump-diffusion settings.

Consider a standard Brownian motion  $W$  in  $\mathbb{R}^m$ ,  $m \leq N$ , a bounded deterministic nonnegative scalar function  $\alpha(s)$ ,  $s \in [0, T_*]$ , and a random measure  $\mu$  on  $[0, T_*] \times \mathbb{R}^m$  with  $\mathbb{P}_*$ -compensator  $F(s, dx)ds$ , where  $\mu$  and  $W$  are mutually independent. Let  $H = (H(t))_{0 \leq t \leq T_*}$  be a time-inhomogeneous Lévy process with canonical decomposition

$$H(t) = \int_0^t \sqrt{\alpha(s)} dW(s) + \int_0^t \int_{\mathbb{R}^m} x(\mu(ds, dx) - F(s, dx)ds). \quad (2.2)$$

We denote by  $\tilde{\mu}$  the compensated random measure of the jumps of  $H$ , that is  $\tilde{\mu}(ds, dx) := \mu(ds, dx) - F(s, dx)ds$ . In order to avoid truncation conventions

---

<sup>1</sup>In a previous unpublished manuscript by the first and third author (Papapantoleon and Skovmand 2010) the efficiency of the standard Euler approach was improved to some extent also, but there was still a costly random drift involved.

we assume that  $F$  satisfies the (stronger than usual) integrability condition

$$\int_0^{T_*} \int_{\mathbb{R}^m} (\|x\| \wedge \|x\|^2) F(s, dx) ds < \infty.$$

We further assume that

$$\int_0^{T_*} \int_{\|x\|>1} \exp(u^\top x) F(s, dx) ds < \infty, \quad (2.3)$$

for all  $\|u\| \leq (1 + \varepsilon)\overline{M}$ , with  $\overline{M}, \varepsilon > 0$  constants. Thus, by construction, the process  $(H(t))_{0 \leq t \leq T_*}$  is a  $\mathbb{P}_*$ -martingale. The cumulant generating function of  $H(t)$ ,  $t \in [0, T_*]$ , is provided by

$$\ln \mathbb{E}[e^{u^\top H(t)}] = \kappa_t(u) = \frac{\alpha(t)}{2} \|u\|^2 + \int_{\mathbb{R}^m} (e^{u^\top x} - 1 - u^\top x) F(t, dx). \quad (2.4)$$

Along with the Lévy martingale (2.2) we introduce a set of bounded deterministic vector-valued functions  $\lambda_i(s) \in \mathbb{R}^m$ ,  $i = 1, \dots, N$ , usually called *loading factors*. In order to avoid local redundances we assume that the matrix  $[\lambda_1, \dots, \lambda_N](s)$  has full rank  $m$  for all  $s \in [0, T_*]$ . Moreover, we assume that  $\|\lambda_i(s)\| \leq \overline{M}$ , for all  $i$ , and  $\|\sum_i \lambda_i(s)\| \leq \overline{M}$ , for all  $s \in [0, T_*]$ .

The Lévy martingale and the set of loading factors then constitute an arbitrage free LIBOR system consistent with (2.1), whose dynamics under the terminal measure  $\mathbb{P}_*$  are given by

$$L_i(t) = L_i(0) \exp \left( \int_0^t b_i(s) ds + \int_0^t \lambda_i^\top(s) dH(s) \right), \quad (2.5)$$

$i = 1, \dots, N$ , where the drift terms in the exponent are given by

$$\begin{aligned} b_i = & -\frac{1}{2} \alpha |\lambda_i|^2 - \sum_{j=i+1}^N \frac{\delta_j L_{j-}}{1 + \delta_j L_{j-}} \alpha \lambda_i^\top \lambda_j \\ & - \int_{\mathbb{R}^m} \left( (e^{\lambda_i^\top x} - 1) \prod_{j=i+1}^N \left( 1 + \frac{\delta_j L_{j-} (e^{\lambda_j^\top x} - 1)}{1 + \delta_j L_{j-}} \right) - \lambda_i^\top x \right) F(\cdot, dx); \end{aligned} \quad (2.6)$$

for details see Eberlein and Özkan (2005). For notational convenience, we set  $L_{j-}(s) := L_j(s-)$  in (2.6), while the time variable is suppressed.

Due to the drift term (2.6), a straightforward Monte Carlo simulation of (2.5) would involve a numerical integration at each time step, since the random terms  $\frac{\delta_j L_{j-}}{1 + \delta_j L_{j-}}$  appear under the integral sign. In order to overcome this problem, we will re-express the drift in terms of random quotients multiplied

with cumulants of the driving process. We have that

$$\begin{aligned}
b_i &= -\kappa(\lambda_i) - \sum_{j=i+1}^N \frac{\delta_j L_{j-}}{1 + \delta_j L_{j-}} \alpha \lambda_i^\top \lambda_j \\
&\quad - \sum_{p=1}^{N-i} \sum_{i < j_1 < \dots < j_p \leq N} \frac{\delta_{j_1} L_{j_1-}}{1 + \delta_{j_1} L_{j_1-}} \dots \frac{\delta_{j_p} L_{j_p-}}{1 + \delta_{j_p} L_{j_p-}} \\
&\quad \times \sum_{q=1}^{p+1} (-1)^{p+q+1} \sum_{0 \leq r_1 < \dots < r_q \leq p} \widehat{\kappa}(\lambda_{j_{r_1}} + \dots + \lambda_{j_{r_q}}); \quad (2.7)
\end{aligned}$$

the derivation is deferred to Appendix A, for brevity. Here  $\widehat{\kappa}$  denotes the part of the cumulant  $\kappa$  stemming from the jumps of  $L$ , that is

$$\widehat{\kappa}_s(u) = \int_{\mathbb{R}^m} (e^{u^\top x} - 1 - u^\top x) F(s, dx). \quad (2.8)$$

Therefore, we can now avoid the numerical integration when simulating LIBOR rates. However, another problem becomes apparent in this representation: the number of terms to be computed in (2.7) grows exponentially fast as a function of the number of LIBOR rates  $N$ , namely it has order  $O(2^N)$ .

**Remark 2.1.** In a practically applicable model, the loading factors  $\lambda_i$  may be decomposed as follows:

$$\begin{aligned}
\lambda_i(t) &= c_i g(T_i - t) e_{i-m(t)} \in \mathbb{R}^m, \\
m(t) &:= \inf\{i : T_i \geq t\}, \quad \|e_i\| = 1, \quad e_i^\top e_j = \rho_{ij}, \quad 1 \leq i, j \leq N,
\end{aligned}$$

for constants  $c_i > 0$ , some (e.g. parametric) scalar function  $g > 0$ , and a correlation structure  $(\rho_{ij})$  which resembles the correlations between forward LIBORs observed in the market. For instance,  $(\rho_{ij})$  may be obtained as a rank- $m$  approximation of a suitably parameterized full rank- $N$  correlation structure; see Schoenmakers (2005) for details. Further, the scalar function  $\alpha$  may be taken as a constant that controls the influence of the Wiener noise with respect to the jump noise.

**Remark 2.2.** The Lévy-driven LIBOR model is constructed under the terminal measure  $\mathbb{P}_{N+1}$  in this paper, for definiteness. As an alternative, for products with shorter maturity for instance, one may consider for some  $T_{\widetilde{N}} < T_{N+1}$ , a Lévy-driven LIBOR model for  $t \leq T_{\widetilde{N}}$  under the measure  $\mathbb{P}_{\widetilde{N}}$ , with respect to the numeraire bond  $B_{\widetilde{N}}$ . Another possibility is to consider as numeraire the spot LIBOR rolling over account

$$\begin{aligned}
B_o(0) &:= 1, \quad B_o(t) := \frac{B_{m(t)}(t)}{B_1(0)} \prod_{i=1}^{m(t)-1} (1 + \delta_i L_i(T_i)), \\
m(t) &:= \min\{m : T_m \geq t\}, \quad 0 < t \leq T_{N+1},
\end{aligned}$$

and the numeraire measure  $\mathbb{P}_o$  associated with it. If one prefers to work in one of these other measures, the drift term (2.6) has to be modified in the following way: for the Libor model in the measure  $\mathbb{P}_{\widetilde{N}}$ , replace in (2.6), if  $i \leq \widetilde{N}$ , the sum  $-\sum_{j=i+1}^N$  and the product  $\prod_{j=i+1}^N$  by  $-\sum_{j=i+1}^{\widetilde{N}-1}$  and  $\prod_{j=i+1}^{\widetilde{N}-1}$

respectively, and if  $i > \tilde{N}$ , by  $\sum_{j=\tilde{N}}^i$  and  $1/\prod_{j=\tilde{N}}^i$  respectively. Likewise, for a LIBOR model in the measure  $\mathbb{P}_\circ$ , replace in (2.6)  $-\sum_{j=i+1}^N$  by  $\sum_{j=m(t)}^i$  and the product  $\prod_{j=i+1}^N$  by  $1/\prod_{j=m(t)}^i$ . We refer to Jamshidian (1999) for more details. The proper choice of a numeraire measure under which the Lévy-driven LIBOR model is constructed may depend on the set of LIBORs involved in a particular (structured) product which has to be evaluated by simulation. In principle, one should choose the measure in such a way that the respective sum and product in the drift (2.6) involve as few terms as possible.

### 3. EFFICIENT AND ACCURATE LOG-LÉVY APPROXIMATIONS

The aim of this section is to derive efficient and accurate log-Lévy approximations for the dynamics of the LIBOR rates under the terminal measure. This is based on an appropriate approximation of the drift term, cf. (2.6), which has two pillars:

- (1) expansion and truncation of the drift term,
- (2) Picard approximation of suitably defined processes.

We will first provide an overview of the approximation argument, and then present the full details in some particular cases.

**3.1. Outline of the method.** Let us denote the log-LIBOR rates by  $G_i$ . They are defined via

$$G_i(t) := \log L_i(t),$$

and satisfy the integrated linear SDE, see (2.5),

$$G_i(t) = G_i(0) + \int_0^t b_i(s) ds + \int_0^t \lambda_i^\top(s) dH(s), \quad (3.1)$$

$0 \leq t \leq T_i$ ,  $1 \leq i \leq N$ . The semimartingale characteristics of  $G_i$  are

$$\begin{aligned} B^i &= \int_0^\cdot b_i(s) ds \\ C^i &= \int_0^\cdot |\lambda_i|^2(s) \alpha(s) ds \\ \int_0^\cdot \int_{\mathbb{R}} 1_A(x) F^i(s, dx) ds &= \int_0^\cdot \int_{\mathbb{R}^m} 1_A(\lambda_i^\top(s)x) F(s, dx) ds, \end{aligned} \quad (3.2)$$

where  $A \in \mathcal{B}(\mathbb{R} \setminus \{0\})$ .

Inspired by the lognormal approximation developed by Kurbanmuradov et al. (2002) in the context of the lognormal LIBOR market model, we will derive log-Lévy approximations for the dynamics of  $L_i$ , or equivalently Lévy approximations for the dynamics of  $G_i$ . The standard remedy for the numerical problems arising in LMMs is to “freeze the drift”, that is to replace the random terms in (2.6) – or (2.7) – by their deterministic initial values. In the present model, this obviously leads to a log-Lévy approximation, which however is not accurate enough.

The method for deriving *efficient* and *accurate* log-Lévy approximations we propose can be summarized in the following steps:

- consider the different product terms  $\frac{\delta_{j_1} L_{j_1}}{1+\delta_{j_1} L_{j_1}} \cdots \frac{\delta_{j_p} L_{j_p}}{1+\delta_{j_p} L_{j_p}} =: X_{j_1 \dots j_p}$  in (2.7), where  $i+1 \leq j_1 < \cdots < j_p \leq N$ ;
- define functions  $h : \mathbb{R}^{j_p} \rightarrow \mathbb{R}$  such that

$$h(G_{j_1}, \dots, G_{j_p}) = X_{j_1 \dots j_p};$$

- apply Itô's formula to  $X_{j_1 \dots j_p}$ , which leads to an SDE of the form

$$\begin{aligned} dX_{j_1 \dots j_p}(s) &= A_{j_1 \dots j_p}(s, L(s))ds + B_{j_1 \dots j_p}(s, L(s))^\top dW(s) \\ &\quad + \int_{\mathbb{R}^m} C_{j_1 \dots j_p}(s, x, L(s)) \tilde{\mu}(ds, dx), \end{aligned} \quad (3.3)$$

with  $L = [L_1, \dots, L_N]$ ;

- use the first step of a Picard iteration to approximate  $X_{j_1 \dots j_p}$  by the Lévy process

$$\begin{aligned} X_{j_1 \dots j_p}^{(1)}(t) &= X_{j_1 \dots j_p}(0) + \int_0^t A_{j_1 \dots j_p}(s, L(0))ds \\ &\quad + \int_0^t B_{j_1 \dots j_p}(s, L(0))^\top dW_s + \int_0^t \int_{\mathbb{R}^m} C_{j_1 \dots j_p}(s, x, L(0)) \tilde{\mu}(ds, dx); \end{aligned} \quad (3.4)$$

- plug the Lévy processes  $X_{j_1 \dots j_p}^{(1)}$  into  $b_i$ , cf. (2.7), which leads to a Lévy approximation for  $b_i$ ;
- finally, integrate by parts to deduce a Lévy approximation for  $G_i$ .

**Remark 3.1.** Note that the “frozen drift” approximation can be easily embedded in this scheme. It corresponds to using just the initial values  $X_{j_1 \dots j_p}(0)$  instead of the Lévy process  $X_{j_1 \dots j_p}^{(1)}$  in (3.4).

**3.2. Log-Lévy approximation schemes.** In the sequel, we are going to follow this recipe for deriving efficient and accurate log-Lévy approximations, and present the full details of the method. However, we will first truncate the drift terms at the second order, in order to reduce the number of terms that need to be calculated.

**1.** The first step is to expand and truncate the drift term at the second order; these computations have been deferred to Appendix A for brevity, see (A.5). We will approximate  $b_i$  by  $b_i''$ , where

$$\begin{aligned} b_i'' &= -\theta_i - \sum_{i+1 \leq j \leq N} \frac{\delta_j L_{j-}}{1 + \delta_j L_{j-}} \eta_{ij} \\ &\quad - \sum_{i+1 \leq k < l \leq N} \frac{\delta_k L_{k-}}{1 + \delta_k L_{k-}} \frac{\delta_l L_{l-}}{1 + \delta_l L_{l-}} \zeta_{ikl}, \end{aligned} \quad (3.5)$$

where

$$\theta_i = \kappa(\lambda_i), \quad \eta_{ij} = \kappa(\lambda_i + \lambda_j) - \kappa(\lambda_i) - \kappa(\lambda_j) \quad (3.6)$$

and

$$\begin{aligned} \zeta_{ikl} &= \widehat{\kappa}(\lambda_i + \lambda_k + \lambda_l) - \widehat{\kappa}(\lambda_i + \lambda_k) - \widehat{\kappa}(\lambda_i + \lambda_l) \\ &\quad - \widehat{\kappa}(\lambda_k + \lambda_l) + \widehat{\kappa}(\lambda_i) + \widehat{\kappa}(\lambda_k) + \widehat{\kappa}(\lambda_l). \end{aligned} \quad (3.7)$$

The number of terms to be calculated is thus reduced from  $O(2^N)$  to  $O(N^2)$ , while the error induced is

$$b_i = b_i'' + O(N^2 \delta^3 \|L\|^3). \quad (3.8)$$

Therefore, the gain in computational time is significant, while the loss in accuracy is usually relatively small. The numerical examples verify this, see section 5.1 for more details.

**2.** The second step is to approximate the random terms

$$Z_j(t) := \frac{\delta_j L_j(t)}{1 + \delta_j L_j(t)} \quad \text{and} \quad Y_{kl}(t) := \frac{\delta_k L_k(t)}{1 + \delta_k L_k(t)} \frac{\delta_l L_l(t)}{1 + \delta_l L_l(t)} \quad (3.9)$$

in (3.5) by a time-inhomogeneous Lévy process. Define the functions

$$f(x) = \frac{\delta_j e^x}{1 + \delta_j e^x} \quad \text{and} \quad g(x_k, x_l) = \frac{\delta_k e^{x_k}}{1 + \delta_k e^{x_k}} \frac{\delta_l e^{x_l}}{1 + \delta_l e^{x_l}},$$

where

$$f'(x) = \frac{\delta_j e^x}{(1 + \delta_j e^x)^2} \quad \text{and} \quad f''(x) = \frac{\delta_j e^x (1 - \delta_j e^x)}{(1 + \delta_j e^x)^3}.$$

The partial derivatives of  $g$  can be computed equally easily, and are denoted

$$g_k = \frac{\partial}{\partial x_k} g, \quad g_l = \frac{\partial}{\partial x_l} g, \quad g_{kl} = \frac{\partial^2}{\partial x_k \partial x_l} g, \quad (3.10)$$

and so forth. We obviously have that

$$Z_j(t) = f(G_j(t)) \quad \text{and} \quad Y_{kl}(t) = g(G_k(t), G_l(t)). \quad (3.11)$$

The functions  $f$  and  $g$  are  $C^2$ -differentiable, hence we can apply Itô's formula for semimartingales (cf. e.g. Jacod and Shiryaev 2003, Theorem I.4.57) to  $Z_j$  and  $Y_{kl}$ . Using (3.1) we may derive (with time variable  $s$  suppressed or denoted by  $\cdot$  in the integrands)

$$\begin{aligned} dZ_j = & \left( \int_{\mathbb{R}^m} \left( f(G_j + \lambda_j^\top x) - f(G_j) - f'(G_j) \lambda_j^\top x \right) F(\cdot, dx) \right. \\ & \left. + f'(G_j) b_j'' + \frac{1}{2} f''(G_j) |\lambda_j|^2 \alpha \right) ds + f'(G_j) \sqrt{\alpha} \lambda_j^\top dW \\ & + \int_{\mathbb{R}^m} \left( f(G_{j-} + \lambda_j^\top x) - f(G_{j-}) \right) (\mu(ds, dx) - F(\cdot, dx) ds). \end{aligned} \quad (3.12)$$

The derivation is given in Appendix B. Hence, we have that

$$\begin{aligned} dZ_j(s) = & A_j(s, L(s)) ds + B_j^\top(s, L_j(s)) dW(s) \\ & + \int_{\mathbb{R}^m} C_j(s, L_j(s), x) (\mu(ds, dx) - F(\cdot, dx) ds), \end{aligned} \quad (3.13)$$

with obvious definitions of the *deterministic* functions  $A_j$ ,  $B_j$ , and  $C_j$ . Due to the drift term  $b_j''$ , the function  $A_j$  depends on the whole LIBOR vector  $L$  rather than  $L_j$  only.



Similarly, we have for  $Y_{kl}$  that

$$\begin{aligned} dY_{kl}(s) &= A_{kl}(s, L(s))ds + B_{kl}^\top(s, L_{kl}(s))dW(s) \\ &\quad + \int_{\mathbb{R}^m} C_{kl}(s, L_{kl}(s), x) (\mu(ds, dx) - F(\cdot, dx)ds), \end{aligned} \quad (3.14)$$

where  $A_{kl}$ ,  $B_{kl}$ , and  $C_{kl}$  are deterministic functions; see Appendix C for all the details. Analogously to (3.13),  $A_{kl}$  depends on the whole LIBOR vector  $L$ , while  $B_{kl}$  and  $C_{kl}$  depend on  $L_k$  and  $L_l$  only; this is denoted by  $L_{kl}$ .

**3.** The next step is to approximate  $Z_j$  and  $Y_{kl}$  by suitable Lévy processes. This approximation is based on a Picard iteration for the SDEs in (3.13) and (3.14). Regarding  $Z$ , the initial value of the Picard iteration is

$$Z_j^{(0)} = Z_j(0) = \frac{\delta_j L_j(0)}{1 + \delta_j L_j(0)}, \quad (3.15)$$

while the first order Picard iteration is provided by

$$\begin{aligned} Z_j^{(1)}(t) &= Z_j(0) + \int_0^t A_j(s, L(0))ds + \int_0^t B_j^\top(s, L_j(0))dW(s) \\ &\quad + \int_0^t \int_{\mathbb{R}^m} C_j(s, L_j(0), x) (\mu(ds, dx) - F(\cdot, dx)ds). \end{aligned} \quad (3.16)$$

We can easily deduce that  $Z^{(1)}$  is a *time-inhomogeneous Lévy process*, since the coefficients  $A_j(\cdot, L(0))$ ,  $B_j(\cdot, L_j(0))$ , and  $C_j(\cdot, L_j(0), \cdot)$  in (3.16) are *deterministic*. Indeed, we have that

$$\begin{aligned} A_j(s, L(0)) &= f'(G_j(0)) b_j^{(0)}(s) + \frac{1}{2} f''(G_j(0)) |\lambda_j|^2(s) \alpha(s) \\ &\quad + \int_{\mathbb{R}^m} \left( f(G_j(0) + \lambda_j^\top(s)x) - f(G_j(0)) - f'(G_j(0)) \lambda_j^\top(s)x \right) F(\cdot, dx), \end{aligned} \quad (3.17)$$

where

$$\begin{aligned} b_j^{(0)}(s) &:= -\theta_i(s) - \sum_{i+1 \leq j \leq N} \frac{\delta_j L_{j-}(0)}{1 + \delta_j L_{j-}(0)} \eta_{ij}(s) \\ &\quad - \sum_{i+1 \leq k < l \leq N} \frac{\delta_k L_{k-}(0)}{1 + \delta_k L_{k-}(0)} \frac{\delta_l L_{l-}(0)}{1 + \delta_l L_{l-}(0)} \zeta_{ikl}(s), \end{aligned}$$

and

$$B_j(s, L_j(0)) = f'(G_j(0)) \sqrt{\alpha(s)} \lambda_j(s), \quad (3.18)$$

$$C_j(s, L_j(0), x) = f(G_j(0) + \lambda_j^\top(s)x) - f(G_j(0)). \quad (3.19)$$

Analogously, the initial value of the Picard iteration for (3.14) is

$$Y_{kl}^{(0)} = Y_{kl}(0) = \frac{\delta_k L_k(0)}{1 + \delta_k L_k(0)} \frac{\delta_l L_l(0)}{1 + \delta_l L_l(0)}, \quad (3.20)$$

and the first order iteration is

$$\begin{aligned} Y_{kl}^{(1)}(t) &= Y_{kl}(0) + \int_0^t A_{kl}(s, L(0))ds + \int_0^t B_{kl}^\top(s, L_{kl}(0))dW(s) \\ &\quad + \int_0^t \int_{\mathbb{R}^m} C_{kl}(s, L_{kl}(0), x) (\mu(ds, dx) - F(\cdot, dx)ds), \end{aligned} \quad (3.21)$$

and we can again deduce that  $Y_{kl}^{(1)}$  is an additive Lévy process.

4. The fourth step is to apply the Lévy approximations of the random terms to (3.5). Let us denote by  $\widehat{b}_i$  the resulting approximate drift term; we have that

$$b_i'' \approx \widehat{b}_i := -\theta_i - \sum_{i+1 \leq j \leq N} \eta_{ij} Z_j^{(1)} - \sum_{i+1 \leq k < l \leq N} \zeta_{ikl} Y_{kl}^{(1)}. \quad (3.22)$$

Keeping in mind that  $\widehat{b}_i$  will be integrated over time, we define

$$V_{ij}(s, t) = \int_s^t \eta_{ij}(r)dr, \quad \text{and} \quad \bar{V}_{ikl}(s, t) = \int_s^t \zeta_{ikl}(r)dr,$$

which are obviously deterministic processes of finite variation. Now, for fixed  $t > 0$ , we can apply integration by parts, which yields

$$\begin{aligned} \int_0^t \eta_{ij}(s) Z_j^{(1)}(s) ds &\stackrel{(3.16)}{=} V_{ij}(0, t) Z_j(0) + \int_0^t V_{ij}(s, t) A_j(s, L(0)) ds \\ &\quad + \int_0^t V_{ij}(s, t) B_j^\top(s, L_j(0)) dW(s) \\ &\quad + \int_0^t V_{ij}(s, t) \int_{\mathbb{R}^m} C_j(s, L_j(0), x) \tilde{\mu}(ds, dx). \end{aligned} \quad (3.23)$$

Similarly for the other term we get

$$\begin{aligned} \int_0^t \zeta_{ikl}(s) Y_{kl}^{(1)}(s) ds &\stackrel{(3.21)}{=} \bar{V}_{ikl}(0, t) Y_{kl}(0) + \int_0^t \bar{V}_{ikl}(s, t) A_{kl}(s, L(0)) ds \\ &\quad + \int_0^t \bar{V}_{ikl}(s, t) B_{kl}^\top(s, L_{kl}(0)) dW(s) \\ &\quad + \int_0^t \bar{V}_{ikl}(s, t) \int_{\mathbb{R}^m} C_{kl}(s, L_{kl}(0), x) \tilde{\mu}(ds, dx). \end{aligned} \quad (3.24)$$

5. Finally, collecting all the pieces together we can derive a Lévy approximation for the log-LIBOR rates. The *approximate* log-LIBOR is denoted

by  $\widehat{G}_i$  and has the following dynamics

$$\widehat{G}_i(t) = G_i(0) + \int_0^t \widehat{b}_i(s) ds + \int_0^t \lambda_i^\top(s) dH(s), \quad (3.25)$$

which using (3.22), (3.23) and (3.24) leads to

$$\begin{aligned} \widehat{G}_i(t) = & \widehat{G}_i(0, t) - \int_0^t \left[ \theta_i(s) + \sum_{i+1 \leq j \leq N} V_{ij}(s, t) A_j(s, L(0)) \right. \\ & \left. + \sum_{i+1 \leq k < l \leq N} \bar{V}_{ikl}(s, t) A_{kl}(s, L(0)) \right] ds \\ & + \int_0^t \left[ \sqrt{\alpha(s)} \lambda_i^\top(s) - \sum_{i+1 \leq j \leq N} V_{ij}(s, t) B_j^\top(s, L_j(0)) \right. \\ & \left. - \sum_{i+1 \leq k < l \leq N} \bar{V}_{ikl}(s, t) B_{kl}^\top(s, L_{kl}(0)) \right] dW(s) \\ & + \int_0^t \int_{\mathbb{R}^m} \left[ \lambda_i^\top(s) x - \sum_{i+1 \leq j \leq N} V_{ij}(s, t) C_j(s, L_j(0), x) \right. \\ & \left. - \sum_{i+1 \leq k < l \leq N} \bar{V}_{ikl}(s, t) C_{kl}(s, L_{kl}(0), x) \right] \tilde{\mu}(ds, dx), \quad (3.26) \end{aligned}$$

with

$$\widehat{G}_i(0, t) := G_i(0) - \sum_{i+1 \leq j \leq N} V_{ij}(0, t) Z_j(0) - \sum_{i+1 \leq k < l \leq N} \bar{V}_{ikl}(0, t) Y_{kl}(0).$$

Let us abbreviate (3.26) by

$$\begin{aligned} \widehat{G}_i(t) &= \widehat{G}_i(0, t) + \int_0^t H_i(t, s) ds + \int_0^t \Theta_i^\top(t, s) dW(s) + \int_0^t I_i(t, s, x) \tilde{\mu}(ds, dx) \\ &=: X_i^{(t)}(t), \end{aligned}$$

where the process  $X_i^{(t)}(r)$ ,  $0 \leq r \leq t$  is defined by

$$X_i^{(t)}(r) := \widehat{G}_i(0, r) + \int_0^r H_i(t, s) ds + \int_0^r \Theta_i^\top(t, s) dW(s) + \int_0^r I_i(t, s, x) \tilde{\mu}(ds, dx).$$

Obviously,  $X_i^{(t)}(r)$ ,  $0 \leq r \leq t$  is a time-inhomogeneous Lévy process whose characteristic function may be expressed by the Lévy–Khintchine formula in terms of  $H_i$ ,  $\Theta_i$  and  $I_i$  in a straightforward manner.

**Remark 3.2.** We will call the approximation in (3.26) the *second order log-Lévy approximation* of the LIBOR rate. If we ignore the second order terms (i.e. those depending on  $L_k$  and  $L_l$ ), we immediately arrive at the

*first order approximation.* The numerical results in section 5 document the improvement from the first to the second order approximation.

**Remark 3.3.** If we restrict our model to the Brownian motion case, the approximation in (3.26) coincides with the “fully lognormal model” of Daniluk and Gařtarek (2005); see also Kurbanmuradov et al. (2002).

**Remark 3.4.** Note that the approximation methods developed in the previous sections do not depend crucially on the choice of the measure. If we work under the spot measure, cf. Remark 2.2, then the Picard approximations can be carried out similarly. However, an additional approximation is required to represent the drift in terms of cumulants as in eq. (2.7) (because of the  $1/\prod_j$  terms).

**3.3. Efficient simulation of the log-Lévy approximation.** In this section, we outline how simulation of the Lévy approximation

$$\widehat{G}_i(t) = \widehat{G}_i(0, t) + \int_0^t H_i(t, s) ds + \int_0^t \Theta_i^\top(t, s) dW(s) + \int_0^t I_i(t, s, x) \widetilde{\mu}(ds, dx) \quad (3.27)$$

can be carried out in an effective way due to the fact that  $\widehat{G}_i(0, t)$  and the integrands in (3.27) are explicitly known deterministic functions.

**(I)** The terms  $\widehat{G}_i(0, t)$  and  $\int_0^t H_i(t, s) ds$  are deterministic integrals which may be computed outside any Monte Carlo loop using some quadrature formula.

**(II)** The Gaussian part

$$\varsigma_i(t) := \int_0^t \Theta_i^\top(t, s) dW(s) \quad (3.28)$$

may be computed either by usual Euler stepping, or even directly at some fixed time  $t$  if only the distribution of  $\widehat{G}(t)$  matters. In this respect, the distribution of any vector  $(\varsigma_{i_1}(t), \dots, \varsigma_{i_k}(t))$  — for simulating a set of log-LIBORs  $(\widehat{G}_{i_1}(t), \dots, \widehat{G}_{i_k}(t))$  — is Gaussian with explicitly known covariance structure, and thus can be simulated straightforwardly.

**(III)** Finally, consider the practically important case where the Lévy measure itself is time homogeneous, i.e.  $F(dx) \equiv F(\cdot, dx)$ . After truncating this measure with respect to jumps with size smaller than some  $\epsilon > 0$  (if needed), simulation of a realization of the jump term in (3.27) may effectively be carried out as follows. First sample on the interval  $(0, t)$  the number  $N_t$  (of jump times) according to a Poisson distribution with intensity  $tF(\{\|x\| > \epsilon\})$ . Next distribute  $N_t$  jump points  $\{s_1, \dots, s_{N_t}\}$  uniformly over the interval  $(0, t)$ , and sample independently for each jump point  $s_l$  a jump  $x_l$ ,  $1 \leq l \leq N_t$  from the probability measure

$$\frac{F(dx \cap \{\|x\| > \epsilon\})}{F(\{\|x\| > \epsilon\})}.$$

Then a realization of the (compensated) jump term is obtained as

$$\varsigma_i^J(t) := \sum_{l=1}^{N_t} \mathbb{I}_i(t, s_{l-}, x_l) - \int_0^t \int_{\|x\|>\epsilon} \mathbb{I}_i(t, s, x) F(dx) ds, \quad (3.29)$$

where the deterministic integral term can be computed outside any Monte Carlo loop by standard methods. Note that a realization of the whole log-LIBOR vector  $(\varsigma_1^J(t), \dots, \varsigma_N^J(t))$  will be computed using the same set of jumps  $(s_l, x_l)$ ,  $l = 1, \dots, N_t$ .

The main benefit from the log-Lévy approximation as outlined above, is the fact that for the simulation of a log-LIBOR vector  $(\widehat{G}_i(t), \dots, \widehat{G}_N(t))$ , the computation of the terms in (2.5) via (2.7) or (3.5) based on each realization of the Brownian motion and the jump process on a fine enough time grid is *not* required. This is in clear contrast to the Euler (or predictor-corrector) discretization of (2.5) and (2.7). It is obvious that in view of the complex structure of (3.5) only, such a simulation would require the (accurate enough) construction of a whole log-LIBOR system  $(\widehat{G}_i(t_j), \dots, \widehat{G}_N(t_j))$  for  $0 < t_1 < \dots < t_n := t$  involving the evaluation of the function  $b''$  at each grid point  $t_j$ . In contrast, simulation of the log-Lévy LIBOR approximation only involves the evaluation of (3.29) at the jump times and the relatively efficient simulation of the Wiener integral (3.28) inside a Monte Carlo loop.

#### 4. ERROR ESTIMATES

In this section, we will provide some error estimates for the log-Lévy approximations in order to offer a theoretical justification for the proposed approximations. The error estimates are rather qualitative in nature, however they allow for useful conclusions.

In view of (3.25) we have for the pathwise error of the (log-)LIBOR approximation,

$$\left| \frac{\widehat{L}_i(t)}{L_i(t)} \right| \leq \exp \left| \widehat{G}_j(t) - G_j(t) \right| \leq \exp \left( \int_0^t |\widehat{b}_i(s) - b_i(s)| ds \right),$$

thus we need to study the difference  $|\widehat{b}_i - b_i|$ . Since the main contribution of this error is due to the first and second order term in (2.7), we consider instead (see (3.5))

$$|\widehat{b}_i - b_i''| \leq \sum_{i+1 \leq j \leq N} |Z_j - Z_j^{(1)}| |\eta_{ij}| + \sum_{i+1 \leq k < l \leq N} |Y_{kl} - Y_{kl}^{(1)}| |\zeta_{ikl}|.$$

Let us assume for simplicity that  $\alpha(s) \equiv 1$ , and that  $K_\eta$  and  $K_\zeta$  are (dimensionless) constants such that

$$\begin{aligned} \max_{1 \leq i < j \leq N} |\eta_{ij}| &\leq K_\eta \max_{1 \leq i \leq N} \sup_{0 \leq t \leq T} \|\lambda_i(t)\|_2^2 =: K_\eta \lambda_{\max}^2, \\ \max_{1 \leq i < k < l \leq N} |\zeta_{ikl}| &\leq K_\zeta \max_{1 \leq i \leq N} \sup_{0 \leq t \leq T} \|\lambda_i(t)\|_2^2 =: K_\zeta \lambda_{\max}^2. \end{aligned}$$

We then have

$$\begin{aligned} \left\| \log \left| \frac{\widehat{L}_i(t)}{L_i(t)} \right| \right\|_{L_2(\mathbb{P}_*)} &\leq K_\eta \lambda_{\max}^2 \max_{i+1 \leq j \leq N} \int_0^t \left\| Z_i^{(1)}(s) - Z_i(s) \right\|_{L_2(\mathbb{P}_*)} ds \\ &+ K_\zeta \lambda_{\max}^2 \max_{i+1 \leq k < l \leq N} \int_0^t \left\| Y_{kl}^{(1)}(s) - Y_{kl}(s) \right\|_{L_2(\mathbb{P}_*)} ds =: (I) + (II). \end{aligned}$$

For the term (I) we get from (3.13) and (3.16)

$$\begin{aligned} \left\| Z_j^{(1)}(s) - Z_j(s) \right\|_{L_2(\mathbb{P}_*)} &\leq \int_0^s |A_j(u, L(0)) - A_j(u, L(u))|_{L_2(\mathbb{P}_*)} du \\ &+ \left( \int_0^s E \|B_j(u, L_j(0)) - B_j(u, L_j(u))\|_2^2 du \right)^{1/2} \\ &+ \left( \int_0^s \int_{\mathbb{R}^m} E (C_j(u, L_j(0), x) - C_j(u, L_j(u), x))^2 F(u, dx) du \right)^{1/2}. \end{aligned}$$

In view of (3.17), (3.18) and (3.19), let  $K_A$ ,  $K_B$ ,  $K_C$  be dimensionless Lipschitz constants such that for all  $1 \leq j \leq N$  and  $0 \leq u \leq T_*$ ,

$$\begin{aligned} |A_j(u, y) - A_j(u, y')| &\leq K_A \lambda_{\max}^2 \|y - y'\|_2, \\ \|B_j(u, y_j) - B_j(u, y'_j)\|_2 &\leq K_B \lambda_{\max} |y_j - y'_j|, \\ \int_{\mathbb{R}^m} (C_j(u, y_j, x) - C_j(u, y'_j, x))^2 F(u, dx) &\leq K_C^2 \lambda_{\max}^2 |y_j - y'_j|^2. \end{aligned}$$

Then, using

$$\begin{aligned} \left\| Z_j^{(1)}(s) - Z_j(s) \right\|_{L_2(\mathbb{P}_*)} &\leq K_A \lambda_{\max}^2 \int_0^s \|L(0) - L(u)\|_{2, L_2(\mathbb{P}_*)} du \\ &+ (K_B + K_C) \lambda_{\max} \left( \int_0^s E |L_j(0) - L_j(u)|^2 du \right)^{1/2}, \end{aligned}$$

we obtain the estimate

$$\begin{aligned} (I) &\leq \lambda_{\max}^4 K_\eta K_A \int_0^t \left( \int_0^s \|L(0) - L(u)\|_{2, L_2(\mathbb{P}_*)} du \right) ds \\ &+ \lambda_{\max}^3 K_\eta (K_B + K_C) \int_0^t \max_{i+1 \leq j \leq N} \left( \int_0^s E |L_j(0) - L_j(u)|^2 du \right)^{1/2} ds, \end{aligned}$$

and a similar expression may be obtained for the second term (II).

On an intuitive level we may interpret the estimates (I) and (II) in the following way: if we roughly consider that (the approximate squared variance)  $E |L_j(0) - L_j(u)|^2 \lesssim \lambda_{\max}^2 u$ , then for (I) we obtain

$$\begin{aligned} (I) &\lesssim \lambda_{\max}^5 K_\eta K_A \int_0^t \int_0^s \sqrt{u} \, du \, ds \\ &\quad + \lambda_{\max}^4 K_\eta (K_B + K_C) \int_0^t \left( \int_0^s u \, du \right)^{1/2} \, ds \\ &= \frac{4}{15} \lambda_{\max}^5 K_\eta K_A t^{5/2} + \frac{\sqrt{2}}{4} \lambda_{\max}^4 K_\eta (K_B + K_C) t^2, \end{aligned}$$

and a similar result for (II). Hence, for some dimensionless constants  $K_1$  and  $K_2$ ,

$$\left\| \log \left| \frac{\widehat{L}_i(t)}{L_i(t)} \right| \right\|_{L_2(\mathbb{P}_*)} \lesssim K_1 (\lambda_{\max}^2 t)^{5/2} + K_2 (\lambda_{\max}^2 t)^2.$$

Concluding, the log-Lévy LIBOR approximations are extremely good as long as  $\lambda_{\max}^2 t$  is small enough but, may become poor as soon as this product grows very large. This issue is confirmed in our numerical experiments.

## 5. NUMERICAL ILLUSTRATIONS

Throughout this section, we will consider a simple example with a flat and constant volatility structure. Similarly zero coupon rates are generated from a flat term structure of interest rates:  $B(0, T_i) = \exp(-0.04 \cdot T_i)$ . We consider a tenor structure with 6 month increments (i.e.  $\delta_i = \frac{1}{2}$ ). As stated in the introduction, the Brownian motion case is already well studied; therefore we set  $\alpha = 0$ , thus limiting ourselves to the case where  $H$  is a pure jump Lévy process. We consider two univariate specifications, for simplicity. The first is a tempered stable or CGMY process (cf. Carr, Geman, Madan, and Yor 2002 and Madan and Yor 2008) with parameters  $M = G = 13$ ,  $Y = 0.25$  and  $C = 48.4201$ , resulting in a process with mean zero and variance 1 (at  $t = 1$ ), infinite activity and finite variation. The CGMY process has cumulant generating function defined for all  $u \in \mathbb{C}$  with  $|\Re u| \leq \min(G, M)$ ,

$$\begin{aligned} \kappa_{\text{CGMY}}(u) &= \Gamma(-Y) G^Y \left\{ \left(1 - \frac{u}{G}\right)^Y - 1 + \frac{uY}{G} \right\} \\ &\quad + \Gamma(-Y) M^Y \left\{ \left(1 + \frac{u}{G}\right)^Y - 1 - \frac{uY}{M} \right\}. \end{aligned} \quad (5.1)$$

The necessary conditions are then satisfied for term structures up to at least 10 years of length because  $\overline{M} = \min(G, M)$ , hence  $\sum_{i=1}^{20} |\lambda_i| \leq 12 < \overline{M}$ . Exact simulation of the increments can be performed without approximation using the approach in Poirot and Tankov (2006). This approach can be used when simulating from (3.1) with or without drift expansions, but cannot be employed in the case of the log-Lévy approximation in (3.26) where jump sizes are transformed in a non-linear fashion. Instead we employ an approximation where we replace jumps smaller than  $\epsilon$  with their expectation which

is zero since the jumps are compensated. This means that jumps bigger than  $\epsilon$  follow a compound Poisson process which can be easily simulated using the so-called Rosinski rejection method (see Rosiński 2001 and Asmussen and Glynn 2007, p. 338). We set the truncation point sufficiently low, at  $\epsilon = 10^{-3}$ , thus making the variance of the truncated term  $\int_{-\epsilon}^{\epsilon} x^2 \nu(dx) = 3.11 \times 10^{-4}$ , which can be considered small enough to safely disregard. To be consistent, we employ this procedure everywhere we simulate from the CGMY process.

The second specification is a compound Poisson process with normally distributed jump sizes — often referred to as the Merton model. The cumulant generating function for  $u \in \mathbb{C}$  is

$$\kappa_{\text{Merton}}(u) = \bar{\lambda} (\exp(\bar{\mu}u + \bar{\sigma}^2 u^2) - 1 - \bar{\mu}u). \quad (5.2)$$

We set  $\bar{\lambda} = 5$ ,  $\bar{\mu} = 0$  and  $\bar{\sigma} = \sqrt{1/\bar{\lambda}}$  yielding a process with mean zero and variance 1 (at  $t = 1$ ), as before.

In order to verify the validity of our approximations we consider both linear and nonlinear payoffs; in particular, forward rate agreements (FRAs), caplets and swaptions. To price FRAs and caplets with strike  $K$  maturing at time  $T_i$ , we compute the following expectations:

$$\text{FRA}_0 = \delta B(0, T_*) \mathbb{E}_{\mathbf{P}^*} \left[ \prod_{l=i+1}^N (1 + \delta L_l(T_{l+1})) (L_i(T_i) - K) \right], \quad (5.3)$$

$$\text{C}_0 = \delta B(0, T_*) \mathbb{E}_{\mathbf{P}^*} \left[ \prod_{l=i+1}^N (1 + \delta L_l(T_{l+1})) (L_i(T_i) - K)^+ \right]. \quad (5.4)$$

Following Kluge (2005, pp. 78), we have that the price of a payer swaption with strike rate  $K$ , where the underlying swap starts at time  $T_i$  and matures at  $T_m$  ( $i < m \leq N$ ) is given by

$$\text{S}_0 = B(0, T_*) \mathbb{E}_{\mathbf{P}^*} \left[ \left( - \sum_{k=i}^m \left( c_k \prod_{l=k}^N (1 + \delta L_l(T_l)) \right) \right)^+ \right], \quad (5.5)$$

where

$$c_k = \begin{cases} -1, & k = i, \\ \delta_k K, & i + 1 \leq k \leq m - 1, \\ 1 + \delta_k K, & k = m. \end{cases} \quad (5.6)$$

**5.1. Performance of the drift expansion.** As we have argued in section 3.2, the truncation of the drift term in equation (2.6) is necessary in order to build a model that is computationally tractable. This section illustrates the effect of this truncation using the standard Euler discretization of the actual dynamics, i.e. equations (2.5) and (2.7).

Due to the complexity of calculating the true drift we limit ourselves to setting  $N = 10$ , corresponding to a 5 year term structure. Furthermore we consider volatility structures constant and flat at  $\lambda_i = 0.2$  and  $\lambda_i = 0.6$  respectively. We simulate 10000 paths and plot the absolute difference between the prices from the drift expansions and the price without expansion (i.e. the full drift in (2.6)) in Figures 5.1 and 5.2. Each Monte Carlo simulation is done using the same random shocks for each method, thus eliminating the Monte Carlo noise as an error source. The figures demonstrate that the



effect of the truncation depends mostly on the level of volatility  $\lambda_i$  and less in the choice of product to price or the driving process. Furthermore, we




FIGURE 5.1. Drift expansion: low volatility scenario.

notice that for low volatility even the first order expansion can be considered adequate, since the maximum of the absolute error is smaller than 0.2 bp. Conversely, for the high volatility case, the second order expansion is necessary to get proper accuracy. However, going to the third order expansion or beyond appears to be unnecessary as there is no visible gain in accuracy ( $< 10^{-5}$  bp). Hence, in the next sections we will use the second order drift expansion as our benchmark case since any resulting error is small enough to be disregarded.

In Table 5.1, CPU times are shown when simulating 10000 paths on an Intel i7 PC running Matlab. Here we can see that highly significant speed-up is achieved when truncating the higher order drift terms, whereas the decrease in speed when taking higher order approximations into account is relatively negligible. The CGMY is slower than the Merton model due to the much higher jump intensity needed in its approximation. We conjecture that the efficiency can be improved using the methods of Kohatsu-Higa and Tankov (2010), but this lies outside the focus of this article.

Finally, to conclude the subsection we should also mention that pricing errors for swaptions (not shown here) are of similar order of magnitude as in case of caplets.

**5.2. Performance of the log-Lévy approximations.** Next we study the performance of the log-Lévy approximations. We increase the number of




FIGURE 5.2. Drift expansion: high volatility scenario.

	Full Drift	1st order	2nd order	3rd order
Merton	358.5	3.95	4.48	4.79
CGMY	471.9	16.29	16.59	16.74

TABLE 5.1. CPU Times (secs) for 10000 paths

rates to the more realistic setting of  $N = 20$  and consider the pricing of FRAs, caplets and swaptions; the latter are maturing at time  $T_i$  and ending at  $(T_i + 3)$  years. Since we have established that errors from the drift expansion can be disregarded, we consider as the benchmark case the second order drift expansion studied in the previous section. In Figure 5.3 we plot prices from the frozen drift, the first and second order log-Lévy approximations of section 3, and include the annuity approximation of the following section for completeness. We use both the Merton and the CGMY model. We can observe that the frozen drift is consistently beaten by both the 1st and 2nd order approximation in both models and for all three products. The 1st and 2nd order log-Lévy approximations have a quite similar performance suggesting that second order approximation may not be necessary. Note that other parameter values (higher/lower intensity for Merton and fatter tails/slower tail decay for CGMY) have also been studied and again the results are qualitatively the same.




FIGURE 5.3. Prices for the Merton and CGMY models.

Concluding, the log-Lévy approximations offer an alternative to the Euler (or predictor-corrector) discretization of the actual dynamics which can be simulated faster and yields almost as accurate options prices.

## 6. APPROXIMATION OF ANNUITIES

In the lognormal LIBOR market model, it is well documented that problems may occur for high volatilities due to a proportionally large Monte Carlo variance in the annuity term used for discounting under the terminal measure, see Beveridge (2010) and Gatarek et al. (2006). Motivated by this numerical problem, we will derive an approximation of the annuity term in the spirit of Gatarek et al. (2006, §10.13).

Let us define the annuity term

$$A_i(t) = \prod_{j=i+1}^N (1 + \delta L_j(t)), \quad (6.1)$$

and consider the vector of log-LIBOR rates  $G = [G_{i+1}, \dots, G_N]$ . We define a function  $f : \mathbb{R}^{N-(i+1)} \rightarrow \mathbb{R}$  such that

$$(x_{i+1}, \dots, x_n) = x \mapsto \prod_{j=i+1}^N (1 + \delta e^{x_j}).$$

The partial derivatives of  $f$  are provided by

$$f_k(x) = \frac{\partial}{\partial x_k} f(x) = \prod_{\substack{j=i+1 \\ j \neq k}}^N (1 + \delta e^{x_j}) \delta e^{x_k} = f(x) \frac{\delta e^{x_k}}{1 + \delta e^{x_k}},$$

for all  $i+1 \leq k \leq N$ , while we obviously have that

$$f(G(t)) = A_i(t). \quad (6.2)$$

Applying Itô's formula to  $f(G)$ , we have that

$$\begin{aligned} f(G(t)) = A_i(t) &= A_i(0) + \sum_{j=i+1}^N \int_0^t f_j(G(s-)) dG_j(s) \\ &\quad + \frac{1}{2} \sum_{j,k=i+1}^N \int_0^t f_{j,k}(G(s-)) d\langle G_k, G_j(s) \rangle^c(s) \\ &\quad + \sum_{s \leq t} \left\{ \Delta f(G(s)) - \sum_{j=i+1}^N f_j(G(s-)) \Delta G_j(s) \right\}. \end{aligned} \quad (6.3)$$

Noting that the annuity is a  $\mathbb{P}_*$ -martingale, we will focus on the martingale parts of (6.3) in the sequel. Using (3.1) and the fact that  $H$  is also a  $\mathbb{P}_*$ -martingale, we get that the *martingale* part of the first summand is

$$\begin{aligned} \sum_{j=i+1}^N \int_0^t f_j(G(s-)) \lambda_j(s) dH(s) &= \sum_{j=i+1}^N \int_0^t \frac{\delta L_j(s-)}{1 + \delta L_j(s-)} f(G(s-)) \lambda_j(s) dH(s) \\ &= \int_0^t A_i(s-) \sum_{j=i+1}^N \frac{\delta L_j(s-)}{1 + \delta L_j(s-)} \lambda_j(s) dH(s). \end{aligned}$$

The second summand is omitted, while the final summands yields that

$$\begin{aligned}
& \sum_{s \leq t} \left\{ \Delta f(G(s)) - \sum_{j=i+1}^N f_j(G(s-)) \Delta G_j(s) \right\} \\
&= \sum_{s \leq t} \left\{ \Delta A_i(s) - A_i(s-) \sum_{j=i+1}^N \frac{\delta L_j(s-)}{1 + \delta L_j(s-)} \Delta G_j(s) \right\} \\
&= \sum_{s \leq t} \left\{ \Delta A_i(s) - A_i(s-) \sum_{j=i+1}^N \frac{\delta L_j(s-)}{1 + \delta L_j(s-)} \lambda_j(s) \Delta H(s) \right\} \\
&= \int_0^t \int_{\mathbb{R}^m} \left\{ A_i(s) - A_i(s-) - A_i(s-) \sum_{j=i+1}^N \frac{\delta L_j(s-)}{1 + \delta L_j(s-)} \lambda_j(s) x \right\} \tilde{\mu}(ds, dx) \\
&\quad - \int_0^t \int_{\mathbb{R}^m} \left\{ A_i(s) - A_i(s-) - A_i(s-) \sum_{j=i+1}^N \frac{\delta L_j(s-)}{1 + \delta L_j(s-)} \lambda_j(s) x \right\} F(s, dx) ds,
\end{aligned} \tag{6.4}$$

where the quantity  $A_i$  in the last two integrals should be understood as

$$A_i(s) = \prod_{j=i+1}^N \left( 1 + \delta \exp \{ G_j(s-) + \lambda_j^\top(s) x \} \right). \tag{6.5}$$

Collecting all the pieces together, we have that the annuity  $A_i$  satisfies the following integrated SDE

$$\begin{aligned}
A_i(t) &= A_i(0) + \int_0^t A_i(s-) \Lambda_i(s-) dH(s) \\
&\quad + \int_0^t \int_{\mathbb{R}^m} \{ A_i(s) - A_i(s-) - A_i(s-) \Lambda_i(s-) \} \tilde{\mu}(ds, dx) \\
&= A_i(0) + \int_0^t A_i(s-) \Lambda_i(s-) dH(s) \\
&\quad + \int_0^t \int_{\mathbb{R}^m} A_i(s-) \left\{ \frac{A_i(s)}{A_i(s-)} - 1 - \Lambda_i(s-) x \right\} \tilde{\mu}(ds, dx),
\end{aligned} \tag{6.6}$$

where

$$\Lambda_i(s-) = \sum_{j=i+1}^N \frac{\delta L_j(s-)}{1 + \delta L_j(s-)} \lambda_j(s). \tag{6.7}$$

The solution of the SDE (6.6) is the stochastic exponential, thus we get that

$$\begin{aligned}
A_i(t) = A_i(0) \exp & \left( \int_0^t \Lambda_i(s-) dW(s) - \frac{1}{2} \int_0^t (\Lambda_i(s-))^2 ds \right. \\
& + \int_0^t \int_{\mathbb{R}^m} \left\{ \frac{A_i(s)}{A_i(s-)} - 1 \right\} \tilde{\mu}(ds, dx) \\
& \left. - \int_0^t \int_{\mathbb{R}^m} \left( \log \left\{ \frac{A_i(s)}{A_i(s-)} \right\} - \frac{A_i(s)}{A_i(s-)} + 1 \right) \mu(ds, dx) \right), \tag{6.8}
\end{aligned}$$

where again  $A_i(s)$  should be understood as in (6.5). By freezing the random terms in the drifts and jump sizes in the above dynamics we get an alternative approximation for the annuity term. Note that the resulting approximation is also a *log-Lévy approximation*.

We can now use this approximation to price caplets and swaptions, noting that their respective payoffs can be written in terms of annuities:

$$\mathbb{C}_0 = \delta B(0, T_*) \mathbb{E}_{\mathbf{P}_*} \left[ (A_{i-1}(T_{i+1}) + (1 - \delta_i K) A_i(T_{i+1}))^+ \right], \tag{6.9}$$

$$\mathbb{S}_0 = B(0, T_*) \mathbb{E}_{\mathbf{P}_*} \left[ \left( - \sum_{k=i}^m c_k A_{k-1}(T_i) \right)^+ \right], \tag{6.10}$$

where the  $c_k$ 's are defined in (5.6). Any other payoff can be expressed in terms of annuities in a similar fashion.

**6.1. Performance of the annuity approximation.** In Figure 6.1, the quality of the various approximations is studied for a number of at-the-money caplets as a function of the volatility. As before we set the number of rates to  $N = 20$ , and simulate 50000 paths for each volatility level. The plot is for the Merton model while the results are similar for CGMY. Using that at-the-money call option prices are increasing and roughly linear functions of volatility (see for example Wilmott 1998, Brenner and Subrahmanyam 1994 and Backus, Foresi, and Wu 2004 for the case of non-Gaussian distributions), we can observe that only the annuity approximation produces sensible option prices at all levels of volatility. Moreover, even the benchmark case fails when volatility grows beyond 30%, meaning that the Monte Carlo simulation has failed to converge. The frozen drift fails at even lower levels of volatility, while the log-Lévy approximations fail at a higher level, similar to the benchmark case. The annuity approximation works for all (higher) levels and also, as we have seen in Figure 5.3, for the low levels. One should therefore be careful when the average (across maturity) at-the-money implied volatilities are above 30% which is indeed the case in the current market for USD denominated LIBOR caplets where volatilities range from roughly 80% in the short end to 25% in the long end (source: Bloomberg).

Moreover, in Figure 6.2 we observe that this problem becomes significantly less severe when limiting the number of rates to 10 with  $\delta_i = 1$  instead of 20 with  $\delta_i = 0.5$ . Needless to say, limiting the number of rates is rarely a possibility in practice.




FIGURE 6.1. Caplet prices as a function of volatility ( $N = 20$ ).




FIGURE 6.2. Caplet prices as a function of volatility ( $N = 10$ ).

In order to intuitively understand why this approximation performs better in the high volatility case than the other methods (e.g. the standard Euler scheme or the log-Lévy approximations), let us just concentrate on

the lognormal case. We have from (6.8) that

$$\log A_i(t) \cong \sum_{j=i+1}^N \frac{\delta L_j(0)}{1 + \delta L_j(0)} \lambda_j \cdot \sqrt{t} \mathcal{N} + \text{deterministic terms}, \quad (6.11)$$

where  $\mathcal{N}$  denotes a standard normal random variate. On the other hand, from (6.1), we get that

$$\log A_i(t) \cong \sum_{j=i+1}^N \delta \exp\left(\lambda_j \cdot \sqrt{t} \mathcal{N} + \text{random terms}\right), \quad (6.12)$$

where actually the method of approximation will only affect the random terms. We can easily conclude from (6.11) and (6.12) that the variance of the annuity approximation is significantly lower than the variance of the standard representation, which results in the faster convergence of the Monte Carlo method. Thus, the annuity log-Lévy approximation should be interpreted as a *variance reduction* technique for the LIBOR market model.

## 7. ECONOMICALLY MEANINGFUL MULTI-DIMENSIONAL LÉVY MEASURES VIA SUBORDINATION

Next, we reflect on the properties the driving process should have for practical applications and provide some recommendations. In an economically realistic Lévy LIBOR model the very structure of the Lévy measure is important. Since, from an economic point of view, any jump in the daily rate typically affects all segments of the yield curve, we require in our modeling that, *at a jump time, all the LIBORs jump, not only the first or second half of the LIBOR curve for example*. Moreover, this requirement should be fulfilled regardless of the structure of the loading factors  $\lambda_i$ ; the latter may be inferred from some calibration procedure for instance. A natural way to meet this property is to take Lévy measures which are *absolutely continuous*. In a jump-diffusion setting this can be easily established by taking as Lévy measure the product of one dimensional absolutely continuous probability measures  $p_i$ , i.e.

$$\nu(dx) = p_1(dx_1) \cdots p_m(dx_m); \quad (7.1)$$

see Belomestny and Schoenmakers (2011). In this paper we consider LIBOR models based on Lévy processes with possibly infinite activity, thus having available flexible and realistic LIBOR models possibly without Wiener part (i.e.  $\alpha \equiv 0$ ). However, Lévy measures of infinite activity cannot be obtained by simply taking the product of a set of one-dimensional Lévy measures of infinite activity. Nonetheless, we seek for absolutely continuous infinite activity Lévy measures such that the entailed jump processes maintain certain (weak) independence properties. Such measures may be constructed by *Brownian subordination* (see e.g. Cont and Tankov 2004) as outlined below.

Let  $W$  be a Wiener process on  $\mathbb{R}^m$ . The characteristic function of  $W(t)$  is given by

$$\mathbb{E} \left[ e^{izW(t)} \right] = e^{-\frac{t}{2} \|z\|^2} =: e^{t\Psi(z)}, \quad z \in \mathbb{R}^m.$$



We now consider a subordinator  $(S_t)_{t \geq 0}$  on  $\mathbb{R}_+$ , with Lévy triplet  $(0, 0, \rho)$ , and with Laplace exponent  $\Xi$ , i.e.

$$\mathbb{E} [e^{uS_t}] = e^{t\Xi(u)} := \exp \left( t \int_{(0, \infty)} (e^{su} - 1) \rho(ds) \right), \quad u \leq 0.$$

Then the  $m$ -dimensional process  $Y$  defined by

$$Y(t) := W(S_t)$$

has characteristic function

$$\begin{aligned} \mathbb{E} [e^{iz^\top Y(t)}] &= \mathbb{E} \left[ \mathbb{E} [e^{iz^\top W(S_t)} | S_t] \right] = \mathbb{E} [e^{S_t \Psi(z)}] = e^{t\Xi(\Psi(z))} \\ &= \exp \left[ t \int_{(0, \infty)} (e^{s\Psi(z)} - 1) \rho(ds) \right] \\ &= \exp \left[ t \int_{(0, \infty)} (e^{-\frac{s}{2} \|z\|^2} - 1) \rho(ds) \right] =: \exp [t\Phi(z)] \end{aligned}$$

As a result,  $Y$  is a pure jump martingale Lévy process with Lévy measure  $\nu^Y$  satisfying

$$\Phi(z) = \int_{(0, \infty)} (e^{-\frac{s}{2} \|z\|^2} - 1) \rho(ds) = \int_{\mathbb{R}^m} (e^{iz^\top x} - 1 - iz^\top x) \nu^Y(dx). \quad (7.2)$$

It is easily checked that

$$\nu^Y(dx) = \int_0^\infty \frac{1}{(\sqrt{2\pi s})^m} e^{-\frac{1}{2s} \|x\|^2} \rho(ds) dx, \quad (7.3)$$

which is a measure with absolutely continuous support.

**Example 7.1.** Let  $(S_t)_{t \geq 0}$  be the inverse Gaussian subordinator with

$$\rho(ds) = \frac{ce^{-\lambda s}}{s^{3/2}} 1_{\{s > 0\}} ds, \quad \text{and} \quad \mathbb{E} [e^{uS_t}] = e^{-2ct\sqrt{\pi}(\sqrt{\lambda - u} - \sqrt{\lambda})}.$$

Then, (7.2) is known explicitly as

$$\Phi(z) = -2c\sqrt{\pi} \left( \sqrt{\lambda + \frac{\sigma^2}{2} \|z\|^2} - \sqrt{\lambda} \right),$$

e.g. see Cont and Tankov (2004).

**Example 7.2.** Let  $(S_t)_{t \geq 0}$  be a Lévy subordinator with the following properties:

$$\rho(dt) = Ce^{-\frac{t}{4}G} D_{-Y}(G) 1_{\{t > 0\}} dt,$$

$$\Xi(u) = 2C\Gamma(-Y) \left[ (G^2 - 2u)^{Y/2} \cos \left( Y \arctan \left( \frac{\sqrt{-2u}}{G} \right) \right) - G^Y \right],$$

where  $D$  is the parabolic cylinder function. Then, (7.2) is known explicitly as the Lévy exponent of the CGMY process, cf. (5.1), with  $G = M$ ; see Madan and Yor (2008).

**Remark 7.3.** By taking in (2.2)  $F(s, dx) := \nu^Y(dx)$  with  $\nu^Y$  given by (7.3), the jump-part of (2.2) is represented by the process  $Y$  constructed above. It is easy to see that  $Y$  has uncorrelated components, although they are generally not independent. Indeed,  $Y(t)$  has mean zero and we have that

$$\mathbb{E} \left[ Y^{(k)}(t) Y^{(l)}(t) \right] = \mathbb{E} \left[ \mathbb{E} \left[ W^{(k)}(S_t) W^{(l)}(S_t) \mid S_t \right] \right] = 0, \quad 1 \leq k < l \leq m.$$

Thus in contrast to the jump-diffusion situation in Belomestny and Schoenmakers (2011) where all components jump at the same time independently, here the components of  $Y$  still jump at the same time but in an uncorrelated rather than in an independent way.

## APPENDIX A. COMPUTATION OF THE DRIFT

**A.1. Full expansion in terms of cumulants.** We will derive a representation for the integral term of the drift (2.6) which does not involve an integration over random terms. Let us denote the integral term by

$$\mathbb{B}_i := \int_{\mathbb{R}^m} \left( \left( e^{\lambda_i^\top x} - 1 \right) \prod_{j=i+1}^N \left( 1 + \frac{\delta_j L_{j-} \left( e^{\lambda_i^\top x} - 1 \right)}{1 + \delta_j L_{j-}} \right) - \lambda_i^\top x \right) F(\cdot, dx).$$

Observe that

$$\begin{aligned} \prod_{j=1}^l (1 + w_j) &= 1 + \sum_{1 \leq j \leq l} w_j + \sum_{1 \leq j_1 < j_2 \leq l} w_{j_1} w_{j_2} \\ &\quad + \sum_{1 \leq j_1 < j_2 < j_3 \leq l} w_{j_1} w_{j_2} w_{j_3} + \dots + w_1 \cdots w_l \\ &= 1 + \sum_{p=1}^l S_p^l(w_1, \dots, w_l), \end{aligned}$$

where  $S_p^l$  denotes the elementary symmetric polynomial of degree  $p$  in  $l$  variables, i.e.

$$S_p^l(w_1, \dots, w_l) := \sum_{1 \leq j_1 < \dots < j_p \leq l} w_{j_1} \cdots w_{j_p}, \quad 1 \leq p \leq l.$$

Thus  $\mathbb{B}_i$  may be rearranged as follows:

$$\begin{aligned} \mathbb{B}_i &= \int \left( e^{\lambda_i^\top x} - 1 - \lambda_i^\top x \right) F(\cdot, dx) + \sum_{p=1}^{N-i} \int \left( e^{\lambda_i^\top x} - 1 \right) \times \\ &\quad S_p^{N-i} \left( \frac{\delta_{i+1} L_{i+1-} \left( e^{\lambda_{i+1}^\top x} - 1 \right)}{1 + \delta_{i+1} L_{i+1-}}, \dots, \frac{\delta_N L_{N-} \left( e^{\lambda_N^\top x} - 1 \right)}{1 + \delta_N L_{N-}} \right) F(\cdot, dx) \\ &:= (I) + (II). \end{aligned}$$

Let us consider in (II) for  $p \geq 1$  the term

$$\begin{aligned} & \int \left( e^{\lambda_i^\top x} - 1 \right) S_p^{N-i} \left( \frac{\delta_{i+1} L_{i+1-} \left( e^{\lambda_{i+1}^\top x} - 1 \right)}{1 + \delta_{i+1} L_{i+1-}}, \dots, \frac{\delta_N L_{N-} \left( e^{\lambda_N^\top x} - 1 \right)}{1 + \delta_N L_{N-}} \right) F(\cdot, dx) \\ &= \sum_{i < j_1 < \dots < j_p \leq N} \frac{\delta_{j_1} L_{j_1-}}{1 + \delta_{j_1} L_{j_1-}} \dots \frac{\delta_{j_p} L_{j_p-}}{1 + \delta_{j_p} L_{j_p-}} \\ & \quad \times \int \left( e^{\lambda_i^\top x} - 1 \right) \left( e^{\lambda_{j_1}^\top x} - 1 \right) \dots \left( e^{\lambda_{j_p}^\top x} - 1 \right) F(\cdot, dx). \end{aligned}$$

With  $j_0 := i$ , we may write

$$\begin{aligned} & \left( e^{\lambda_i^\top x} - 1 \right) \left( e^{\lambda_{j_1}^\top x} - 1 \right) \dots \left( e^{\lambda_{j_p}^\top x} - 1 \right) \tag{A.1} \\ &= (-1)^{p+1} \left( 1 - e^{\lambda_{j_0}^\top x} \right) \left( 1 - e^{\lambda_{j_1}^\top x} \right) \dots \left( 1 - e^{\lambda_{j_p}^\top x} \right) \\ &= (-1)^{p+1} \left[ 1 + \sum_{q=1}^{p+1} S_q^{p+1} \left( -e^{\lambda_{j_0}^\top x}, \dots, -e^{\lambda_{j_p}^\top x} \right) \right] = (-1)^{p+1} [1 + (*)] \end{aligned}$$

where

$$\begin{aligned} (*) &= \sum_{q=1}^{p+1} (-1)^q S_q^{p+1} \left( e^{\lambda_{j_0}^\top x}, \dots, e^{\lambda_{j_p}^\top x} \right) = \sum_{q=1}^{p+1} (-1)^q \sum_{0 \leq r_1 < \dots < r_q \leq p} e^{\lambda_{j_{r_1}}^\top x} \dots e^{\lambda_{j_{r_q}}^\top x} \\ &= \sum_{q=1}^{p+1} (-1)^q \sum_{0 \leq r_1 < \dots < r_q \leq p} \underbrace{\left( e^{\lambda_{j_{r_1}}^\top x + \dots + \lambda_{j_{r_q}}^\top x} - 1 - (\lambda_{j_{r_1}}^\top x + \dots + \lambda_{j_{r_q}}^\top x) \right)}_{O(\|x\|^2)} \\ & \quad + \sum_{q=1}^{p+1} (-1)^q \sum_{0 \leq r_1 < \dots < r_q \leq p} \left( 1 + \lambda_{j_{r_1}}^\top x + \dots + \lambda_{j_{r_q}}^\top x \right). \end{aligned}$$

Obviously, expression (A.1) is of order  $O(\|x\|^2)$  for any  $p \geq 1$ , hence (!) it must hold

$$1 + \sum_{q=1}^{p+1} (-1)^q \sum_{0 \leq r_1 < \dots < r_q \leq p} \left( 1 + \lambda_{j_{r_1}}^\top x + \dots + \lambda_{j_{r_q}}^\top x \right) = 0.$$

Therefore, we can deduce the following representation for the integral term

$$\begin{aligned}
\mathbb{B}_i &= \int \left( e^{\lambda_i^\top x} - 1 - \lambda_i^\top x \right) F(\cdot, dx) \\
&\quad + \sum_{p=1}^{N-i} \sum_{i < j_1 < \dots < j_p \leq N} \frac{\delta_{j_1} L_{j_1-}}{1 + \delta_{j_1} L_{j_1-}} \dots \frac{\delta_{j_p} L_{j_p-}}{1 + \delta_{j_p} L_{j_p-}} \sum_{q=1}^{p+1} (-1)^{p+q+1} \\
&\quad \times \sum_{0 \leq r_1 < \dots < r_q \leq p} \int \left( e^{(\lambda_{j_{r_1}} + \dots + \lambda_{j_{r_q}})^\top x} - 1 - (\lambda_{j_{r_1}} + \dots + \lambda_{j_{r_q}})^\top x \right) F(\cdot, dx) \\
&= \widehat{\kappa}(\lambda_i) + \sum_{p=1}^{N-i} \sum_{i < j_1 < \dots < j_p \leq N} \frac{\delta_{j_1} L_{j_1-}}{1 + \delta_{j_1} L_{j_1-}} \dots \frac{\delta_{j_p} L_{j_p-}}{1 + \delta_{j_p} L_{j_p-}} \\
&\quad \times \sum_{q=1}^{p+1} (-1)^{p+q+1} \sum_{0 \leq r_1 < \dots < r_q \leq p} \widehat{\kappa}(\lambda_{j_{r_1}} + \dots + \lambda_{j_{r_q}}). \tag{A.2}
\end{aligned}$$

**A.2. First order expansion of (A.2).** Let us consider the first order expansion of  $\mathbb{B}_i$ ; we get

$$\begin{aligned}
\mathbb{B}_i &= \widehat{\kappa}(\lambda_i) + \sum_{i < j < n} \frac{\delta_j L_{j-}}{1 + \delta_j L_{j-}} \sum_{q=1}^2 (-1)^q \sum_{0 \leq r_1 < \dots < r_q \leq 1} \widehat{\kappa}(\lambda_{j_{r_1}} + \dots + \lambda_{j_{r_q}}) \\
&\quad + O(\|L\|^2).
\end{aligned}$$

Note that

$$\begin{aligned}
&\sum_{q=1}^2 (-1)^q \sum_{0 \leq r_1 < \dots < r_q \leq 1} \widehat{\kappa}(\lambda_{j_{r_1}} + \dots + \lambda_{j_{r_q}}) \\
&= - \sum_{0 \leq r_1 \leq 1} \widehat{\kappa}(\lambda_{j_{r_1}}) + \sum_{0 \leq r_1 < r_2 \leq 1} \widehat{\kappa}(\lambda_{j_{r_1}} + \lambda_{j_{r_2}}) \\
&= -\widehat{\kappa}(\lambda_{j_0}) - \widehat{\kappa}(\lambda_{j_1}) + \widehat{\kappa}(\lambda_{j_0} + \lambda_{j_1}).
\end{aligned}$$

Thus we obtain the following expression for the first order expansion of the integral term  $\mathbb{B}_i$

$$\mathbb{B}'_i = \widehat{\kappa}(\lambda_i) + \sum_{i < j \leq N} \frac{\delta_j L_{j-}}{1 + \delta_j L_{j-}} (\widehat{\kappa}(\lambda_i + \lambda_j) - \widehat{\kappa}(\lambda_i) - \widehat{\kappa}(\lambda_j)), \tag{A.3}$$

which leads to the following approximation for the drift term  $b_i$  in (2.6)

$$b'_i = \kappa(\lambda_i) + \sum_{i < j \leq N} \frac{\delta_j L_{j-}}{1 + \delta_j L_{j-}} (\kappa(\lambda_i + \lambda_j) - \kappa(\lambda_i) - \kappa(\lambda_j)), \tag{A.4}$$

taking also the terms stemming from the diffusion into account.

A.3. **Second order expansion of (A.2).** Analogously, we can also derive a second order expansion of  $\mathbb{B}_i$ ; we get

$$\begin{aligned} \mathbb{B}_i &= \widehat{\kappa}(\lambda_i) + \sum_{i < j \leq N} \frac{\delta_j L_{j-}}{1 + \delta_j L_{j-}} (\widehat{\kappa}(\lambda_i + \lambda_j) - \widehat{\kappa}(\lambda_i) - \widehat{\kappa}(\lambda_j)) \\ &\quad + \sum_{i+1 \leq k < l \leq N} \frac{\delta_k L_{k-}}{1 + \delta_k L_{k-}} \frac{\delta_l L_{l-}}{1 + \delta_l L_{l-}} \left( \widehat{\kappa}(\lambda_i + \lambda_k + \lambda_l) - \widehat{\kappa}(\lambda_i + \lambda_k) \right. \\ &\quad \left. - \widehat{\kappa}(\lambda_i + \lambda_l) - \widehat{\kappa}(\lambda_k + \lambda_l) + \widehat{\kappa}(\lambda_i) + \widehat{\kappa}(\lambda_k) + \widehat{\kappa}(\lambda_l) \right) \\ &\quad + O(\|L\|^3), \end{aligned}$$

which leads to the following second order expansion of  $b_i$  in (2.6)

$$\begin{aligned} b_i'' &= \kappa(\lambda_i) + \sum_{i < j \leq N} \frac{\delta_j L_{j-}}{1 + \delta_j L_{j-}} (\kappa(\lambda_i + \lambda_j) - \kappa(\lambda_i) - \kappa(\lambda_j)) \\ &\quad + \sum_{i+1 \leq k < l \leq N} \frac{\delta_k L_{k-}}{1 + \delta_k L_{k-}} \frac{\delta_l L_{l-}}{1 + \delta_l L_{l-}} \left( \widehat{\kappa}(\lambda_i + \lambda_k + \lambda_l) - \widehat{\kappa}(\lambda_i + \lambda_k) \right. \\ &\quad \left. - \widehat{\kappa}(\lambda_i + \lambda_l) - \widehat{\kappa}(\lambda_k + \lambda_l) + \widehat{\kappa}(\lambda_i) + \widehat{\kappa}(\lambda_k) + \widehat{\kappa}(\lambda_l) \right). \quad (\text{A.5}) \end{aligned}$$

#### APPENDIX B. DERIVATION OF (3.12)

Using the Itô formula for general semimartingales (cf. Jacod and Shiryaev 2003, Theorem I.4.57) we have

$$\begin{aligned} Z_j &= Z_j(0) + \int_0^\cdot f'(G_j(s-)) dG_j + \frac{1}{2} \int_0^\cdot f''(G_j) d\langle G_j^c, G_j^c \rangle \\ &\quad + \sum_{0 < s \leq \cdot} (f(G_j(s)) - f(G_j(s-)) - f'(G_j(s-)) \Delta G_j(s)), \quad (\text{B.1}) \end{aligned}$$

where  $\langle G_j^c, G_j^c \rangle$  denotes the quadratic variation of the continuous martingale part of  $G_j$ , that is

$$d\langle G_j^c, G_j^c \rangle(s) = |\lambda_j|^2(s) \alpha(s) ds. \quad (\text{B.2})$$

The sum in (B.1), using (3.2), may be written as

$$\begin{aligned} &\sum_{0 < s \leq \cdot} (f(G_j(s-) + \Delta G_j(s)) - f(G_j(s-)) - f'(G_j(s-)) \Delta G_j(s)) \quad (\text{B.3}) \\ &= \int_0^\cdot \int_{\mathbb{R}^m} \left( f(G_j(s-) + \lambda_j^\top x) - f(G_j(s-)) - f'(G_j(s-)) \lambda_j^\top x \right) \mu(ds, dx) \\ &= \int_0^\cdot \int_{\mathbb{R}^m} \left( f(G_j(s-) + \lambda_j^\top x) - f(G_j(s-)) - f'(G_j(s-)) \lambda_j^\top x \right) F(s, dx) ds \\ &\quad + \int_0^\cdot \int_{\mathbb{R}^m} \left( f(G_j(s-) + \lambda_j^\top x) - f(G_j(s-)) - f'(G_j(s-)) \lambda_j^\top x \right) \tilde{\mu}(ds, dx). \end{aligned}$$

Moreover,

$$\begin{aligned} \int_0^\cdot f'(G_j(s-))dG_j &= \int_0^\cdot f'(G_j(s-))b_j ds + \int_0^t f'(G_j(s-))\sqrt{\alpha}\lambda_j^\top dW \\ &\quad + \int_0^\cdot \int_{\mathbb{R}^m} f'(G_j(s-))\lambda_j^\top x \tilde{\mu}(ds, dx). \end{aligned} \quad (\text{B.4})$$

Finally, by plugging (B.2), (B.3) and (B.4) into (B.1), (3.12) follows.

### APPENDIX C. DERIVATION OF (3.14)

Applying Itô's formula for general semimartingales again, we have

$$\begin{aligned} Y_{kl} &= Y_{kl}(0) + \sum_{i=k,l} \int_0^\cdot g_i(G_k(s-), G_l(s-))dG_i \\ &\quad + \frac{1}{2} \sum_{i,j=k,l} \int_0^\cdot g_{ij}(G_k(s-), G_l(s-))d\langle G_i^c, G_j^c \rangle \\ &\quad + \sum_{0 < s \leq \cdot} \left( \Delta g(G_k(s), G_l(s)) - \sum_{i=k,l} g_i(G_k(s-), G_l(s-))\Delta G_i(s) \right), \end{aligned} \quad (\text{C.1})$$

where

$$d\langle G_i^c, G_j^c \rangle(s) = \lambda_i(s)\lambda_j(s)\alpha(s)ds. \quad (\text{C.2})$$

The sum in (C.1), using (3.2), may be written as

$$\begin{aligned} &\sum_{0 < s \leq \cdot} \left( \Delta g(G_k(s), G_l(s)) - \sum_{i=k,l} g_i(G_k(s-), G_l(s-))\Delta G_i(s) \right) \\ &= \int_0^\cdot \int_{\mathbb{R}^m} \left( g(G_k(s-), G_l(s-)) - g(G_k(s), G_l(s)) \right. \\ &\quad \left. - \sum_{i=k,l} g_i(G_k(s-), G_l(s-))\lambda_i^\top x \right) \mu(ds, dx) \\ &= \int_0^\cdot \int_{\mathbb{R}^m} \left( g(G_k(s-), G_l(s-)) - g(G_k(s), G_l(s)) \right. \\ &\quad \left. - \sum_{i=k,l} g_i(G_k(s-), G_l(s-))\lambda_i^\top x \right) F(s, dx) ds \\ &+ \int_0^\cdot \int_{\mathbb{R}^m} \left( g(G_k(s-), G_l(s-)) - g(G_k(s), G_l(s)) \right. \\ &\quad \left. - \sum_{i=k,l} g_i(G_k(s-), G_l(s-))\lambda_i^\top x \right) \tilde{\mu}(ds, dx). \end{aligned} \quad (\text{C.3})$$

Moreover

$$\begin{aligned} \int_0^\cdot g_i(G_k, G_l) dG_i &= \int_0^\cdot g_i(G_k, G_l) b_i ds + \int_0^t g_i(G_k, G_l) \sqrt{\alpha} \lambda_j^\top dW \\ &\quad + \int_0^\cdot \int_{\mathbb{R}^m} g_i(G_k, G_l) \lambda_j^\top x \tilde{\mu}(ds, dx). \end{aligned} \quad (\text{C.4})$$

Finally, putting all the pieces together we have that

$$\begin{aligned} Y_{kl}(t) &= Y_{kl}(0) + \int_0^t A_{kl}(s, L(s)) ds + \int_0^t B_{kl}^\top(s, L_{kl}(s)) dW(s) \\ &\quad + \int_0^t \int_{\mathbb{R}^m} C_{kl}(s, L_{kl}(s), x) (\mu(ds, dx) - F(\cdot, dx) ds), \end{aligned} \quad (\text{C.5})$$

where

$$\begin{aligned} A_{kl}(s, L(s)) &= \sum_{i=k,l} g_i(G_k(s-), G_l(s-)) b_i(s) \\ &\quad + \frac{1}{2} \sum_{i,j=k,l} \int_0^\cdot g_{ij}(G_k(s-), G_l(s-)) \lambda_i(s) \lambda_j(s) \alpha(s) \\ &\quad + \int_{\mathbb{R}^m} \left( g(G_k(s-) + \lambda_k^\top x, G_l(s-) + \lambda_l^\top x) - g(G_k(s), G_l(s)) \right. \\ &\quad \left. - \sum_{i=k,l} g_i(G_k(s-), G_l(s-)) \lambda_i^\top x \right) F(s, dx), \end{aligned} \quad (\text{C.6})$$

$$B_{kl}^\top(s, L_{kl}(s)) = \sum_{i=k,l} g_i(G_k(s-), G_l(s-)) \sqrt{\alpha(s)} \lambda_i^\top(s) \quad (\text{C.7})$$

and

$$\begin{aligned} C_{kl}(s, L_{kl}(s), x) &= \sum_{i=k,l} g_i(G_k(s-), G_l(s-)) \lambda_i^\top(s) x \\ &\quad + g(G_k(s-) + \lambda_k^\top x, G_l(s-) + \lambda_l^\top x) - g(G_k(s), G_l(s)) \\ &\quad - \sum_{i=k,l} g_i(G_k(s-), G_l(s-)) \lambda_i^\top x. \end{aligned} \quad (\text{C.8})$$

Here  $L_{kl}(s) := (L_k(s), L_l(s))$  and denotes that  $B_{kl}$  and  $C_{kl}$  depend on  $L_k$  and  $L_l$  (via  $G_k$  and  $G_l$ ).

## REFERENCES

- Andersen, L. and R. Brotherton-Ratcliffe (2005). Extended LIBOR market models with stochastic volatility. *J. Comput. Finance* 9, 1–40.
- Asmussen, S. and P. W. Glynn (2007). *Stochastic Simulation: Algorithms and Analysis*. Springer.

- Backus, D., S. Foresi, and L. Wu (2004). Accounting for biases in Black-Scholes. [SSRN/585623](#).
- Belomestny, D., S. Mathew, and J. Schoenmakers (2009). Multiple stochastic volatility extension of the LIBOR market model and its implementation. *Monte Carlo Methods Appl.* *15*, 285–310.
- Belomestny, D. and J. Schoenmakers (2011). A jump-diffusion LIBOR model and its robust calibration. *Quant. Finance* *11*, 529–546.
- Beveridge, C. (2010). Very long-stepping in the spot measure of the LIBOR market model. *Wilmott J.* *2*(6), 289–299.
- Black, F. (1976). The pricing of commodity contracts. *J. Financ. Econ.* *3*, 167–179.
- Brace, A., T. Dun, and G. Barton (2001). Towards a central interest rate model. In E. Jouini, J. Cvitanić, and M. Musiela (Eds.), *Option Pricing, Interest Rates and Risk Management*, pp. 278–313. Cambridge University Press.
- Brace, A., D. Gątarek, and M. Musiela (1997). The market model of interest rate dynamics. *Math. Finance* *7*, 127–155.
- Brenner, M. and M. Subrahmanyam (1994). A simple approach to option valuation and hedging in the Black-Scholes model. *Financial Analysts J.* *50*(2), 25–28.
- Carr, P., H. Geman, D. B. Madan, and M. Yor (2002). The fine structure of asset returns: An empirical investigation. *J. Business* *75*, 305–332.
- Cont, R. and P. Tankov (2004). *Financial Modelling with Jump Processes*. Chapman and Hall/CRC Press.
- Daniluk, A. and D. Gątarek (2005). A fully log-normal LIBOR market model. *Risk* *18*(9), 115–118.
- Dun, T., G. Barton, and E. Schlögl (2001). Simulated swaption delta-hedging in the lognormal forward LIBOR model. *Int. J. Theor. Appl. Finance* *4*, 677–709.
- Eberlein, E. and F. Özkan (2005). The Lévy LIBOR model. *Finance Stoch.* *9*, 327–348.
- Gatarek, D., P. Bachert, and R. Maksymiuk (2006). *The LIBOR Market Model in Practice*. Wiley.
- Glasserman, P. and S. G. Kou (2003). The term structure of simple forward rates with jump risk. *Math. Finance* *13*, 383–410.
- Glasserman, P. and N. Merener (2003a). Cap and swaption approximations in LIBOR market models with jumps. *J. Comput. Finance* *7*, 1–36.
- Glasserman, P. and N. Merener (2003b). Numerical solution of jump-diffusion LIBOR market models. *Finance Stoch.* *7*, 1–27.
- Glasserman, P. and X. Zhao (2000). Arbitrage-free discretization of log-normal forward LIBOR and swap rate models. *Finance Stoch.* *4*, 35–68.
- Hunter, C., P. Jäckel, and M. Joshi (2001). Getting the drift. *Risk* *14*, 81–84.
- Jacod, J. and A. N. Shiryaev (2003). *Limit Theorems for Stochastic Processes* (2nd ed.). Springer.
- Jamshidian, F. (1997). LIBOR and swap market models and measures. *Finance Stoch.* *1*, 293–330.



- Jamshidian, F. (1999). LIBOR market model with semimartingales. Working Paper, NetAnalytic Ltd.
- Joshi, M. and A. Stacey (2008). New and robust drift approximations for the LIBOR market model. *Quant. Finance* 8, 427–434.
- Kluge, W. (2005). *Time-inhomogeneous Lévy processes in interest rate and credit risk models*. Ph. D. thesis, Univ. Freiburg.
- Kohatsu-Higa, A. and P. Tankov (2010). Jump-adapted discretization schemes for Lévy-driven SDEs. *Stochastic Process. Appl.* 120, 2258–2285.
- Kurbanmuradov, O., K. Sabelfeld, and J. Schoenmakers (2002). Lognormal approximations to LIBOR market models. *J. Comput. Finance* 6, 69–100.
- Madan, D. B. and M. Yor (2008). Representing the CGMY and Meixner processes as time changed Brownian motions. *J. Comput. Finance* 12, 27–47.
- Miltersen, K. R., K. Sandmann, and D. Sondermann (1997). Closed form solutions for term structure derivatives with log-normal interest rates. *J. Finance* 52, 409–430.
- Papapantoleon, A. and D. Skovmand (2010). Picard approximation of stochastic differential equations and application to LIBOR models. Preprint, [arXiv/1007:3362](https://arxiv.org/abs/1007.3362).
- Poirot, J. and P. Tankov (2006). Monte Carlo option pricing for tempered stable (CGMY) processes. *Asia-Pac. Finan. Markets* 13, 327–344.
- Rosiński, J. (2001). Series representations of Lévy processes from the perspective of point processes. In O. E. Barndorff-Nielsen, T. Mikosch, and S. I. Resnick (Eds.), *Lévy Processes: Theory and Applications*, pp. 401–415. Birkhäuser.
- Schlögl, E. (2002). A multicurrency extension of the lognormal interest rate market models. *Finance Stoch.* 6, 173–196.
- Schoenmakers, J. (2005). *Robust LIBOR Modelling and Pricing of Derivative Products*. Chapman & Hall/CRC Press.
- Wilmott, P. (1998). *Derivatives: The Theory and Practice of Financial Engineering*. Wiley.
- Wu, L. and F. Zhang (2006). LIBOR market model with stochastic volatility. *J. Industr. Manag. Optim.* 2, 199–227.

INSTITUTE OF MATHEMATICS, TU BERLIN, STRASSE DES 17. JUNI 136, 10623 BERLIN, GERMANY

*E-mail address:* [papapan@math.tu-berlin.de](mailto:papapan@math.tu-berlin.de)

WEIERSTRASS INSTITUTE FOR APPLIED ANALYSIS AND STOCHASTICS, MOHRENSTR. 39, 10117 BERLIN, GERMANY

*E-mail address:* [schoenma@wias-berlin.de](mailto:schoenma@wias-berlin.de)

AARHUS SCHOOL OF BUSINESS, AARHUS UNIVERSITY, FUGLESANGS ALLÉ 4, 8210 AARHUS V, DENMARK

*E-mail address:* [davids@asb.dk](mailto:davids@asb.dk)