

**EDP-convergence for a linear reaction-diffusion system  
with fast reversible reaction**

Artur Stephan

submitted: December 5, 2020

Weierstrass Institute  
Mohrenstr. 39  
10117 Berlin  
Germany  
E-Mail: [artur.stephan@wias-berlin.de](mailto:artur.stephan@wias-berlin.de)

No. 2793  
Berlin 2020



---

2010 *Mathematics Subject Classification.* 49S05, 47J30, 35A15, 35K57, 92E20, 35Q84.

*Key words and phrases.* Markov process with detailed balance, linear reaction-diffusion system, gradient systems, gradient flows, evolutionary convergence, Energy-Dissipation-Balance, coarse-graining, microscopic equilibrium, Gamma-convergence.

Research supported by DFG via SFB 1114 (project no.235221301, subproject C05).

Edited by  
Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS)  
Leibniz-Institut im Forschungsverbund Berlin e. V.  
Mohrenstraße 39  
10117 Berlin  
Germany

Fax: +49 30 20372-303  
E-Mail: [preprint@wias-berlin.de](mailto:preprint@wias-berlin.de)  
World Wide Web: <http://www.wias-berlin.de/>

# EDP-convergence for a linear reaction-diffusion system with fast reversible reaction

Artur Stephan

## Abstract

We perform a fast-reaction limit for a linear reaction-diffusion system consisting of two diffusion equations coupled by a linear reaction. We understand the linear reaction-diffusion system as a gradient flow of the free energy in the space of probability measures equipped with a geometric structure, which contains the Wasserstein metric for the diffusion part and cosh-type functions for the reaction part. The fast-reaction limit is done on the level of the gradient structure by proving EDP-convergence with tilting. The limit gradient system induces a diffusion system with Lagrange multipliers on the linear slow-manifold. Moreover, the limit gradient system can be equivalently described by a coarse-grained gradient system, which induces a diffusion equation with a mixed diffusion constant for the coarse-grained slow variable.

## 1 Introduction

Considering two species  $X_1$  and  $X_2$ , which diffuse in a bounded medium  $\Omega \subset \mathbb{R}^d$  and react linearly  $X_1 \rightleftharpoons X_2$ , the evolution of their concentrations  $c = (c_1, c_2)$  can be described by the linear reaction-diffusion system

$$\begin{aligned}\dot{c}_1 &= \delta_1 \Delta c_1 - (\tilde{\alpha} c_1 - \tilde{\beta} c_2) \\ \dot{c}_2 &= \delta_2 \Delta c_2 + (\tilde{\alpha} c_1 - \tilde{\beta} c_2)\end{aligned}\tag{1.1}$$

complemented with no-flux boundary conditions and initial conditions, where  $\delta_1, \delta_2 > 0$  are diffusion coefficients for species  $X_1$  and  $X_2$ , respectively, and  $\tilde{\alpha}, \tilde{\beta} > 0$  are reaction rates describing the reaction speed of the linear reaction  $X_1 \rightleftharpoons X_2$ . The aim of the paper is to investigate system (1.1) if the reaction is much faster than the diffusion. To do this, we introduce a small parameter  $\varepsilon > 0$  and assume that the reaction rates are given by  $\tilde{\alpha} = \frac{1}{\varepsilon} \sqrt{\frac{\alpha}{\beta}}$ ,  $\tilde{\beta} = \frac{1}{\varepsilon} \sqrt{\frac{\beta}{\alpha}}$ . Then, the system (1.1) can be rewritten in an  $\varepsilon$ -dependent reaction-diffusion system

$$\begin{aligned}\dot{c}_1^\varepsilon &= \delta_1 \Delta c_1^\varepsilon - \frac{1}{\varepsilon} \left( \sqrt{\frac{\alpha}{\beta}} c_1^\varepsilon - \sqrt{\frac{\beta}{\alpha}} c_2^\varepsilon \right) \\ \dot{c}_2^\varepsilon &= \delta_2 \Delta c_2^\varepsilon + \frac{1}{\varepsilon} \left( \sqrt{\frac{\alpha}{\beta}} c_1^\varepsilon - \sqrt{\frac{\beta}{\alpha}} c_2^\varepsilon \right).\end{aligned}\tag{1.2}$$

Reaction systems and reaction-diffusion systems with slow and fast time scales have attracted a lot of attention in the last years [Eva80, HvdHP00, Bot03, BoP10, BoP11, BPR12, MuN11, MiS20, DDJ20, MPS20, PeR20]. Bothe and Hilhorst proved a fast-reaction limit  $\varepsilon \rightarrow 0$  for (1.2) in the following form.

**Theorem 1.1** ([BoH02]). *Let  $\Omega \subset \mathbb{R}^d$  be a domain with Lipschitz boundary. Let  $c_1^\varepsilon$  and  $c_2^\varepsilon$  be weak solutions of (1.2) with no-flux boundary conditions  $\nabla c_i^\varepsilon \cdot \nu = 0$  on  $\partial\Omega$ . Then  $c_1^\varepsilon \rightarrow c_1$  and  $c_2^\varepsilon \rightarrow c_2$  in  $L^2([0, T] \times \Omega)$  as  $\varepsilon \rightarrow 0$  and we have  $\frac{c_1}{\beta} = \frac{c_2}{\alpha}$ . Moreover, defining the coarse-grained concentration  $\hat{c} = c_1 + c_2$ , then  $\hat{c}$  solves the diffusion equation  $\dot{\hat{c}} = \hat{\delta} \Delta \hat{c}$  with a new mixed diffusion coefficient  $\hat{\delta} = \frac{\beta\delta_1 + \alpha\delta_2}{\alpha + \beta}$ .*

Essentially, the proof uses the free-energy as a Lyapunov function to derive  $\varepsilon$ -uniform bounds on the concentrations  $c_i^\varepsilon$  and their gradients  $\nabla c_i^\varepsilon$ , which is then used to prove convergence towards the slow manifold  $\{c \in [0, \infty]^2 \mid \alpha c_1 = \beta c_2\}$ . This proof also works for nonlinear reactions once  $\varepsilon$ -uniform  $L^\infty$ -estimates are established (see [BoH02]). On the linear slow manifold, one easily verifies that the coarse grained concentration  $\hat{c} := \frac{\alpha + \beta}{\beta} c_1 = \frac{\alpha + \beta}{\alpha} c_2 = c_1 + c_2$  solves  $\dot{\hat{c}} = \hat{\delta} \Delta \hat{c}$  where  $\hat{\delta} = \frac{\beta\delta_1 + \alpha\delta_2}{\alpha + \beta}$  is the effective mixed diffusion coefficient.

In this work, we are not primarily interested in convergence of solutions of system (1.2). Instead, we perform the fast-reaction limit on the level of the underlying variational structure, which then implies convergence of solutions as a byproduct. Our starting point is that reaction-diffusion systems such as (1.2) can be written as a gradient flow equation induced by a gradient system  $(Q, \mathcal{E}, \mathcal{R}_\varepsilon^*)$ , where the state space  $Q$  is the space of probability measures  $Q = \text{Prob}(\Omega \times \{1, 2\})$  and the driving functional is the free-energy  $\mathcal{E}(\mu) = \int_\Omega \sum_{j=1}^2 E_B\left(\frac{c_j}{w_j}\right) w_j dx$  for measures  $\mu = c dx$ , with the Boltzmann function  $E_B(r) = r \log r - r + 1$  and the (in general space dependent) stationary measure  $w = (w_1, w_2)^\top$ . The dissipation potential  $\mathcal{R}_\varepsilon^*$  that determines the geometry of the underlying space is given by two parts  $\mathcal{R}_\varepsilon^* = \mathcal{R}_{\text{diff}}^* + \mathcal{R}_{\text{react}, \varepsilon}^*$  describing the diffusion and reaction separately. Since the pioneering work of Otto and coauthors [JKO98, Ott01] it is known that diffusion has to be understood as a gradient system driven by the free-energy in the space of probability measures equipped with the Wasserstein distance. The corresponding dissipation potential  $\mathcal{R}_{\text{diff}}^*$  is quadratic and given by

$$\mathcal{R}_{\text{diff}}^*(\mu, \xi) = \frac{1}{2} \int_\Omega \sum_{i=1}^2 \delta_i |\nabla \xi_i|^2 d\mu_i.$$

Later Mielke [Mie11] proposed a quadratic gradient structure also for reaction-diffusion systems with the same driving functional. Geometric properties of that gradient structure were investigated in [LiM13, GK\*20]. Here, we are not interested in that gradient structure, but use a different, the so-called *cosh-type gradient structure*, where the reaction part is given by

$$\mathcal{R}_{\text{react}, \varepsilon}^*(\mu, \xi) = \frac{1}{\varepsilon} \int_\Omega C^*(\xi_1(x) - \xi_2(x)) \sqrt{d\mu_1 d\mu_2},$$

with  $C^*(r) = 4(\cosh(r/2) - 1)$ . Setting  $\mathcal{R}_\varepsilon^* = \mathcal{R}_{\text{diff}}^* + \mathcal{R}_{\text{react}, \varepsilon}^*$ , the reaction-diffusion system (1.2) can now be written as a gradient flow equation

$$\dot{\mu} = \partial_\xi \mathcal{R}_\varepsilon^*(\mu, -D\mathcal{E}(\mu)).$$

Although there are many gradient structures for (1.2) (see e.g. [MiS20, Sect. 4]) and the cosh-type gradient structure entails several technical difficulties as defining a nonlinear kinetic relation and not inducing metric on  $Q$ , it nevertheless has several significant features. Historically, it has its origin in [Mar15] where, following thermodynamic considerations, chemical reactions are written in exponential terms. In recent years, the cosh-gradient structure has been derived via a large-deviation principle [MPR14, MP\*17], and it was shown that it is stable under limit processes [LM\*17] that are similar to our approach. Moreover, it does not explicitly depend on the stationary measure  $w$ , which allows for an

rigorous distinction between the energetic and dissipative part [MiS20]. This is physically reasonable because a change of the energy by an external field should not influence the geometric structure of the underlying space.

The goal of the paper is to construct an effective gradient system  $(Q, \mathcal{E}, \mathcal{R}_{\text{eff}}^*)$  and perform the limit  $(Q, \mathcal{E}, \mathcal{R}_{\varepsilon}^*) \rightarrow (Q, \mathcal{E}, \mathcal{R}_{\text{eff}}^*)$  as  $\varepsilon \rightarrow 0$ . For this, we use the notion of *convergence of gradient systems in the sense of the energy-dissipation principle*, shortly called *EDP-convergence*. EDP-convergence was introduced in [DFM18] and further developed in [MMP20, MiS20] and is based on the dissipation functional

$$\mathfrak{D}_{\varepsilon}^{\eta}(\mu) = \int_0^T \mathcal{R}_{\varepsilon}(\mu, \dot{\mu}) + \mathcal{R}_{\varepsilon}^*(\mu, \eta - D\mathcal{E}(\mu)) dt$$

which, for solutions  $\mu$  of the gradient flow equation describes the total dissipation between initial time  $\mathcal{E}(\mu(0))$  and final time  $\mathcal{E}(\mu(T))$ , and can now be defined for general trajectories  $\mu \in L^1([0, T], Q)$ . The notion of EDP-convergence with tilting requires  $\Gamma$ -convergences of the energies  $\mathcal{E}_{\varepsilon} \xrightarrow{\Gamma} \mathcal{E}_0$  and of the dissipation functionals  $\mathfrak{D}_{\varepsilon}^{\eta} \xrightarrow{\Gamma} \mathfrak{D}_0^{\eta}$  in suitable topologies, such that for all tilts  $\eta$  the limit  $\mathfrak{D}_0^{\eta}$  has the form  $\mathfrak{D}_0^{\eta}(\mu) = \int_0^T \mathcal{R}_{\text{eff}}(\mu, \dot{\mu}) + \mathcal{R}_{\text{eff}}^*(\mu, \eta - D\mathcal{E}_0(\mu)) dt$ , see Section 2.2 for a precise definition. Importantly, the effective dissipation potential  $\mathcal{R}_{\text{eff}}$  in the  $\Gamma$ -limit is independent of the tilts, hence allowing for extended energies. In our situation, the tilts  $\eta$  correspond to an external potential  $V = (V_1, V_2)$  added to the energy  $\mathcal{E}$ . On the level of the PDE, the starting reaction-diffusion system is extended to a reaction-drift-diffusion system of the form

$$\frac{d}{dt} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \text{div} \left( \begin{pmatrix} \delta_1 \nabla c_1 \\ \delta_2 \nabla c_2 \end{pmatrix} + \begin{pmatrix} \delta_1 c_1 \nabla V_1 \\ \delta_2 c_2 \nabla V_2 \end{pmatrix} \right) + \frac{1}{\varepsilon} \begin{pmatrix} -\sqrt{\frac{\alpha}{\beta}} e^{\frac{V_1 - V_2}{2}} & \sqrt{\frac{\beta}{\alpha}} e^{\frac{V_2 - V_1}{2}} \\ \sqrt{\frac{\alpha}{\beta}} e^{\frac{V_1 - V_2}{2}} & -\sqrt{\frac{\beta}{\alpha}} e^{\frac{V_2 - V_1}{2}} \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix}.$$

The main result of the paper is Theorem 4.3 which asserts tilt EDP-convergence of  $(Q, \mathcal{E}, \mathcal{R}_{\varepsilon}^*)$  to  $(Q, \mathcal{E}, \mathcal{R}_{\text{eff}}^*)$  as  $\varepsilon \rightarrow 0$  where the effective dissipation potential is given by

$$\mathcal{R}_{\text{eff}}^* = \mathcal{R}_{\text{diff}}^* + \chi_{\{\xi_1 = \xi_2\}},$$

where  $\chi_A$  is the characteristic function of convex analysis taking values zero in  $A$  and infinity otherwise. The effective dissipation potential describes diffusion but restricts the chemical potential  $\xi = (\xi_1, \xi_2)$  to a linear submanifold. The induced gradient flow equation of the gradient system  $(Q, \mathcal{E}, \mathcal{R}_{\text{eff}}^*)$  is then given by a system of drift-diffusion equations on a linear submanifold with a space and time dependent Lagrange multiplier  $\lambda$

$$\begin{aligned} \dot{c}_1 &= \text{div}(\delta_1 \nabla c_1 + \delta_1 c_1 \nabla V_1) - \lambda \\ \dot{c}_2 &= \text{div}(\delta_2 \nabla c_2 + \delta_2 c_2 \nabla V_2) + \lambda \end{aligned} \quad , \quad \frac{c_1}{\beta e^{-V_1}} = \frac{c_2}{\alpha e^{-V_2}}.$$

Moreover, as an immediate consequence of Theorem 4.3, we obtain that the effective gradient flow equation can be equivalently described as a drift-diffusion equation of the coarse-grained concentration  $\hat{c}$ , see Proposition 4.5. Introducing the mixed diffusion coefficient  $\hat{\delta}^V = \frac{\delta_1 \beta e^{-V_1} + \delta_2 \alpha e^{-V_2}}{\beta e^{-V_1} + \alpha e^{-V_2}}$  and the mixed potential  $\hat{V} = -\log\left(\frac{\beta}{\alpha + \beta} e^{-V_1} + \frac{\alpha}{\alpha + \beta} e^{-V_2}\right)$ , we obtain

$$\hat{c} = \text{div} \left( \hat{\delta}^V \nabla \hat{c} + \hat{\delta}^V \hat{c} \nabla \hat{V} \right),$$

which is in accordance with [BoH02] in the potential-free case  $V = \text{const}$ . Moreover, we obtain a natural coarse-grained gradient structure  $(\hat{Q}, \hat{\mathcal{E}}, \hat{\mathcal{R}}^*)$ , where  $\hat{Q} = \text{Prob}(\Omega)$  is the coarse-grained state space and  $\hat{\mathcal{E}}, \hat{\mathcal{R}}^*$  are the coarse-grained energy functional and dissipation potential, respectively.

Interestingly, this coarse-grained gradient structure  $(\hat{Q}, \hat{\mathcal{E}}, \hat{\mathcal{R}}^*)$  contains the same information as the effective gradient structure  $(Q, \mathcal{E}, \mathcal{R}_{\text{eff}}^*)$ , although defined on a smaller state space, see Proposition 4.5.

The result on tilt EDP-convergence is an immediate consequence of the  $\Gamma$ -convergence result of the dissipation functional  $\mathcal{D}_\varepsilon^\eta$  (Theorem 5.12). The primal dissipation potential  $\mathcal{R}_\varepsilon$  is given by an infimal sum consisting of diffusion fluxes and reaction fluxes coupled via a generalized continuity equation, see Section 3.3. Theorem 5.12 follows from the following observations:  $\mathcal{R}_\varepsilon^*$  converges monotonically to a singular limit  $\mathcal{R}_{\text{eff}}^*$ , the primal dissipation potentials  $\mathcal{R}_\varepsilon$  degenerate. It is not possible to control the rates of  $\dot{\mu}_1$  and  $\dot{\mu}_2$  separately by  $\mathcal{R}_\varepsilon$ , since the reaction flux between both species may become unbounded. Instead, it is possible to prove compactness for the sum (or slow variable)  $\mu_1 + \mu_2$  by  $\mathcal{R}_\varepsilon$ , and proving convergence towards the slow-manifold where an equilibration takes place, i.e.  $\alpha c_1^\varepsilon - \beta c_2^\varepsilon \rightarrow 0$ . The two pieces of complementary information provide strong convergence of the densities  $c^\varepsilon$  in  $L^1([0, T] \times \Omega)$ . This procedure has been already successfully applied for linear and nonlinear reaction systems [MiS20, MPS20] and is here applied to a space-dependent evolution system. A posteriori we conclude that the limit measure  $\mu^0$  has indeed an absolutely continuous representative using results from [AGS05]. The construction of the recovery sequence relies on the fact that the limit dissipation functional can be equivalently written as a functional of coarse-grained variables. Only the reaction flux, which is present for positive  $\varepsilon > 0$  and hidden for  $\varepsilon = 0$ , has to be reconstructed. One observes that diffusion causes the reaction flux on an infinitesimally small scale. Since the dissipation functional considers also fluctuations which may be not strictly positive and not smooth in contrast to the solution of the linear reaction diffusion system (1.2), the construction of a recovery sequence is completed by a suitable approximation argument.

Let us finally mention, that the same results can also be established for reaction-diffusion systems, where more than two species are involved. Applying the coarse-graining and reconstruction machinery as developed in [MiS20], a similar  $\Gamma$ -convergence result for the dissipation functional can be proved. For notational convenience we restrict to the two-species situation and briefly discuss the multi-species case in Section 6. We refer also to [Ste21], where coarse-graining and reconstruction for concentrations as well as the fluxes is developed.

## 2 Gradient structures

### 2.1 Gradient systems and the energy-dissipation principle

Let us briefly recall what we mean with a gradient system. Following [Mie16], we call a triple  $(Q, \mathcal{E}, \mathcal{R})$  a *gradient system* if

- (1)  $Q$  is a closed convex subset of a Banach space  $X$
- (2)  $\mathcal{E} : Q \rightarrow \mathbb{R}_\infty := \mathbb{R} \cup \{\infty\}$  is a functional (such as the free energy)
- (3)  $\mathcal{R} : Q \times X \rightarrow \mathbb{R}_\infty$  is a dissipation potential, which means that for any  $u \in Q$  the functional  $\mathcal{R}(u, \cdot) : X \rightarrow \mathbb{R}_\infty$  is lower semicontinuous (lsc), nonnegative and convex, and it satisfies  $\mathcal{R}(u, 0) = 0$ .

We define the dual dissipation potential  $\mathcal{R}^* : Q \times X^* \rightarrow [0, \infty]$  using the Legendre transform via

$$\mathcal{R}^*(u, \xi) = (\mathcal{R}(u, \cdot))^*(\xi) = \sup_{v \in X} \{ \langle v, \xi \rangle - \mathcal{R}(u, v) \} .$$

The gradient system is uniquely described by  $(Q, \mathcal{E}, \mathcal{R})$  or, equivalently by  $(Q, \mathcal{E}, \mathcal{R}^*)$  and, in particular, in this paper we use the second representation.

The dynamics of a gradient system can be formulated in different ways as an equation in  $X$ , in  $\mathbb{R}$  or in  $X^*$  (the dual Banach space of  $X$ ), respectively:

- (1) **Force balance in  $X^*$ :**  $0 \in \partial_{\dot{u}} \mathcal{R}(u, \dot{u}) + D\mathcal{E}(u) \in X^*$ ,
- (2) **Power balance in  $\mathbb{R}$ :**  $\mathcal{R}(u, \dot{u}) + \mathcal{R}^*(u, -D\mathcal{E}(u)) = -\langle D\mathcal{E}(u), \dot{u} \rangle$ ,
- (3) **Rate equation in  $X$ :**  $\dot{u} \in \partial_{\xi} \mathcal{R}^*(u, -D\mathcal{E}(u)) \in X$ .

(Here,  $\partial$  denotes the subdifferential of convex analysis.) Equations (1) and (3) are called *gradient flow equation* associated with  $(Q, \mathcal{E}, \mathcal{R}^*)$ . The equivalent formulations rely on the following fact: Let  $X$  be a reflexive Banach space and  $\Psi : X \rightarrow \mathbb{R}_{\infty}$  be a proper, convex and lsc. Then for every  $\xi \in X^*$  and  $v \in X$  the following five statements, the so-called *Legendre-Fenchel-equivalences*, are equivalent:

$$\begin{array}{lll} v \in \operatorname{Argmin}_{w \in X} (\Psi(w) - \langle \xi, w \rangle) & \Leftrightarrow & \Psi(v) + \Psi^*(\xi) = \langle \xi, v \rangle \\ \Leftrightarrow & & \Leftrightarrow \\ v \in \partial \Psi^*(\xi) & \Leftrightarrow & \Psi(v) + \Psi^*(\xi) = \langle \xi, v \rangle \\ & \Leftrightarrow & \xi \in \operatorname{Argmin}_{\eta \in X^*} (\Psi^*(\eta) - \langle \eta, v \rangle) \end{array}$$

Especially the second dynamic formulation, the power balance (2), is interesting for us. Integrating the power balance (2) in time from 0 to  $T$  and using the chain rule for the time-derivative of  $t \mapsto \mathcal{E}(u(t))$ , we get another equivalent formulation of the dynamics of the gradient system, which is called *Energy-Dissipation-Balance*:

$$(EDB) \quad \mathcal{E}(u(T)) + \int_0^T [\mathcal{R}(u, \dot{u}) + \mathcal{R}^*(u, -D\mathcal{E}(u))] dt = \mathcal{E}(u(0)). \quad (2.1)$$

Equation (EDB) compares the energy of the system at time  $t = 0$  and at time  $t = T$ , the difference is described by the total dissipation from  $t = 0$  to  $t = T$ . This gives rise to another definition: We define the *De Giorgi dissipation functional* as

$$\mathfrak{D}(u) = \int_0^T [\mathcal{R}(u, \dot{u}) + \mathcal{R}^*(u, -D\mathcal{E}(u))] dt,$$

for  $u \in W^{1,1}([0, T], Q)$  and extend it to infinity otherwise. The following *energy-dissipation principle* provides the definition for solutions of the gradient flow equation, see e.g. [AGS05, Prop. 1.4.1], [AM\*12, Def. 1.1], [MMP20, Thm 2.5].

**Definition 2.1.** We say  $u \in W^{1,1}([0, T], Q)$  is a solution of the gradient flow equation (1) or (3) induced by the gradient system  $(Q, \mathcal{E}, \mathcal{R}^*)$ , if  $\mathcal{E}(u(0)) < \infty$  and the energy-dissipation balance holds.

## 2.2 Definition of EDP-convergence

The definition of EDP-convergence for gradient systems relies on the notion of  $\Gamma$ -convergence for functionals (cf. [Dal93]). If  $Y$  is a Banach space and  $I_{\varepsilon} : Y \rightarrow \mathbb{R}_{\infty}$  we write  $I_{\varepsilon} \xrightarrow{\Gamma} I_0$  and

$I_\varepsilon \xrightarrow{\Gamma} I_0$  for  $\Gamma$ -convergence in the strong and weak topology, respectively. If both holds this is called Mosco-convergence and written as  $I_\varepsilon \xrightarrow{M} I_0$ .

For families of gradient systems  $(X, \mathcal{E}_\varepsilon, \mathcal{R}_\varepsilon)$ , three different levels of EDP-convergence are introduced and discussed in [DFM18, MMP20], called simple EDP-convergence, EDP-convergence with tilting and contact EDP-convergence with tilting. EDP-convergence with tilting is the strongest notion, since it implies the other two notions. Here we will only use the first two notions. For all three notions the choice of weak or strong topology is still to be decided according to the specific problem.

**Definition 2.2** (Simple EDP-convergence). A family of gradient structures  $(Q, \mathcal{E}_\varepsilon, \mathcal{R}_\varepsilon)$  is said to *EDP-converge* to the gradient system  $(Q, \mathcal{E}_0, \mathcal{R}_{\text{eff}})$  if the following conditions hold:

- 1  $\mathcal{E}_\varepsilon \xrightarrow{\Gamma} \mathcal{E}_0$  on  $Q \subset X$ ;
- 2  $\mathfrak{D}_\varepsilon$  strongly  $\Gamma$ -converges to  $\mathfrak{D}_0$  on  $L^1([0, T], Q)$  conditioned to bounded energies (we write  $\mathfrak{D}_\varepsilon \xrightarrow{\Gamma_{\mathbb{R}}} \mathfrak{D}_0$ ), i.e. we
  - 2.1 (Liminf-estimate) For all strongly converging families  $u_\varepsilon \rightarrow u$  in  $L^1([0, T], Q)$  which satisfy  $\sup_{\varepsilon > 0} \text{ess sup}_{t \in [0, T]} \mathcal{E}_\varepsilon(u_\varepsilon(t)) < \infty$ , we have  $\liminf_{\varepsilon \rightarrow 0^+} \mathfrak{D}_\varepsilon(u_\varepsilon) \geq \mathfrak{D}_0(u)$ ,
  - 2.2 (Limsup-estimate) For all  $\tilde{u} \in L^1([0, T], Q)$  there exists a strongly converging family  $u_\varepsilon \rightarrow \tilde{u}$  in  $L^1([0, T], Q)$  with  $\sup_{\varepsilon > 0} \text{ess sup}_{t \in [0, T]} \mathcal{E}_\varepsilon(\tilde{u}_\varepsilon(t)) < \infty$  such that we have  $\limsup_{\varepsilon \rightarrow 0^+} \mathfrak{D}_\varepsilon(u_\varepsilon) \leq \mathfrak{D}_0(\tilde{u})$ .
- 3 There is an effective dissipation potential  $\mathcal{R}_{\text{eff}} : Q \times X \rightarrow \mathbb{R}_\infty$  such that  $\mathfrak{D}_0$  takes the form of a dual sum, namely  $\mathfrak{D}_0(u) = \int_0^T \{\mathcal{R}_{\text{eff}}(u, \dot{u}) + \mathcal{R}_{\text{eff}}^*(u, -D\mathcal{E}_{\text{eff}}(u))\} dt$ .

Similarly, one can also use weak  $\Gamma$ - or Mosco-convergence conditioned to bounded energy, which we will then write as  $\mathfrak{D}_\varepsilon \xrightarrow{\Gamma_{\mathbb{R}}} \mathfrak{D}_0$  and  $\mathfrak{D}_\varepsilon \xrightarrow{M_{\mathbb{R}}} \mathfrak{D}_0$ . In fact, for our fast-slow reaction systems we are going to prove  $\mathfrak{D}_\varepsilon \xrightarrow{M_{\mathbb{R}}} \mathfrak{D}_0$ .

A general feature of EDP-convergence is that, under suitable conditions, solutions  $u$  of the gradient flow equation  $\dot{u} = \partial_\xi \mathcal{R}_{\text{eff}}^*(u, -D\mathcal{E}_0(u))$  of the effective gradient system  $(X, \mathcal{E}_0, \mathcal{R}_{\text{eff}})$  are indeed limits of solutions  $u^\varepsilon$  of the gradient flow equation  $\dot{u} = \partial_\xi \mathcal{R}_\varepsilon^*(u, -D\mathcal{E}_\varepsilon(u))$ , see e.g. [Bra13, Thm. 11.3], [Mis20, Lem. 3.4] and [MMP20, Lem. 2.8].

A strengthening of simple EDP-convergence is the so-called *EDP-convergence with tilting*. This notion involves the tilted energy functionals  $\mathcal{E}_\varepsilon^\eta : Q \ni u \mapsto \mathcal{E}_\varepsilon(u) - \langle \eta, u \rangle$ , where the tilt  $\eta$  (also called *external loading*) varies through the whole dual space  $X^*$ .

**Definition 2.3** (EDP-convergence with tilting (cf. [MMP20, Def. 2. 14])). A family of gradient structures  $(Q, \mathcal{E}_\varepsilon, \mathcal{R}_\varepsilon)$  is said to *EDP-converge with tilting* to the gradient system  $(Q, \mathcal{E}_0, \mathcal{R}_{\text{eff}})$ , if for all tilts  $\eta \in X^*$  we have  $(Q, \mathcal{E}_\varepsilon^\eta, \mathcal{R}_\varepsilon)$  EDP-converges to  $(Q, \mathcal{E}_0^\eta, \mathcal{R}_{\text{eff}})$ .

Clearly, we have that  $\mathcal{E}_\varepsilon \xrightarrow{\Gamma} \mathcal{E}_0$  implies  $\mathcal{E}_\varepsilon^\eta \xrightarrow{\Gamma} \mathcal{E}_0^\eta$  for all  $\eta \in X^*$  (and similarly for weak  $\Gamma$ -convergence), since the linear tilt  $u \mapsto -\langle \eta, u \rangle$  is weakly continuous. The main and nontrivial assumption is that additionally

$$\mathfrak{D}_\varepsilon^\eta : u \mapsto \int_0^T \{\mathcal{R}_\varepsilon(u, \dot{u}) + \mathcal{R}_\varepsilon^*(u, \eta - D\mathcal{E}_\varepsilon(u))\} dt$$



$\Gamma$ -converges in  $L^1([0, T], Q)$  to  $\mathfrak{D}_0^\eta$  for all  $\eta \in X^*$  and that this limit  $\mathfrak{D}_0^\eta$  is given in  $\mathcal{R} \oplus \mathcal{R}^*$ -form with  $\mathcal{R}_{\text{eff}}$  via

$$\mathfrak{D}_0^\eta(u) = \int_0^T \{ \mathcal{R}_{\text{eff}}(u, \dot{u}) + \mathcal{R}_{\text{eff}}^*(u, \eta - D\mathcal{E}_{\text{eff}}(u)) \} dt.$$

The main point is that  $\mathcal{R}_{\text{eff}}$  remains independent of  $\eta \in X^*$ . We refer to [MMP20] for a discussion of this and the other two notions of EDP-convergence.

### 3 Gradient system of reaction-diffusion systems

In this section, we present the gradient system  $(Q, \mathcal{E}, \mathcal{R}_\varepsilon^*)$ , which induces the reaction-diffusion system (1.2). In Section 3.2 we derive the gradient flow equation of the gradient system including general tilts of the energy. In Section 3.3 we compute the primal dissipation potential  $\mathcal{R}_\varepsilon$ , which is only implicitly given via a infimal-convolution, and the total dissipation functional  $\mathfrak{D}_\varepsilon^\eta$ , which will be the main object of interest in Section 4. In Section 3, the computations are basically formal; the precise functional analytic setting is presented in Section 4 which also includes the  $\Gamma$ -convergence and EDP-convergence result.

#### 3.1 Gradient structure for the linear reaction system

Although a gradient system induces a unique gradient flow equation, a general evolution equation can often be described by many different gradient systems. The choice of the gradient structure is a question of modeling since it adds thermodynamic information to the system, which is not inherent in the evolution equation itself. Here, we follow the pioneering work of Otto and coauthors [JKO98, Ott01] who showed that certain diffusion type equations can be understood as a gradient flow equation of the free energy in the space of probability measures equipped with the Wasserstein distance. Later Mielke proposed a gradient structure for a reaction diffusion system satisfying detailed balance [Mie11]. For a system with two species with a reversible reaction detailed balance is always satisfied. For the reaction part, we use the gradient structure which has been derived via a large-deviation principle from a microscopic Markov process in [MPR14]. We refer also to [Ren18], where our choice of gradient structure has been formally derived.

The gradient system  $(Q, \mathcal{E}, \mathcal{R}_\varepsilon^*)$  is defined as follows: The state space is the space of probability measure on  $Q \times \{1, 2\}$

$$Q := \text{Prob}(\Omega \times \{1, 2\}) = \{ \mu = (\mu_1, \mu_2) \in \mathbb{R}^2 : \mu_i \in \mathcal{M}(\Omega), \mu_i \geq 0, \sum_{i=1}^2 \mu_i(\Omega) = 1 \},$$

where we assume that  $\Omega \subset \mathbb{R}^d$  is a compact domain with normalized mass  $|\Omega| = 1$ . The driving energy functional  $\mathcal{E} : Q \rightarrow \mathbb{R}_\infty := \mathbb{R} \cup \{\infty\}$  is the free-energy of the reaction-diffusion system. It is finite for measures  $\mu = (\mu_1, \mu_2)$  with Lebesgue density  $c = (c_1, c_2)$  only and has the form

$$\mathcal{E}(\mu) := \begin{cases} \int_\Omega \sum_{j=1}^2 E_B \left( \frac{c_j}{w_j} \right) w_j dx, & \text{if } \mu = c \cdot dx \\ \infty, & \text{otherwise.} \end{cases} \quad (3.1)$$

where the Boltzmann function is defined as  $E_B(r) = r \log r - r + 1$  and the positive stationary measure is given by  $w = \frac{1}{\alpha + \beta} (\beta, \alpha)^T$ . Note that the stationary measure  $w$  as well as the energy  $\mathcal{E}$

is  $\varepsilon$ -independent. The derivative of the energy  $\mathcal{E}$  is only defined in its domain, i.e. for measures with Lebesgue density  $c$ , and has the form

$$D\mathcal{E}(\mu) = \sum_{j=1}^2 (\log c_j - \log w_j) = \sum_{j=1}^2 \left( \log \frac{c_j}{w_j} \right).$$

As the equation splits into a diffusion and reaction part, so does the dual dissipation functional. We define

$$\mathcal{R}_\varepsilon^*(\mu, \xi) := \mathcal{R}_{\text{diff}}^*(\mu, \xi) + \mathcal{R}_{\text{react},\varepsilon}^*(\mu, \xi)$$

where

$$\begin{aligned} \mathcal{R}_{\text{diff}}^*(\mu, \xi) &:= \frac{1}{2} \int_{\Omega} \sum_{j=1}^2 \delta_j |\nabla \xi_j(x)|^2 d\mu_j, \\ \mathcal{R}_{\text{react},\varepsilon}^*(\mu, \xi) &:= \frac{1}{\varepsilon} \int_{\Omega} C^*(\xi_1(x) - \xi_2(x)) d\sqrt{\mu_1 \mu_2}, \end{aligned}$$

where we use the cosh-function  $C^*(x) = 4(\cosh(x/2) - 1)$  and for measures  $\mu$  with Lebesgue density  $c$  we have  $d\sqrt{\mu_1 \mu_2} := \sqrt{c_1 c_2} dx$ .

The diffusion part  $\mathcal{R}_{\text{diff}}^*$  induces the Wasserstein distance on  $Q$ . The  $\varepsilon$ -dependent reaction part  $\mathcal{R}_{\text{react},\varepsilon}^*$  forces the evolution close to a linear submanifold given by

$$\mathcal{R}_{\text{react},\varepsilon}^*(\mu, -D\mathcal{E}(\mu)) = 0 \Leftrightarrow \alpha c_1 - \beta c_2 = 0.$$

Note, that since  $\mathcal{R}_{\text{react},\varepsilon}^*$  is not 2-homogeneous, it does not define a metric on  $Q$ . We refer to [PR\*20] which treats similar and general dissipation potentials and understands them as generalized transport costs on discrete spaces. Note that  $\mathcal{R}_\varepsilon^*$  does not depend on the stationary measure  $w$  explicitly, as highlighted in [MiS20].

### 3.2 The tilted gradient flow equation

In this section, we derive the gradient flow equation of the gradient system  $(Q, \mathcal{E}, \mathcal{R}_\varepsilon^*)$ . To exploit the full information of the dissipation potential, we consider general tilted energies. First, we present how a change of energy by a linear tilt corresponds to a change of stationary measure, and secondly, we compute the induced gradient flow equation.

Let us first consider two free energies (3.1) with different stationary measures  $w, \tilde{w}$ , which may be space dependent but are assumed to be positive. Assuming a density  $\mu = c dx$  and using  $\sum_i \int_{\Omega} w_i dx = \sum_i \int_{\Omega} c_i dx = 1$  (where we used  $|\Omega| = 1$ ), we have

$$\mathcal{E}(\mu) = \sum_{i=1}^2 \int_{\Omega} E_B \left( \frac{c_i}{w_i} \right) w_i dx = \sum_{i=1}^2 \int_{\Omega} \{c_i \log c_i - c_i \log w_i\} dx.$$

In particular, we conclude that  $\tilde{\mathcal{E}}(\mu) + \sum_{i=1}^2 \int_{\Omega} c_i \log \tilde{w}_i dx = \mathcal{E}(\mu) + \sum_{i=1}^2 \int_{\Omega} c_i \log w_i dx$  which implies

$$\tilde{\mathcal{E}}(\mu) = \mathcal{E}(\mu) + \sum_{i=1}^2 \int_{\Omega} c_i \log \left( \frac{w_i}{\tilde{w}_i} \right) dx.$$

Hence, changing the underlying stationary measure corresponds to a linear tilt of the energy by a two component potential  $V = (V_1, V_2)$  where  $V_i = \log\left(\frac{w_i}{\bar{w}_i}\right)$ . On the other hand, a tilted energy has a different stationary measure as its minimum. To compute the new stationary measure, we introduce tilted energies

$$\mathcal{E}^V(\mu) := \mathcal{E}(\mu) + \sum_{i=1}^2 \int_{\Omega} V_i d\mu_i,$$

where  $V \in C^1(\Omega, \mathbb{R}^2)$  is a two component smooth potential. Moreover, we introduce  $\eta_i := e^{-V_i}$  and clearly, we have  $\eta_i > 0$  on  $\Omega \subset \mathbb{R}^d$ . We compute the stationary state  $w^V$  by minimizing  $\mathcal{E}^V$  on the space  $Q = \text{Prob}(\Omega \times \{1, 2\})$ . We obtain the space dependent stationary measure

$$w_i^V = \frac{1}{Z} w_i e^{-V_i}, \text{ where } Z := \sum_{i=1}^2 \int_{\Omega} w_i e^{-V_i} dx. \quad (3.2)$$

Next, we compute the tilted gradient flow equation  $\dot{c} = \partial_{\xi} \mathcal{R}_{\varepsilon}^*(\mu, -D\mathcal{E}^V(\mu))$ , which is induced by the gradient system  $(Q, \mathcal{E}^V, \mathcal{R}_{\varepsilon}^* = \mathcal{R}_{\text{diff}}^* + \mathcal{R}_{\text{react}, \varepsilon}^*)$ . First, we observe that  $\mathcal{E}(\mu) < \infty$  if and only if  $\mathcal{E}^V(\mu) < \infty$ . Inserting  $\xi_i = (-D\mathcal{E}^V(\mu))_i$  into  $\partial_{\xi} \mathcal{R}_{\text{diff}, \varepsilon}^*(\mu, \xi)$ , we see that

$$\partial_{\xi} \mathcal{R}_{\text{diff}}^*(\mu, \cdot)|_{\xi = -D\mathcal{E}^V(\mu)} = -(\text{div}(\delta_i c_i \nabla(-\log(c_i/w_i) - V_i)))_{i=1,2} = \text{div}(\delta_i \nabla c_i + \delta_i c_i \nabla V_i)_{i=1,2},$$

which is a system of two uncoupled drift-diffusion equations or Fokker-Planck equations for the concentrations  $c_i$  where the fluxes are given by a diffusion part  $-\delta_i \nabla c_i$  and a drift part  $-\delta_i c_i \nabla V_i$ .

For the reaction part of the dual dissipation potential, we insert  $\xi_i = (-D\mathcal{E}^V(\mu))_i$  into the term  $\partial_{\xi} \mathcal{R}_{\text{react}, \varepsilon}^*(\mu, \cdot)$ . It readily verifies the identity  $(C^*)'(\log p - \log q) = \frac{p-q}{\sqrt{pq}}$  for the cosh-function and conclude

$$\sqrt{c_1 c_2} (C^*)'(\xi_1(x) - \xi_2(x))|_{\xi = -D\mathcal{E}^V(\mu)} = \sqrt{c_1 c_2} \frac{\frac{c_2}{w_2 \eta_2} - \frac{c_1}{w_1 \eta_1}}{\sqrt{\frac{c_1}{w_1 \eta_1} \frac{c_2}{w_2 \eta_2}}} = \sqrt{w_1 \eta_1 w_2 \eta_2} \left( \frac{c_2}{w_2 \eta_2} - \frac{c_1}{w_1 \eta_1} \right).$$

Hence, we get

$$\partial_{\xi_1} \mathcal{R}_{\text{react}, \varepsilon}^*(\mu, \cdot)|_{\xi = -D\mathcal{E}^V(\mu)} = -\partial_{\xi_2} \mathcal{R}_{\text{react}}^*(\mu, \cdot)|_{\xi = -D\mathcal{E}^V(\mu)} = \frac{1}{\varepsilon} \left( \sqrt{\frac{\beta}{\alpha}} \sqrt{\frac{\eta_1}{\eta_2}} c_2 - \sqrt{\frac{\alpha}{\beta}} \sqrt{\frac{\eta_2}{\eta_1}} c_1 \right),$$

which is linear in  $c = (c_1, c_2)$ . In vector notation, we get a tilted Markov generator of the form

$$\partial_{\xi} \mathcal{R}_{\text{react}, \varepsilon}^*(\mu, -D\mathcal{E}^V(\mu)) = \frac{1}{\varepsilon} \begin{pmatrix} -\sqrt{\frac{\alpha}{\beta}} \frac{\eta_2}{\eta_1} & \sqrt{\frac{\beta}{\alpha}} \frac{\eta_1}{\eta_2} \\ \sqrt{\frac{\alpha}{\beta}} \frac{\eta_2}{\eta_1} & -\sqrt{\frac{\beta}{\alpha}} \frac{\eta_1}{\eta_2} \end{pmatrix} c = \frac{1}{\varepsilon} \begin{pmatrix} -\sqrt{\frac{\alpha}{\beta}} e^{\frac{V_1 - V_2}{2}} & \sqrt{\frac{\beta}{\alpha}} e^{\frac{V_2 - V_1}{2}} \\ \sqrt{\frac{\alpha}{\beta}} e^{\frac{V_1 - V_2}{2}} & -\sqrt{\frac{\beta}{\alpha}} e^{\frac{V_2 - V_1}{2}} \end{pmatrix} c$$

which has the space dependent stationary measure

$$w^V = \frac{1}{Z(\alpha + \beta)} (\beta \eta^1, \alpha \eta^2)^{\text{T}} = \frac{1}{Z} (w_1 e^{-V_1}, w_2 e^{-V_2})^{\text{T}}.$$

Summarizing, the tilted evolution equation has the form

$$\frac{d}{dt} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \text{div} \left( \begin{pmatrix} \delta_1 \nabla c_1 \\ \delta_2 \nabla c_2 \end{pmatrix} + \begin{pmatrix} \delta_1 c_1 \nabla V_1 \\ \delta_2 c_2 \nabla V_2 \end{pmatrix} \right) + \frac{1}{\varepsilon} \begin{pmatrix} -\sqrt{\frac{\alpha}{\beta}} e^{\frac{V_1 - V_2}{2}} & \sqrt{\frac{\beta}{\alpha}} e^{\frac{V_2 - V_1}{2}} \\ \sqrt{\frac{\alpha}{\beta}} e^{\frac{V_1 - V_2}{2}} & -\sqrt{\frac{\beta}{\alpha}} e^{\frac{V_2 - V_1}{2}} \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix}, \quad (3.3)$$

which is a linear reaction-drift-diffusion system with space dependent reaction rates. In the special case without external forcing  $V = \text{const}$ , we get the linear reaction diffusion system (1.2). Note that the reaction part still inherits symmetry since the product of the off-diagonal elements is constant in space. In particular, not all general linear reaction-drift-diffusion system with space dependent reaction rates for two species can be expressed in the form (3.3) and are induced by the gradient system  $(Q, \mathcal{E}^V, \mathcal{R}_\varepsilon^* = \mathcal{R}_{\text{diff}}^* + \mathcal{R}_{\text{react},\varepsilon}^*)$ .

### 3.3 The dissipation functional

In this section, we compute the dissipation functional  $\mathfrak{D}_\varepsilon$ , which consists of two parts: the velocity part given by the primal dissipation potential  $\mathcal{R}_\varepsilon$  and the slope-part (sometimes also called *Fisher information*)  $\mathcal{R}_\varepsilon^*(\mu, -D\mathcal{E}(\mu))$ . Here, all computations are formal and we always assume that the measure  $\mu$  has a Lebesgue density  $c$ . The precise functional analytic setting is presented in the Section 4.

The primal dissipation potential  $\mathcal{R}_\varepsilon$ , given by the Legendre transform of the dual dissipation potential  $\mathcal{R}_\varepsilon^* = \mathcal{R}_{\text{diff}}^* + \mathcal{R}_{\varepsilon,\text{react}}^*$ , can be computed via inf-convolution of  $\mathcal{R}_{\text{diff}}^*$  and  $\mathcal{R}_{\text{react},\varepsilon}^*$ . First, we compute both primal dissipation potentials separately. To do this, we introduce the following notation: For a convex, lsc. function  $F : X \rightarrow [0, \infty]$  on a reflexive and separable Banach space  $X$  with Legendre dual  $F^*$ , we define the function  $\tilde{F} : [0, \infty] \times X \rightarrow [0, \infty]$  by

$$\tilde{F}(a, x) := (a F^*(\cdot))^*(x) = \begin{cases} a F\left(\frac{1}{a}x\right) & \text{for } a > 0, \\ \chi_0(x) & \text{for } a = 0. \end{cases}$$

Introducing the quadratic function  $Q(x) = \frac{1}{2}|x|^2$  on  $\mathbb{R}^d$ , the primal dissipation potential of the diffusion part  $\mathcal{R}_{\text{diff}}^*$  is given by

$$\mathcal{R}_{\text{diff}}(\mu, \nu) = \sum_{j=1}^2 \int_{\Omega} \tilde{Q}(\delta_j c_j, J_j) dx,$$

where  $J_j$  is, by definition, the unique solution of the elliptic equation  $v_j + \text{div} J_j = 0$  with  $J \cdot \nu = 0$  on  $\partial\Omega$ . For positive  $c_j$ , we have  $\tilde{Q}(\delta_j c_j, J_j) = \frac{1}{2} \frac{|J_j|^2}{\delta_j c_j}$ .

The primal dissipation potential of the reaction part is

$$\mathcal{R}_{\text{react},\varepsilon}(\mu, b) = \begin{cases} \int_{\Omega} \tilde{C}\left(\frac{\sqrt{c_1 c_2}}{\varepsilon}, b_2\right) dx, & \text{for } b_1 + b_2 = 0 \\ \infty & \text{for } b_1 + b_2 \neq 0 \end{cases},$$

where  $\tilde{C} = (C^*)^*$  is the Legendre transform of the cosh-function  $C^*(x) = 4(\cosh(x/2) - 1)$ . In the following, we use the inequality

$$\frac{1}{2}|r| \cdot \log(|r| + 1) \leq C(r) \leq 2|r| \cdot \log(|r| + 1), \quad (3.4)$$

which, in particular, implies that the Orlicz class for  $A \subset \mathbb{R}^d$  given by

$$\tilde{L}^C(A) := \{u \in L^1(A) : \int_A C(u) dx < \infty\},$$

is, in fact, a Banach space  $\tilde{L}^C(A) = L^C(A)$  with the norm  $\|u\|_C = \sup_{\int_A C^*(v) \leq 1} \left| \int_A uv dx \right|$ .

Importantly, functions  $\tilde{Q}$ ,  $\tilde{C}$  as well as the functionals  $\mathcal{R}_{\text{diff}}$ ,  $\mathcal{R}_{\text{react},\varepsilon}$  are convex on their domain of definition.

The primal dissipation potential  $\mathcal{R}_\varepsilon$  is the inf-convolution of  $\mathcal{R}_{\text{diff}}$  and  $\mathcal{R}_{\text{react},\varepsilon}$ , and is given by

$$\begin{aligned} \mathcal{R}_\varepsilon(\mu, v) &= \inf_{v=u_1+u_2} \{ \mathcal{R}_{\text{diff}}(\mu, u_1) + \mathcal{R}_{\text{react},\varepsilon}(\mu, u_2) \} \\ &= \inf_{J,b} \left\{ \sum_{j=1}^2 \int_{\Omega} \tilde{Q}(\delta_j c_j, J_j) dx + \int_{\Omega} \tilde{C} \left( \frac{\sqrt{c_1 c_2}}{\varepsilon}, b_2(x) \right) dx : \begin{cases} v_1 = -\text{div} J_1 + b_1 \\ v_2 = -\text{div} J_2 + b_2 \\ b_1 + b_2 = 0 \end{cases} \right\}. \end{aligned}$$

In time-integrated form we get for  $v = \dot{\mu}$  that

$$\begin{aligned} \int_0^T \mathcal{R}_\varepsilon(\mu, \dot{\mu}) dt &= \inf_{v=v_1+v_2} \int_0^T \{ \mathcal{R}_{\text{diff}}(\mu, v_1) + \mathcal{R}_{\text{react},\varepsilon}(\mu, v_2) \} dt \\ &= \inf_{J,b} \left\{ \int_0^T \left\{ \sum_{j=1}^2 \int_{\Omega} \tilde{Q}(\delta_j c_j, J_j) dx + \int_{\Omega} \tilde{C} \left( \frac{\sqrt{c_1 c_2}}{\varepsilon}, b_2(x) \right) \right\} dx dt : (c, J, b) \in (\text{gCE}) \right\}. \end{aligned}$$

where we introduce the notation of a (linear) generalized continuity equation

$$(c, J, b) \in (\text{gCE}) \Leftrightarrow \left\{ b_1 + b_2 = 0 \text{ and } \begin{cases} \dot{c}_1 = -\text{div} J_1 + b_1 \\ \dot{c}_2 = -\text{div} J_2 + b_2 \end{cases} \right\}.$$

Without the reaction part,  $\int_0^T \mathcal{R}_\varepsilon dt$  is the dynamic formulation à la Benamou-Brenier of the Wasserstein distance in  $Q$  [BeB00], which can be equivalently written in the form

$$\mathcal{W}_2(\mu_0, \mu_1)^2 = \inf \left\{ \int_0^1 \int_{\Omega} \sum_{j=1}^2 \delta_j |v_j|^2 d\mu_j : \dot{\mu}_j + \text{div}(\mu_j v_j) = 0, \mu_{j,0} = \mu_0, \mu_{j,1} = \mu_1 \right\}$$

expressed in terms of transport velocities  $v_j = J_j/c_j$ . The Wasserstein distance can be interpreted as a cost in transporting mass from one measure  $\mu_0$  to  $\mu_1$ . In our situation  $\int_0^T \mathcal{R}_\varepsilon dt$  is jointly convex in  $c$ ,  $J$  and  $b$  and corresponds to modified cost function which also takes the reaction fluxes into account. The optimal diffusion fluxes  $J_j$  and reaction fluxes  $b_j$  have to satisfy the generalized continuity equation. Note that  $\int_0^T \mathcal{R}_\varepsilon dt$  does not induce a metric on  $Q$  since the reaction part is not quadratic.

Next, we compute the tilted slope part  $\mathcal{R}_\varepsilon^*(\mu, -D\mathcal{E}^V(\mu))$ . To do this, we introduce the relative densities  $\rho_j^V$  of  $\mu$  w.r.t. the stationary measure  $w^V dx$ , i.e.  $\rho_j^V = \frac{d\mu}{w_j^V dx} = \frac{c_j}{w_j^V}$ , where by (3.2) the stationary measure is  $w_j^V = \frac{1}{Z} w_i e^{-V_j}$ . Since  $V \in C^1(\Omega, \mathbb{R}^2)$  and  $\Omega \subset \mathbb{R}^d$  is compact,  $\mu$  is absolutely continuous w.r.t. the Lebesgue measure  $dx$  if and only if it is w.r.t. the stationary measure  $w^V dx$ . Inserting  $\xi = -D\mathcal{E}^V(\mu) = -(\log(c_i/w_i) + V_i)_{i=1,2}$  in the dual dissipation potential  $\mathcal{R}_\varepsilon^*$ , we get for the diffusive part

$$\mathcal{R}_{\text{diff}}^*(\mu, -D\mathcal{E}^V(\mu)) = \frac{1}{2} \int_{\Omega} \sum_{j=1}^2 \delta_j c_j |\nabla (\log c_j/w_j + V_j)|^2 dx.$$

Using  $w_j^V = \frac{1}{Z} w_j e^{-V_j}$ , a short calculation shows  $\delta_j c_j |\nabla (\log c_j/w_j + V_j)|^2 = \delta_j w_j^V \frac{|\nabla \rho_j^V|^2}{\rho_j^V}$ . Hence, we have

$$\mathcal{R}_{\text{diff}}^*(\mu, -D\mathcal{E}^V(\mu)) = \frac{1}{2} \int_{\Omega} \sum_{j=1}^2 \delta_j w_j^V \frac{|\nabla \rho_j^V|^2}{\rho_j^V} dx.$$

For the reaction part, we use the identity  $C^*(\log p - \log q) = 2 \frac{(\sqrt{p} - \sqrt{q})^2}{\sqrt{pq}}$  and get

$$\begin{aligned} \mathcal{R}_{\text{react},\varepsilon}^*(\mu, -D\mathcal{E}^V(\mu)) &= 2 \int_{\Omega} \frac{1}{\varepsilon} \sqrt{c_1 c_2} \frac{\left(\sqrt{c_1/\eta_1 w_1} - \sqrt{c_2/\eta_2 w_2}\right)^2}{\sqrt{c_1 c_2/\eta_1 w_1 \eta_2 w_2}} dx \\ &= \frac{2}{\varepsilon} \int_{\Omega} \sqrt{w_1^V w_2^V} \left(\sqrt{\rho_1^V} - \sqrt{\rho_2^V}\right)^2 dx. \end{aligned}$$

Summarizing, the total dissipation functional is

$$\begin{aligned} \mathfrak{D}_{\varepsilon}^V(\mu) &= \int_0^T \mathcal{R}_{\varepsilon}(\mu, \dot{\mu}) + \mathcal{R}_{\varepsilon}^*(\mu, -V - D\mathcal{E}(\mu)) dt \tag{3.5} \\ &= \inf_{(c,J,b) \in (\text{gCE})} \left\{ \int_0^T \left\{ \int_{\Omega} \sum_{j=1}^2 \tilde{\mathcal{Q}}(\delta_j c_j, J_j) dx + \int_{\Omega} \tilde{\mathcal{C}}\left(\frac{\sqrt{c_1 c_2}}{\varepsilon}, b_2(x)\right) dx \right\} dt \right\} + \\ &\quad + \int_0^T \left\{ \frac{1}{2} \int_{\Omega} \sum_{j=1}^2 \delta_j w_j^V \frac{|\nabla \rho_j^V|^2}{\rho_j^V} dx + \frac{2}{\varepsilon} \int_{\Omega} \sqrt{w_1^V w_2^V} \left(\sqrt{\rho_1^V} - \sqrt{\rho_2^V}\right)^2 dx \right\} dt. \end{aligned}$$

## 4 EDP-convergence result

In this section we state the EDP-convergence result for the gradient systems  $(Q, \mathcal{E}, \mathcal{R}_{\varepsilon}^*)$  to  $(Q, \mathcal{E}, \mathcal{R}_{\text{eff}}^*)$  and discuss the properties of the effective gradient system  $(Q, \mathcal{E}, \mathcal{R}_{\text{eff}}^*)$ . Since the energy  $\mathcal{E}$  is  $\varepsilon$ -independent the major challenge is to prove  $\Gamma$ -convergence of the dissipation functional  $\mathfrak{D}_{\varepsilon}^V$ , which is a functional defined on the space of trajectories in the state space  $Q$ . To be mathematical precise, we first fix the functional analytic setting.

The state space  $Q = \text{Prob}(\Omega \times \{1, 2\})$  is equipped with the  $p$ -Wasserstein distance  $d_{\mathcal{W}_p}$ , where in our situation either  $p = 1$  or  $p = 2$ . Recall that for any compact euclidean subspace  $E \subset \mathbb{R}^k$  the  $p$ -Wasserstein distance is defined on the space of probability measures  $\text{Prob}(E)$  by

$$d_{\mathcal{W}_p}(\mu^1, \mu^2)^p = \min_{\gamma \in \Gamma(\mu^1, \mu^2)} \int_E |x - y|^p d\gamma(x, y),$$

where  $\Gamma(\mu^1, \mu^2)$  is the set of all transport plans with marginals  $\mu^1$  and  $\mu^2$  (see e.g. [AGS05]). The  $p$ -Wasserstein distance  $d_{\mathcal{W}_p}$  metrizes the weak\*-topology of measures, i.e. convergence tested against continuous functions on  $E$ . In the following we will consider either  $E = \Omega$  or  $E = \Omega \times \{1, 2\}$ .

To define the topology in the space of trajectories on  $Q$ , we start very coarse, where we understand the trajectories on  $Q$  as measures in space and time. We denote the space of trajectories by  $L_w^{\infty}([0, T], Q)$  equipped with the weak\*-measureability. The weak\*-convergence is defined as usual by

$$\mu^{\varepsilon}(\cdot) \rightarrow \mu^0(\cdot) \quad :\Leftrightarrow \quad \forall i \in \{1, 2\}, \forall \phi \in C^{\infty}(\Omega \times [0, T]) : \int_0^T \int_{\Omega} \phi d\mu_i^{\varepsilon}(x) dt \rightarrow \int_0^T \int_{\Omega} \phi d\mu_i^0(x) dt.$$

A finer topology, which enables to prove compactness and evaluate the effective dissipation functional is then given by the a priori bounds

$$\sup_{\varepsilon \in ]0, 1]} \mathfrak{D}_{\varepsilon}^V(\mu^{\varepsilon}) \leq C, \quad \sup_{\varepsilon \in ]0, 1]} \text{ess sup}_{t \in [0, T]} \mathcal{E}(\mu^{\varepsilon}(t)) \leq C.$$

In fact, as presented in Section 5.1, these bounds provide that the measures  $\mu^\varepsilon$  have Lebesgue densities  $c^\varepsilon$  which converge strongly in  $L^1([0, T] \times \Omega, \mathbb{R}_{\geq 0}^2)$ . Moreover, the limiting coarse-grained measure  $\hat{\mu}^0 = \mu_1^0 + \mu_2^0$  has an representative which is absolutely continuous in time with values in  $(\text{Prob}(\Omega), d_{\mathcal{W}_2})$ , i.e. there is a function  $m \in L^1([0, T])$  such that for all  $t, s \in [0, T]$  with  $s \leq t$  we have

$$d_{\mathcal{W}_2}(\hat{\mu}^0(s), \hat{\mu}^0(t)) \leq \int_s^t m(r) dr.$$

Each component  $\mu_i^0$ ,  $i = 1, 2$  is not a trajectory in the space of probability measure, but in the space of non-negative Radon measures. Proposition 5.11 shows that  $\mu_i^0$  is absolutely continuous in time with values in  $(\mathcal{M}_+(\Omega), d_{\mathcal{W}_1})$  exploiting the dual formulation of the 1-Wasserstein distance (see e.g. [Edw11]). This compactness result is comparable to the result of Bothe and Hilhorst [BoH02], where also strong convergence of solutions  $c = (c_1, c_2)$  is proved. In particular, similar to the space independent situation in [MiS20, Ste19, MPS20] one cannot guarantee that  $\mu^\varepsilon(t) \rightarrow \mu^0(t)$  in  $Q$  for all times  $t \in [0, T]$  as jumps in time cannot be excluded. Instead the limit  $\mu^0 = c^0 dx$  has an absolutely continuous representative.

## 4.1 Main theorem

Let us state our main EDP-convergence result. For doing this, we define for  $V \in C^1(\Omega, \mathbb{R}^2)$  the total dissipation functional on  $L_w^\infty([0, T], Q)$  as

$$\mathfrak{D}_\varepsilon^V(\mu) = \begin{cases} \int_0^T \{ \mathcal{R}_\varepsilon(\mu, \dot{\mu}) + \mathcal{R}_\varepsilon^*(\mu, -D\mathcal{E}^V(\mu)) \} dt, & \mu \in \text{AC}([0, T], Q), \mu = c dx \text{ a.e. in } [0, T] \\ \infty & \text{otherwise.} \end{cases}$$

If  $\mu = c dx$  a.e. in  $[0, T]$ , then the dissipation functional is given by

$$\begin{aligned} & \int_0^T \mathcal{R}_\varepsilon(\mu, \dot{\mu}) + \mathcal{R}_\varepsilon^*(\mu, -V - D\mathcal{E}(\mu)) dt \\ &= \inf_{(c, J, b) \in (\text{gCE})} \left\{ \int_0^T \left\{ \int_\Omega \sum_{j=1}^2 \tilde{Q}(\delta_j c_j, J_j) dx + \int_\Omega \tilde{C} \left( \frac{\sqrt{c_1 c_2}}{\varepsilon}, b_2(x) \right) dx \right\} dt \right\} + \\ &+ \int_0^T \left\{ \frac{1}{2} \int_\Omega \sum_{j=1}^2 \delta_j w_j^V \frac{|\nabla \rho_j^V|^2}{\rho_j^V} dx + \frac{2}{\varepsilon} \int_\Omega \sqrt{w_1^V w_2^V} \left( \sqrt{\rho_1^V} - \sqrt{\rho_2^V} \right)^2 dx \right\} dt, \quad (4.1) \end{aligned}$$

where the infimum is taken over all Borel fluxes  $J_j \in \mathcal{M}([0, T] \times \Omega, \mathbb{R}^d)$ ,  $b_j \in \mathcal{M}([0, T] \times \Omega, \mathbb{R})$  which satisfy the generalized continuity equation (gCE) in the sense of distributions, i.e.

$$\begin{aligned} \forall j = 1, 2 \forall \phi \in C_c^\infty([0, T] \times \Omega) : & \int_0^T \int_\Omega \dot{\phi} c_j - \nabla \phi \cdot J_j dx dt = - \int_0^T \int_\Omega b_j \phi dx dt, \\ & J \cdot \nu = 0 \text{ on } \partial\Omega. \end{aligned}$$

*Remark 4.1.* Strictly speaking, the functions  $\tilde{Q}$  and  $\tilde{C}$  are not defined for measures  $J_j, b_j$  and hence, the formula for the dissipation functional (3.5) is a priori not correct. In fact, introducing the related functional as in Lemma 5.2, the dissipation functional can be expressed via the densities of  $J_j, b_j$ . These densities are in  $L^1([0, T] \times \Omega)$  as Lemma 5.1 shows. For notational convenience, we identify the measures with their Lebesgue densities and stick to the above expression (3.5). In Lemma 5.7, we, in fact, show compactness for the sequence of measures  $J_i^\varepsilon$ .

The main result is the  $\Gamma$ -convergence of  $\mathfrak{D}_\varepsilon^V$  to the effective dissipation functional  $\mathfrak{D}_0^V$  which is defined by

$$\mathfrak{D}_0^V(\mu) = \begin{cases} \int_0^T \mathcal{R}_{\text{eff}}(\mu, \dot{\mu}) + \mathcal{R}_{\text{eff}}^*(\mu, -D\mathcal{E}^V(\mu)) dt, & \text{if } \mu \in \text{AC}([0, T], Q), \mu = c \, dx \text{ a.e. in } [0, T] \\ \infty & \text{otherwise.} \end{cases}$$

where

$$\mathcal{R}_{\text{eff}}^*(\mu, \xi) = \mathcal{R}_{\text{diff}}^*(\mu, \xi) + \chi_{\{\xi_1 = \xi_2\}}(\xi), \quad \mathcal{R}_{\text{eff}}(\mu, v) = (\mathcal{R}_{\text{eff}}(\mu, \cdot))^*(v). \quad (4.2)$$

**Theorem 4.2.** *Let  $V \in C^1(\Omega, \mathbb{R}^2)$ . On  $L_w^\infty([0, T], Q)$ , we have  $\Gamma$ -convergence constraint to bounded energies of  $\mathfrak{D}_\varepsilon^V$ , i.e.  $\mathfrak{D}_\varepsilon^V \xrightarrow{\Gamma_E} \mathfrak{D}_0^V$  where*

$$\mathfrak{D}_0^V(\mu) = \begin{cases} \int_0^T \mathcal{R}_{\text{eff}}(\mu, \dot{\mu}) + \mathcal{R}_{\text{eff}}^*(\mu, -V - D\mathcal{E}(\mu)) dt, & \mu \in \text{AC}([0, T], Q), \mu = c \, dx \text{ a.e. } [0, T] \\ \infty & \text{otherwise} \end{cases} \quad (4.3)$$

with

$$\begin{aligned} \mathcal{R}_{\text{eff}}^*(\mu, \xi) &= \mathcal{R}_{\text{diff}}^*(\mu, \xi) + \chi_{\{\xi_1 = \xi_2\}}(\xi), \\ \mathcal{R}_{\text{eff}}(\mu, v) &= \inf_{u+\tilde{u}=v} \{ \mathcal{R}_{\text{diff}}(\mu, \tilde{u}) + \chi_0(u_1 + u_2) \} = \\ &= \inf \left\{ \sum_{j=1}^2 \int_{\Omega} \tilde{\mathcal{Q}}(\delta_j c_j, J_j) dx : u_1 + u_2 = 0, \left\{ \begin{array}{l} v_1 = -\text{div} J_1 + u_1 \\ v_2 = -\text{div} J_2 + u_2 \end{array} \right\} \right\}. \end{aligned}$$

The theorem states that the limit dissipation functional is again of  $\mathcal{R} \oplus \mathcal{R}^*$ -form with an effective dissipation potential  $\mathcal{R}_{\text{eff}}^*$ . The effective dissipation potential  $\mathcal{R}_{\text{eff}}^*$  consists again of two terms describing the diffusion and a coupling, which forces the chemical potential  $-D\mathcal{E}^V$  to equilibration. This equilibration provides the microscopic equilibria of the densities  $\rho^V$  defining the slow manifold of the evolution.

In the Section 5, we will present the detailed proof of this  $\Gamma$ -convergence result. In this section, we discuss the effective gradient system and its induced gradient flow equation. As we will see the associated gradient flow equation can be understood as an evolution equation on  $Q$  and also on a smaller state space  $\hat{Q} := \text{Prob}(\Omega)$  of coarse-grained variables.

As an immediate consequence, Theorem 4.2 implies that  $(Q, \mathcal{E}, \mathcal{R}_\varepsilon^*)$  EDP-converges with tilting to  $(Q, \mathcal{E}, \mathcal{R}_{\text{eff}}^*)$ .

**Theorem 4.3.** *Let  $\mathcal{R}_{\text{eff}}^*$  be defined by (4.2). Then the gradient system  $(Q, \mathcal{E}, \mathcal{R}_\varepsilon^*)$  EDP-converges with tilting to  $(Q, \mathcal{E}, \mathcal{R}_{\text{eff}}^*)$ .*

*Proof.* The energy  $\mathcal{E}$  is  $\varepsilon$ -independent and lsc. on  $Q$ . Hence, it  $\Gamma$ -converges to itself. Theorem 4.2 implies that  $\mathfrak{D}_\varepsilon^V$   $\Gamma$ -converges to  $\mathfrak{D}_0^V$  and  $\mathfrak{D}_0^V$  is of  $\mathcal{R} \oplus \mathcal{R}^*$  structure, where the effective dissipation potential  $\mathcal{R}_{\text{eff}}$  is independent of the tilts  $\eta = V$ . Hence, EDP-convergence with tilting is established.  $\square$

## 4.2 Effective gradient flow equation

In this section, we discuss the effective gradient flow equation that is associated by the limit gradient structure  $(Q, \mathcal{E}, \mathcal{R}_{\text{eff}}^*)$ . Similar to the space-independent situation in [MiS20, MPS20] the limit gradient



structure can also be equivalently understood as a gradient structure on a smaller coarse-grained space of slow variables  $\hat{Q}$ . In particular, we obtain an effective gradient flow equation on the original state space  $Q$  with a Lagrange multiplier ensuring the projection on the slow manifold, and, moreover, an effective gradient flow equation in coarse-grained variables. First, we discuss the effective gradient flow equation with Lagrange multipliers, and secondly, the coarse-grained gradient structure and its induced gradient flow equation. Throughout the section the potential  $V \in C^1(\Omega, \mathbb{R}^2)$  is fixed.

#### 4.2.1 Gradient flow equation with Lagrange multipliers

For being brief, the calculations in this section are rather formal. The effective dissipation potential  $\mathcal{R}_{\text{eff}}^* = \mathcal{R}_{\text{diff}}^* + \chi_{\{\xi_1 = \xi_2\}}$  consists of two parts: the first describes the dissipation of the evolution and the second provides the linear constraint of being on the slow manifold and also the corresponding Lagrange multiplier. The evolution equation is given by

$$\dot{\mu} \in \partial_{\xi} \mathcal{R}_{\text{eff}}^*(\mu, -V - D\mathcal{E}(\mu)) = \partial_{\xi} \left\{ \mathcal{R}_{\text{diff}}^*(\mu, -V - D\mathcal{E}(\mu)) + \chi_{\{\xi_1 = \xi_2\}}(-V - D\mathcal{E}(\mu)) \right\}.$$

Following [EkT76], the subdifferential of a sum is given by the sum of the subdifferential, if one term is continuous, which holds for the first term. For the second term, the subdifferential of the characteristic function is only definite in its domain, i.e. if

$$-V_1 - D\mathcal{E}(\mu)_1 = -V_2 - D\mathcal{E}(\mu)_2,$$

which implies that  $\mu = cdx$  and that their densities satisfy the relation  $\frac{c_1}{\beta e^{-V_1}} = \frac{c_2}{\alpha e^{-V_2}}$  defining the linear slow manifold. Moreover, on its domain we have for the subdifferential that  $\partial \chi_{\{\xi_1 = \xi_2\}} = \mathcal{M}(\Omega) \begin{pmatrix} 1 \\ -1 \end{pmatrix}$ . Hence, we conclude that

$$\dot{\mu} \in \partial_{\xi} \left\{ \mathcal{R}_{\text{diff}}^*(\mu, -V - D\mathcal{E}(\mu)) \right\} + \mathcal{M}(\Omega) \begin{pmatrix} 1 \\ -1 \end{pmatrix}, \quad \frac{c_1}{\beta e^{-V_1}} = \frac{c_2}{\alpha e^{-V_2}},$$

which implies the gradient flow equation on the slow manifold with a Lagrange multiplier  $\lambda = (\lambda_1, \lambda_2)$  for the densities of the form

$$\begin{cases} \dot{c}_1 = \text{div} \{ \delta_1 \nabla c_1 + \delta_1 c_1 \nabla V_1 \} + \lambda_1(t, x) \\ \dot{c}_2 = \text{div} \{ \delta_2 \nabla c_2 + \delta_2 c_2 \nabla V_2 \} + \lambda_2(t, x) \end{cases}, \quad \lambda_1 + \lambda_2 = 0, \quad \frac{c_1}{\beta e^{-V_1}} = \frac{c_2}{\alpha e^{-V_2}}. \quad (4.4)$$

#### 4.2.2 Coarse-grained gradient structure and its gradient flow equation

Now, we discuss the effective gradient structure  $(Q, \mathcal{E}, \mathcal{R}_{\text{eff}}^*)$  in the slow coarse-grained variables. To do this, we introduce the coarse grained probability measure  $\hat{\mu} = \mu_1 + \mu_2$  on  $\Omega$  and the corresponding concentrations  $\hat{c} := c_1 + c_2$ . Moreover, we define the equilibrated densities  $\hat{\rho}^V = \rho_1^V = \rho_2^V$  and the coarse-grained stationary measure  $\hat{w}^V = w_1^V + w_2^V$ , for which we get  $\hat{c} = \rho_1^V w_1^V + \rho_2^V w_2^V = \hat{\rho}^V (w_1^V + w_2^V) = \hat{\rho}^V \hat{w}^V$ . We introduce the coarse-grained diffusion coefficient  $\hat{\delta}^V = \frac{\delta_1 w_1^V + \delta_2 w_2^V}{w_1^V + w_2^V}$ .

With this notation, we may define the coarse-grained gradient structure  $(\hat{Q}, \hat{\mathcal{E}}, \hat{\mathcal{R}}^*)$ . On the state space  $\hat{Q} = \text{Prob}(\Omega)$ , we define

$$\begin{aligned}\hat{\mathcal{R}}^*(\hat{\mu}, \hat{\xi}) &:= \mathcal{R}_{\text{eff}}^* \left( \left( \frac{w_1^V}{w_1^V + w_2^V} \hat{\mu}, \frac{w_2^V}{w_1^V + w_2^V} \hat{\mu} \right), (\hat{\xi}, \hat{\xi}) \right) = \frac{1}{2} \int_{\Omega} \delta^V |\nabla \hat{\xi}|^2 d\hat{\mu}, \quad (4.5) \\ \hat{\mathcal{E}}(\hat{\mu}) &:= \mathcal{E}^V \left( \frac{w_1^V}{w_1^V + w_2^V} \hat{\mu}, \frac{w_2^V}{w_1^V + w_2^V} \hat{\mu} \right).\end{aligned}$$

Introducing the coarse-grained potential  $\hat{V} = -\log(w_1^V + w_2^V) - \log Z = -\log(\hat{w}^V) - \log Z = -\log(w_1 e^{-V_1} + w_2 e^{-V_2})$ , for which the exponential is given by the weighted arithmetic mean of the exponentials  $e^{-V_1}$  and  $e^{-V_2}$ , i.e.  $e^{-\hat{V}} = w_1 e^{-V_1} + w_2 e^{-V_2}$  (we used that  $w_1 + w_2 = 1$ ). Easy calculations show that the energy has for the explicit form

$$\hat{\mathcal{E}}(\hat{\mu}) = \int_{\Omega} \hat{\mu} \log \hat{\mu} + \hat{\mu} \hat{V} dx.$$

The coarse-grained dissipation functional is defined by

$$\hat{\mathcal{D}}(\hat{\mu}) = \int_0^T \hat{\mathcal{R}}(\hat{\mu}, \dot{\hat{\mu}}) + \hat{\mathcal{R}}^*(\hat{\mu}, -D\hat{\mathcal{E}}(\hat{\mu})) dt,$$

which incorporates the tilt via the coarse-grained variables. Note, that the coarse-grained dissipation potential  $\hat{\mathcal{R}}^*$  depends explicitly on the tilt  $V$  via the diffusion coefficient  $\delta^V$ . This is not a contradiction to tilt-EDP convergence (Theorem 4.3), because in original variables the effective dissipation potential (4.3) is indeed independent of the tilts. The tilts dependence of  $\hat{\mathcal{R}}^*$  originates from the energy and tilt dependent slow manifold.

To relate the dissipation functional  $\mathfrak{D}_0^V$  with the coarse grained dissipation functional  $\hat{\mathcal{D}}$ , we first show that also an equilibration of the fluxes occurs. To do this the following convexity property is important.

**Lemma 4.4.** *Let  $X$  a separable and reflexive Banach space and let  $F : X \rightarrow \mathbb{R}_{\infty}$  be convex and lsc. Then for the function  $\tilde{F} : [0, \infty[ \times X \rightarrow \mathbb{R}_{\infty}$ , we have*

$$\tilde{F} \left( \sum_{i=1}^I a_i, \sum_{i=1}^I x_i \right) \leq \sum_{i=1}^I \tilde{F}(a_i, x_i).$$

*If  $F$  is strictly convex then equality holds if and only if  $(a_i, x_i) = (0, 0)$  whenever  $a_i = 0$  or  $x_i/a_i = x_j/a_j$  whenever  $a_i, a_j > 0$ . Moreover, if  $F(0) = 0$ , we have the following monotonicity property*

$$\tilde{F}(a_1, x) \leq \tilde{F}(a_2, x), \quad \text{if } a_1 \geq a_2.$$

*Proof.* Let pairs  $(a_i, x_i)$  for  $i = 1, \dots, I$  be given. If  $a_i = 0$ , then either  $x_i = 0$  and the claim has to be shown for  $I - 1$ -number of pairs, or  $x_i \neq 0$  and the right-hand side is  $\infty$  meaning that the claim is trivial. So let us assume that  $a_i > 0$  for all  $i = 1 \dots, I$ . Then  $\tilde{F}(a_i, x_i) = a_i F(x_i/a_i)$  and the claim is equivalent to

$$\sum_{i=1}^I \frac{a_i}{\sum_{i=1}^I a_i} F(x_i/a_i) \geq F \left( \sum_{i=1}^I \frac{a_i}{\sum_{i=1}^I a_i} \frac{x_i}{a_i} \right) = F \left( \frac{\sum_{i=1}^I x_i}{\sum_{i=1}^I a_i} \right),$$

which holds since  $F$  is convex. If  $F$  is strictly convex then we immediately observe that whenever  $a_i, a_j > 0$  we have  $\frac{x_i}{a_i} = \frac{x_j}{a_j}$ , and whenever  $a_i = 0$  that also  $x_i = 0$ .

To see the monotonicity property, we observe that  $\tilde{F}(a, 0) = 0$  for all  $a \geq 0$ . Hence, we have

$$\tilde{F}(a_1 + a_2, x) \leq \tilde{F}(a_1, x) + \tilde{F}(a_2, 0) = \tilde{F}(a_1, x),$$

which proves the claim.  $\square$

Recalling formula (4.3) of the effective dissipation functional  $\mathfrak{D}_0^V$  and using the above lemma, we observe that the velocity part of the dissipation functional  $\mathfrak{D}_0^V$  can now be estimated. In particular, we will see that the limit dissipation functional  $\mathfrak{D}_0^V$  can be equivalently expressed in coarse-grained variables  $(\hat{\mu}, \hat{J})$  by using that an equilibration of concentrations also provides an equilibration of the corresponding fluxes. In the reconstruction strategy in Section 5.3 this equilibration is explicitly used (see 4.10 and (5.3)).

**Proposition 4.5.** *Let  $\mu \in AC([0, T], Q)$  with  $\mathfrak{D}_0^V(\mu) < \infty$  and  $\text{ess sup}_{t \in [0, T]} \mathcal{E}(\mu(t)) < \infty$ . Then the following holds:*

1 We have  $\mathfrak{D}_0^V(\mu) = \hat{\mathfrak{D}}(\hat{\mu})$  where  $\hat{\mu} = \mu_1 + \mu_2$  and

$$\begin{aligned} \hat{\mathfrak{D}}(\hat{\mu}) &= \int_0^T \hat{\mathcal{R}}(\hat{\mu}, \dot{\hat{\mu}}) + \hat{\mathcal{R}}^*(\hat{\mu}, -D\hat{\mathcal{E}}(\hat{\mu})) dt \\ &= \inf_{\hat{J}: \dot{\hat{\mu}} + \text{div} \hat{J} = 0} \left\{ \int_0^T \int_{\Omega} \tilde{Q}(\delta^V \hat{c}, \hat{J}) + \frac{\delta \hat{w}^V}{2} \frac{|\nabla \hat{\rho}^V|^2}{\hat{\rho}^V} dx dt \right\}. \end{aligned}$$

2 The chain-rule holds for  $[0, T] \ni t \mapsto \hat{\mathcal{E}}(\hat{\mu}(t)) \in \mathbb{R}$ , i.e. we have

$$\frac{d}{dt} \hat{\mathcal{E}}(\hat{\mu}(t)) = \langle D\hat{\mathcal{E}}(\hat{\mu}(t)), \dot{\hat{\mu}}(t) \rangle.$$

3 The gradient flow equation of the gradient system  $(\hat{Q}, \hat{\mathcal{E}}, \hat{\mathcal{R}}^*)$  is given by

$$\dot{\hat{c}} = -\text{div} \left( \delta^V \hat{c} \nabla \left( -D\hat{\mathcal{E}}(\hat{\mu}) \right) \right) = \text{div} \left( \delta^V \nabla \hat{c} + \delta^V \hat{c} \nabla \hat{V} \right), \quad (4.6)$$

with the potential  $\hat{V} = -\log \hat{w}^V$  and stationary measure  $\hat{w}^V$ .

Equation (4.6) shows that the coarse-grained gradient flow equation induced by  $(\hat{Q}, \hat{\mathcal{E}}, \hat{\mathcal{R}}^*)$  is a drift-diffusion equation of the coarse-grained concentration  $\hat{c}$  with mixed diffusion constant  $\delta^V$ . In particular, in the tilt free case we have  $\delta^V = \text{const} = \frac{\beta \delta_1 + \alpha \delta_2}{\alpha + \beta}$ , and we recover the result of [BoH02].

*Proof.* To prove Part 1, we first observe that the bounded energy and dissipation for the trajectory  $\mu$  implies that we have  $\frac{c_1}{w_1^V} = \frac{c_2}{w_2^V}$  a.e. in  $[0, T] \times \Omega$ . Using  $\hat{c} = c_1 + c_2$  for the densities, we get  $\hat{c} = \frac{w_1^V + w_2^V}{w_1^V} c_1 = \frac{w_1^V + w_2^V}{w_2^V} c_2$ . The Fisher information  $\mathcal{S}_0^V(\mu) := \mathcal{R}_{\text{eff}}^*(\mu, -V - D\mathcal{E}(\mu))$  has the form

$$\mathcal{S}_0^V(\mu) = \frac{1}{2} \int_{\Omega} \sum_{j=1}^2 \delta_j w_j^V \frac{|\nabla \rho_j^V|^2}{\rho_j^V} dx = \frac{1}{2} \int_{\Omega} \left( \frac{\delta_1 w_1^V + \delta_2 w_2^V}{w_1^V + w_2^V} \right) (w_1^V + w_2^V) \frac{|\nabla \hat{\rho}^V|^2}{\hat{\rho}^V} dx \quad (4.7)$$

$$= \frac{1}{2} \int_{\Omega} \delta^V \hat{w}^V \frac{|\nabla \hat{\rho}^V|^2}{\hat{\rho}^V} dx = \hat{\mathcal{R}}^*(\hat{\mu}, -D\hat{\mathcal{E}}(\hat{\mu})). \quad (4.8)$$

Lemma 4.4 provides that also an equilibration of the fluxes occurs. Indeed, defining the coarse-grained flux  $\hat{J} = J_1 + J_2$ , we conclude

$$\frac{|J_1|^2}{\delta_1 c_1} + \frac{|J_2|^2}{\delta_2 c_2} \geq \frac{|J_1 + J_2|^2}{\delta_1 c_1 + \delta_2 c_2} = \frac{|\hat{J}|^2}{\frac{\delta_1 w_1^V + \delta_2 w_2^V}{w_1^V + w_2^V} \hat{c}} = \frac{|\hat{J}|^2}{\delta \hat{c}}, \quad (4.9)$$

where  $\hat{\delta}^V := \frac{\delta_1 w_1^V + \delta_2 w_2^V}{w_1^V + w_2^V}$ . Equality holds if and only if  $(J_1, c_1) = 0$  or  $(J_2, c_2) = 0$  or  $J_1/\delta_1 c_1 = J_2/\delta_2 c_2 = \hat{J}/\hat{\delta}^V \hat{c}$ . The last condition is equivalent to

$$\hat{J} = \frac{\delta_1 w_1^V}{\delta_1 w_1^V + \delta_2 w_2^V} J_1 = \frac{\delta_2 w_2^V}{\delta_1 w_1^V + \delta_2 w_2^V} J_2, \quad (4.10)$$

which provides an explicit formula for the coarse-grained diffusion flux.

For the dissipation functional that means

$$\begin{aligned} \mathfrak{D}_0^V(\mu) &= \inf_{(c, J, b) \in (\text{gCE})} \int_0^T \left\{ \int_{\Omega} \sum_{j=1}^2 \tilde{\mathcal{Q}}(\delta_j c_j, J_j) dx + \frac{1}{2} \int_{\Omega} \sum_{j=1}^2 \delta_j w_j^V \frac{|\nabla \rho_j|^2}{\rho_j} dx \right\} dt \\ &\geq \inf_{\hat{c} + \text{div} \hat{J} = 0} \int_0^T \left\{ \int_{\Omega} \tilde{\mathcal{Q}}(\hat{\delta} \hat{c}, \hat{J}) dx + \frac{1}{2} \int_{\Omega} \hat{\delta} \hat{w}^V \frac{|\nabla \hat{\rho}|^2}{\hat{\rho}} dx \right\} dt. \end{aligned} \quad (4.11)$$

To prove equality, we first observe that  $\hat{J}, J_1, J_2$  satisfy the same boundary conditions. Moreover, the explicitly derived reaction flux  $b_1, b_2$  from (5.3) shows that the reconstructed fluxes  $(J_1, J_2)$  from coarse-grained flux  $\hat{J}$  is admissible. Hence, we obtain equality

$$\mathfrak{D}_0^V(\mu) = \inf_{\hat{c} + \text{div} \hat{J} = 0} \int_0^T \left\{ \int_{\Omega} \tilde{\mathcal{Q}}(\hat{\delta} \hat{c}, \hat{J}) dx + \frac{1}{2} \int_{\Omega} \hat{\delta} \hat{w}^V \frac{|\nabla \hat{\rho}|^2}{\hat{\rho}} dx \right\} dt,$$

which proves the first part.

For the chain-rule in Part 2, we refer to the proof [FrL19, Lem 4.8] since we consider the pure diffusive situation. The proof uses a time-regularization argument and convexity of the Fisher-information following the ideas of [MRS13, Prop. 2.4].

For Part 3, we compute the evolution equation that is induced by the gradient system is  $(\hat{Q}, \hat{\mathcal{E}}, \hat{\mathcal{R}}^*)$ . We have

$$\begin{aligned} \partial_{\hat{\xi}} \hat{\mathcal{R}}^*(\hat{\mu}, \hat{\xi}) &= -\text{div} \left( \hat{\delta}^V \hat{c} \nabla \hat{\xi} \right), \\ D \hat{\mathcal{E}}(\hat{\mu}) &= \log \hat{\mu} + 1 - \log \hat{w}^V - \log Z, \\ \nabla \left( -D \hat{\mathcal{E}}(\hat{\mu}) \right) &= -\frac{\nabla \hat{\mu}}{\hat{\mu}} + \frac{\nabla \hat{w}^V}{\hat{w}^V} = -\frac{\nabla \hat{\mu}}{\hat{\mu}} - \nabla \hat{V}, \end{aligned}$$

which results in

$$\hat{c} = -\text{div} \left( \hat{\delta}^V \hat{c} \nabla (-D \hat{\mathcal{E}}(\hat{\mu})) \right) = \text{div} \left( \hat{\delta}^V \nabla \hat{c} + \hat{\delta}^V \hat{c} \nabla \hat{V} \right).$$

□

Note that the coarse-grained gradient flow equation (4.6) is equivalent to the gradient flow equation with Lagrange multipliers (4.4). Indeed, adding both equations in (4.4) together and using that the original concentrations can be expressed by the coarse-grained concentrations via

$$c_i = \frac{w_i e^{-V_i}}{w_1 e^{-V_1} + w_2 e^{-V_2}} \hat{c}, \quad (4.12)$$

the coarse-grained gradient flow equation (4.6) with the drift term  $\hat{\delta}^V \hat{c} \nabla \hat{V}$  can be readily derived. Conversely, using (4.12), we see that  $c = (c_1, c_2)$  are on the slow manifold and satisfy (4.4). The corresponding Lagrange multipliers  $\lambda = (\lambda_1, \lambda_2)$  can be explicitly calculated. Introducing the difference of the diffusion constants  $\bar{\delta} = \delta_1 - \delta_2$  and the potentials  $\bar{V} = V_1 - V_2$ , we have

$$\lambda_1 = \frac{w_2 e^{-V_2}}{w_1 e^{-V_1} + w_2 e^{-V_2}} \left( -\bar{\delta} \Delta c_1 + (\delta_2 \nabla \bar{V} - \bar{\delta} \nabla V_1) \cdot \nabla c_1 + c_1 \{ \delta_2 \nabla \bar{V} \nabla V_1 - \bar{\delta} \Delta V_1 \} \right)$$

$$\lambda_2 = \frac{w_1 e^{-V_1}}{w_1 e^{-V_1} + w_2 e^{-V_2}} \left( \bar{\delta} \Delta c_2 + (-\delta_1 \nabla \bar{V} + \bar{\delta} \nabla V_2) \cdot \nabla c_2 + c_2 \{ -\delta_1 \nabla \bar{V} \nabla V_2 + \bar{\delta} \Delta V_2 \} \right).$$

We observe that the Lagrange multiplier  $\lambda_i$  has the same regularity as the right-hand side of the evolution of  $c_i$ . Moreover, both evolution equations are completely uncoupled but contain a linear annihilation/creation term, which depends on the potential  $V = (V_1, V_2)$  and the diffusion coefficient  $\delta = (\delta_1, \delta_2)$ . A lengthy calculation shows that indeed we have  $\lambda_1 + \lambda_2 = 0$ .

## 5 Proof of $\Gamma$ -convergence

In this section we prove the  $\Gamma$ -convergence result of Theorem 4.2. As usual, we prove  $\Gamma$ -convergence in three steps: First deriving compactness, secondly establishing the liminf-estimate by exploiting the compactness, thirdly constructing the recovery sequence for the limsup-estimate.

In the following the next lemma will be useful.

**Lemma 5.1.** *Let  $F : \mathbb{R}^k \rightarrow [0, \infty[$  be a convex, lsc. function of superlinear growth, i.e.  $F(r)/r \rightarrow \infty$  as  $r \rightarrow \infty$ . Then there is a constant  $k_F > 0$  such that for any measurable functions  $W : \Omega \rightarrow \mathbb{R}^k$  and  $\rho : \Omega \rightarrow \mathbb{R}_{\geq 0}$  it holds*

$$\int_{\Omega} |W| dx \leq \int_{\Omega} \tilde{F}(\rho, W) dx + k_F \int_{\Omega} \rho dx.$$

*Proof.* Let  $W$  and  $\rho$  be given. We define three measurable subsets of  $\Omega$ :

$$\Omega_0 = \{x : \rho(x) = 0\}, \quad \Omega_1 = \{x : \rho \neq 0, \frac{1}{\rho} |W| \leq F(\frac{1}{\rho} W)\}, \quad \Omega_2 = \{x : \rho \neq 0, \frac{1}{\rho} |W| > F(\frac{1}{\rho} W)\}.$$

Since  $F$  is superlinear, there is a constant  $k_F > 0$  such that on  $\Omega_2$  it holds  $W/\rho \leq k_F$ . Hence we can estimate

$$\begin{aligned} \int_{\Omega} |W| dx &\leq \int_{\Omega_0} |W| dx + \int_{\Omega_1} \frac{|W|}{\rho} \rho dx + \int_{\Omega_2} \frac{|W|}{\rho} \rho dx \\ &\leq \int_{\Omega_0} \tilde{F}(\rho, W) dx + \int_{\Omega_1} F(\frac{1}{\rho} W) \rho dx + k_F \int_{\Omega_2} \rho dx \\ &\leq \int_{\Omega} \tilde{F}(\rho, W) dx + k_F \int_{\Omega} \rho dx. \end{aligned}$$

□

Moreover, we need the following classical lemma. It guarantees the necessary regularity for the limits, and moreover, it provides the desired liminf-estimate.

**Lemma 5.2** (AGS05-Lemma 9.4.3, [AGS05]). *Let  $F : [0, \infty[ \rightarrow [0, \infty]$  be a proper, lsc, convex function with superlinear growth. We define the related functional*

$$\mathcal{F}(\mu, \gamma) = \begin{cases} \int_A F\left(\frac{d\mu}{d\gamma}\right) d\gamma, & \text{if } \mu \ll \gamma, \\ \infty, & \text{otherwise.} \end{cases}$$

*Let  $\mu^\varepsilon, \gamma^\varepsilon \in \text{Prob}(A)$  be two sequences with  $\mu^\varepsilon \xrightarrow{*} \mu^0$  and  $\gamma^\varepsilon \xrightarrow{*} \gamma^0$ . Then*

$$\liminf_{\varepsilon \rightarrow 0} \mathcal{F}(\mu^\varepsilon, \gamma^\varepsilon) \geq \mathcal{F}(\mu^0, \gamma^0).$$

*In particular, if the left-hand side is finite then for the limits it holds  $\mu^0 \ll \gamma^0$ .*

## 5.1 Compactness

In this section, we derive the required compactness for proving the liminf-estimate in Section 5.2.

Recall that for given potential  $V \in C^1(\Omega, \mathbb{R}^2)$  the dissipation functional  $\mathfrak{D}_\varepsilon^V$  is defined on the space of trajectories equipped with the weak topology, i.e.  $\mu^\varepsilon \rightarrow \mu^0 \in L_w^\infty([0, T], Q)$  if and only if it holds

$$\forall i \in \{1, 2\}, \forall \phi \in C_0^\infty(\Omega \times [0, T]) : \int_0^T \int_\Omega \phi d\mu_i^\varepsilon(x) dt \rightarrow \int_0^T \int_\Omega \phi d\mu_i^0(x) dt.$$

In the following we want to derive compactness for a sequence  $(\mu^\varepsilon)_{\varepsilon > 0}$  of trajectories, satisfying the a priori bounds

$$\sup_{\varepsilon > 0} \text{ess sup}_{t \in [0, T]} \mathcal{E}(\mu^\varepsilon(t)) \leq C, \quad \sup_{\varepsilon > 0} \mathfrak{D}_\varepsilon^V(\mu^\varepsilon) \leq C, \quad (5.1)$$

where the total dissipation functional is

$$\mathfrak{D}_\varepsilon^V(\mu) = \begin{cases} \int_0^T \mathcal{R}_\varepsilon(\mu, \dot{\mu}) + \mathcal{R}_\varepsilon^*(\mu, -D\mathcal{E}^V(\mu)) dt, & \mu \in \text{AC}([0, T], Q), \mu = c dx \text{ a.e. in } [0, T] \\ \infty & \text{otherwise,} \end{cases}$$

and for  $\mu = c dx$  we have

$$\begin{aligned} & \int_0^T \mathcal{R}_\varepsilon(\mu, \dot{\mu}) + \mathcal{R}_\varepsilon^*(\mu, -V - D\mathcal{E}(\mu)) dt \\ &= \inf_{(c, J, b) \in (\text{gCE})} \left\{ \int_0^T \left\{ \int_\Omega \sum_{j=1}^2 \tilde{Q}(\delta_j c_j, J_j) dx + \int_\Omega \tilde{C}\left(\frac{\sqrt{c_1 c_2}}{\varepsilon}, b_2(x)\right) dx \right\} dt \right\} +, \\ &+ \int_0^T \frac{1}{2} \int_\Omega \sum_{j=1}^2 \delta_j w_j^V \frac{|\nabla \rho_j^V|^2}{\rho_j^V} dx + \frac{2}{\varepsilon} \int_\Omega \sqrt{w_1^V w_2^V} \left( \sqrt{\rho_1^V} - \sqrt{\rho_2^V} \right)^2 dx dt. \end{aligned} \quad (5.2)$$

Using the bound of the dissipation functional  $\mathfrak{D}_\varepsilon^V(\mu^\varepsilon) \leq C$ , we conclude that there are diffusive fluxes  $J^\varepsilon = (J_1^\varepsilon, J_2^\varepsilon)$  and reaction fluxes  $b^\varepsilon = (b_1^\varepsilon, b_2^\varepsilon)$  such that  $J_i^\varepsilon \in \mathcal{M}(\Omega, \mathbb{R}^d)$ ,  $b_i^\varepsilon \in \mathcal{M}(\Omega, \mathbb{R})$  and  $(\mu^\varepsilon, J^\varepsilon, b^\varepsilon)$  satisfies the continuity equation

$$(c, J, b) \in (\text{gCE}) \Leftrightarrow \left\{ b_1 + b_2 = 0 \text{ and } \left\{ \begin{array}{l} \dot{c}_1 = -\text{div} J_1 + b_1 \\ \dot{c}_2 = -\text{div} J_2 + b_2 \end{array} \right\} \right\}.$$

Moreover, we get bounds:

$$i = 1, 2 : \int_0^T \int_{\Omega} \frac{|\nabla \rho_i^{V,\varepsilon}|^2}{\rho_i^{V,\varepsilon}} dx dt \leq C, \quad \int_0^T \int_{\Omega} \tilde{Q}(c_i^\varepsilon, J_i^\varepsilon) dx dt \leq C,$$

$$\int_0^T \int_{\Omega} \tilde{C}\left(\frac{\sqrt{c_1^\varepsilon c_2^\varepsilon}}{\varepsilon}, b_2^\varepsilon(x)\right) dx dt \leq C, \quad \frac{1}{\varepsilon} \int_0^T \int_{\Omega} \left(\sqrt{\rho_1^{V,\varepsilon}} - \sqrt{\rho_2^{V,\varepsilon}}\right)^2 dx dt \leq C.$$

*Remark 5.3.* Following [Man07] a distributional solution  $(\mu, J, B)$  of the generalized continuity equation  $\dot{\mu} = -\operatorname{div} J + B$  satisfying  $\int_0^T \int_{\Omega} |B| + |J| dx < \infty$  can be assumed to be absolutely continuous. The bounds can be obtained easily using Lemma 5.1 for fixed  $\varepsilon > 0$ .

Although the functional is convex in the concentration  $c$  and in the fluxes  $J$  and  $b$ , weak convergence would be sufficient to prove a liminf-estimate using a Joffe-type argument. But, comparing the situation with the evolution equation, we aim in proving even strong convergence for the densities  $c^\varepsilon \rightarrow c^0$  in  $L^1([0, T] \times \Omega, \mathbb{R}_{\geq 0}^2)$ . This is done in two steps: First, compactness of coarse-grained variables, and secondly, convergence towards the slow manifold is shown, which together implies strong compactness. This strategy has successfully been applied already in the space-independent case in [MiS20, MPS20]. Moreover, we show that the limit trajectory  $\mu^0 = c^0 dx$  has a representative which is in  $\operatorname{AC}([0, T], Q)$ . Note that it is not possible to prove pointwise convergence  $\mu^\varepsilon(t) \xrightarrow{*} \mu^0(t)$  for all  $t \in [0, T]$ . Instead, pointwise convergence is only shown for the coarse-grained variables  $\hat{\mu}^\varepsilon := \mu_1^\varepsilon + \mu_2^\varepsilon$ .

First, we derive weak compactness in space-time, which immediately follows from the uniform bound in  $\varepsilon$  and time on the energy.

**Lemma 5.4** (Very Weak compactness in space-time). *Let  $(\mu^\varepsilon)_{\varepsilon>0}$ ,  $\mu^\varepsilon \in L_w^\infty([0, T], Q)$  satisfy  $\sup_{\varepsilon>0} \operatorname{ess\,sup}_{t \in [0, T]} \mathcal{E}(\mu^\varepsilon(t)) \leq C$ . Then for a.e.  $t \in [0, T]$  the measure  $\mu^\varepsilon(t, \cdot)$  has a Lebesgue density  $c^\varepsilon(t, \cdot)$ . Moreover, there is a subsequence (not relabeled), such that their densities  $c^\varepsilon$  are uniformly integrable in  $\Omega \times [0, T] \times \{1, 2\}$  and hence,  $c_i^\varepsilon$  converges weakly in  $L^1([0, T] \times \Omega)$  to  $c_i^0$  for  $i = 1, 2$ .*

*Proof.* The bound on the energy implies that a.e.  $t \in [0, T]$  the measure  $\mu^\varepsilon(t, \cdot)$  has a Lebesgue density  $c^\varepsilon(t, \cdot)$ . Moreover, the functional  $\mu \mapsto \int_0^T \mathcal{E}(\mu) dt$  is superlinear and convex. Hence, it follows by the Theorem of de Vallée -Poussin that  $\mu^\varepsilon$  are uniformly integrable and hence,  $\mu_i^\varepsilon$  converges weakly in  $L^1([0, T] \times \Omega)$  to  $\mu_i^0$ .  $\square$

In the following, we are going to derive compactness for the concentrations  $c_i^\varepsilon$  and the diffusive fluxes  $J_i^\varepsilon$ . It is not possible to get compactness for the fast reaction flux  $b_2^\varepsilon$  by bounding the dissipation functional. In particular, pointwise convergence for the measures  $\mu^\varepsilon(t)$  cannot be achieved.

*Remark 5.5.* To see that compactness for the fast reaction flux  $b_2^\varepsilon$  is not possible obtain, we set  $\rho^\varepsilon = 1$  constant in  $[0, T] \times \Omega \times \{1, 2\}$  and  $b_2^\varepsilon = b^\varepsilon$  constant in  $[0, T] \times \Omega$ . Then, a bound means on the dissipation functional implies a bound

$$\infty > \int_0^T \int_{\Omega} C\left(\frac{\sqrt{c_1^\varepsilon c_2^\varepsilon}}{\varepsilon}, b_2^\varepsilon(x)\right) dx \approx C\left(\frac{1}{\varepsilon}, b^\varepsilon\right) = \frac{1}{\varepsilon} C(\varepsilon b^\varepsilon) \approx |b^\varepsilon| \log(\varepsilon |b^\varepsilon| + 1).$$

Setting  $b^\varepsilon = -\log \varepsilon$ , we easily see that  $|b^\varepsilon| \log(\varepsilon |b^\varepsilon| + 1) \rightarrow 0$  as  $\varepsilon \rightarrow 0$ , however,  $b^\varepsilon \rightarrow \infty$ . Hence, it is not possible to obtain compactness for the fast reaction flux  $b_2^\varepsilon$ . Later in Lemma 5.14 the “converse” statement is proved: If  $\iint C(b) dx dt < \infty$  then  $\iint C(\frac{1}{\varepsilon}, b) dx dt \rightarrow 0$ .

Next, we are going to derive time-regularity for the sequence  $(\mu^\varepsilon)_{\varepsilon>0}$  in proving compactness for the coarse-grained trajectories  $\hat{\mu}^\varepsilon = \mu_1^\varepsilon + \mu_2^\varepsilon$ . In particular, we are able to prove pointwise convergence in time.

**Lemma 5.6** (Time Regularity of  $\mu^\varepsilon$ ). *Let  $(\mu^\varepsilon)_{\varepsilon>0}$ ,  $\mu^\varepsilon \in L_w^\infty([0, T], Q)$  satisfying the a priori bounds (5.1). Then the curves  $t \mapsto \hat{\mu}^\varepsilon := \mu_1^\varepsilon(t) + \mu_2^\varepsilon(t)$  have  $\varepsilon$ -uniform bounded total variation in the space  $\text{Prob}(\Omega)$  equipped with the 1-Wasserstein distance, i.e.*

$$\|\hat{\mu}^\varepsilon\|_{TV} := \sup \left\{ \sum_{k=1}^K \mathcal{W}_1(\hat{\mu}^\varepsilon(t_k), \hat{\mu}^\varepsilon(t_{k-1})) : 0 = t_0 < \dots < t_k < \dots < t_K = T \right\}.$$

In particular, by Helly's selection principle, we conclude pointwise convergence  $\hat{\mu}^\varepsilon(t) := \mu_1^\varepsilon(t) + \mu_2^\varepsilon(t) \xrightarrow{*} \hat{\mu}^0(t) := \mu_1^0(t) + \mu_2^0(t)$  for all  $t \in [0, T]$  in  $\text{Prob}(\Omega)$  along a suitable subsequence.

*Proof.* We exploit the dual formulation of the  $L^1$ -Wasserstein distance, i.e. integrating against Lipschitz functions (see e.g. [AGS05]). Take  $\phi \in C^1(\Omega \times \{1, 2\})$  with  $\|\phi\|_{W^{1,\infty}(\Omega)} \leq 1$  and  $\phi = (\hat{\phi}, \hat{\phi})$ . Using the continuity equations  $\left\{ b_1 + b_2 = 0 \text{ and } \begin{cases} \dot{c}_1 = -\text{div} J_1 + b_1 \\ \dot{c}_2 = -\text{div} J_2 + b_2 \end{cases} \right\}$ , we conclude for all  $[t_1, t_2] \subset [0, T]$  that

$$\begin{aligned} \int_{\Omega} \phi \cdot (d\mu^\varepsilon(t_2) - d\mu^\varepsilon(t_1)) &= \int_{t_1}^{t_2} \langle \phi, \dot{\mu}^\varepsilon \rangle dt \\ &\leq \int_{t_1}^{t_2} \int_{\Omega} \nabla \hat{\phi} \cdot (J_1^\varepsilon + J_2^\varepsilon) \, dx \, dt \leq \int_{t_1}^{t_2} \int_{\Omega} \sum_{i=1}^2 |J_i^\varepsilon| \, dx \, dt. \end{aligned}$$

By the bound on the dissipation functional, we obtain  $\varepsilon$ -uniform bounds on the term  $\int_{t_1}^{t_2} \int_{\Omega} Q(\delta_j c_j, J_j) dx dt$ . Moreover, we have  $\int_{\Omega} \rho_j^\varepsilon \leq 1$  for all  $t \in [0, T]$ . Hence by Lemma 5.1 we conclude an  $\varepsilon$ -uniform bound on each addend  $\int_{t_1}^{t_2} \int_{\Omega} \sum_{i=1}^2 |J_i^\varepsilon| \, dx \, dt$ , which by summing up implies that  $\|\hat{\mu}^\varepsilon\|_{TV}$  is  $\varepsilon$ -uniformly bounded.  $\square$

Next, the compactness result from Lemma 5.2 is used in order to prove compactness for the fluxes and spatial regularity.

**Lemma 5.7** (Regularity for the fluxes and spatial regularity). *Let  $(\mu^\varepsilon)_{\varepsilon>0}$  with  $\mu^\varepsilon \in L_w^\infty([0, T], Q)$  satisfying the a priori bounds (5.1). Then the corresponding diffusive fluxes  $J^\varepsilon : [0, T] \times \Omega \rightarrow \mathbb{R}^2$  converge weakly-star  $J^\varepsilon \xrightarrow{*} J^0$  in  $\mathcal{M}([0, T] \times \Omega \times \{1, 2\})$  and  $J_j^0 \ll \mu_j^0$ . Moreover  $\nabla \rho^{V,\varepsilon} \xrightarrow{*} \nabla \rho^{V,0}$  in  $\mathcal{M}([0, T] \times \Omega \times \{1, 2\})$  and  $\nabla \rho_j^0 \ll \mu_j^0$ . In particular, we conclude that  $\rho_j^\varepsilon$  is uniformly bounded in  $L^1([0, T], W^{1,1}(\Omega))$ , which also implies that  $\hat{c}^\varepsilon$  is uniformly in  $L^1([0, T], W^{1,1}(\Omega))$ .*

*Proof.* By the bound on the dissipation functional, we get (after extracting a suitable subsequence of  $\varepsilon \rightarrow 0$ ) that  $J^\varepsilon \xrightarrow{*} J^0$ . Moreover, we have  $\int_0^T \int_{\Omega} \frac{|J_j^\varepsilon|^2}{\rho_j^\varepsilon} dx \leq C$  and  $\rho_j^\varepsilon \xrightarrow{*} \rho_j^0$ . Hence applying the Lemma 5.2, we conclude that  $J_j^0 \ll \mu_j^0$ . Similarly, we conclude compactness for the gradients  $\nabla \rho^{V,\varepsilon}$ . The only thing that remains is to identify the limit. But this is clear by definition of the weak derivatives, i.e. integrating against smooth test functions, because this is captured in the weak star convergence. Lemma 5.1 implies that  $\rho_j^\varepsilon$  is uniformly bounded in  $L^1([0, T], W^{1,1}(\Omega))$ .  $\square$

The spatial regularity and the temporal regularity provides a compactness result by a BV-generalization of the Aubin-Lions-Simon Lemma.



**Theorem 5.8** ([BaP86, HPR18]). *Let  $X, Y, Z$  be Banach spaces such that  $X$  is compactly embedded in  $Y$ , and  $Y$  is continuously in  $Z^*$ . Let  $u^\varepsilon$  be a bounded sequence in  $L^1([0, T], X)$  and in  $BV([0, T], Z^*)$ . Then (up to a sequence)  $u^\varepsilon$  strongly converges in  $L^1([0, T], Y)$ .*

In our situation we immediately conclude that  $\hat{c}^\varepsilon$  converges strongly.

**Corollary 5.9** (Strong convergence of coarse-grained variables). *Let  $(\mu^\varepsilon)_{\varepsilon>0}$  with  $\mu^\varepsilon \in L_w^\infty([0, T], Q)$  satisfying the a priori bounds (5.1). Then the coarse-grained densities  $\hat{c}^\varepsilon$  converge strongly in  $L^1([0, T] \times \Omega)$ .*

*Proof.* Lemma 5.6 provides that  $\hat{c}^\varepsilon$  is bounded in  $BV([0, T], W^{1,\infty}(\Omega)^*)$  and Lemma 5.7 provides that  $\hat{c}^\varepsilon$  is bounded in  $L^1([0, T], W^{1,1}(\Omega))$ . Since the embedding  $W^{1,1}(\Omega) \subset L^1(\Omega)$  is compact and the embedding  $L^1(\Omega) \subset W^{1,\infty}(\Omega)^*$  is continuous, Theorem 5.8 yields that the sequence  $\hat{c}^\varepsilon$  is compact in  $L^1([0, T] \times \Omega)$ .  $\square$

It is also clear that we get convergence towards the fast manifold, which results from the Fisher information of the fast reaction.

**Lemma 5.10** (Convergence towards microscopic equilibrium and strong compactness). *Let  $(\mu^\varepsilon)_{\varepsilon>0}$ ,  $\mu^\varepsilon \in L_w^\infty([0, T], Q)$  satisfying the a priori bounds (5.1). Then there is a subsequence such that  $c^\varepsilon \rightarrow c^0$  strongly in  $L^1([0, T] \times \Omega)$  and, moreover, it holds  $\rho_1^{V,0} = \rho_2^{V,0}$  a.e. in  $[0, T] \times \Omega$ .*

*Proof.* The bound on the dissipation functional provides  $\int_0^T \int_\Omega \left( \sqrt{\rho_1^{V,\varepsilon}} - \sqrt{\rho_2^{V,\varepsilon}} \right)^2 dxdt \leq C\varepsilon$ .

Hence, we conclude  $\|\sqrt{\rho_1^{V,\varepsilon}} - \sqrt{\rho_2^{V,\varepsilon}}\|_{L^2([0,T] \times \Omega)} \rightarrow 0$  as  $\varepsilon \rightarrow 0$ . In particular, we conclude that  $\rho_1^{V,0} = \rho_2^{V,0}$ . The strong convergence towards the slow manifold provides strong convergence for the whole sequence. Indeed, using Cauchy-Schwartz inequality and  $x - y = (\sqrt{x} - \sqrt{y})(\sqrt{x} + \sqrt{y})$ , we have

$$\|\rho_1^{V,\varepsilon} - \rho_2^{V,\varepsilon}\|_{L^1([0,T] \times \Omega)} \leq \|\sqrt{\rho_1^{V,\varepsilon}} - \sqrt{\rho_2^{V,\varepsilon}}\|_{L^2([0,T] \times \Omega)} \|\sqrt{\rho_1^{V,\varepsilon}} + \sqrt{\rho_2^{V,\varepsilon}}\|_{L^2([0,T] \times \Omega)}.$$

The last term can be estimated by the AM-GM inequality

$$\|\sqrt{\rho_1^{V,\varepsilon}} + \sqrt{\rho_2^{V,\varepsilon}}\|_{L^2([0,T] \times \Omega)}^2 = \int_0^T \int_\Omega \rho_1^{V,\varepsilon} + \rho_2^{V,\varepsilon} + 2\sqrt{\rho_1^{V,\varepsilon}\rho_2^{V,\varepsilon}} dxdt \leq 2 \int_0^T \int_\Omega (\rho_1^{V,\varepsilon} + \rho_2^{V,\varepsilon}) dxdt,$$

and the right-hand side is bounded since  $\mu(t) \in Q$  for  $t \in [0, T]$ . Hence, we conclude that  $\|\rho_1^{V,\varepsilon} - \rho_2^{V,\varepsilon}\|_{L^1([0,T] \times \Omega)} \rightarrow 0$  as  $\varepsilon \rightarrow 0$ .

Using this convergence, we have that also  $c_i^\varepsilon \rightarrow c_i^0$  strongly in  $L^1([0, T] \times \Omega)$ . Indeed, we have

$$\begin{aligned} c_i^\varepsilon - w_i^V \frac{c_1^0 + c_2^0}{w_1^V + w_2^V} &= w_i^V \left( \frac{c_i^\varepsilon}{w_i^V} - \frac{c_1^0 + c_2^0}{w_1^V + w_2^V} \right) \\ &= w_i^V \left( \frac{c_i^\varepsilon}{w_i^V} - \frac{c_1^\varepsilon + c_2^\varepsilon}{w_1^V + w_2^V} + \frac{c_1^\varepsilon + c_2^\varepsilon}{w_1^V + w_2^V} - \frac{c_1^0 + c_2^0}{w_1^V + w_2^V} \right) \\ &= (-1)^i \left( \frac{\frac{c_2^\varepsilon}{w_2^V} - \frac{c_1^\varepsilon}{w_1^V}}{w_1^V w_2^V (w_1^V + w_2^V)} \right) + \frac{w_i^V}{w_1^V + w_2^V} (c_1^\varepsilon + c_2^\varepsilon - (c_1^0 + c_2^0)), \end{aligned}$$

and both terms converge strongly to zero as  $\varepsilon \rightarrow 0$  by convergence of  $\hat{c}^\varepsilon \rightarrow \hat{c}^0$  and  $\rho_1^{V,\varepsilon} - \rho_2^{V,\varepsilon} \rightarrow 0$ .  $\square$

Finally, we show that the limit  $\mu^0 = c^0 dx$  has an absolutely continuous representative in the space of probability measures. To do this, we exploit the characterization of absolutely continuous curves as solutions of the continuity equation following [AGS05].

**Proposition 5.11.** *Let  $(\mu^\varepsilon)_{\varepsilon>0}$ ,  $\mu^\varepsilon \in L^\infty([0, T], Q)$  satisfying the a priori bounds (5.1) and let  $c^0$  be the limit of the densities  $c^\varepsilon$ . Then the coarse-grained slow variable  $\hat{\mu} = \mu_1^0 + \mu_2^0 = (c_1^0 + c_2^0) dx \in L_w^\infty([0, T], \text{Prob}(\Omega))$  has a representative (in time), which is absolutely continuous in the space of probability measures equipped with the 2-Wasserstein metric. Moreover, each component  $\mu_i^0$  has an absolutely continuous representative (in time), which is absolutely continuous in the space of non-negative Radon measures equipped with the 1-Wasserstein metric.*

*Proof.* The coarse-grained measures  $\hat{\mu}^\varepsilon$  satisfy continuity equation  $\dot{\hat{\mu}}^\varepsilon + \text{div}(\hat{J}^\varepsilon) = 0$  in the sense of distributions where  $\hat{J}^\varepsilon = J_1^\varepsilon + J_2^\varepsilon$  is the coarse-grained diffusion flux. Since the linear continuity equation is stable under weak convergence, we conclude that also the limits satisfy the same continuity equation  $\dot{\hat{\mu}}^0 + \text{div}(\hat{J}^0) = 0$ , where  $\hat{J}^0$  is the weak\*-limit of  $\hat{J}^\varepsilon$  (see Lemma 5.7). Using (4.9), the bound on the dissipation functional implies a bound  $\int_0^T \int_\Omega Q(\hat{c}^0, \hat{J}^0) dx dt < \infty$ . Let us define the transport velocity  $\hat{v} \in \mathcal{M}([0, T] \times \Omega, \mathbb{R})$

$$\hat{v} = \begin{cases} \frac{\hat{J}}{\hat{c}} & \text{for } \hat{c} > 0 \\ 0 & \text{for } \hat{c} = 0 \end{cases}.$$

Then  $\int_0^T \int_\Omega Q(\hat{c}, \hat{J}) dx dt = \frac{1}{2} \int_0^T \int_\Omega |\hat{v}|^2 \hat{c} dx dt = \frac{1}{2} \int_0^T \int_\Omega |\hat{v}|^2 d\hat{\mu} dt$  and the bound on the dissipation functional implies the bound on the Borel velocity field  $\|\hat{v}\|_{L^2(\hat{\mu})} < \infty$ . Hence, by Theorem 8.3.1 from [AGS05] it follows that  $t \mapsto \hat{\mu}(t) \in (\text{Prob}(\Omega), d_{W_2})$  has a continuous representative which is absolutely continuous.

To prove time-regularity for  $\mu_i^0$  for  $i = 1, 2$ , we first observe that  $\mu_i^0 = \frac{w_i^V}{w_1^V + w_2^V} \hat{\mu}$  is a non-negative Radon measure. To show that it has an absolutely continuous representative, we proceed as in Lemma 5.6 and exploit the dual formulation of the 1-Wasserstein distance on the space of non-negative Radon measures, i.e. integrating against Lipschitz functions (see e.g. [Edw11]). Let  $\phi \in C^1(\Omega)$  with  $\|\phi\|_{W^{1,\infty}(\Omega)} \leq 1$ . Using the continuity equations  $\dot{\mu}_i^0 + \text{div}(\hat{J}^0) = 0$ , we conclude for all  $[t_1, t_2] \subset [0, T]$  that

$$\begin{aligned} \int_\Omega \phi \cdot (d\mu_i^0(t_2) - d\mu_i^0(t_1)) &= \int_{t_1}^{t_2} \langle \phi, \dot{\mu}_i^0 \rangle dt = \int_{t_1}^{t_2} \langle \phi \frac{w_i^V}{w_1^V + w_2^V}, \dot{\hat{\mu}} \rangle dt \\ &\leq \int_{t_1}^{t_2} \int_\Omega \nabla \left( \phi \frac{w_i^V}{w_1^V + w_2^V} \right) \cdot \hat{J} dx dt \leq C \int_{t_1}^{t_2} \int_\Omega |\hat{J}| dx dt, \end{aligned}$$

where  $C = C(w, V)$ . The bound on the dissipation functional provides again that the right-hand side is bounded for each interval  $[t_1, t_2] \subset [0, T]$ . Summing up, we conclude that  $\mu_i^0$  has an absolutely continuous representative, which proves the claim.  $\square$

## 5.2 Liminf-estimate

In this section, we state and prove the liminf-estimate of the  $\Gamma$ -convergence result Theorem 4.2. Once the compactness is established the proof of the liminf-estimate is comparatively easy.

**Theorem 5.12.** *Let  $\mu^\varepsilon \rightarrow \mu^0$  in  $L_w^\infty([0, T], Q)$  such that  $\sup_{\varepsilon \in ]0,1]} \sup_{t \in [0,T]} \mathcal{E}(\mu^\varepsilon) < \infty$ . Then, we have the liminf-estimate*

$$\liminf_{\varepsilon \rightarrow 0} \mathfrak{D}_\varepsilon(\mu^\varepsilon) \geq \mathfrak{D}_0(\mu^0),$$

where the limit dissipation functional is defined by

$$\mathfrak{D}_0^V(\mu) = \begin{cases} \int_0^T \mathcal{R}_{\text{eff}}(\mu, \dot{\mu}) + \mathcal{R}_{\text{eff}}^*(\mu, -D\mathcal{E}^V(\mu)) dt, & \mu \in \text{AC}([0, T], Q), \mu = c dx \text{ a.e. in } [0, T] \\ \infty & \text{otherwise} \end{cases}$$

with

$$\begin{aligned} \mathcal{R}_{\text{eff}}^*(\mu, \xi) &= \mathcal{R}_{\text{diff}}^*(\mu, \xi) + \chi_{\{\xi_1 = \xi_2\}}(\xi) \\ \mathcal{R}_{\text{eff}}(\mu, v) &= \inf \left\{ \sum_{j=1}^2 \int_{\Omega} \tilde{\mathcal{Q}}(\delta_j c_j, J_j) dx : u_1 + u_2 = 0, \left\{ \begin{array}{l} v_1 = -\text{div} J_1 + u_1 \\ v_2 = -\text{div} J_2 + u_2 \end{array} \right\} \right\}. \end{aligned}$$

*Proof.* We may assume that  $\mathfrak{D}_\varepsilon(\mu^\varepsilon) \leq C < \infty$  (otherwise the claim is trivial). For the given curves  $t \mapsto \mu^\varepsilon(t) \in Q$  take diffusive fluxes  $J^\varepsilon$  and reactive fluxes  $b^\varepsilon$ , which satisfy the generalized continuity equation

$$(c, J, b) \in (\text{gCE}) \Leftrightarrow \left\{ b_1 + b_2 = 0 \text{ and } \left\{ \begin{array}{l} \dot{c}_1 = -\text{div} J_1 + b_1 \\ \dot{c}_2 = -\text{div} J_2 + b_2 \end{array} \right\} \right\},$$

and approximate the infimum in  $\mathfrak{D}_\varepsilon(\mu^\varepsilon)$  arbitrarily close, i.e.

$$\mathfrak{D}_\varepsilon(\mu_\varepsilon) + \varepsilon \geq \int_0^T \mathcal{D}_\varepsilon(\mu_\varepsilon, J^\varepsilon, b^\varepsilon) dt.$$

The integrand  $\mathcal{D}_\varepsilon$  consists of a velocity and a slope part and both of them split into a reaction and a diffusion part:

$$\begin{aligned} \mathcal{D}_\varepsilon(\mu, J, b) &= \int_{\Omega} \sum_{j=1}^2 \tilde{\mathcal{Q}}(\delta_j c_j, J_j) dx + \int_{\Omega} \tilde{\mathcal{C}} \left( \frac{\sqrt{c_1 c_2}}{\varepsilon}, b_2(x) \right) dx + \\ &\quad + \frac{1}{2} \int_{\Omega} \sum_{j=1}^2 \delta_j w_j^V \frac{|\nabla \rho_j^V|^2}{\rho_j^V} dx + \frac{2}{\varepsilon} \int_{\Omega} \sqrt{w_1^V w_2^V} \left( \sqrt{\rho_1^V} - \sqrt{\rho_2^V} \right)^2 dx \\ &=: \mathcal{V}_{\text{diff}}(\mu, J) + \mathcal{V}_{\text{react}, \varepsilon}(\mu, b) + \mathcal{S}_{\text{diff}}(\mu) + \mathcal{S}_{\text{react}, \varepsilon}(\mu). \end{aligned}$$

Clearly, we also have  $\int_0^T \mathcal{D}_\varepsilon(\mu_\varepsilon, J^\varepsilon, b^\varepsilon) dt \leq C + \varepsilon < \infty$ . By Lemma 5.10, we conclude compactness for the densities  $c^\varepsilon \rightarrow c^0$  in  $L^1([0, T] \times \Omega \times \{1, 2\})$  and by Lemma 5.7 that  $J^\varepsilon \xrightarrow{*} J^0$  in  $\mathcal{M}([0, T] \times \Omega \times \{1, 2\})$ . Using the lower-semicontinuity result from Lemma 5.2 (which implies the liminf-estimates for  $\mathcal{V}_{\text{diff}}$  and  $\mathcal{S}_{\text{diff}}$ ) and that  $\mathcal{V}_{\text{react}, \varepsilon}, \mathcal{S}_{\text{react}, \varepsilon} \geq 0$ , we obtain the estimate

$$\liminf_{\varepsilon \rightarrow 0} \int_0^T \mathcal{D}_\varepsilon(\mu^\varepsilon) dt \geq \int_0^T \left\{ \int_{\Omega} \sum_{j=1}^2 \tilde{\mathcal{Q}}(\delta_j c_j^0, J_j^0) dx + \frac{1}{2} \int_{\Omega} \sum_{j=1}^2 \delta_j w_j^V \frac{|\nabla \rho_j^0|^2}{\rho_j^0} dx \right\} dt.$$

Let us define  $A_i^\varepsilon = \dot{c}_i^\varepsilon + \text{div} J_i^\varepsilon$ . We conclude convergence for  $A_i^\varepsilon \rightarrow A_i^0$  in the sense of distributions, and, moreover, we have  $A_1^\varepsilon + A_2^\varepsilon \rightarrow \dot{c}_1^0 + \dot{c}_2^0 + \text{div} J_1^0 + \text{div} J_2^0 = 0$ . Let us define  $u_1 := A_0^2 = \dot{c}_2^0 + \text{div} J_2^0$  and  $u_2 := A_0^1 = \dot{c}_1^0 + \text{div} J_1^0$ . Then  $u_1 + u_2 = 0$ ,  $A_1^0 + u_1 = 0$  and  $A_2^0 + u_2 = 0$ . In particular, we conclude the pointwise estimate

$$\int_{\Omega} \sum_{j=1}^2 \tilde{\mathcal{Q}}(\delta_j c_j^0, J_j^0) dx \geq \inf_{(J, u)} \left\{ \int_{\Omega} \sum_{j=1}^2 \tilde{\mathcal{Q}}(\delta_j c_j^0, J_j^0) dx : \left\{ \begin{array}{l} \dot{c}_1 + \text{div} J_1 + u_1 = 0 \\ \dot{c}_2 + \text{div} J_2 + u_2 = 0 \\ u_1 + u_2 = 0 \end{array} \right\} \right\},$$

which finally establishes the liminf-estimate

$$\liminf_{\varepsilon \rightarrow 0} \int_0^T \mathcal{D}_\varepsilon(\mu^\varepsilon) dt \geq \mathfrak{D}_0^V(\mu).$$

□

### 5.3 Construction of the recovery sequence

In this section, we construct the recovery sequence for the functional  $\mathfrak{D}_0^V$  to finish the  $\Gamma$ -convergence result in Theorem 4.2. To be precise, we will show the following:

**Theorem 5.13.** *Let  $\mu^0 \in L_w^\infty([0, T] \times \Omega, \mathbb{R}_{\geq 0}^2)$  such that the a priori bounds  $\mathfrak{D}_0^V(\mu^0) < \infty$  and  $\text{ess sup}_{t \in [0, T]} \mathcal{E}(\mu^0(t)) < \infty$  hold. Then there is a sequence  $(\mu^\varepsilon)_{\varepsilon > 0}$ ,  $\mu^\varepsilon \in \text{AC}([0, T], Q)$ ,  $\sup_{\varepsilon > 0} \text{ess sup}_{t \in [0, T]} \mathcal{E}(\mu^\varepsilon(t)) < \infty$ , such that the densities converge  $c^\varepsilon \rightarrow c^0$  strongly in  $L^1([0, T] \times \Omega \times \{1, 2\})$  and we have  $\mathfrak{D}_\varepsilon^V(\mu^\varepsilon) \rightarrow \mathfrak{D}_0^V(\mu^0)$ .*

Using Proposition 4.5, we see that the limit functionals do not contain more information than the functionals in coarse-grained variables and it holds  $\mathfrak{D}_0^V(\mu^0) = \hat{\mathfrak{D}}(\hat{\mu}^0)$ . Hence, we may reconstruct the dissipation functional with the corresponding diffusion and reaction flux  $(J, b)$  from the coarse-grained variables  $(\hat{c}, \hat{J})$ .

The dissipation functional  $\mathfrak{D}_0^V$  is defined on the space of general fluctuations around the solution of the evolution equation (1.2). These fluctuations are neither strictly positive nor smooth. The proof of the limsup-estimate is done in several steps, which are elaborated in the next lemmas. The bound  $\mathfrak{D}_0^V(\mu^0) < \infty$  can be assumed without loss of generality because the other case is already treated in the liminf-estimate.

*Proof of Theorem 5.13.* We do the reconstruction in three steps using different approximation methods. We will do the following steps:

- 1 Proposition 5.21 shows that for  $\mu^0 = c^0 dx$  with sufficiently smooth and positive density  $c^0$  the constant sequence  $\mu^{\varepsilon, \gamma} = \mu^0$  satisfies  $|\mathfrak{D}_\varepsilon(\mu^{\varepsilon, \gamma}) - \mathfrak{D}_0(\mu^{\varepsilon, \gamma})| \rightarrow 0$ .
- 2 Lemma 5.15 overcomes the positivity assumption, i.e. it shows that for all  $\mu^0 = c^0 dx$  there is a positive  $c^\gamma$  such that  $c^\gamma \rightarrow c^0$  and  $\mathfrak{D}_0(\mu^\gamma) \rightarrow \mathfrak{D}_0(\mu^0)$  as  $\gamma \rightarrow 0$ .
- 3 A mollification argument as in [AGS05, Lemma 8.1.10] and stated in Lemma 5.20 allows us to overcome regularity by smoothing, which shows  $\mathfrak{D}_0(\mu^{\varepsilon, \gamma}) \rightarrow \mathfrak{D}_0(\mu^\gamma)$  as  $\varepsilon \rightarrow 0$ .

Hence, defining the recovery sequence  $\mu^\varepsilon := \mu^{\varepsilon, \gamma}$ , we have

$$|\mathfrak{D}_\varepsilon(\mu^\varepsilon) - \mathfrak{D}_0(\mu^0)| \leq |\mathfrak{D}_\varepsilon(\mu^{\varepsilon, \gamma}) - \mathfrak{D}_0(\mu^{\varepsilon, \gamma})| + |\mathfrak{D}_0(\mu^{\varepsilon, \gamma}) - \mathfrak{D}_0(\mu^\gamma)| + |\mathfrak{D}_0(\mu^\gamma) - \mathfrak{D}_0(\mu^0)|,$$

where the first term tends to zero by the first reconstruction step, the second term tends to zero by the third reconstruction step and the third term tends to zero by the second reconstruction step, which in total proves the desired convergence. □

Before performing the three recovery steps in Section 5.3.2, we first illustrate the general idea of constructing the recovery sequence by forgetting about positivity and regularity issues for the first moment.

### 5.3.1 Construction of recovery sequence for smooth and positive measures

To show the general idea, let us firstly assume that the density of  $\hat{\mu}$  is sufficiently smooth and positive, i.e. we assume that its Lebesgue density satisfies  $\hat{c} \geq \frac{1}{C} > 0$  on  $\Omega \times [0, T]$  and has a enough regularity that will be specified below. Let  $\hat{J}$  be the diffusion flux which provides the minimum in  $\hat{\mathcal{D}}(\hat{\mu}^0) = \mathcal{D}_0^V(\mu)$  and satisfies  $\hat{c} + \operatorname{div}(\hat{J}) = 0$ . We define the reconstructed variables by

$$\begin{aligned} c_1 &= \frac{w_1^V}{w_1^V + w_2^V} \hat{c}, & c_2 &= \frac{w_2^V}{w_1^V + w_2^V} \hat{c}, & J_1 &= \frac{\delta_1 w_1^V}{\delta_1 w_1^V + \delta_2 w_2^V} \hat{J}, & J_2 &= \frac{\delta_2 w_2^V}{\delta_1 w_1^V + \delta_2 w_2^V} \hat{J}. \\ b_1 &= \left( \frac{\delta_1 - \delta_2}{\delta_1 w_1^V + \delta_2 w_2^V} \frac{w_1^V w_2^V}{w_1^V + w_2^V} \right) \operatorname{div} \hat{J} + \hat{J} \cdot \nabla \left( \frac{\delta_1 w_1^V}{\delta_1 w_1^V + \delta_2 w_2^V} \right), & b_2 &= -b_1. \end{aligned} \quad (5.3)$$

The reconstructed concentrations  $c$  and diffusion fluxes  $J = (J_1, J_2)$  are proportional to the coarse grained concentration  $\hat{c}$  and diffusion flux  $\hat{J}$ , respectively. On the coarse-grained level, which considers only one species there is no reaction flux anymore. (This changes when considering large reaction-diffusion system as explained in Section 6). The reactive flux  $b = (b_1, b_2)$  is given as a function of the coarse-grained diffusion flux  $\hat{J}$ , which means that in the limit the diffusion determines the hidden reaction.

Concerning regularity issues, we immediately observe the following. Since  $w^V$  is smooth and positive,  $c_1, c_2$  have the same regularity as  $\hat{c}$  and also  $J_1, J_2$  have the same regularity as  $\hat{J}$ . Only the reaction fluxes  $b_i$  are a priori not well-defined (e.g. in  $L^1$ ) if  $\operatorname{div} \hat{J}$  and  $\hat{J}$  are not regular enough. This means, that the reaction flux between the fast-connected species is not well-defined for general  $\hat{J} \in \mathcal{M}(\Omega \times [0, T], \mathbb{R}^3)$ . Note, that regularity assumptions for  $\operatorname{div} \hat{J}$  are not needed if  $\delta_1 = \delta_2$ , i.e. if both species diffuse with the same diffusion constant. In particular, in this situation no regularization argument as in Lemma 5.20 is necessary. Moreover, no additional regularity for  $\hat{J}$  is needed if  $\frac{\delta_1 w_1^V}{\delta_1 w_1^V + \delta_2 w_2^V} = \theta \in ]0, 1[$  is constant. This is equivalent to

$$V_1(x) - V_2(x) = \log \left( \frac{1 - \theta \delta_1 \beta}{\theta \delta_2 \alpha} \right) = \text{const},$$

which means that the potentials  $V_1, V_2$  differ in a constant on  $\Omega$ . In particular, this implies that for the coarse-grained potential  $\hat{V}$  we have  $\nabla \hat{V} = \nabla V_1 = \nabla V_2$ . As we will see, enough regularity for  $\hat{J}$  is already obtained from bounds on the dissipation functional. Of course, regularity properties for  $\operatorname{div} \hat{J}$  and  $\hat{J}$  are independent of each other.

So let us assume for the moment that  $b_i$  is well-defined. Then we conclude  $(c, J, b) \in (\text{gCE})$ , because we have

$$\begin{aligned} \dot{c}_1 + \operatorname{div} J_1 &= \frac{w_1^V}{w_1^V + w_2^V} \dot{\hat{c}} + \operatorname{div} \left( \frac{\delta_1 w_1^V}{\delta_1 w_1^V + \delta_2 w_2^V} \hat{J} \right) = \\ &= \frac{w_1^V}{w_1^V + w_2^V} \dot{\hat{c}} + \hat{J} \cdot \nabla \left( \frac{\delta_1 w_1^V}{\delta_1 w_1^V + \delta_2 w_2^V} \right) + \left( \frac{\delta_1 w_1^V}{\delta_1 w_1^V + \delta_2 w_2^V} \right) \operatorname{div} \hat{J} \\ &= -\frac{w_1^V}{w_1^V + w_2^V} \operatorname{div} \hat{J} + \hat{J} \cdot \nabla \left( \frac{\delta_1 w_1^V}{\delta_1 w_1^V + \delta_2 w_2^V} \right) + \left( \frac{\delta_1 w_1^V}{\delta_1 w_1^V + \delta_2 w_2^V} \right) \operatorname{div} \hat{J} = b_1, \end{aligned}$$

where we used that  $(\hat{c}, \hat{J})$  solves  $\dot{\hat{c}} + \operatorname{div} \hat{J} = 0$ . Similarly, we see that  $\dot{c}_2 + \operatorname{div} J_2 = b_2$  and, by definition, we have  $b_1 + b_2 = 0$ . Moreover, boundary properties of  $\hat{J}$  remain for  $J = (J_1, J_2)$ .

Since  $\frac{c_1}{w_1^V} = \frac{c_2}{w_2^V}$ , we conclude that  $\mathcal{D}_\varepsilon^V(\mu) \leq \mathcal{D}_0(\mu) + \int_0^T \int_\Omega \tilde{C} \left( \frac{\sqrt{c_1 c_2}}{\varepsilon}, b_2 \right) dx dt$ . That means, that for proving that the constant sequence  $\mu^\varepsilon = \mu$  is a recovery sequence, it suffice to show that

$\int_0^T \int_\Omega \tilde{\mathcal{C}}\left(\frac{\sqrt{c_1 c_2}}{\varepsilon}, b_2\right) dxdt \rightarrow 0$  as  $\varepsilon \rightarrow 0$ . This is, in fact, shown in the next lemma under the assumption that  $\operatorname{div} \hat{J}, \hat{J} \in L^C([0, T] \times \Omega)$ . The proof basically uses the monotonicity property of the Legendre dual function  $\tilde{\mathcal{C}}(a_1, b) \leq \tilde{\mathcal{C}}(a_2, b)$  as  $a_1 \geq a_2$  (see Lemma 4.4), its superlinear growth and the dominated convergence theorem.

**Lemma 5.14.** *Let  $\hat{c} \in L^1(\Omega \times [0, T])$  with  $\hat{c} \geq \frac{1}{C}$  a.e. in  $[0, T] \times \Omega$  for a constant  $C > 0$ , and let  $\hat{J} : \Omega \rightarrow \mathbb{R}^d$  satisfy  $\operatorname{div} \hat{J}, |\hat{J}| \in L^C([0, T] \times \Omega)$ . Then  $\int_0^T \int_\Omega \tilde{\mathcal{C}}\left(\frac{\sqrt{c_1 c_2}}{\varepsilon}, b_2\right) dxdt \rightarrow 0$  as  $\varepsilon \rightarrow 0$ .*

*Proof.* Since  $\hat{c} \geq \frac{1}{C}$  a.e., Lemma 4.4 yields  $\tilde{\mathcal{C}}\left(\frac{\sqrt{c_1 c_2}}{\varepsilon}, b_2\right) \leq \tilde{\mathcal{C}}\left(\frac{1}{C\varepsilon}, b_2\right) = \frac{1}{C\varepsilon} \mathcal{C}(C\varepsilon b_2)$ . Moreover, we have the estimate

$$\frac{1}{2}|r| \log(|r| + 1) \leq \mathcal{C}(r) \leq 2|r| \log(|r| + 1),$$

which implies

$$\int_0^T \int_\Omega \tilde{\mathcal{C}}\left(\frac{\sqrt{c_1 c_2}}{\varepsilon}, b_2\right) dxdt \leq \frac{1}{C\varepsilon} \int_0^T \int_\Omega \mathcal{C}(C\varepsilon b_2) dxdt \leq \int_0^T \int_\Omega 2|b_2| \log(C\varepsilon|b_2| + 1) dxdt.$$

By assumption, we have that  $\operatorname{div} \hat{J}, |\hat{J}| \in L^C([0, T] \times \Omega)$ . Since  $V \in C^1(\bar{\Omega})$  and the Orlicz space  $L^C([0, T] \times \Omega)$  is a Banach space, we conclude that  $b_2 \in L^C([0, T] \times \Omega)$ . By the inequality for  $\mathcal{C}$ , this implies that for  $\varepsilon < \frac{1}{C}$ , the right-hand side is bounded. We show that (for a subsequence) the integrand converges to zero pointwise a.e. in  $[0, T] \times \Omega$ . By the dominated convergence theorem, this would imply that  $\int_0^T \int_\Omega \tilde{\mathcal{C}}\left(\frac{\sqrt{c_1 c_2}}{\varepsilon}, b_2\right) dxdt \rightarrow 0$  as  $\varepsilon \rightarrow 0$ .

To see that the integrand converges to zero pointwise, we firstly observe that  $b_2 \in L^C([0, T] \times \Omega) \subset L^1([0, T] \times \Omega)$ , which means that

$$\int_0^T \int_\Omega \log(\varepsilon|b_2| + 1) dxdt \leq \varepsilon \int_0^T \int_\Omega |b_2| dxdt \rightarrow 0.$$

Hence, (for a subsequence)  $\log(\varepsilon|b_2| + 1)$  converges pointwise to zero and, thus, also  $|b_2| \log(C\varepsilon|b_2| + 1)$ .  $\square$

In fact, the above proof is quite robust and already suggests that the same convergence holds even if  $\hat{c}^\varepsilon \rightarrow 0$  not too fast somewhere in  $\Omega \times [0, T]$  and  $\|\hat{b}_\varepsilon^2\| \approx \varepsilon^{-\alpha}$  for some  $\alpha > 0$ . This is proved in Proposition 5.21.

In the following, we need to overcome the positivity assumption  $\hat{c} \geq \frac{1}{C}$  and the regularity assumption for  $\hat{J}$  and  $\operatorname{div} \hat{J}$ . The first is done by shifting the density  $\hat{c}^\delta := \frac{1}{Z_\delta}(\hat{c} + \delta)$ ,  $\delta > 0$ ; the necessary regularity of  $\hat{J}$  is provided immediately by the bound on the dissipation functional; the regularity of  $\operatorname{div} \hat{J}$  is achieved by smoothing using that  $(\hat{c}, \hat{J})$  is a solution of the coarse-grained continuity equation  $\hat{c} + \operatorname{div} \hat{J} = 0$ .

### 5.3.2 Auxiliary results for construction of recovery sequence for general measures

First, we show how to overcome the positivity assumption. This is done by a controlled positive shift.

**Lemma 5.15.** *For all  $\mu^0 = c^0 dx$  satisfying  $\mathfrak{D}_0^V(\mu^0) = \hat{\mathfrak{D}}(\hat{\mu}^0) < \infty$  there is a sequence  $(\hat{c}^\gamma)$  of densities satisfying  $\hat{c}^\gamma \geq \gamma$ ,  $\hat{c}^\gamma \rightarrow \hat{c}$  in  $L^1([0, T] \times \Omega)$  such that for their corresponding measures we have  $\hat{\mathfrak{D}}(\hat{\mu}^\gamma) = \mathfrak{D}_0(\mu^\gamma) \rightarrow \mathfrak{D}_0(\mu^0) = \hat{\mathfrak{D}}(\mu^0)$  as  $\gamma \rightarrow 0$  and  $\sup_{\gamma \in [0, 1]} \operatorname{ess\,sup}_{t \in [0, T]} \mathcal{E}(\mu^\gamma(t)) < \infty$ .*

*Proof.* For small  $\gamma > 0$ , we define  $\hat{c}^\gamma := \frac{1}{Z_\gamma}(\hat{c} + 2\gamma)$ , where  $Z_\gamma = 1 + 2\gamma|\Omega| = 1 + 2\gamma > 0$  is the normalization factor such that  $\int_\Omega \hat{c}^\gamma dx = 1$ . Hence,  $Z_\gamma \searrow 1$ ,  $\hat{c}^\gamma \rightarrow \hat{c}$ ,  $\sup_{\gamma \in [0,1]} \text{ess sup}_{t \in [0,T]} \mathcal{E}(\mu^\gamma(t)) < \infty$  and w.l.o.g. we assume that  $\hat{c}^\gamma \geq \gamma$ . Moreover, we define  $\hat{J}^\gamma := \frac{1}{Z_\gamma} \hat{J}$ . Clearly,  $\hat{J}^\gamma \cdot \nu = 0$  on  $\partial\Omega$  and  $(\hat{c}^\gamma, \hat{J}^\gamma)$  solves the continuity equation  $\dot{\hat{c}}^\gamma + \text{div} \hat{J}^\gamma = 0$ . We compute the terms in the dissipation functional  $\mathfrak{D}_0^V(\mu^\gamma)$ . We have  $\hat{\rho}^{V,\gamma} = \frac{\hat{c}^\gamma}{w^V} = \frac{1}{Z_\gamma} \frac{\hat{c} + 2\gamma}{w^V}$ . Using the bounds  $\sup_{x \in \bar{\Omega}} \{|\nabla(1/w^V)|, w^V\} \leq C$ , we get

$$\nabla \hat{\rho}^{V,\gamma} = \frac{1}{Z_\gamma} \left\{ \nabla \left( \frac{\hat{c}}{w^V} \right) + 2\gamma \nabla \left( \frac{1}{w^V} \right) \right\} \Rightarrow |\nabla \hat{\rho}^{V,\gamma}| \leq \frac{1}{Z_\gamma} \{ |\nabla \hat{\rho}^V| + 2\gamma C \}$$

Using the estimates  $\frac{1}{a+\delta} \leq \frac{1}{a}$ ,  $\frac{1}{Z_\gamma} \leq 1$  and the inequality  $2xy \leq \sqrt{\gamma}x^2 + \frac{1}{\sqrt{\gamma}}y^2$  in the second estimate, we get the pointwise estimate

$$\begin{aligned} \frac{|\nabla \hat{\rho}^{V,\gamma}|^2}{\hat{\rho}^{V,\gamma}} &= \frac{1}{Z_\gamma} \frac{(|\nabla \hat{\rho}^V| + 2\gamma C)^2}{\hat{\rho}^V + \frac{2\gamma}{w^V}} = \frac{1}{Z_\gamma} \frac{|\nabla \hat{\rho}^V|^2 + 4\gamma C |\nabla \hat{\rho}^V| + 4\gamma^2 C^2}{\hat{\rho}^V + \frac{2\gamma}{w^V}} \\ &\leq \frac{|\nabla \hat{\rho}^V|^2}{\hat{\rho}^V} + 2 \frac{2\gamma C |\nabla \hat{\rho}^V|}{\hat{\rho}^V + \frac{2\gamma}{w^V}} + \frac{4\gamma^2 C^2}{\hat{\rho}^V + \frac{2\gamma}{w^V}} \\ &\leq \frac{|\nabla \hat{\rho}^V|^2}{\hat{\rho}^V} + \frac{1}{\hat{\rho}^V + \frac{2\gamma}{w^V}} \left( \sqrt{\gamma} |\nabla \hat{\rho}^V|^2 + \frac{1}{\sqrt{\gamma}} \{2\gamma C\}^2 \right) + 2\gamma C^2 w^V \\ &\leq \frac{|\nabla \hat{\rho}^V|^2}{\hat{\rho}^V} (1 + \sqrt{\gamma}) + 2\sqrt{\gamma} w^V C^2 (1 + \sqrt{\gamma}). \end{aligned}$$

Hence,  $\int_0^T \int_\Omega \delta^V w^V \frac{|\nabla \hat{\rho}^{V,\gamma}|^2}{\hat{\rho}^{V,\gamma}} dx dt \rightarrow \int_0^T \int_\Omega \delta^V w^V \frac{|\nabla \hat{\rho}^V|^2}{\hat{\rho}^V} dx dt$  as  $\gamma \rightarrow 0$ .

Similarly, we get

$$\int_0^T \int_\Omega \frac{|\hat{J}^\gamma|^2}{\hat{\delta} \hat{c}^\gamma} dx dt = \frac{1}{Z_\gamma} \int_0^T \int_\Omega \frac{|\hat{J}|^2}{\hat{\delta}(\hat{c} + 2\gamma)} dx dt \leq \int_0^T \int_\Omega \frac{|\hat{J}|^2}{\hat{\delta} \hat{c}} dx dt,$$

which implies that  $\int_0^T \int_\Omega \tilde{\mathbf{Q}}(\hat{\delta} \hat{c}^\gamma, \hat{J}^\gamma) dx dt \leq \int_0^T \int_\Omega \tilde{\mathbf{Q}}(\hat{\delta} \hat{c}, \hat{J}) dx dt$ . Hence, we conclude that  $\hat{c}^\gamma \geq \gamma$ ,  $\hat{c}^\gamma \rightarrow \hat{c}$  and  $\hat{\mathfrak{D}}(\hat{c}^\gamma) \rightarrow \hat{\mathfrak{D}}(\hat{c})$  as  $\gamma \rightarrow 0$ .  $\square$

Next, we are going to show that the flux  $b_1 = -b_2$  can be made sufficiently smooth, i.e. at least in  $L^C$  which would allow us to proceed similar as in Lemma 5.14.

Recalling the formula for the reconstructed flux, we have

$$b_2 = \left( \frac{\delta_1 - \delta_2}{\delta_1 w_1^V + \delta_2 w_2^V} \frac{w_1^V w_2^V}{w_1^V + w_2^V} \right) \text{div} \hat{J} + \hat{J} \cdot \nabla \left( \frac{\delta_1 w_1^V}{\delta_1 w_1^V + \delta_2 w_2^V} \right) =: a_1 \text{div} \hat{J} + \hat{J} \cdot a_2$$

and  $b_2 = -b_1$ , where  $a_1 \in C^1(\bar{\Omega}, \mathbb{R})$ ,  $a_2 \in C^0(\bar{\Omega}, \mathbb{R}^d)$  such that  $\sup_{x \in \bar{\Omega}} \{|a_1(x)|, |a_2(x)|\} \leq C$ . In particular, the regularity of  $b_1 = -b_2$  does not depend on  $a_1, a_2$ . We are going to prove that  $\text{div} \hat{J}$  and  $\hat{J}$  have enough regularity. The regularity of  $\hat{J}$  follows from the bound on the dissipation functional. The regularity of  $\text{div} \hat{J}$  is achieved by mollification.

First, we show that  $\hat{J} \in L^{\tilde{p}}([0, T] \times \Omega)$ , for some  $\tilde{p} > 1$ . Clearly, we have  $\hat{J} \in L^1([0, T] \times \Omega)$  by Lemma 5.1 and the bound on the dissipation functional. To improve the regularity of  $\hat{J}$ , we firstly improve the regularity of  $\hat{c} \in L^p([0, T] \times \Omega)$  which is provided from the bound on the dissipation

functional  $\mathfrak{D}_0$  and the energy functional  $\mathcal{E}$ . The bound on the energy yields  $\hat{c} \in L^\infty([0, T], L^1(\Omega))$ . The bound on the Fisher-information yields again by Lemma 5.1 that  $\hat{c} \in L^1([0, T], W^{1,1}(\Omega))$  as in Lemma 5.7. By the Sobolev embedding theorem, we have the compact embedding  $W^{1,1}(\Omega) \subset L^q(\Omega)$ , where  $1 - \frac{1}{d} > \frac{1}{q} \Leftrightarrow q < \frac{d}{d-1}$ . Thus  $\hat{c} \in L^\infty([0, T], L^1(\Omega)) \cap L^1([0, T], L^{\frac{d}{d-1}}(\Omega))$ . Using the next classical interpolation result we get  $\hat{c} \in L^p([0, T] \times \Omega)$  for some  $p > 1$ .

**Theorem 5.16** (Theorem 5.1.2, [BeL76]). *Let  $X_1, X_2$  be Banach spaces. Then for the complex interpolation spaces, it holds for any  $\theta \in ]0, 1[$*

$$[L^{p_1}([0, T], X_1), L^{p_2}([0, T], X_2)]_\theta \simeq L^{p_\theta}([0, T], [X, Y]_\theta),$$

$$\text{where } \frac{1}{p_\theta} = \frac{1-\theta}{p_1} + \frac{\theta}{p_2}.$$

In our particular situation, we have the following.

**Lemma 5.17.** *Let  $\mu^0 \in L_w^\infty([0, T], Q)$  such that  $\mathfrak{D}_0^V(\mu^0) < \infty$  and  $\text{ess sup}_{t \in [0, T]} \mathcal{E}(\mu^0(t)) < \infty$ . Then the density  $\hat{c}^0$  is in  $L^p([0, T] \times \Omega)$  with  $p = \frac{d+1}{d} > 1$ .*

*Proof.* To apply Theorem 5.16, we first observe that the Lebesgue spaces form interpolation couples.

In our situation we have  $p_1 = 1, p_2 = \infty, X_1 = L^{\frac{d}{d-1}}(\Omega), X_2 = L^1(\Omega)$ . Hence,  $p_\theta = \frac{1}{1-\theta} > 1$ . Moreover,  $[X, Y]_\theta = [L^q(\Omega), L^1(\Omega)]_\theta \simeq L^{q_\theta}(\Omega)$ , where  $\frac{1}{q_\theta} = \frac{1-\theta}{q} + \frac{\theta}{1}$ . Setting  $p_\theta = q_\theta$ , we conclude  $\frac{1-\theta}{1-\theta} = q = \frac{d}{d-1}$ . Solving  $p_\theta = q_\theta$  for  $\theta$ , we obtain  $\theta = \frac{1}{1+d}$ , and hence  $p_\theta = q_\theta = \frac{d+1}{d} > 1$ . Summarizing, we conclude

$$\hat{c} \in L^\infty([0, T], L^1(\Omega)) \cap L^1([0, T], W^{1,1}(\Omega)) \subset L^{\frac{d+1}{d}}([0, T] \times \Omega). \quad (5.4)$$

□

**Remark 5.18.** In particular, if  $d = 2$ , then  $\hat{c} \in L^{3/2}([0, T] \times \Omega)$  and if  $d = 3$ , then  $\hat{c} \in L^{4/3}([0, T] \times \Omega)$ . Iterating the procedure, it is even possible to obtain  $\hat{c} \in L^{(d+2)/d}([0, T] \times \Omega)$  for  $d \geq 2$ .

Knowing integrability of  $\hat{c}$ , we get also better integrability of the fluxes  $\hat{J} \in L^{\tilde{p}}([0, T] \times \Omega)$  for some  $\tilde{p} > 1$ , which follows from the next lemma.

**Lemma 5.19.**

1 We have for all  $J \in \mathbb{R}^d$  and  $c > 0$  that

$$\frac{|J|^2}{c} + \frac{1}{p} c^p \geq \left(1 + \frac{1}{p}\right) |J|^{\frac{2p}{p+1}}.$$

2 Let  $\mu^0 \in L_w^\infty([0, T] \times \Omega, \mathbb{R}_{\geq 0}^2)$  such that  $\mathfrak{D}_0^V(\mu^0) < \infty$  and  $\text{ess sup}_{t \in [0, T]} \mathcal{E}(\mu^0(t)) < \infty$  and let  $\hat{J} \in \mathcal{M}([0, T] \times \Omega, \mathbb{R}^d)$  be the corresponding diffusion flux satisfying the continuity equation. Then  $\hat{J} \in L^{\tilde{p}}([0, T] \times \Omega, \mathbb{R}^d)$  for  $\tilde{p} = \frac{2d+2}{2d+1} > 1$ .

*Proof.* For proving the first part, let us define for fixed  $J \in \mathbb{R}^d$  the function  $F : ]0, \infty[ \rightarrow \mathbb{R}, F(c) := \frac{|J|^2}{c} + \frac{1}{p} c^p$ . Clearly,  $F \geq 0$  and  $F(c) \rightarrow \infty$  as  $c \rightarrow 0$  or  $c \rightarrow \infty$ . We compute the minimum. We have  $F'(c) = -|J|^2 c^{-2} + c^{p-1}$  and hence the critical point is at  $c_0 = |J|^{2/(p+1)}$ . Inserting  $c_0$  into  $F$  we get  $F(c) \geq F(c_0) = |J|^2 |J|^{-2/(p+1)} + \frac{1}{p} |J|^{2p/(p+1)} = (1 + \frac{1}{p}) |J|^{2p/(p+1)}$ , which proves the claim.

For second part, we use that by Lemma 5.17 we have  $\hat{c} \in L^p([0, T] \times \Omega)$  for  $p = \frac{d+1}{d}$ . This implies by the first part that  $\hat{J} \in L^{\tilde{p}}([0, T] \times \Omega, \mathbb{R}^d)$  for  $\tilde{p} = \frac{2 \frac{d+1}{d} + 2}{2 \frac{d+1}{d} + 1} = \frac{2d+2}{2d+1}$ . □



To obtain regularity for the whole reaction flux  $b_1 = -b_2$ , we have to get regularity also for  $\operatorname{div} \hat{J}$ . This is done by mollifying the solution  $(\hat{c}, \hat{J})$  of the continuity equation  $\hat{c} + \operatorname{div} \hat{J} = 0$  in time. We already now that  $\hat{c} : [0, T] \rightarrow \operatorname{Prob}(\Omega)$  is continuous by Lemma 5.11. With a slight abuse of notation, we denote by  $\hat{c}$  also a continuous continuation on  $\mathbb{R}$  such that  $\hat{c} \in L^p(\mathbb{R}, L^p(\Omega))$ . Now, we mollify in time and define  $\hat{c}^\epsilon(t) = \int_{\mathbb{R}} \hat{c}(s) \psi_\epsilon(t-s) ds$  where  $\psi_\epsilon$  is a positive and symmetric mollifier. Analogously, we define  $\hat{J}^\epsilon$  by convolution, i.e.  $\hat{J}^\epsilon(t) = \int_{\mathbb{R}} \hat{J}(s) \psi_\epsilon(t-s) ds$ . Since the continuity equation is linear, the smoothed functions  $(\hat{c}^\epsilon, \hat{J}^\epsilon)$  satisfy again the continuity equation with the same no-flux boundary conditions.

The next lemma shows that the dissipation functional can be approximated by mollifying  $(\hat{c}, \hat{J})$ . This basically uses the convexity of  $\hat{\mathfrak{D}}$  in  $(\hat{c}, \hat{J})$ .

**Lemma 5.20.** *Let  $\mu^0 \in L_w^\infty([0, T] \times \Omega, \mathbb{R}_{\geq 0}^2)$  such that the a priori bounds  $\mathfrak{D}_0^V(\mu^0) < \infty$  and  $\operatorname{ess\,sup}_{t \in [0, T]} \mathcal{E}(\mu^0(t)) < \infty$  hold. Let  $\hat{J} \in \mathcal{M}([0, T] \times \Omega, \mathbb{R}^d)$  be the corresponding diffusion flux satisfying the continuity equation. Let  $\psi^\epsilon : \mathbb{R} \rightarrow \mathbb{R}$  be a positive and symmetric mollifier. Define  $\hat{c}^\epsilon(t) = \int_{\mathbb{R}} \hat{c}(s) \psi_\epsilon(t-s) ds$  and  $\hat{J}^\epsilon(t) = \int_{\mathbb{R}} \hat{J}(s) \psi_\epsilon(t-s) ds$ . Then, we have the energy bound  $\sup_{\epsilon \in ]0, 1[} \operatorname{ess\,sup}_{t \in [0, T]} \mathcal{E}(\mu^\epsilon(t)) < \infty$  and*

$$\begin{aligned} & \int_0^T \left\{ \int_{\Omega} \sum_{j=1}^2 \tilde{\mathfrak{Q}}(\delta_j c_j^\epsilon, J_j^\epsilon) dx + \frac{1}{2} \int_{\Omega} \sum_{j=1}^2 \delta_j w_j^V \frac{|\nabla \rho_j^\epsilon|^2}{\rho_j^\epsilon} dx \right\} dt \\ & \rightarrow \int_0^T \left\{ \int_{\Omega} \sum_{j=1}^2 \tilde{\mathfrak{Q}}(\delta_j c_j^0, J_j^0) dx + \frac{1}{2} \int_{\Omega} \sum_{j=1}^2 \delta_j w_j^V \frac{|\nabla \rho_j^0|^2}{\rho_j^0} dx \right\} dt, \end{aligned} \quad (5.5)$$

which in particular, implies that  $\mathfrak{D}_\epsilon^V(\mu^\epsilon) \rightarrow \mathfrak{D}_0^V(\mu^0)$ .

*Proof.* The energy bound on  $\mu^\epsilon$  is trivially satisfied. The convergence for the dissipation functional (5.5) follows directly as the proof of Lemma 8.1.10 in [AGS05] since the integrand is convex in  $(\hat{c}, \hat{J})$ .  $\square$

With these preparations, we are able to show the remaining step in the proof of Theorem 5.13. The next proposition shows in analogy to Lemma 5.14 that the contribution by the reaction flux  $b_2^\epsilon$  to the dissipation functional  $\mathfrak{D}_\epsilon^V$  converges to zero as  $\epsilon \rightarrow 0$ .

**Proposition 5.21.** *Let  $\mu^0 \in L_w^\infty([0, T] \times \Omega, \mathbb{R}_{\geq 0}^2)$  satisfy the a priori bounds  $\mathfrak{D}_0^V(\mu^0) = \hat{\mathfrak{D}}_0(\hat{\mu}) < \infty$  and  $\operatorname{ess\,sup}_{t \in [0, T]} \mathcal{E}(\mu^0(t)) < \infty$ , and let  $\hat{J} \in \mathcal{M}([0, T] \times \Omega, \mathbb{R}^d)$  be the corresponding diffusion flux satisfying the continuity equation. Let  $\psi^\epsilon : \mathbb{R} \rightarrow \mathbb{R}$  be a positive and symmetric mollifier, which is specified below. Let  $\hat{c}^\epsilon, \hat{J}^\epsilon$  the mollified functions as in Lemma 5.20 and  $\hat{c}^{\epsilon, \gamma}, \hat{J}^{\epsilon, \gamma}$  the mollified and shifted functions as in Lemma 5.15.*

*Let  $\psi^\epsilon$  be such that  $\|\hat{c}^\epsilon\|_{L^{\tilde{p}}([0, T] \times \Omega)} \lesssim \frac{1}{\epsilon^\alpha}$  and  $\gamma \geq C\epsilon^{1-\lambda}$  for  $\epsilon \rightarrow 0$ , where  $\tilde{p}$  is the integrability exponent of the fluxes as in Lemma 5.19,  $C > 0$  is a positive constant and  $\lambda \in [0, 1[$ ,  $\alpha \in [0, 1[$  satisfies the inequality  $d + 1 \leq \frac{\lambda}{2\alpha}$ . Then, we have  $|\mathfrak{D}_\epsilon(\mu^{\epsilon, \gamma}) - \mathfrak{D}_0(\mu^{\epsilon, \gamma})| \rightarrow 0$ .*

*Proof.* First, we observe that for given  $\hat{c}$  such a mollifier and these constants  $\alpha, \lambda$  satisfying all the conditions can be easily constructed.

To prove the convergence, we follow the same strategy as in the proof of Lemma 5.14. Defining the reconstructed concentrations and fluxes as in (5.3), we observe that

$$|\mathfrak{D}_\epsilon(\mu^{\epsilon, \gamma}) - \mathfrak{D}_0(\mu^{\epsilon, \gamma})| \leq \int_0^T \int_{\Omega} \tilde{\mathfrak{C}} \left( \frac{\sqrt{c_1^{\epsilon, \gamma} c_2^{\epsilon, \gamma}}}{\epsilon}, b_2^{\epsilon, \gamma} \right) dx dt.$$

Using the bound from below on  $\hat{c}^\varepsilon$ , and the inequality  $\log(x+1) \leq C_{\tilde{p}} x^{\tilde{p}-1}$ , we get the estimate

$$\begin{aligned} \tilde{C} \left( \frac{\sqrt{c_1^{\varepsilon,\gamma} c_2^{\varepsilon,\gamma}}}{\varepsilon}, b_2^{\varepsilon,\gamma} \right) &\leq \tilde{C} (C\varepsilon^{-\lambda}, b_2^{\varepsilon,\gamma}) \leq C\varepsilon^{-\lambda} C (C^{-1}\varepsilon^\lambda b_2^{\varepsilon,\gamma}) \\ &\leq 2C\varepsilon^{-\lambda} |b_2^{\varepsilon,\gamma}| C^{-1}\varepsilon^\lambda \log(C^{-1}\varepsilon^\lambda |b_2^{\varepsilon,\gamma}| + 1) \\ &\leq 2C_{\tilde{p}} |b_2^{\varepsilon,\gamma}| C^{-\tilde{p}+1} \varepsilon^{\lambda(\tilde{p}-1)} |b_2^{\varepsilon,\gamma}|^{\tilde{p}-1} \leq \tilde{C} |b_2^{\varepsilon,\gamma}|^{\tilde{p}} \varepsilon^{\lambda(\tilde{p}-1)}, \end{aligned}$$

where  $\tilde{C} = \tilde{C}(C_{\tilde{p}}, C)$ . By  $\|\hat{c}^\varepsilon\|_{L^{\tilde{p}}([0,T] \times \Omega)} \lesssim \frac{1}{\varepsilon^\alpha}$ , we conclude that  $\|\operatorname{div} \hat{J}^\varepsilon\|_{\tilde{p}} \lesssim \frac{1}{\varepsilon^\alpha}$ , and by Lemma 5.19, we have  $\hat{J}^\varepsilon \in L^{\tilde{p}}([0, T] \times \Omega)$ . Together this implies that  $\|b_2^\varepsilon\|_{L^{\tilde{p}}([0,T] \times \Omega)} \lesssim \frac{1}{\varepsilon^\alpha}$ . Hence, we get

$$|\mathfrak{D}_\varepsilon(\mu^{\varepsilon,\gamma}) - \mathfrak{D}_0(\mu^{\varepsilon,\gamma})| \lesssim \varepsilon^{\lambda(\tilde{p}-1)} \varepsilon^{-\tilde{p}\alpha} = \varepsilon^{-\frac{1}{2d+1}\{(2d+2)\alpha-\lambda\}}.$$

Choosing,  $\lambda, \alpha \in [0, 1[$  such that  $d+1 \leq \frac{\lambda}{2\alpha}$ , we conclude that the right-hand side converges to zero, which proves the claim.  $\square$

## 6 Remarks for reaction-diffusion systems involving more species

In the last section, we comment on linear reaction-diffusion systems involving more species. The evolution equation for concentrations  $c \in \mathbb{R}_{\geq 0}^I$ ,  $I \in \mathbb{N}$  is given by

$$\dot{c} = \operatorname{diag}(\delta_1, \dots, \delta_I) \Delta c + A^\varepsilon c$$

where  $A^\varepsilon = A^S + \frac{1}{\varepsilon} A^F$  is a Markov generator (preserving positivity and total mass), which consists of a slow part and a fast part. The main assumption is the  $A^\varepsilon$  satisfies detailed balance with respect to its stationary measure  $w^\varepsilon$ . Similar to [MiS20, MPS20], we are going to assume that the stationary vector  $w^\varepsilon$  satisfies  $w^\varepsilon \rightarrow w^0$  as  $\varepsilon \rightarrow 0$  and that  $w^0 > 0$ . The positivity of the limit stationary measure  $w^0$  means that in the limit the evolution respects all concentrations  $c_i$  and is not degenerate.

The gradient structure is defined on the state space

$$Q = \operatorname{Prob}(\Omega \times \{1, \dots, I\}) := \{\mu = (\mu_1, \dots, \mu_I) \in \mathbb{R}^I : \mu_i \in \mathcal{M}(\Omega), \mu_i \geq 0, \mu_i(\Omega) = 1\}.$$

The driving energy functional  $\mathcal{E}_\varepsilon : X \rightarrow \mathbb{R}_\infty$  has the form

$$\mathcal{E}_\varepsilon(\mu) = \begin{cases} \int_\Omega \sum_{j=1}^I E_B \left( \frac{c_j}{w_j^\varepsilon} \right) w_j^\varepsilon dx, & \text{if } \mu = c dx \\ \infty, & \text{otherwise.} \end{cases}$$

and the dual dissipation potential splits into two parts

$$\begin{aligned} \mathcal{R}^*(\mu, \xi) &= \mathcal{R}_{\text{diff}}^*(\mu, \xi) + \mathcal{R}_{\text{react}}^*(\mu, \xi) \\ \mathcal{R}_{\text{diff}}^*(\mu, \xi) &= \frac{1}{2} \int_\Omega \sum_{j=1}^I \delta_j |\nabla \xi_j(x)|^2 d\mu_j, \quad \mathcal{R}_{\text{react}, \varepsilon}^*(\mu, \xi) = \int_\Omega \sum_{i < j} \kappa_{ij}^\varepsilon C^*(\xi_i(x) - \xi_j(x)) d\sqrt{\mu_i \mu_j}, \end{aligned}$$

where  $\kappa_{ij}^\varepsilon := A_{ij}^\varepsilon \left( \frac{w_j^\varepsilon}{w_i^\varepsilon} \right)^{1/2}$ . In particular, the reaction part of the dissipation potential splits into a fast part and a slow part

$$\begin{aligned} \mathcal{R}_{\text{react}, \varepsilon}^*(\mu, \xi) &= \mathcal{R}_{\text{slow}, \varepsilon}^*(\mu, \xi) + \frac{1}{\varepsilon} \mathcal{R}_{\text{fast}, \varepsilon}^*(\mu, \xi) \\ \mathcal{R}_{\text{xy}, \varepsilon}^*(\mu, \xi) &= \int_\Omega \sum_{i < j} \tilde{\kappa}_{ij}^\varepsilon C^*(\xi_i(x) - \xi_j(x)) d\sqrt{\mu_i \mu_j}, \quad \text{xy} \in \{\text{slow}, \text{fast}\}, \end{aligned}$$

where  $\tilde{\kappa}_{ij}^\varepsilon$  are bounded and positive uniformly in  $\varepsilon > 0$ . In particular, we call a reaction and its flux  $b_{ij}$  slow if  $A_{ij}^\varepsilon = O(1)$  and fast if  $A_{ij}^\varepsilon = O(\varepsilon^{-1})$ . Due to the detailed balance assumption and by  $w^0 > 0$ , the distinction between fast and slow reactions is indeed well-defined.

In the remainder of the section, we briefly explain how to generalize the proof of the EDP-convergence result also for this situation. Major differences occur at two stages, namely in 1) deriving compactness for slow reaction fluxes, 2) proving the limsup-estimate. The reaction fluxes of the fast reactions are not seen in the limit and have to be reconstructed in an analogous way as in the 2-species situation. Firstly, we explain the compactness result and the liminf-estimate. Secondly, we comment on the limsup-estimate.

## 6.1 Compactness for slow reaction fluxes and liminf-estimate

Here, we comment on proving compactness and the liminf-estimate for the multi species case. In comparison to the previous situation, compactness for the concentrations, by using strong compactness for coarse-grained variables and convergence towards the slow manifold, can be derived (cf. Lemmas 5.6, 5.10). Moreover, compactness of diffusion fluxes and spatial regularity follows, too (cf. Lemma 5.19).

In contrast to the situation of two species connected with one fast reaction, where no slow reaction fluxes exists, compactness for slow reaction fluxes  $b_{ij}^\varepsilon$  has to be derived in the multi species case. This follows immediately from Lemma 5.2, once compactness of  $\sqrt{c_i^\varepsilon c_j^\varepsilon}$  is obtained. At this point it is clear that weak convergence of  $c^\varepsilon \rightharpoonup c^0$  is not sufficient. Instead the previously derived strong convergence of  $c^\varepsilon \rightarrow c^0$  implies by dominated convergence also strong convergence of  $\sqrt{c_i^\varepsilon c_j^\varepsilon} \rightarrow \sqrt{c_i^0 c_j^0}$  in  $L^1([0, T] \times \Omega)$ , and hence, compactness for the slow fluxes  $b_{ij}^\varepsilon$ . Compactness for fast reaction fluxes can not be obtained as already mentioned in Remark 5.5. Having proved compactness, the proof of the liminf-estimate is exactly the same as for Theorem 5.12, since the functional  $\mathcal{D}_\varepsilon$  is jointly convex in all variables  $(c, J, b)$ .

## 6.2 Equilibration and reconstruction of reaction fluxes und recovery sequence

A crucial observation throughout the proof of the  $\Gamma$ -convergence was Lemma 4.4, which provides an equilibration of fluxes assuming microscopic equilibria for the concentrations. In Lemma 4.5, we derived equilibration for the diffusion fluxes. Similarly, also an equilibration of the slow reaction fluxes can be derived. In [MiS20] a general operator-theoretic coarse-graining and reconstruction procedure has been developed. This method can also be applied to derive coarse-grained fluxes and a coarse-grained continuity equation, see [Ste21]. Importantly for us, as in (5.3) the reconstructed slow reaction fluxes depend linearly on the coarse-grained reaction fluxes. The fast reaction fluxes are then of the form

$$b_{ij} = a_1 \operatorname{div} \hat{J}_i + a_2 \hat{J}_i + \sum_{j=3}^{k(i)} a_{ij} \hat{b}_{ij},$$

where all functions  $a_j$  are  $C^0(\Omega, \mathbb{R}^{k_j})$ , where  $k_1 = 1$ ,  $k_2 = d$  and  $k_{ij} = 1$ .

In order to prove that the constant sequence for smooth and positive concentrations is indeed a recovery sequence, we follow the same reasoning as in the Lemmas 5.14 and 5.21. The only difference comes from the explicit depends on the coarse-grained reaction flux  $\hat{b}_{ij}$ . Using the bound on the limit dissipation functional (which provides bounds on  $\int_0^T \int_\Omega \tilde{C}(\sqrt{\hat{c}_i \hat{c}_j}, \hat{b}_{ij}) dx dt$ ) and the next Lemma 6.1,

we obtain that  $\hat{b}_{ij} \in L^C([0, T] \times \Omega)$  (we refer to [FrM21] for a proof). Since  $L^C$  is an Orlicz-space, we conclude that the reconstructed fluxes  $b_i$  are in  $L^C$ . This allows to proceed as in Lemma 5.21 and proves the existence of a recovery sequence.

**Lemma 6.1** ([FrM21]). *Let  $p > 1$ . Then, for all  $a \geq 0$  and  $B \in \mathbb{R}$  we have*

$$\tilde{C}(a, B) \geq \left(1 - \frac{1}{p}\right) \tilde{C}(B) - \frac{2}{p} a^p.$$

*In particular, setting  $a = \sqrt{c_i c_j}$  and  $B = b_{ij}$  we have*

$$\int_0^T \int_{\Omega} \tilde{C}(\sqrt{c_i c_j}, b_{ij}) dx dt + \|\sqrt{c_i c_j}\|_{L^p([0, T] \times \Omega)}^p \gtrsim \int_0^T \int_{\Omega} \tilde{C}(b_{ij}) dx dt,$$

*which proves that  $b_{ij} \in L^C([0, T] \times \Omega)$  if  $c_i, c_j \in L^p([0, T] \times \Omega)$  for some  $p > 1$ .*

**Acknowledgement:** The research was supported by Deutsche Forschungsgemeinschaft (DFG) through the Collaborative Research Center SFB 1114 “Scaling Cascades in Complex Systems” (Project no. 235221301), subproject C05 “Effective models for materials and interfaces with multiple scales”. The author is grateful to Alexander Mielke for many helpful discussions and for having proposed the problem.

## References

- [AGS05] L. AMBROSIO, N. GIGLI, and G. SAVARÉ. *Gradient flows in metric spaces and in the space of probability measures*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 2005.
- [AM\*12] S. ARNRICH, A. MIELKE, M. A. PELETIER, G. SAVARÉ, and M. VENERONI. Passing to the limit in a Wasserstein gradient flow: from diffusion to reaction. *Calc. Var. Part. Diff. Eqns.*, 44, 419–454, 2012.
- [BaP86] V. BARBU and T. PRECUPANU. *Convexity and optimization in Banach spaces*. D. Reidel Publishing Co., Dordrecht, third edition, 1986.
- [BeB00] J.-D. BENAMOU and Y. BRENIER. A computational fluid mechanics solution to the Monge-Kantorovich mass transfer problem. *Numer. Math.*, 84(3), 375–393, 2000.
- [BeL76] J. BERGH and J. LÖFSTRÖM. *Interpolation spaces - an introduction*. Springer-Verlag, 1976.
- [BoH02] D. BOTHE and D. HILHORST. A reaction-diffusion system with fast reversible reaction. *J. Math. Anal. Appl.*, 2002.
- [BoP10] D. BOTHE and M. PIERRE. Quasi-steady-state approximation for a reaction-diffusion system with fast intermediate. *J. Math. Anal. Appl.*, 368(1), 120–132, 2010.
- [BoP11] D. BOTHE and M. PIERRE. The instantaneous limit for reaction-diffusion systems with a fast irreversible reaction. *Discr. Cont. Dynam. Systems Ser. S*, 8(1), 49–59, 2011.
- [Bot03] D. BOTHE. Instantaneous limits of reversible chemical reactions in presence of macroscopic convection. *J. Differential Equations*, 193(1), 27–48, 2003.
- [BPR12] D. BOTHE, M. PIERRE, and G. ROLLAND. Cross-diffusion limit for a reaction-diffusion system with fast reversible reaction. *Comm. Partial Differential Equations*, 37(11), 1940–1966, 2012.
- [Bra13] A. BRAIDES. *Local minimization, Variational Evolution and Gamma-convergence*. Lect. Notes Math. Vol. 2094. Springer, 2013.
- [Dal93] G. DAL MASO. *An Introduction to  $\Gamma$ -Convergence*. Birkhäuser Boston Inc., Boston, MA, 1993.
- [DDJ20] E. S. DAUS, L. DESVILLETES, and A. JÜNGEL. Cross-diffusion systems and fast-reaction limits. *Bull. Sci. math.*, 159(102824), 2020.

- [DFM18] P. DONDL, T. FRENZEL, and A. MIELKE. A gradient system with a wiggly energy and relaxed EDP-convergence. *ESAIM Control Optim. Calc. Var.*, 2018. To appear. WIAS preprint 2459.
- [Edw11] D. EDWARDS. On the kantorovich-rubinstein theorem. *Exposition. Math.*, 29, 387–398, 2011.
- [EKT76] I. EKELAND and R. TEMAM. *Convex Analysis and Variational Problems*. North Holland, 1976.
- [Eva80] L. C. EVANS. A convergence theorem for a chemical diffusion-reaction system. *Houston Journal of Mathematics*, 6(2), 259–267, 1980.
- [FrL19] T. FRENZEL and M. LIERO. Effective diffusion in thin structures via generalized gradient systems and EDP-convergence. *WIAS Preprint 2601*, 2019.
- [FrM21] T. FRENZEL and A. MIELKE. Deriving the kinetic relation for the flux through a membrane via EDP-convergence. *in preparation*, 2021.
- [GK\*20] P. GLADBACH, E. KOPFER, J. MAAS, and L. PORTINALE. Homogenisation of one-dimensional discrete optimal transport. *Journal de Mathématiques Pures et Appliquées*, 139(7), 204–234, 2020.
- [HPR18] M. HEIDA, R. I. A. PATTERSON, and D. R. M. RENGER. Topologies and measures on the space of functions of bounded variation taking values in a Banach or metric space. *J. Evol. Eqns.*, 19, 111–152, 2018.
- [HvdHP00] D. HILHORST, R. VAN DER HOUT, and L. PELETIER. Nonlinear diffusion in the presence of fast reaction. *Nonlinear Analysis*, 41, 803–823, 2000.
- [JKO98] R. JORDAN, D. KINDERLEHRER, and F. OTTO. The variational formulation of the Fokker-Planck equation. *SIAM J. Math. Analysis*, 29(1), 1–17, 1998.
- [LiM13] M. LIERO and A. MIELKE. Gradient structures and geodesic convexity for reaction-diffusion systems. *Phil. Trans. Royal Soc. A*, 371(2005), 20120346, 28, 2013.
- [LM\*17] M. LIERO, A. MIELKE, M. A. PELETIER, and D. R. M. RENGER. On microscopic origins of generalized gradient structures. *Discr. Cont. Dynam. Systems Ser. S*, 10(1), 1–35, 2017.
- [Man07] S. MANIGLIA. Probabilistic representation and uniqueness results for measure-valued solutions of transport equations. *J. Math. Pures Appl. (9)*, 87(6), 601–626, 2007.
- [Mar15] R. MARCELIN. Contribution a l'étude de la cinétique physico-chimique. *Annales de Physique*, III, 120–231, 1915.
- [Mie11] A. MIELKE. A gradient structure for reaction-diffusion systems and for energy-drift-diffusion systems. *Nonlinearity*, 24, 1329–1346, 2011.
- [Mie16] A. MIELKE. On evolutionary  $\Gamma$ -convergence for gradient systems (Ch. 3). In A. Muntean, J. Rademacher, and A. Zagaris, editors, *Macroscopic and Large Scale Phenomena: Coarse Graining, Mean Field Limits and Ergodicity*, Lecture Notes in Applied Math. Mechanics Vol. 3, pages 187–249. Springer, 2016. Proc. of Summer School in Twente University, June 2012.
- [MiS20] A. MIELKE and A. STEPHAN. Coarse-graining via EDP-convergence for linear fast-slow reaction systems. *M3AS: Math. Models Meth. Appl. Sci.*, 2020.
- [MMP20] A. MIELKE, A. MONTEFUSCO, and M. A. PELETIER. Exploring families of energy-dissipation landscapes via tilting — three types of EDP convergence. *WIAS preprint 2668*, 2020.
- [MP\*17] A. MIELKE, R. I. A. PATTERSON, M. A. PELETIER, and D. R. M. RENGER. Non-equilibrium thermodynamical principles for chemical reactions with mass-action kinetics. *SIAM J. Appl. Math.*, 77(4), 1562–1585, 2017.
- [MPR14] A. MIELKE, M. A. PELETIER, and D. R. M. RENGER. On the relation between gradient flows and the large-deviation principle, with applications to Markov chains and diffusion. *Potential Analysis*, 41(4), 1293–1327, 2014.
- [MPS20] A. MIELKE, M. A. PELETIER, and A. STEPHAN. EDP convergence for nonlinear fast slow reaction systems with detailed balance. *arXiv:2010.08458*, 2020.
- [MRS13] A. MIELKE, R. ROSSI, and G. SAVARÉ. Nonsmooth analysis of doubly nonlinear evolution equations. *Calc. Var. Part. Diff. Eqns.*, 46(1-2), 253–310, 2013.
- [MuN11] H. MURAKAWA and H. NINOMIYA. Fast reaction limit of a three-component reaction-diffusion system. *J. Math. Anal. Appl.*, 379(1), 150–170, 2011.
- [Ott01] F. OTTO. The geometry of dissipative evolution equations: the porous medium equation. *Comm. Partial Differential Equations*, 26, 101–174, 2001.
- [PeR20] M. A. PELETIER and M. RENGER. Fast reaction limits via  $\gamma$ -convergence of the flux rate functional. *arXiv 2009.14546*, 2020.
- [PR\*20] M. A. PELETIER, R. ROSSI, G. SAVARÉ, and O. TSE. Jump processes as generalized gradient flows. *arXiv:2006.10624*, 2020.

- [Ren18] D. R. M. RENGER. Gradient and GENERIC systems in the space of fluxes, applied to reacting particle systems. *Entropy*, 20(8), 2018.
- [Ste19] A. STEPHAN. On EDP-convergence for gradient systems with different time scales. *Proc. Appl. Math. Mech.*, 2019.
- [Ste21] A. STEPHAN. Coarse-graining and reconstruction for markov operators. *in preparation*, 2021.