

## **Infinite-dimensional diffusions and depletion interaction for a model of colloids**

Myriam Fradon<sup>1</sup>, Alexander Zass<sup>2</sup>

submitted: March 3, 2025

<sup>1</sup> Université de Lille  
CNRS UMR 8524  
Laboratoire P. Painlevé  
Bât. M2  
59655 Villeneuve d'Ascq  
France  
E-Mail: myriam.fradon@univ-lille.fr

<sup>2</sup> Weierstrass Institute  
Mohrenstr. 39  
10117 Berlin  
Germany  
E-Mail: alexander.zass@wias-berlin.de

No. 3181  
Berlin 2025



---

2020 *Mathematics Subject Classification.* 60K35, 60H10, 60J55, 82D99.

*Key words and phrases.* Stochastic differential equation, hard core interaction, reversible measure, collision local time, colloids, depletion interaction, Gibbs point process.

Edited by  
Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS)  
Leibniz-Institut im Forschungsverbund Berlin e. V.  
Mohrenstraße 39  
10117 Berlin  
Germany

Fax: +49 30 20372-303  
E-Mail: [preprint@wias-berlin.de](mailto:preprint@wias-berlin.de)  
World Wide Web: <http://www.wias-berlin.de/>

# Infinite-dimensional diffusions and depletion interaction for a model of colloids

Myriam Fradon, Alexander Zass

## Abstract

We consider infinite-dimensional random diffusion dynamics for the Asakura–Oosawa model of interacting hard spheres of two different sizes. We construct a solution to the corresponding SDE with collision local times, analyse its reversible measures, and observe the emergence of an attractive short-range depletion interaction between the large spheres. We study the Gibbs measures associated to this new interaction, exploring connections to percolation and optimal packing.

## 1 Introduction

In this work, we consider colloidal dynamics, that is the motion of large colloidal particles within a medium formed by much smaller solvent particles. The model is inspired by the Asakura–Oosawa (AO) model [AO54] of Chemical Physics; there, the authors introduce a size-asymmetric binary mixture of hard spheres to model colloidal suspensions, successfully explaining the origin of an attractive interaction that is observed when looking at large colloidal particles in a solution of non-adsorbing polymers, and which is referred to as “attraction through repulsion” [LT11]. We consider a modification of the model, due to A. Vrij [Vri76], in which the colloids are represented as large hard spheres, and the polymers as small penetrable spheres, which we will call particles. We present here a two-fold study of this model – both from the dynamic (infinite-dimensional SDEs) point of view, and from the static (infinite-volume Gibbs point processes) one.

Indeed, already in the early 20th century, J. Perrin observed that these objects are characterised by a Brownian-type motion [Per09]. To preserve shift invariance, we are therefore modelling infinitely-many hard spheres in Euclidean space, randomly-diffusing through a random medium of infinitely-many randomly-diffusing very small particles. Since spheres cannot overlap, and spheres and particles do not overlap either, the model is an infinite-dimensional SDE with normal reflection on the boundary of the set of forbidden, overlapping configurations.

Since the pioneering work of A. V. Skorokhod [Sko61, Sko62] on half-spaces, pathwise solutions of reflected stochastic differential equations have been constructed on finite-dimensional domains under boundary conditions of increasing generality: see H. Tanaka [Tan79] for convex domains, P. L. Lions and A. S. Sznitman [LS84] for admissible domains, H. Saisho [Sai87] for domains satisfying exterior sphere and interior cone conditions, P. Dupuis and H. Ishii [DI93, DI08] for non-smooth domains, and [Fra13] for domains defined by a finite set of constraints. On the other hand, infinite-dimensional Brownian diffusions, without boundary condition, have been first studied by R. Lang [Lan77] and H.-O. Georgii [Geo79]. Our present work combines the difficulties of infinite-dimensional Brownian systems and those induced by the reflection on the non-smooth boundary of the domain of non-overlapping configurations.

H. Tanemura [Tan96] first used such an infinite-dimensional reflected SDE to study a system of infinitely-many identical Brownian hard balls. His results have been extended to identical hard balls that also have an additional smooth interaction with potentially-infinite range (see [FR00, FRT00, FR07, Tan22]). Recently, we extended these ideas in [FKRZ24] to two-type hard-core diffusions, in order to construct dynamics for the Asakura–Oosawa model. However, the results therein were limited to a finite number of large hard spheres in a bath of a infinitely-many particles; as a result, the model in [FKRZ24], though isotropic, does not have the spatial invariance which can be expected from a colloidal suspension at such a microscopic scale.

The aim of the present paper is then to construct a more natural, shift-invariant Asakura–Oosawa dynamics, where both the hard spheres and the particles are infinitely many. This natural generalisation is clearly not straightforward: since spheres, contrary to particles, are not penetrable, an infinite number of spheres induces infinite-range non-overlap conditions; this requires precise estimates (namely, to control so-called *bad paths*) which lead to new, highly non-trivial, technical difficulties.

Colloids and the phenomenon of depletion interaction have recently been investigated using the tools and language of Statistical Mechanics. For example: in [JT20], the authors use the emergence of the effective depletion interaction to show that cluster expansion converges in an improved activity regime; in [JKSZ24], the Gibbs variational principle is shown to hold also when the size of hard spheres is unbounded; in [WJL23], the authors analyse geometric criteria for the absence of multi-body depletion interactions. In this framework, our work shows how the Gibbs measures at the heart of the above studies come up naturally as the reversible equilibrium measures of the colloid-polymer dynamics.

The core of the paper is dedicated to the proof of existence of a unique strong solution to the two-type infinite-dimensional SDE that is introduced in Section 1.2: in Section 2, we construct a sequence of approximating processes via penalisation, and in Section 3, we prove that such a sequence converges to the unique solution of the SDE. In Section 4, we analyse the emergence of the depletion interaction. More precisely, we consider the projection of the reversible measures onto the subsystem of hard spheres. We prove a correspondence between the two-type and the one-type Gibbs measures, and present an infinite-dimensional gradient diffusion associated to these measures. Finally, we show that in the low-intensity regime there is absence of percolation, whereas the high-intensity regime leads to the phenomenon of optimal packing.

## 1.1 Allowed configurations of spheres and particles

We consider an infinite number of large spheres with radius  $\hat{r} > 0$  and centres  $\hat{x}_1, \hat{x}_2, \dots$  in  $\mathbb{R}^d$ ,  $d \geq 2$ , along with an infinite number of smaller spherical particles with radius  $0 < \hat{r} < \hat{r}$  and centres  $\hat{x}_1, \hat{x}_2, \dots$  in  $\mathbb{R}^d$ . This two-type system is subject to the following *non-overlap constraints*: the large spheres are not allowed to overlap; the small particles can overlap each other, but are not allowed to overlap the large ones.

The configuration space is the set  $\mathcal{M}$  of locally-finite Radon point measures on  $\mathbb{X} = \mathbb{R}^d \times \{\circ, \cdot\}$ , i.e. those of the form  $\mathbf{x} = \hat{\mathbf{x}}\hat{\mathbf{x}} = \sum_{i \in I} \delta_{\hat{x}_i} + \sum_{k \in K} \delta_{\hat{x}_k}$ , with  $\hat{x}_i \in \mathbb{R}^d \times \{\circ\}$ ,  $\hat{x}_k \in \mathbb{R}^d \times \{\cdot\}$  for  $I, K \subset \mathbb{N}^* = \mathbb{N} \setminus \{0\} = \{1, 2, \dots\}$ , such that  $\mathbf{x}(\Lambda) < +\infty$  for any compact subset  $\Lambda \subset \mathbb{X}$ .  $\mathcal{M}$  is endowed with the topology of vague convergence.

The sum of two point measures is denoted by the juxtaposition  $\mathbf{xy} := \mathbf{x} + \mathbf{y}$ . As the point measures we consider are a.s. simple, we identify them with their support. For any  $\Lambda \subset \mathbb{R}^d$ ,  $\mathbf{y}_\Lambda := \mathbf{y} \cap \Lambda$  is the restriction of the sphere and particle configuration  $\mathbf{y}$  to  $\Lambda$ . We denote by  $\mathcal{M}_\Lambda$  the corresponding set of point measures on  $\Lambda \times \{\circ, \cdot\}$  and by  $\mu_\Lambda$  the restriction to  $\mathcal{M}_\Lambda$  of a measure  $\mu$  on  $\mathcal{M}$ . The above

notations are, of course, analogous when considering only sphere or particle configurations, leading to  $\mathring{\mathcal{M}}_\Lambda$  and  $\mathring{\mathcal{M}}_\Lambda$ , respectively.

According to the non-overlap constraints, the subset  $\mathcal{D} \subset \mathcal{M}$  of so-called *allowed configurations* is

$$\mathcal{D} = \left\{ \mathbf{x} = \dot{\mathbf{x}}\dot{\mathbf{x}} \in \mathcal{M} : \begin{array}{l} \forall i \neq j, \quad |\dot{x}_i - \dot{x}_j| \geq 2\dot{r}, \\ \forall i, k, \quad |\dot{x}_i - \dot{x}_k| \geq \dot{r} + \dot{r} \end{array} \right\}.$$

The second constraint describes a *depletion* shell of thickness  $\dot{r}$  around each sphere, that is forbidden for the centres of the small particles. Given an allowed configuration  $\dot{\mathbf{x}}$  of hard spheres, in order for the configuration  $\dot{\mathbf{x}}\dot{\mathbf{x}}$  to also be allowed, the particles  $\dot{\mathbf{x}}$  cannot be placed within the forbidden area  $\mathbb{B}(\dot{\mathbf{x}}) := \bigcup_{\dot{x} \in \dot{\mathbf{x}}} B(\dot{x}, \dot{r}^\circ)$ , where  $\dot{r}^\circ := \dot{r} + \dot{r}$  is the depletion radius.

Here and in the sequel,  $B(x, R)$  denotes the open ball with centre  $x$  and radius  $R$ ,  $|A|$  is the  $d$ -dimensional volume of  $A \subset \mathbb{R}^d$ , and  $v_d := |B(0, 1)|$  the volume of the  $d$ -dimensional unit sphere.

## 1.2 Dynamics for the AO model

Having fixed notations, we can now introduce our dynamical model. Fix a probability space  $(\Omega, \mathcal{F}, P)$ ; the system is described as follows:

- Infinitely-many *hard spheres* with radius  $\dot{r} > 0$ , whose centres at time  $t$  are denoted by  $(\dot{X}_i(t))_{i \in \mathbb{N}^*}$ , move according to independent Brownian motions  $(\dot{W}_i)_{i \in \mathbb{N}^*}$ .
- The hard spheres evolve in a time-inhomogeneous *random medium* consisting of infinitely-many *small particles* with radius  $\dot{r} \in (0, \dot{r})$ , whose centres at time  $t$  are denoted by  $(\dot{X}_k(t))_{k \in \mathbb{N}^*}$ , themselves moving according to independent Brownian motions  $(\dot{W}_k)_{k \in \mathbb{N}^*}$ .
- The only interactions are due to the non-overlap constraints. The *local times*  $(L_{ij})_{i, j \in \mathbb{N}^*}$  ensure that there is no pairwise overlap between the hard spheres: in case of a collision, they induce an instantaneous repulsion given by a normal reflection at the boundary of the set of allowed configurations. Similarly, the local times  $(\ell_{ik})_{i, k \in \mathbb{N}^*}$  model the non-overlap constraint between small particles and hard spheres. As a result, at each time, the two-type configuration should belong to the set  $\mathcal{D}$  of allowed configurations.

This two-type dynamics can be described by the following doubly-infinite SDE with reflection. We are only interested in a solution on  $[0, 1]$ ; Markov property then yields a solution on  $\mathbb{R}_+$ .

$$\left\{ \begin{array}{l} \text{for any } i \in \mathbb{N}^*, k \in \mathbb{N}^*, t \in [0, 1], \\ d\dot{X}_i(t) = d\dot{W}_i(t) + \sum_{j=1}^{+\infty} (\dot{X}_i - \dot{X}_j)(t) dL_{ij}(t) + \sum_{k=1}^{+\infty} (\dot{X}_i - \dot{X}_k)(t) d\ell_{ik}(t) \\ d\dot{X}_k(t) = \dot{\sigma} d\dot{W}_k(t) + \dot{\sigma}^2 \sum_{j=1}^{+\infty} (\dot{X}_k - \dot{X}_j)(t) d\ell_{ik}(t) \\ \forall j \in \mathbb{N}^*, L_{ij}(0) = \ell_{ik}(0) = 0, L_{ij} = L_{ji}, L_{ii} \equiv 0, \\ \int_0^t \mathbb{1}_{|\dot{X}_i(s) - \dot{X}_j(s)| \neq 2\dot{r}} dL_{ij}(s) = 0, \int_0^t \mathbb{1}_{|\dot{X}_i(s) - \dot{X}_k(s)| \neq \dot{r}^\circ} d\ell_{ik}(s) = 0. \end{array} \right. \quad (\mathcal{S}_\infty)$$

The *diffusion coefficient*  $\dot{\sigma} > 0$  is fixed, related to the temperature and to the mass of the small particles. We do not introduce a corresponding coefficient  $\dot{\sigma}$  for the hard spheres, i.e. we simply set

it equal to 1; this is not a restriction, as it suffices to choose, as the time unit of the model, the time at which the standard deviation of each coordinate of the Brownian motions  $\dot{W}_i$  is equal to the length unit of the model.

### 1.3 Main results

The following set of Gibbs measures will be the reversible measures for the above system.

**Definition 1.1.** For a probability measure  $\mu$  on  $\mathcal{M}$ , we write  $\mu \in \mathcal{G}_{\dot{z}, \dot{z}}$  and say that it is a *Gibbs measure* on  $\mathcal{M}$ , with activities  $\dot{z}, \dot{z}$ , associated to the above hard-core interaction, if, for any bounded subset  $\Lambda \subset \mathbb{R}^d$  and any positive measurable function  $F$  on  $\mathcal{D}$ ,

$$\int_{\mathcal{D}} F(\mathbf{x}) \mu(d\mathbf{x}) := \int_{\mathcal{D}} \frac{1}{Z_{\Lambda}(\mathbf{y})} \int_{\dot{\mathcal{M}}} \int_{\dot{\mathcal{M}}} F(\mathbf{y}_{\Lambda^c} \dot{\mathbf{x}} \dot{\mathbf{x}}) \mathbb{1}_{\mathcal{D}}(\mathbf{y}_{\Lambda^c} \dot{\mathbf{x}} \dot{\mathbf{x}}) \pi_{\Lambda}^{\dot{z}}(d\dot{\mathbf{x}}) \pi_{\Lambda}^{\dot{z}}(d\dot{\mathbf{x}}) \mu(d\mathbf{y}), \quad (1.1)$$

where  $\pi_{\Lambda}^{\dot{z}}(d\dot{\mathbf{x}})$  denotes the Poisson point process on  $\dot{\mathcal{M}}_{\Lambda}$  with intensity  $\dot{z} > 0$ ,  $\pi_{\Lambda}^{\dot{z}}(d\dot{\mathbf{x}})$  denotes the Poisson point process on  $\dot{\mathcal{M}}_{\Lambda}$  with intensity  $\dot{z} > 0$ , and the normalisation factor  $Z_{\Lambda}(\mathbf{y}) = Z_{\Lambda}(\mathbf{y}_{\Lambda^c}) = \iint \mathbb{1}_{\mathcal{D}}(\mathbf{y}_{\Lambda^c} \dot{\mathbf{x}} \dot{\mathbf{x}}) \pi_{\Lambda}^{\dot{z}}(d\dot{\mathbf{x}}) \pi_{\Lambda}^{\dot{z}}(d\dot{\mathbf{x}})$  is the partition function.

Note that  $\mu \in \mathcal{G}_{\dot{z}, \dot{z}}$  is concentrated on the set  $\mathcal{D}$  of admissible configuration. Also note that  $Z_{\Lambda}(\mathbf{y}) \leq 1$ , since  $\mathbb{1}_{\mathcal{D}} \leq 1$ , and  $Z_{\Lambda}(\mathbf{y}) > 0$ , since  $\mathbf{y} \in \mathcal{D}$  and  $\dot{\mathbf{x}} \dot{\mathbf{x}}$  is a finite configuration.

While existence of at least one such measure is a classical result (see, e.g. [Rue99]), it is conjectured [LT11, Chapter 3] that a non-uniqueness phase transition, that is  $\text{Card}(\mathcal{G}_{\dot{z}, \dot{z}}) > 1$ , occurs for large enough hard-sphere activities.

A main result of this paper is the following theorem, whose proof is split in Sections 2 and 3.

**Theorem 1.2.** Let  $\mu \in \mathcal{G}_{\dot{z}, \dot{z}}$ ,  $\dot{z}, \dot{z} > 0$ . The two-type infinite-dimensional SDE  $(\mathcal{S}_{\infty})$  admits a unique  $\mathcal{D}$ -valued strong solution, for  $\mu$ -almost every deterministic initial condition. Moreover, this solution is time-reversible if the initial condition is random and distributed according to  $\mu$ .

In Section 4, we investigate the emergence of an effective depletion interaction. The main results can be summarised in the following two theorems.

**Theorem 1.3.** The sphere marginal probability measure  $\dot{\mu}$  of  $\mu \in \mathcal{G}_{\dot{z}, \dot{z}}$  is an element of  $\mathcal{G}_{\dot{z}}(\dot{z}\mathcal{E})$ , i.e. it solves, for any bounded  $\Lambda \subset \mathbb{R}^d$  and any positive measurable  $F$  on  $\dot{\mathcal{M}}$ ,

$$\int_{\mathcal{D}} F(\dot{\mathbf{x}}) \dot{\mu}(d\dot{\mathbf{x}}) = \int_{\mathcal{D}} \frac{1}{Z_{\Lambda}(\dot{\mathbf{y}})} \int_{\dot{\mathcal{M}}} F(\dot{\mathbf{x}}_{\Lambda^c}) \exp(-\dot{z}\mathcal{E}_{\Lambda}(\dot{\mathbf{x}}_{\Lambda^c})) \mathbb{1}_{\mathcal{D}}(\dot{\mathbf{y}}_{\Lambda^c} \dot{\mathbf{x}}) \pi_{\Lambda}^{\dot{z}}(d\dot{\mathbf{x}}) \dot{\mu}(d\dot{\mathbf{y}}), \quad (1.2)$$

where  $\mathcal{E}_{\Lambda}(\dot{\mathbf{x}}) := |\mathbb{B}(\dot{\mathbf{x}}_{\Lambda}) \setminus \mathbb{B}(\dot{\mathbf{x}}_{\Lambda^c})|$ . Moreover, there exists a critical activity  $\dot{z}_c > 0$  such that, for any  $\dot{z} < \dot{z}_c$ , there is no percolation of interacting hard spheres  $\dot{\mu}$ -almost surely; and  $\dot{\mu}$  attains the closest packing density as  $\dot{z} \rightarrow \infty$ , for any  $\dot{z} > 0$ .

**Theorem 1.4.** Let  $\mathcal{E}(\dot{\mathbf{x}}) := |\mathbb{B}(\dot{\mathbf{x}})|$ . For  $d \geq 3$  and  $\dot{r}/\dot{r} \leq \frac{2}{3}\sqrt{3} - 1$ , the following gradient dynamics admits a unique strong solution, and  $\dot{\mu} \in \mathcal{G}_{\dot{z}}(\dot{z}\mathcal{E})$  is reversible for this dynamics for any  $\dot{z}, \dot{z} > 0$ :

$$\left\{ \begin{array}{l} \text{for any } i \in \mathbb{N}^*, t \in [0, 1], \\ d\dot{X}_i(t) = d\dot{W}_i(t) - \frac{\dot{z}}{2} \nabla_i \mathcal{E}(\dot{X}_1, \dot{X}_2, \dots) dt + \sum_{j=1}^{+\infty} (\dot{X}_i - \dot{X}_j)(t) dL_{ij}(t), \\ \forall j \in \mathbb{N}^*, L_{ij}(0) = 0, L_{ij} = L_{ji}, \int_0^t \mathbb{1}_{|\dot{X}_i(s) - \dot{X}_j(s)| \neq 2\dot{r}} dL_{ij}(s) = 0, L_{ii} \equiv 0. \end{array} \right.$$

Note that the solutions of the above theorems are unique in the sense of [Tan96, Lemma 5.4], that is as elements of a set of regular paths, see Remark 3.9.

## 2 Construction of the approximating process

Let  $\mu \in \mathcal{G}_{\dot{z}, \dot{z}}$ ,  $\dot{z}, \dot{z} > 0$ . In this section, for  $\mu$ -almost every initial condition  $\mathbf{y} \in \mathcal{D}$ , we construct, via a penalisation procedure, a sequence of approximating processes that start from an approximation  $\mu_R^{\mathbf{y}}$  of the candidate reversible measure for  $(\mathcal{S}_\infty)$ , and that mostly stay in  $B(0, R)$ . The solution of  $(\mathcal{S}_\infty)$  will then be shown, in Section 3, to be the limit as  $R \rightarrow \infty$  of such a sequence.

### 2.1 Penalisation functions

For every fixed radius  $R > 0$  and outside configuration  $\mathbf{y} = \dot{\mathbf{y}}\dot{\mathbf{y}} \in \mathcal{D}$ , we define  $\mathring{B}_R^{\mathbf{y}}$  (resp.  $\mathring{B}_R^{\dot{\mathbf{y}}}$ ) as the part of  $B(0, R)$  in which the centre of a sphere (resp. a particle) can be put without conflicting with the  $\mathbf{y}$  spheres and particles whose centres are outside of  $B(0, R)$ , that is:

$$\mathring{B}_R^{\mathbf{y}} := B(0, R) \setminus B(\dot{\mathbf{y}}_{B(0,R)^c}, 2\dot{r}) \setminus B(\dot{\mathbf{y}}_{B(0,R)^c}, \dot{r}), \quad \mathring{B}_R^{\dot{\mathbf{y}}} := B(0, R) \setminus \mathbb{B}(\dot{\mathbf{y}}_{B(0,R)^c}). \quad (2.1)$$

The sphere-penalisation function  $\mathring{\psi}_R^{\mathbf{y}} : \mathbb{R}^d \rightarrow \mathbb{R}$  and the particle-penalisation function  $\mathring{\psi}_R^{\dot{\mathbf{y}}} : \mathbb{R}^d \rightarrow \mathbb{R}$  are chosen among non-negative functions of class  $C^2$  with bounded derivatives, such that  $\nabla \mathring{\psi}_R^{\mathbf{y}}$  vanishes on  $\mathring{B}_R^{\mathbf{y}}$ ,  $\nabla \mathring{\psi}_R^{\dot{\mathbf{y}}}$  vanishes on  $\mathring{B}_R^{\dot{\mathbf{y}}}$ , and they are small enough outside of  $\mathring{B}_R^{\mathbf{y}}$  and  $\mathring{B}_R^{\dot{\mathbf{y}}}$  respectively, in the sense that:

$$\sum_{R=1}^{\infty} \int_{(\mathring{B}_R^{\mathbf{y}})^c} e^{-\mathring{\psi}_R^{\mathbf{y}}(x)} dx < +\infty \quad \text{and} \quad \sum_{R=1}^{\infty} \int_{(\mathring{B}_R^{\dot{\mathbf{y}}})^c} e^{-\mathring{\psi}_R^{\dot{\mathbf{y}}}(x)} dx < +\infty. \quad (2.2)$$

That is, for the finite measures on  $\mathbb{R}^d$  defined as  $\mathring{\lambda}_R^{\mathbf{y}}(d\dot{x}) := e^{-\mathring{\psi}_R^{\mathbf{y}}(\dot{x})} d\dot{x}$ , and  $\mathring{\lambda}_R^{\dot{\mathbf{y}}}(d\dot{x}) := e^{-\mathring{\psi}_R^{\dot{\mathbf{y}}}(\dot{x})} d\dot{x}$ , where  $d\dot{x}$  denotes the Lebesgue measure, we assume that  $\sum_{R=1}^{\infty} \mathring{\lambda}_R^{\mathbf{y}}((\mathring{B}_R^{\mathbf{y}})^c) + \mathring{\lambda}_R^{\dot{\mathbf{y}}}((\mathring{B}_R^{\dot{\mathbf{y}}})^c) < +\infty$ . An example of such functions  $\mathring{\psi}_R^{\mathbf{y}}$  and  $\mathring{\psi}_R^{\dot{\mathbf{y}}}$ , having linear growth with respect to the Euclidean norm at infinity, is given in Section A.1.

These penalisation functions will be used as drifts in a finite-dimensional penalised dynamics whose solution will be shown to approximate the solution of  $(\mathcal{S}_\infty)$ .

## 2.2 Penalised SDE

We consider the penalised SDE around  $B(0, R)$  for a finite number  $n$  of spheres and a finite number  $m$  of particles given by

$$\left\{ \begin{array}{l} \text{for any } i \in \{1, \dots, n\}, k \in \{1, \dots, m\}, t \in [0, 1], \\ \dot{X}_i(t) = \dot{X}_i(0) + \dot{W}_i(t) - \frac{1}{2} \int_0^t \nabla \psi_R^y(\dot{X}_i(s)) ds \\ \quad + \sum_{j=1}^n \int_0^t (\dot{X}_i - \dot{X}_j)(s) dL_{ij}(s) + \sum_{k=1}^m \int_0^t (\dot{X}_i - \dot{X}_k)(s) dl_{ik}(s) \\ \dot{X}_k(t) = \dot{X}_k(0) + \dot{W}_k(t) - \frac{\dot{\sigma}^2}{2} \int_0^t \nabla \psi_R^y(\dot{X}_k(s)) ds + \dot{\sigma}^2 \sum_{i=1}^n \int_0^t (\dot{X}_k - \dot{X}_i)(s) dl_{ik}(s) \\ \forall j \in \{1, \dots, n\}, L_{ij}(0) = \ell_{ik}(0) = 0, L_{ij} \equiv L_{ji}, L_{ii} \equiv 0 \\ \int_0^t \mathbb{1}_{|\dot{X}_i(s) - \dot{X}_j(s)| \neq 2\hat{r}} dL_{ij}(s) = 0, \int_0^t \mathbb{1}_{|\dot{X}_i(s) - \dot{X}_k(s)| \neq \hat{r}} dl_{ik}(s) = 0. \end{array} \right. \quad (\mathcal{S}_R^y)$$

The existence of a unique solution to the above system follows from [FKRZ24, Proposition 2.2]:

**Proposition 2.1.** For any fixed outside condition  $\mathbf{y} \in \mathcal{D}$ , any fixed radius  $R > 0$ , and any number  $n \in \mathbb{N}$  of large spheres and  $m \in \mathbb{N}$  of small particles, the penalised finite-dimensional SDE with reflection  $(\mathcal{S}_R^y)$  admits a unique  $\mathcal{D}$ -valued strong solution  $\mathbf{X}^{\mathbf{y}, R, n, m}(\mathbf{x}) = (\mathbf{X}^{\mathbf{y}, R, n, m}(\mathbf{x}, t), t \in [0, 1])$ ,

$$\mathbf{X}^{\mathbf{y}, R, n, m}(\mathbf{x}, t) := (\dot{X}_i^{\mathbf{y}, R, n, m}(\mathbf{x}, t), \dot{X}_k^{\mathbf{y}, R, n, m}(\mathbf{x}, t), L_{ij}^{\mathbf{y}, R, n, m}(\mathbf{x}, t), \ell_{ik}^{\mathbf{y}, R, n, m}(\mathbf{x}, t))_{1 \leq i, j \leq n, 1 \leq k \leq m},$$

for  $\nu_{R, n, m}^y$ -almost every deterministic initial condition  $\mathbf{x} = \dot{\mathbf{x}} = (\dot{x}_1, \dots, \dot{x}_n, \dot{x}_1, \dots, \dot{x}_m) \in \mathcal{D}$ , where the measure  $\nu_{R, n, m}^y$ , concentrated on the admissible configurations with  $n$  spheres and  $m$  particles, is defined, for any positive measurable function  $F$  on  $\mathcal{D}$ , by

$$\int_{\mathcal{D}} F(\mathbf{x}) d\nu_{R, n, m}^y(\mathbf{x}) := \int_{\mathbb{R}^{nd}} \int_{\mathbb{R}^{md}} F(\dot{\mathbf{x}}) \mathbb{1}_{\mathcal{D}}(\dot{\mathbf{x}}) \otimes_{k=1}^m \dot{\lambda}_R^y(d\dot{x}_k) \otimes_{i=1}^n \dot{\lambda}_R^y(d\dot{x}_i). \quad (2.3)$$

Moreover, the finite measure  $\nu_{R, n, m}^y$  is reversible for the dynamics  $(\mathcal{S}_R^y)$ .

Note that  $\nu_{R, n, m}^y$  only depends on  $\mathbf{y}$  via  $\psi_R^y$  and  $\dot{\psi}_R^y$ , which only depend on  $\mathbf{y}_{B(0, R)^c}$ . The initial configurations  $\mathbf{x} = \dot{\mathbf{x}}$  given by  $\nu_{R, n, m}^y$  belong to  $\mathcal{D}$ , but the superposition  $\mathbf{y}_{B(0, R)^c} \dot{\mathbf{x}}$  is in general not in  $\mathcal{D}$ .

**Remark 2.2.** For  $n = 0$ , there is no sphere process and no local time process  $L_{ij}$  and  $\ell_{ik}$ ; the process  $\mathbf{X}^{\mathbf{y}, R, 0, m}(\mathbf{x}, \cdot) = (\dot{X}_k^{\mathbf{y}, R, 0, m}(\mathbf{x}, \cdot))_{1 \leq k \leq m}$  only involves  $m$  independent particles. Then (2.3) holds with the convention that  $\dot{\mathbf{x}} = \emptyset$ ,  $\int_{\mathbb{R}^0} C^{\text{st}} \otimes_{i=1}^0 \dot{\lambda}_R^y(d\dot{x}_i) = C^{\text{st}}$ , and  $\int F(\mathbf{x}) d\nu_{R, 0, m}^y(\mathbf{x}) := \int_{\mathbb{R}^{md}} F(\dot{\mathbf{x}}) \otimes_{k=1}^m \dot{\lambda}_R^y(d\dot{x}_k)$ .

Similarly, for  $m = 0$ , there is no particle process and no local time process  $\ell_{ik}$ ; the process  $\mathbf{X}^{\mathbf{y}, R, n, 0}(\mathbf{x}, t) = (\dot{X}_i^{\mathbf{y}, R, n, 0}(\mathbf{x}, \cdot), L_{ij}^{\mathbf{y}, R, n, 0}(\mathbf{x}, \cdot))_{1 \leq i, j \leq n}$  only involves  $n$  hard spheres. Then (2.3) holds with the convention that  $\dot{\mathbf{x}} = \emptyset$  and  $\int F(\mathbf{x}) d\nu_{R, n, 0}^y(\mathbf{x}) := \int_{\mathbb{R}^{nd}} F(\dot{\mathbf{x}}) \mathbb{1}_{\mathcal{D}}(\dot{\mathbf{x}}) \otimes_{i=1}^n \dot{\lambda}_R^y(d\dot{x}_i)$ .

For  $n = 0$  and  $m = 0$ , the process is empty. For the sake of completeness, in the sequel we let  $\int F(\emptyset) d\nu_{R, 0, 0}^y(\mathbf{x}) := F(\emptyset)$ .



**Definition 2.3.** Let  $Q_{R,n,m}^y$  denote the time-reversible pushforward measure of the solution of the penalised SDE  $(\mathcal{S}_R^y)$  with initial measure  $\nu_{R,n,m}^y$ :

$$Q_{R,n,m}^y(\cdot) := \int P\left(\left(\mathbf{X}^{y,R,n,m}(\mathbf{x}, t)\right)_{t \in [0,1]} \in \cdot\right) d\nu_{R,n,m}^y(\mathbf{x}). \quad (2.4)$$

## 2.3 Mixture of penalised processes

We construct an approximation of the expected reversible measure  $\mu$  as a Poissonian mixture of the reversible measures of the solutions of  $(\mathcal{S}_R^y)$  for different numbers of spheres and particles:

$$\mu_R^y := \frac{1}{\mathcal{Z}_R^y} \sum_{n=0}^{+\infty} \frac{\dot{z}^n}{n!} \sum_{m=0}^{+\infty} \frac{\dot{z}^m}{m!} \nu_{R,n,m}^y. \quad (2.5)$$

For any  $y$ , the following mixture  $Q_R^y$  of penalised processes distributions starting from  $\mu_R^y$  is reversible:

$$Q_R^y := \frac{1}{\mathcal{Z}_R^y} \sum_{n=0}^{+\infty} \frac{\dot{z}^n}{n!} \sum_{m=0}^{+\infty} \frac{\dot{z}^m}{m!} Q_{R,n,m}^y. \quad (2.6)$$

Finally, we define the approximate process  $X^{y,R}$  as the solution of  $(\mathcal{S}_R^y)$  with  $n = \sharp(\dot{y}_{B(0,R)})$  and  $m = \sharp(\dot{y}_{B(0,R)})$ , where  $\sharp$  denotes the number of points, and with initial configuration equal to  $\dot{y}_{B(0,R)} \dot{y}_{B(0,R)}$ , that is,

$$X^{y,R}(\cdot) := \mathbf{X}^{y,R,\sharp(\dot{y}_{B(0,R)}),\sharp(\dot{y}_{B(0,R)})}(\dot{y}_{B(0,R)}, \cdot). \quad (2.7)$$

To obtain Theorem 1.2, we will prove that, as  $R$  tends to infinity, the  $Q_R^y$ -distributed reversible process converges, in a locally eventually-constant way, to a solution of  $(\mathcal{S}_\infty)$ , and the distribution of  $X^{y,R}$  is arbitrary close to  $Q_R^y$  when the outside configuration  $y$  is chosen according to  $\mu$ .

## 3 Convergence of the penalised processes

In this section, we show that the penalised process  $X^{y,R}$  converges to the solution of  $(\mathcal{S}_\infty)$ : we first prove, in Section 3.1, that the convergence holds in a certain subset of  $\Omega$ , and then, in Section 3.2, show that this subset has full mass under the reversible measures. Uniqueness is shown in Remark 3.9.

In order to prove the convergence, we have to check that balls (spheres or particles) only interact locally, making sure that the probability of *bad paths*, i.e. those of balls that interact with a great number of other ones, vanishes. The proof relies on the fact that *long chains* of interacting spheres rarely form, and that balls coming *very fast* from far away also are improbable. Let us first define these two events. See Figure 1 for a visualisation of a long chain.

**Definition 3.1** (Long chains). For every positive radius  $\alpha$ , every positive leeway  $\varepsilon$  and every chain length  $\kappa$ , define the set of configurations which have a long sphere chain by

$$\mathcal{B}_{\text{Chain}}(\alpha, \kappa, \varepsilon) := \left\{ \mathbf{x} \in \mathcal{D} \text{ s.t. for some } \{\dot{x}_{j_1}, \dots, \dot{x}_{j_{\kappa+1}}\} \subset \dot{\mathbf{x}} \right. \\ \left. |\dot{x}_{j_1}| \leq \alpha \text{ and } |\dot{x}_{j_1} - \dot{x}_{j_2}| < 2\dot{r} + \varepsilon, \dots, |\dot{x}_{j_\kappa} - \dot{x}_{j_{\kappa+1}}| < 2\dot{r} + \varepsilon \right\}.$$




Figure 1: Highlighted, an example of an  $\varepsilon$ -chain of length  $\kappa = 10$ .

For every positive time step  $\delta$ , the set of configuration paths whose  $\delta$ -discretisation presents an  $\varepsilon$ -chain is

$$\mathcal{B}'_{\text{Chain}}(\delta, \alpha, \kappa, \varepsilon) := \left\{ \mathbf{X}: [0, 1] \rightarrow \mathcal{D} \text{ s.t. } \exists k \in \mathbb{N} \mathbf{X}(k\delta) \in \mathcal{B}_{\text{Chain}}(\alpha, \kappa, \varepsilon) \right\}.$$

**Definition 3.2** (Fast balls). For every positive radius  $\alpha$ , every positive range  $\varepsilon$  and every positive time duration  $\delta$ , define the set of paths which have oscillation  $\varepsilon$  during  $\delta$  as

$$\mathcal{B}'_{\text{Fast}}(\alpha, \delta, \varepsilon) := \left\{ f \in C^0([0, 1]; \mathbb{R}^d) \text{ s.t. } \min_{0 \leq s \leq 1} |f(s)| \leq \alpha \text{ and } \sup_{\substack{|t-s| < \delta \\ 0 \leq s, t \leq 1}} |f(t) - f(s)| > \varepsilon \right\}.$$

The set of configuration paths involving a sphere or particle that moves too fast is defined as

$$\mathcal{B}'_{\text{Fast}}(\alpha, \delta, \varepsilon) := \left\{ \mathbf{X}: [0, 1] \rightarrow \mathcal{D} \text{ s.t. } \exists i, \dot{X}_i(\cdot) \in \mathcal{B}'_{\text{Fast}}(\alpha, \delta, \varepsilon) \text{ or } \exists k, \dot{X}_k(\cdot) \in \mathcal{B}'_{\text{Fast}}(\alpha, \delta, \varepsilon) \right\}.$$

We will show that the approximate processes  $X^{\mathbf{y}, R}$  coincide for  $R$  large enough, as far as their sphere and particles in some bounded area are concerned. This happens as soon as, for any  $\varepsilon$  smaller than some positive threshold  $\varepsilon(\mathbf{y})$  and for all ranges  $R$  large enough, both processes  $X^{\mathbf{y}, R}$  and  $X^{\mathbf{y}, R+1}$  do not belong to the  $\mathcal{B}'_{\text{Chain}}$  and  $\mathcal{B}'_{\text{Fast}}$  events given by  $\varepsilon$  and some suitable parameters  $\delta$ ,  $\alpha$ , and  $\kappa$  depending on  $R$ . Formally, the set on which the convergence holds, and which will be shown to be of full measure, is:

$$\Omega^{\mathbf{y}} := \left\{ \omega \in \Omega : \exists \varepsilon(\mathbf{y}) > 0 \forall \varepsilon < \varepsilon(\mathbf{y}) \exists R(\varepsilon) \in \mathbb{N} \text{ s.t. } \forall R \geq R(\varepsilon), \right. \\ \left. X^{\mathbf{y}, R}(\omega) \notin \mathcal{B}'_{\text{Chain}}(\delta(R), \alpha(R), \kappa(R), \varepsilon), \quad X^{\mathbf{y}, R} \notin \mathcal{B}'_{\text{Fast}}(\alpha(R), \delta(R), \varepsilon/4), \right. \\ \left. X^{\mathbf{y}, R+1}(\omega) \notin \mathcal{B}'_{\text{Chain}}(\delta(R), \alpha(R), \kappa(R), \varepsilon), \quad X^{\mathbf{y}, R+1} \notin \mathcal{B}'_{\text{Fast}}(\alpha(R), \delta(R), \varepsilon/4) \right\},$$

where  $\delta(R) := 1/\lfloor R^{1/3} \rfloor$ ,  $\alpha(R) := R - 2\hat{r}$  and  $\kappa(R) := \lfloor R^{1/3} \rfloor$ .

The construction of a solution to  $(\mathcal{S}_\infty)$  is then split into the two following propositions, proved in Sections 3.1 and 3.2 respectively, and a short Section 3.3 to check the reversibility of the limit process obtained in these propositions.

**Proposition 3.3.** For any fixed  $\mathbf{y}$  in the interior of  $\mathcal{D}$ , for any  $\omega \in \Omega^{\mathbf{y}}$ , the sequence of processes  $(X^{\mathbf{y},R}(\omega), \omega \in \Omega^{\mathbf{y}})_{R \in \mathbb{N}^*}$  converges to a limit process that solves  $(\mathcal{S}_\infty)$  with initial condition  $\mathbf{y}$ .

**Proposition 3.4.** Any Gibbs measure  $\mu \in \mathcal{G}_{z,z}$  has its support in  $\{\mathbf{y} \in \mathcal{D} \text{ s.t. } P(\Omega^{\mathbf{y}}) = 1\}$ , that is  $\int_{\mathcal{D}} P((\Omega^{\mathbf{y}}))\mu(d\mathbf{y}) = 1$ .

### 3.1 Proof of Proposition 3.3 : convergence on a subset

**Definition 3.5.** For each configuration  $\mathbf{x} \in \mathcal{D}$ , each radius  $\rho > 0$ , and each leeway  $\varepsilon > 0$ :

- The set of spheres which belong to  $B(0, \rho)$  or to an  $\varepsilon$ -chain connected to  $B(0, \rho)$  is:

$$\mathring{\mathcal{J}}(\mathbf{x}, \rho, \varepsilon) := \left\{ i \text{ s.t. } |\dot{x}_i| < \rho \text{ or for some } \{\dot{x}_{j_1}, \dots, \dot{x}_{j_k}\} \subset \dot{\mathbf{x}} \quad |\dot{x}_{j_1}| \leq \rho \text{ and} \right. \\ \left. |\dot{x}_{j_1} - \dot{x}_{j_2}| < 2\dot{r} + \varepsilon, \dots, |\dot{x}_{j_{k-1}} - \dot{x}_{j_k}| < 2\dot{r} + \varepsilon, |\dot{x}_{j_k} - \dot{x}_i| < 2\dot{r} + \varepsilon \right\}.$$

- The set of particles which belong to  $B(0, \rho)$  or are close to some  $\varepsilon$ -chain connected to  $B(0, \rho)$  is:

$$\dot{\mathcal{J}}(\mathbf{x}, \rho, \varepsilon) := \left\{ k \text{ s.t. } |\dot{x}_k| < \rho \text{ or } \exists i \in \mathring{\mathcal{J}}(\mathbf{x}, \rho, \varepsilon) \text{ s.t. } |\dot{x}_k - \dot{x}_i| \leq \overset{\circ}{r} + \varepsilon \right\}.$$

The following crucial lemma is proved in Section A.2:

**Lemma 3.6.** Assume that for a chain size  $\kappa \geq 2$ , a time step  $\delta$  (inverse of integer), and a ball size  $\alpha$ , the path  $X : [0, 1] \rightarrow \mathcal{D}$  has neither too large a chain nor too large an oscillation, i.e.

$$X \notin \mathcal{B}'_{\text{Chain}}(\delta, \alpha, \kappa, \varepsilon) \quad \text{and} \quad X \notin \mathcal{B}'_{\text{Fast}}(\alpha, \delta, \varepsilon/4).$$

Fix  $\rho$  such that  $\rho_0 := \rho + \frac{2\kappa}{\delta}(2\dot{r} + \varepsilon) \leq \alpha$ . Then, for any  $a \in \{0, 1, 2, \dots, \frac{1}{\delta}\}$ , for all  $t \in [a\delta, (a+1)\delta]$ , and for  $\rho_a := \rho_0 - 2a\kappa(2\dot{r} + \varepsilon) = \rho + 2\left(\frac{1}{\delta} - a\right)\kappa(2\dot{r} + \varepsilon)$ :

- Spheres in  $\mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon)$  do not bump into the other balls (separation property):
 
$$\forall j \notin \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \forall i \in \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \quad |\dot{X}_j(t) - \dot{X}_i(t)| > 2\dot{r} + \frac{\varepsilon}{2},$$

$$\forall k \notin \dot{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \forall i \in \dot{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \quad |\dot{X}_k(t) - \dot{X}_i(t)| > \overset{\circ}{r} + \frac{\varepsilon}{2}.$$
- Spheres and particles not too far from the origin stay around the origin (localisation property):
 
$$\forall i \in \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \quad |\dot{X}_i(t)| \leq \rho_a + \kappa(2\dot{r} + \varepsilon) + \frac{\varepsilon}{4},$$

$$\forall k \in \dot{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \quad |\dot{X}_k(t)| \leq \rho_a + \kappa(2\dot{r} + \varepsilon) + \overset{\circ}{r} + \frac{5\varepsilon}{4}.$$
- The index sets form a decreasing sequence (nested inclusion property):
 
$$\mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon) \supset \mathring{\mathcal{J}}(X((a+1)\delta), \rho_{a+1}, \varepsilon),$$

$$\dot{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon) \supset \dot{\mathcal{J}}(X((a+1)\delta), \rho_{a+1}, \varepsilon).$$

Fix a configuration  $\mathbf{y} = \dot{\mathbf{y}}\dot{\mathbf{y}} \in \mathcal{D}$ , a radius  $\rho$ , and consider an  $\omega \in \Omega^{\mathbf{y}}$ . By definition, for any fixed  $\varepsilon < \varepsilon(\mathbf{y})$ , there exists  $R(\varepsilon) > 0$  such that, for every  $R \geq R(\varepsilon)$ ,

$$X^{\mathbf{y},R}(\omega) \notin \mathcal{B}'_{\text{Chain}}(\delta(R), \alpha(R), \kappa(R), \varepsilon), \quad X^{\mathbf{y},R}(\omega) \notin \mathcal{B}'_{\text{Fast}}(\alpha(R), \delta(R), \varepsilon/4) \text{ and} \\ X^{\mathbf{y},R+1}(\omega) \notin \mathcal{B}'_{\text{Chain}}(\delta(R), \alpha(R), \kappa(R), \varepsilon), \quad X^{\mathbf{y},R}(\omega) \notin \mathcal{B}'_{\text{Fast}}(\alpha(R), \delta(R), \varepsilon/4).$$

Recall that  $\delta(R) := 1/\lfloor R^{1/3} \rfloor$ ,  $\alpha(R) := R - 2\mathring{r}$  and  $\kappa(R) := \lfloor R^{1/3} \rfloor$ . Since  $\kappa(R)/\delta(R) \sim R^{2/3} \ll \alpha(R) \sim R$ , there exists  $R(\varepsilon, \rho) \geq R(\varepsilon)$  such that for any  $R \geq R(\varepsilon, \rho)$ :

$$\rho + \frac{\varepsilon}{\delta(R)} + \left( \frac{2}{\delta(R)} + 1 \right) \kappa(R)(2\mathring{r} + \varepsilon) + \mathring{r} + \frac{5\varepsilon}{4} \leq \alpha(R). \quad (3.1)$$

We now fix  $R \geq R(\varepsilon, \rho)$ . Let us prove by induction that, for all  $t \in [0, 1]$ : for all  $i$  such that  $|\mathring{y}_i| < \rho$ , all  $k$  such that  $|\mathring{y}_k| < \rho$ , and all  $R \geq R(\varepsilon, \rho)$ ,

$$\mathring{X}_i^{\mathbf{y},R}(\omega, t) = \mathring{X}_i^{\mathbf{y},R+1}(\omega, t) \text{ and } \dot{X}_k^{\mathbf{y},R}(\omega, t) = \dot{X}_k^{\mathbf{y},R+1}(\omega, t).$$

At time  $t = 0$ , the initial configurations  $X^{\mathbf{y},R}(\omega, 0) = \mathbf{y}_{B(0,R)}$  and  $X^{\mathbf{y},R+1}(\omega, 0) = \mathbf{y}_{B(0,R+1)}$  coincide as far as their spheres in  $\mathring{\mathcal{J}}(X^{\mathbf{y},R}(\omega, 0), \rho_0, \varepsilon)$  and their particles in  $\dot{\mathcal{J}}(X^{\mathbf{y},R}(\omega, 0), \rho_0, \varepsilon)$  are concerned, for

$$\rho_0 := \rho + \frac{\varepsilon}{\delta(R)} + \frac{2\kappa(R)}{\delta(R)}(2\mathring{r} + \varepsilon),$$

since  $\rho_0 + \kappa(R)(2\mathring{r} + \varepsilon) + \mathring{r} \leq R$  thanks to (3.1). This provides the initial step for the induction procedure.

We turn to the induction step. Assume that, for a fixed integer  $a$  between 0 and  $1/\delta(R)$ , and for  $\rho_a := \rho + \frac{\varepsilon}{\delta(R)} + 2\left(\frac{1}{\delta(R)} - a\right)\kappa(R)(2\mathring{r} + \varepsilon)$ , the processes  $X^{\mathbf{y},R}(\omega, \cdot)$  and  $X^{\mathbf{y},R+1}(\omega, \cdot)$  and their local times coincide on  $[0, a\delta(R)]$  as far as their spheres in  $\mathring{\mathcal{J}}(X^{\mathbf{y},R}(\omega, a\delta), \rho_a, \varepsilon)$  and their particles in  $\dot{\mathcal{J}}(X^{\mathbf{y},R}(\omega, a\delta), \rho_a, \varepsilon)$  are concerned, and satisfy  $(\mathcal{S}_\infty)$  restricted to these balls up to time  $a\delta(R)$ . We want to prove the same result for  $(a+1)$  instead of  $a$ . The separation property in Lemma 3.6 implies that the spheres in

$$\mathring{\mathcal{J}} := \mathring{\mathcal{J}}(X^{\mathbf{y},R}(\omega, a\delta(R)), \rho_a, \varepsilon) = \mathring{\mathcal{J}}(X^{\mathbf{y},R+1}(\omega, a\delta(R)), \rho_a, \varepsilon)$$

or the particles in

$$\dot{\mathcal{J}} := \dot{\mathcal{J}}(X^{\mathbf{y},R}(\omega, a\delta(R)), \rho_a, \varepsilon) = \dot{\mathcal{J}}(X^{\mathbf{y},R+1}(\omega, a\delta(R)), \rho_a, \varepsilon)$$

do not bump, from time  $a\delta(R)$  to time  $(a+1)\delta(R)$ , into the spheres and particles which do not belong to these index sets. Thanks to  $(\mathcal{S}_R^{\mathbf{y}})$  and its Markov property, this implies:

$$\left\{ \begin{array}{l} \text{for any } i \in \mathring{\mathcal{J}}, \text{ any } k \in \dot{\mathcal{J}}, \text{ and any } t \in [a\delta(R), (a+1)\delta(R)], \\ \mathring{X}_i^{\mathbf{y},R}(\omega, t) = \mathring{X}_i^{\mathbf{y},R}(\omega, a\delta(R)) + \mathring{W}_i(\omega, t) - \mathring{W}_i(\omega, a\delta(R)) - \frac{1}{2} \int_{a\delta(R)}^t \nabla \mathring{\psi}_R^{\mathbf{y}}(\mathring{X}_i^{\mathbf{y},R}(\omega, s)) ds \\ \quad + \sum_{j \in \mathring{\mathcal{J}}} \int_{a\delta(R)}^t (\mathring{X}_i^{\mathbf{y},R} - \mathring{X}_j^{\mathbf{y},R})(\omega, s) dL_{ij}(\omega, s) + \sum_{k \in \dot{\mathcal{J}}} \int_{a\delta(R)}^t (\mathring{X}_i^{\mathbf{y},R} - \mathring{X}_k^{\mathbf{y},R})(\omega, s) dl_{ik}(\omega, s) \\ \dot{X}_k^{\mathbf{y},R}(\omega, t) = \dot{X}_k^{\mathbf{y},R}(\omega, a\delta(R)) + \dot{W}_k^{\mathbf{y},R}(\omega, t) - \dot{W}_k^{\mathbf{y},R}(\omega, a\delta(R)) - \frac{\dot{\sigma}^2}{2} \int_{a\delta(R)}^t \nabla \dot{\psi}_R^{\mathbf{y}}(\dot{X}_k^{\mathbf{y},R}(\omega, s)) ds \\ \quad + \dot{\sigma}^2 \sum_{j \in \dot{\mathcal{J}}} \int_{a\delta(R)}^t (\dot{X}_k^{\mathbf{y},R} - \dot{X}_j^{\mathbf{y},R})(\omega, s) dl_{jk}(\omega, s). \end{array} \right.$$

This is also true with  $R+1$  instead of  $R$ . The  $\mathring{X}_i^{\mathbf{y},R}(\omega, \cdot)$  and  $\mathring{X}_i^{\mathbf{y},R+1}(\omega, \cdot)$  for  $i \in \mathring{\mathcal{J}}$ , as well as the  $\dot{X}_k^{\mathbf{y},R}(\omega, \cdot)$  and  $\dot{X}_k^{\mathbf{y},R+1}(\omega, \cdot)$  for  $k \in \dot{\mathcal{J}}$ , stay in the ball  $B(0, R - 2\mathring{r})$  where  $\nabla \mathring{\psi}_R^{\mathbf{y}}$ ,  $\nabla \mathring{\psi}_{R+1}^{\mathbf{y}}$ ,  $\nabla \dot{\psi}_R^{\mathbf{y}}$  and  $\nabla \dot{\psi}_{R+1}^{\mathbf{y}}$  vanish, as a consequence of the localisation property from Lemma 3.6 and the inequality

$\rho_a + \kappa(2\mathring{r} + \varepsilon) + \mathring{r} + \frac{5\varepsilon}{4} \leq R - 2\mathring{r}$ . The restriction to these balls and this time interval of  $(\mathcal{S}_R^y)$  and  $(\mathcal{S}_{R+1}^y)$  both coincide with the corresponding restriction of  $(\mathcal{S}_\infty)$ .

The equality of  $X^{y,R}(\omega, \cdot)$  and  $X^{y,R+1}(\omega, \cdot)$  at time  $a\delta(R)$  for spheres and particles in  $\mathring{\mathcal{J}}$  and  $\mathring{\mathcal{J}}$ , and the uniqueness of the solution of the above equation imply that  $X^{y,R}(\omega, \cdot)$  and  $X^{y,R+1}(\omega, \cdot)$  – and the corresponding local times – coincide on the time interval  $[a\delta(R), (a+1)\delta(R)]$  as far as their spheres in  $\mathring{\mathcal{J}}$  and their particles in  $\mathring{\mathcal{J}}$  are concerned.

In particular,  $\mathring{X}_i^{y,R}(\omega, (a+1)\delta(R)) = \mathring{X}_i^{y,R+1}(\omega, (a+1)\delta(R))$  and  $\mathring{X}_k^{y,R}(\omega, (a+1)\delta(R)) = \mathring{X}_k^{y,R+1}(\omega, (a+1)\delta(R))$  for spheres  $i \in \mathring{\mathcal{J}}$  and particles  $k \in \mathring{\mathcal{J}}$ . The inclusion property from Lemma 3.6 then yields

$$\begin{aligned} \mathring{\mathcal{J}}(X^{y,R}((a+1)\delta(R)), \rho_{a+1}, \varepsilon) &= \mathring{\mathcal{J}}(X^{y,R+1}((a+1)\delta(R)), \rho_{a+1}, \varepsilon) \subset \mathring{\mathcal{J}}, \\ \mathring{\mathcal{J}}(X^{y,R}((a+1)\delta(R)), \rho_{a+1}, \varepsilon) &= \mathring{\mathcal{J}}(X^{y,R+1}((a+1)\delta(R)), \rho_{a+1}, \varepsilon) \subset \mathring{\mathcal{J}}. \end{aligned}$$

As a result,  $X^{y,R}(\omega, \cdot)$  and  $X^{y,R+1}(\omega, \cdot)$  and their local times coincide on  $[0, (a+1)\delta(R)]$  as far as their spheres in  $\mathring{\mathcal{J}}(X^{y,R}(\omega, (a+1)\delta(R)), \rho_{a+1}, \varepsilon)$  and their particles in  $\mathring{\mathcal{J}}(X^{y,R}(\omega, (a+1)\delta(R)), \rho_{a+1}, \varepsilon)$  are concerned, and the equation  $(\mathcal{S}_R^y)$  they satisfy reduces to  $(\mathcal{S}_\infty)$  for these balls on this time interval. This concludes the induction step.

The above step-by-step coincidence result holds for each integer  $a$  up to  $1/\delta(R)$ . The solutions  $X^{y,R}(\omega, \cdot)$  and  $X^{y,R+1}(\omega, \cdot)$  of the penalised equations starting from configuration  $\mathbf{y}$  are equal up to time 1 for their components which are spheres in  $\mathring{\mathcal{J}}(X^{y,R}(\omega, 1), \rho_{1/\delta(R)}, \varepsilon)$  or particles in  $\mathring{\mathcal{J}}(X^{y,R}(\omega, 1), \rho_{1/\delta(R)}, \varepsilon)$ .

Since  $X^{y,R}(\omega, \cdot)$  and  $X^{y,R+1}(\omega, \cdot)$  both do not have balls that cover a distance larger than  $\frac{\varepsilon}{\delta(R)}$  during the time interval  $[0, 1]$  if they ever enter  $B(0, \alpha(R))$ , hence stay in  $B(0, \rho + \frac{\varepsilon}{\delta(R)}) = B(0, \rho_{1/\delta(R)})$  if they start in  $B(0, \rho)$ , we have

$$\left\{ i : |\mathring{y}_i| < \rho \right\} \subset \mathring{\mathcal{J}}(X^{y,R}(\omega, 1), \rho_{1/\delta(R)}, \varepsilon) \quad \text{and} \quad \left\{ k : |\mathring{y}_k| < \rho \right\} \subset \mathring{\mathcal{J}}(X^{y,R}(\omega, 1), \rho_{1/\delta(R)}, \varepsilon).$$

The penalised processes  $X^{y,R}(\omega, \cdot)$  and  $X^{y,R+1}(\omega, \cdot)$  are then the same process (including local times) for all spheres and particles starting in  $B(0, \rho)$ , and satisfy  $(\mathcal{S}_\infty)$ .

Let us summarise the result of this induction process: For any  $\mathbf{y} = \mathring{\mathbf{y}} \in \mathcal{D}$  and  $\omega \in \Omega^y$ , for  $\varepsilon < \varepsilon(\mathbf{y})$ , for any finite collection of spheres  $\mathring{y}_i$  and particles  $\mathring{y}_k$ , there exists a radius  $\rho$  larger than all the  $|\mathring{y}_i|$ 's and  $|\mathring{y}_k|$ 's, and there exists  $R(\varepsilon, \rho)$  large enough such that, as soon as  $R > R(\varepsilon, \rho)$ , all the  $\mathring{X}_i^{y,R}(\omega, t)$  and  $\mathring{X}_k^{y,R}(\omega, t)$  are equal for  $t \in [0, 1]$ . All the local times for the sphere-sphere collisions and the sphere-particle collisions in this collection also do not depend on  $R$  as soon as  $R > R(\varepsilon, \rho)$ . Moreover, the eventually-constant limit paths  $\mathring{X}_i^{y,\infty}(\omega, \cdot) := \lim_{R \rightarrow +\infty} \mathring{X}_i^{y,R}(\omega, \cdot)$  and  $\mathring{X}_k^{y,\infty}(\omega, \cdot) := \lim_{R \rightarrow +\infty} \mathring{X}_k^{y,R}(\omega, \cdot)$  satisfy  $(\mathcal{S}_\infty)$ . Since  $\mathbf{X}^{y,R}(\omega, 0) = \mathbf{y}$  for all  $R$ 's by construction,  $\mathbf{X}^{y,\infty}(\omega, 0) := \mathbf{y}$ . This concludes the proof of Proposition 3.3.

### 3.2 Proof of Proposition 3.4 : almost sure convergence

We prove here Proposition 3.4, that is that  $\int_{\mathcal{D}} P((\Omega^y)^c) \mu(dy) = 0$  for  $\mu \in \mathcal{G}_{z,z}$ . Recall that, for  $\delta(R) := 1/\lfloor R^{1/3} \rfloor$ ,  $\alpha(R) := R - 2\mathring{r}$  and  $\kappa(R) := \lfloor R^{1/3} \rfloor$ , the complement set of  $\Omega^y$  is

$$(\Omega^y)^c := \bigcap_{\varepsilon(\mathbf{y}) \in 1/\mathbb{N}} \bigcup_{\substack{\varepsilon \in 1/\mathbb{N} \\ \varepsilon < \varepsilon(\mathbf{y})}} \limsup_{R \rightarrow +\infty} \left\{ X^{y,R} \in \mathcal{B}'(\varepsilon, R) \text{ or } X^{y,R+1} \in \mathcal{B}'(\varepsilon, R) \right\},$$

where  $\mathcal{B}'(\varepsilon, R) := \mathcal{B}'_{\text{Chain}}(\delta(R), \alpha(R), \kappa(R), \varepsilon) \cup \mathcal{B}'_{\text{Fast}}(\alpha(R), \delta(R), \varepsilon/4)$ . This can be rewritten as

$$(\Omega^{\mathbf{y}})^c = \bigcap_{\varepsilon(\mathbf{y}) \in 1/\mathbb{N}} \bigcup_{\substack{\varepsilon \in 1/\mathbb{N} \\ \varepsilon < \varepsilon(\mathbf{y})}} \limsup_{R \rightarrow +\infty} \left\{ X^{\mathbf{y}, R} \in \mathcal{B}'(\varepsilon, R) \cup \mathcal{B}'(\varepsilon, R-1) \right\}.$$

Thanks to the Borel–Cantelli lemma,  $\int_{\mathcal{D}} P((\Omega^{\mathbf{y}})^c) \mu(d\mathbf{y}) = 0$  as soon as there exists  $\varepsilon_0 > 0$  such that:

$$\forall \varepsilon < \varepsilon_0, \quad \sum_R \int_{\mathcal{D}} P\left(X^{\mathbf{y}, R} \in \mathcal{B}'(\varepsilon, R) \cup \mathcal{B}'(\varepsilon, R-1)\right) \mu(d\mathbf{y}) < +\infty.$$

A sufficient condition for the above summability to hold is

$$\exists \varepsilon_0 > 0 : \quad \forall \varepsilon < \varepsilon_0, \quad \sum_R \int_{\mathcal{D}} Q_R^{\mathbf{y}}\left(\mathcal{B}'(\varepsilon, R) \cup \mathcal{B}'(\varepsilon, R-1)\right) \mu(d\mathbf{y}) < +\infty, \quad (3.2)$$

$$\text{and } \sum_R \mathbf{d}_{TV}(R) < +\infty, \quad \text{where} \quad (3.3)$$

$$\mathbf{d}_{TV}(R) := \sup_{\Theta \subset \mathcal{C}([0,1], \mathcal{D})} \left| \int_{\mathcal{D}} P\left(X^{\mathbf{y}, R} \in \Theta\right) \mu(d\mathbf{y}) - \int_{\mathcal{D}} Q_R^{\mathbf{y}}(\Theta) \mu(d\mathbf{y}) \right|.$$

The proof of (3.2) relies on the two following results on chains and fast motion, whose proofs are postponed to Sections A.3 and A.4, respectively.

**Lemma 3.7.** For each sphere activity  $\dot{z} > 0$ , each chain length  $\kappa \in \mathbb{N}^*$ , each radius  $\alpha > 0$ , and each chain leeway  $\varepsilon \in (0, 2\dot{r})$ , the probability of a long sphere chain to occur is bounded by:

$$\forall \mathbf{y} \in \mathcal{D}, \forall R > 0, \quad Q_R^{\mathbf{y}}\left(\mathcal{B}'_{\text{Chain}}(\delta, \alpha, \kappa, \varepsilon)\right) \leq \frac{1}{\delta} \left( \dot{z} \left( (2\dot{r} + \varepsilon)^d - (2\dot{r})^d \right) \mathbf{v}_d \right)^{\kappa}.$$

**Lemma 3.8.** There exists a constant  $C_{\text{Fast}} \equiv C_{\text{Fast}}(d, \dot{\sigma}, \dot{z}, \dot{z})$ , depending only on the dimension  $d$ , the particle diffusion coefficient  $\dot{\sigma}$ , and the activities  $\dot{z}$  and  $\dot{z}$ , such that for any oscillation parameters  $0 < \varepsilon \leq 1$ ,  $0 < \delta \leq 1$ , and any radius  $\alpha \geq 1$ :

$$\forall \mathbf{y} \in \mathcal{D}, \forall R > 0, \quad Q_R^{\mathbf{y}}\left(\mathcal{B}'_{\text{Fast}}(\alpha, \delta, \varepsilon)\right) \leq C_{\text{Fast}} \frac{\alpha^d}{\delta^{d+1}} \exp\left(-\frac{\varepsilon^2}{10 d \delta \max(1, \dot{\sigma}^2)}\right).$$

Thanks to Lemma 3.7,

$$\begin{aligned} & \sum_R \int_{\mathcal{D}} Q_R^{\mathbf{y}}\left(\mathcal{B}'_{\text{Chain}}(\delta(R), \alpha(R), \kappa(R), \varepsilon)\right) \mu(d\mathbf{y}) \\ & \leq \sum_R \lfloor R^{1/3} \rfloor \left( \dot{z} \left( (2\dot{r} + \varepsilon)^d - (2\dot{r})^d \right) \mathbf{v}_d \right)^{\lfloor R^{1/3} \rfloor} < +\infty, \end{aligned}$$

as soon as  $\varepsilon$  is small enough for  $\dot{z} \left( (2\dot{r} + \varepsilon)^d - (2\dot{r})^d \right) \mathbf{v}_d < 1$  to hold, which is the case, for example, if  $\varepsilon < \min(2\dot{r}, \left( \dot{z} (4\dot{r} + 1)^d \mathbf{v}_d \right)^{-1})$ . Under the same condition on  $\varepsilon$ , one also obtains

$$\sum_R \int_{\mathcal{D}} Q_R^{\mathbf{y}}\left(\mathcal{B}'_{\text{Chain}}(\delta(R-1), \alpha(R-1), \kappa(R-1), \varepsilon)\right) \mu(d\mathbf{y}) < +\infty.$$

Thanks to Lemma 3.8, for any  $0 < \varepsilon \leq 1$ ,

$$\begin{aligned} & \sum_R \int_{\mathcal{D}} Q_R^{\mathbf{y}} \left( \mathcal{B}'_{\text{Fast}}(\alpha(R), \delta(R), \varepsilon/4) \right) \mu(d\mathbf{y}) \\ & \leq C_{\text{Fast}} \sum_R [R^{1/3}]^{d+1} (R - 2\hat{r})^d \mathbf{v}_d \exp \left( -\frac{\varepsilon^2 [R^{1/3}]}{10 d \max(1, \hat{\sigma}^2)} \right) < +\infty, \end{aligned}$$

and the same summability holds for

$$\sum_R \int_{\mathcal{D}} Q_R^{\mathbf{y}} \left( \mathcal{B}'_{\text{Fast}}(\alpha(R-1), \delta(R-1), \varepsilon/4) \right) \mu(d\mathbf{y}) < +\infty.$$

Thus (3.2) holds. It only remains to prove (3.3), i.e. that the total variation distance between  $\int_{\mathcal{D}} Q_R^{\mathbf{y}}(\cdot) \mu(d\mathbf{y})$  and  $\int_{\mathcal{D}} P(X^{\mathbf{y},R} \in \cdot) \mu(d\mathbf{y})$  converges to zero in a summable way. From (2.7) we have that, for any measurable set  $\Theta$  of continuous paths in  $\mathcal{D}$ :

$$P(X^{\mathbf{y},R} \in \Theta) = P(\mathbf{X}^{\mathbf{y},R, \#(\dot{\mathbf{y}}_{B(0,R)}), \#(\dot{\mathbf{y}}_{B(0,R)})}(\mathbf{y}_{B(0,R)}, \cdot) \in \Theta) =: F_{\Theta}^{\mathbf{y}}(\mathbf{y}_{B(0,R)}),$$

where  $F_{\Theta}^{\mathbf{y}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) := P(\mathbf{X}^{\mathbf{y},R, \#(\dot{\mathbf{x}}), \#(\dot{\mathbf{x}})}(\dot{\mathbf{x}}\dot{\mathbf{x}}, \cdot) \in \Theta)$ . Note that  $F_{\Theta}^{\mathbf{y}}$  only depends on  $\mathbf{y}_{B(0,R)^c}$  and not on  $\mathbf{y}_{B(0,R)}$ . Taking  $\Lambda = B(0, R)$ , with  $\dot{\mathbf{x}} = \{\dot{x}_1, \dots, \dot{x}_n\}$  and  $\dot{\mathbf{x}} = \{\dot{x}_1, \dots, \dot{x}_m\}$ , in the definition (1.1) of  $\mu$ , we find:

$$\begin{aligned} & \int_{\mathcal{D}} P(X^{\mathbf{y},R} \in \Theta) \mu(d\mathbf{y}) = \int_{\mathcal{D}} F_{\Theta}^{\mathbf{y}}(\mathbf{y}_{B(0,R)}) \mu(d\mathbf{y}) \\ & = \int_{\mathcal{D}} \frac{1}{Z_{B(0,R)}(\mathbf{y})} \sum_{n=0}^{\infty} \frac{\dot{z}^n}{n!} \sum_{m=0}^{\infty} \frac{\dot{z}^m}{m!} \\ & \quad \int_{B(0,R)^n} \int_{B(0,R)^m} F_{\Theta}^{\mathbf{y}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) \mathbf{1}_{\mathcal{D}}(\mathbf{y}_{B(0,R)^c} \dot{\mathbf{x}}\dot{\mathbf{x}}) \otimes_{k=1}^m d\dot{x}_k \otimes_{i=1}^n d\dot{x}_i \mu(d\mathbf{y}). \end{aligned}$$

Recall that  $\dot{B}_R^{\mathbf{y}}$  and  $\dot{B}_R^{\mathbf{y}}$  are the parts of  $B(0, R)$  where a sphere or a particle can be put without conflict with the outside configuration  $\mathbf{y}_{B(0,R)}$  (see (2.1)):

$$\begin{aligned} & \int_{\mathcal{D}} P(X^{\mathbf{y},R} \in \Theta) \mu(d\mathbf{y}) \\ & = \int_{\mathcal{D}} \frac{1}{Z_{B(0,R)}(\mathbf{y})} \sum_{n=0}^{\infty} \frac{\dot{z}^n}{n!} \sum_{m=0}^{\infty} \frac{\dot{z}^m}{m!} \int_{(\dot{B}_R^{\mathbf{y}})^n} \int_{(\dot{B}_R^{\mathbf{y}})^m} F_{\Theta}^{\mathbf{y}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) \mathbf{1}_{\mathcal{D}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) \otimes_{k=1}^m d\dot{x}_k \otimes_{i=1}^n d\dot{x}_i \mu(d\mathbf{y}). \end{aligned}$$

According to the definitions (2.6), (2.4), (2.3) of the measures  $Q_R^{\mathbf{y}}$ ,  $Q_{R,n,m}^{\mathbf{y}}$ ,  $\nu_{R,n,m}^{\mathbf{y}}$ , we have

$$\begin{aligned} Q_R^{\mathbf{y}}(\Theta) & = \frac{1}{Z_R^{\mathbf{y}}} \sum_{n=0}^{+\infty} \frac{\dot{z}^n}{n!} \sum_{m=0}^{+\infty} \frac{\dot{z}^m}{m!} \int P(\mathbf{X}^{\mathbf{y},R,n,m}(\mathbf{x}, \cdot) \in \Theta) d\nu_{R,n,m}^{\mathbf{y}}(\mathbf{x}) \\ & = \frac{1}{Z_R^{\mathbf{y}}} \sum_{n=0}^{+\infty} \frac{\dot{z}^n}{n!} \sum_{m=0}^{+\infty} \frac{\dot{z}^m}{m!} \int_{\mathbb{R}^{nd}} \int_{\mathbb{R}^{md}} F_{\Theta}^{\mathbf{y}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) \mathbf{1}_{\mathcal{D}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) \otimes_{i=1}^n \dot{\lambda}_R^{\mathbf{y}}(d\dot{x}_i) \otimes_{k=1}^m \dot{\lambda}_R^{\mathbf{y}}(d\dot{x}_k). \end{aligned}$$

Distributing the terms according to the number of spheres and particles in  $B(0, R)$ , we get the total variation distance:

$$\mathbf{d}_{TV}(R) = \sup_{\Theta} \left| \int_{\mathcal{D}} \sum_{n=0}^{\infty} \frac{\dot{z}^n}{n!} \sum_{m=0}^{\infty} \frac{\dot{z}^m}{m!} \int_{\mathbb{R}^{nd}} \int_{\mathbb{R}^{md}} F_{\Theta}^{\mathbf{y}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) \mathbf{1}_{\mathcal{D}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) \Delta(\mathbf{y}, \dot{\mathbf{x}}\dot{\mathbf{x}}) \otimes_{k=1}^m d\dot{x}_k \otimes_{i=1}^n d\dot{x}_i \mu(d\mathbf{y}) \right|,$$

where

$$\Delta(\mathbf{y}, \dot{\mathbf{x}}\dot{\mathbf{x}}) := \frac{\prod_{i=1}^n \mathbb{1}_{\dot{x}_i \in \dot{B}_R^{\mathbf{y}}}}{Z_{B(0,R)}(\mathbf{y})} \prod_{k=1}^m \mathbb{1}_{\dot{x}_k \in \dot{B}_R^{\mathbf{y}}} - \frac{e^{-\sum_{i=1}^n \dot{\psi}_R^{\mathbf{y}}(\dot{x}_i)} e^{-\sum_{k=1}^m \dot{\psi}_R^{\mathbf{y}}(\dot{x}_k)}}{\mathcal{Z}_R^{\mathbf{y}}}.$$

Using the fact that  $|F_{\Theta}^{\mathbf{y}}(\dot{\mathbf{x}}\dot{\mathbf{x}})|$  is bounded by 1, we get

$$\mathbf{d}_{TV}(R) \leq \int_{\mathcal{D}} \sum_{n=0}^{\infty} \frac{\dot{z}^n}{n!} \sum_{m=0}^{\infty} \frac{\dot{z}^m}{m!} \int_{\mathbb{R}^{nd}} \int_{\mathbb{R}^{md}} \mathbb{1}_{\mathcal{D}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) |\Delta(\mathbf{y}, \dot{\mathbf{x}}\dot{\mathbf{x}})| \otimes_{k=1}^m d\dot{x}_k \otimes_{i=1}^n d\dot{x}_i \mu(d\mathbf{y}).$$

Note that the terms are well-defined even for  $n = 0$  and/or  $m = 0$ , with the convention that  $\prod_{i=1}^0 \cdots = 1$  and  $\sum_{i=1}^0 \cdots = 0$ . We split the difference  $\Delta(\mathbf{y}, \dot{\mathbf{x}}\dot{\mathbf{x}})$  in two:

$$\begin{aligned} |\Delta(\mathbf{y}, \dot{\mathbf{x}}\dot{\mathbf{x}})| &\leq \prod_{i=1}^n \mathbb{1}_{\dot{x}_i \in \dot{B}_R^{\mathbf{y}}} \prod_{k=1}^m \mathbb{1}_{\dot{x}_k \in \dot{B}_R^{\mathbf{y}}} \left| \frac{1}{Z_{B(0,R)}(\mathbf{y})} - \frac{1}{\mathcal{Z}_R^{\mathbf{y}}} \right| \\ &\quad + \frac{1}{\mathcal{Z}_R^{\mathbf{y}}} \left| \prod_{i=1}^n \mathbb{1}_{\dot{x}_i \in \dot{B}_R^{\mathbf{y}}} \prod_{k=1}^m \mathbb{1}_{\dot{x}_k \in \dot{B}_R^{\mathbf{y}}} - e^{-\sum_{i=1}^n \dot{\psi}_R^{\mathbf{y}}(\dot{x}_i)} e^{-\sum_{k=1}^m \dot{\psi}_R^{\mathbf{y}}(\dot{x}_k)} \right|. \end{aligned}$$

Since  $\dot{\psi}_R^{\mathbf{y}}$  vanishes on  $\dot{B}_R^{\mathbf{y}}$  and  $\dot{\psi}_R^{\mathbf{y}}$  vanishes on  $\dot{B}_R^{\mathbf{y}}$ , the argument of the second absolute value is negative:

$$\prod_{i=1}^n \mathbb{1}_{\dot{x}_i \in \dot{B}_R^{\mathbf{y}}} \prod_{k=1}^m \mathbb{1}_{\dot{x}_k \in \dot{B}_R^{\mathbf{y}}} \leq e^{-\sum_{i=1}^n \dot{\psi}_R^{\mathbf{y}}(\dot{x}_i)} e^{-\sum_{k=1}^m \dot{\psi}_R^{\mathbf{y}}(\dot{x}_k)},$$

and the argument of the first absolute value in the bound of  $|\Delta(\mathbf{y}, \dot{\mathbf{x}}\dot{\mathbf{x}})|$  is positive:

$$\begin{aligned} Z_{B(0,R)}(\mathbf{y}) &= \sum_{n=0}^{+\infty} \frac{\dot{z}^n}{n!} \sum_{m=0}^{+\infty} \frac{\dot{z}^m}{m!} \int_{\mathbb{R}^{nd}} \int_{\mathbb{R}^{md}} \prod_{i=1}^n \mathbb{1}_{\dot{x}_i \in \dot{B}_R^{\mathbf{y}}} \prod_{k=1}^m \mathbb{1}_{\dot{x}_k \in \dot{B}_R^{\mathbf{y}}} \mathbb{1}_{\mathcal{D}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) \otimes_{i=1}^n d\dot{x}_i \otimes_{k=1}^m d\dot{x}_k \\ &\leq \sum_{n=0}^{+\infty} \frac{\dot{z}^n}{n!} \sum_{m=0}^{+\infty} \frac{\dot{z}^m}{m!} \int_{\mathbb{R}^{nd}} \int_{\mathbb{R}^{md}} \mathbb{1}_{\mathcal{D}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) \otimes_{i=1}^n \dot{\lambda}_R^{\mathbf{y}}(d\dot{x}_i) \otimes_{k=1}^m \dot{\lambda}_R^{\mathbf{y}}(d\dot{x}_k) = \mathcal{Z}_R^{\mathbf{y}}. \end{aligned}$$

Consequently, the first part in the upper bound of  $|\Delta(\mathbf{y}, \dot{\mathbf{x}}\dot{\mathbf{x}})|$  yields  $Z_{B(0,R)}(\mathbf{y})$  and the second part yields  $\mathcal{Z}_R^{\mathbf{y}} - Z_{B(0,R)}(\mathbf{y})$ :

$$\begin{aligned} \mathbf{d}_{TV}(R) &\leq \int_{\mathcal{D}} \sum_{n=0}^{+\infty} \frac{\dot{z}^n}{n!} \sum_{m=0}^{+\infty} \frac{\dot{z}^m}{m!} \int_{(\dot{B}_R^{\mathbf{y}})^n} \int_{(\dot{B}_R^{\mathbf{y}})^m} \mathbb{1}_{\mathcal{D}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) \left( \frac{1}{Z_{B(0,R)}(\mathbf{y})} - \frac{1}{\mathcal{Z}_R^{\mathbf{y}}} \right) \otimes_{k=1}^m d\dot{x}_k \otimes_{i=1}^n d\dot{x}_i \\ &\quad + \frac{1}{\mathcal{Z}_R^{\mathbf{y}}} \int_{\mathbb{R}^{nd}} \int_{\mathbb{R}^{md}} \mathbb{1}_{\mathcal{D}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) \left( e^{-\sum_{i=1}^n \dot{\psi}_R^{\mathbf{y}}(\dot{x}_i)} e^{-\sum_{k=1}^m \dot{\psi}_R^{\mathbf{y}}(\dot{x}_k)} - \prod_{i=1}^n \mathbb{1}_{\dot{x}_i \in \dot{B}_R^{\mathbf{y}}} \prod_{k=1}^m \mathbb{1}_{\dot{x}_k \in \dot{B}_R^{\mathbf{y}}} \right) \\ &\quad \otimes_{k=1}^m d\dot{x}_k \otimes_{i=1}^n d\dot{x}_i \mu(d\mathbf{y}) \\ &\leq \int_{\mathcal{D}} \left( \frac{1}{Z_{B(0,R)}(\mathbf{y})} - \frac{1}{\mathcal{Z}_R^{\mathbf{y}}} \right) Z_{B(0,R)}(\mathbf{y}) + \frac{1}{\mathcal{Z}_R^{\mathbf{y}}} \left( \mathcal{Z}_R^{\mathbf{y}} - Z_{B(0,R)}(\mathbf{y}) \right) \mu(d\mathbf{y}) \\ &\leq 2 \int_{\mathcal{D}} \frac{\mathcal{Z}_R^{\mathbf{y}} - Z_{B(0,R)}(\mathbf{y})}{\mathcal{Z}_R^{\mathbf{y}}} \mu(d\mathbf{y}). \end{aligned} \tag{3.4}$$



To prove (3.3), we only have find a summable upper bound for the above integral. Since  $\psi_R^y(\dot{x}_i) = 0$  for  $\dot{x}_i \in \dot{B}_R^y$ , and  $\psi_R^y(\dot{x}_k) = 0$  for  $\dot{x}_k \in \dot{B}_R^y$ , we have

$$1 - e^{\sum_{i=1}^n \psi_R^y(\dot{x}_i)} e^{\sum_{k=1}^m \psi_R^y(\dot{x}_k)} \prod_{i=1}^n \mathbb{1}_{\dot{x}_i \in \dot{B}_R^y} \prod_{k=1}^m \mathbb{1}_{\dot{x}_k \in \dot{B}_R^y} \leq \sum_{i=1}^n \mathbb{1}_{\dot{x}_i \notin \dot{B}_R^y} + \sum_{k=1}^m \mathbb{1}_{\dot{x}_k \notin \dot{B}_R^y}.$$

Using the exchangeability of the spheres and particles, we obtain

$$\begin{aligned} & \mathcal{Z}_R^y - Z_{B(0,R)}(\mathbf{y}) \\ & \leq \sum_{n=0}^{+\infty} \frac{z^n}{n!} \sum_{m=0}^{+\infty} \frac{z^m}{m!} \int_{\mathbb{R}^{nd}} \int_{\mathbb{R}^{md}} \mathbb{1}_{\mathcal{D}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) \left( n \mathbb{1}_{\dot{x}_n \notin \dot{B}_R^y} + m \mathbb{1}_{\dot{x}_m \notin \dot{B}_R^y} \right) \otimes_{i=1}^n \dot{\lambda}_R^y(d\dot{x}_i) \otimes_{k=1}^m \dot{\lambda}_R^y(d\dot{x}_k). \end{aligned}$$

Since  $\mathbb{1}_{\mathcal{D}}(\dot{x}_1 \dots \dot{x}_n \dot{x}_1 \dots \dot{x}_m) \leq \mathbb{1}_{\mathcal{D}}(\dot{x}_1 \dots \dot{x}_{n-1} \dot{x}_1 \dots \dot{x}_m)$ , and analogously for the particles,

$$\begin{aligned} & \mathcal{Z}_R^y - Z_{B(0,R)}(\mathbf{y}) \\ & \leq z \sum_{n=1}^{+\infty} \frac{z^{n-1}}{(n-1)!} \sum_{m=0}^{+\infty} \frac{z^m}{m!} \int_{\mathbb{R}^{(n-1)d}} \int_{\mathbb{R}^{md}} \mathbb{1}_{\mathcal{D}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) \otimes_{i=1}^{n-1} \dot{\lambda}_R^y(d\dot{x}_i) \otimes_{k=1}^m \dot{\lambda}_R^y(d\dot{x}_k) \dot{\lambda}_R^y((\dot{B}_R^y)^c) \\ & \quad + z \sum_{n=0}^{+\infty} \frac{z^n}{n!} \sum_{m=1}^{+\infty} \frac{z^{m-1}}{(m-1)!} \int_{\mathbb{R}^{nd}} \int_{\mathbb{R}^{(m-1)d}} \mathbb{1}_{\mathcal{D}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) \otimes_{i=1}^n \dot{\lambda}_R^y(d\dot{x}_i) \otimes_{k=1}^{m-1} \dot{\lambda}_R^y(d\dot{x}_k) \dot{\lambda}_R^y((\dot{B}_R^y)^c) \\ & \leq \mathcal{Z}_R^y \left( z \dot{\lambda}_R^y((\dot{B}_R^y)^c) + z \dot{\lambda}_R^y((\dot{B}_R^y)^c) \right). \end{aligned}$$

Injecting this in (3.4) yields

$$\mathbf{d}_{TV}(R) \leq 2 \int_{\mathcal{D}} z \dot{\lambda}_R^y((\dot{B}_R^y)^c) + z \dot{\lambda}_R^y((\dot{B}_R^y)^c) \mu(d\mathbf{y}). \quad (3.5)$$

Finally, by assumption (2.2), we have that  $\sum_R \mathbf{d}_{TV}(R) < +\infty$ , which in turn implies that

$$\int_{\mathcal{D}} P((\Omega^y)^c) \mu(d\mathbf{y}) = 0, \text{ concluding the proof of Proposition 3.4.}$$

As a result, the solution  $X^{\mathbf{y},\infty}(\omega, \cdot)$  of  $(\mathcal{S}_\infty)$  constructed in Section 3.1 as a limit process exists for  $\mu$ -almost every  $\omega$ , for any fixed  $\mathbf{y}$  in the interior of  $\mathcal{D}$ .

**Remark 3.9.** The process  $X^{\mathbf{y},\infty}(\omega, \cdot)$  is the unique solution of  $(\mathcal{S}_\infty)$  in the sense of [Tan96, Lemma 5.4]. This means that  $X^{\mathbf{y},\infty}$  coincides  $\mu$ -a.s. with any solution  $X$  of  $(\mathcal{S}_\infty)$  starting from  $\mathbf{y}$  in the class of paths for which: there exists  $\varepsilon > 0$  and a rate  $p \in \mathbb{N}^*$  such that, for all  $\rho > 0$  and infinitely-many  $m \in \mathbb{N}^*$ , there exists a finite sequence of rational times  $t_0 = 0 < t_1 < \dots < t_{m'-1} < t_{m'} = 1$ , with  $t_{a+1} - t_a \leq \frac{1}{m}$ , and a sequence of bounded sets  $C_0, \dots, C_{m'-1}$  in  $\mathbb{R}^d$ , with  $B(0, \rho + m) \subset C_{m'-1} \subset \dots \subset C_0 \subset B(0, \rho + m + m^p)$ , with  $B(C_{a+1}, \varepsilon) \subset C_a$  for each  $a$ , such that, for every  $a \in \{0, \dots, m' - 1\}$ , the boundary of  $C_a$  separates balls in such a way that they do not interact:

$$\begin{aligned} \forall a \in \{0, 1, \dots, m' - 1\}, \quad \forall t \in \left[ \frac{a}{m}, \frac{a+1}{m} \right], \quad \inf\{|\dot{X}_i(t) - x| : i \in \mathbb{N}^*, x \in \partial C_a\} > \dot{r} + \frac{\varepsilon}{4} \\ \text{and } \inf\{|\dot{X}_k(t) - x| : k \in \mathbb{N}^*, x \in \partial C_a\} > \dot{r} + \frac{\varepsilon}{4}. \end{aligned}$$

That the limit solution  $X^{\mathbf{y},\infty}$  belongs to this class follows from the computations of Section 3.1. Any solution belonging to this class inherits the uniqueness property of the finite SDE, since every sphere or particle belongs on successive time intervals to finite sets of interacting balls, hence they coincide with  $X^{\mathbf{y},\infty}(\omega, \cdot)$ .

### 3.3 Reversibility of the limit under Gibbs measures

To complete the proof of Theorem 1.2, we only have to note that the limit process  $X^{\mathbf{y},\infty}(\omega, \cdot)$ , constructed in Sections 3.1 and 3.2 for  $\mu$ -almost every  $\omega$ , admits  $\mu$  as a reversible measure, for any Gibbs measure  $\mu \in \mathcal{G}_{\dot{z},\dot{z}}$ .

This is a straightforward consequence of the reversibility of the penalised processes under the approximate measures  $Q_R^{\mathbf{y}}$ . Indeed, for any time  $T$  in  $[0, 1]$ , any subdivision  $0 \leq t_1 < \dots < t_j \leq T$ , and any bounded continuous local functions  $F_1, \dots, F_j$  on  $\mathcal{M}$ , we have

$$\begin{aligned} & \left| \int_{\mathcal{D}} E \left( \prod_{i=1}^j F_i(X^{\mathbf{y},\infty}(T - t_i)) - \prod_{i=1}^j F_i(X^{\mathbf{y},\infty}(t_i)) \right) \mu(d\mathbf{y}) \right| \\ &= \lim_{R \rightarrow +\infty} \left| \int_{\mathcal{D}} E \left( \prod_{i=1}^j F_i(X^{\mathbf{y},R}(T - t_i)) - \prod_{i=1}^j F_i(X^{\mathbf{y},R}(t_i)) \right) \mu(d\mathbf{y}) \right| \\ &\leq \lim_{R \rightarrow +\infty} 2 \prod_{i=1}^j \|F_i\|_{\infty} \mathbf{d}_{TV}(R) = 0, \end{aligned}$$

since, thanks to the reversibility of  $X^{\mathbf{y},R}$  starting from  $\mu_R^{\mathbf{y}}$ ,

$$\int_{\mathcal{D}} E \left( \prod_{i=1}^j F_i(X^{\mathbf{y},R}(T - t_i)) - \prod_{i=1}^j F_i(X^{\mathbf{y},R}(t_i)) \right) \mu_R^{\mathbf{y}}(d\mathbf{y}) = 0.$$

## 4 Occurrence of a depletion interaction between hard spheres

In this section, we study the emergence of an attractive interaction, the *depletion interaction*, between hard spheres, due to the presence of the particle medium. We first identify the projection  $\hat{\mu}$  of the reversible measure  $\mu$  introduced in (1.1), as a Gibbs measure with a new effective interaction. We then construct a gradient dynamics whose reversible measure is given by  $\hat{\mu}$ . Finally, in Sections 4.3 and 4.4, we study the properties of the Gibbs measures  $\hat{\mu}$  in the low- and high-activity regimes, respectively.

### 4.1 The projection of the two-type reversible measure

We study here the projection of the reversible measure  $\mu \in \mathcal{G}_{\dot{z},\dot{z}}$  onto the subsystem of hard spheres. For any finite configuration  $\dot{\mathbf{x}} = \{\dot{x}_1, \dots, \dot{x}_n\}$ ,  $n \geq 1$ , of hard spheres, consider the energy

$$\mathcal{E}(\dot{\mathbf{x}}) := |\mathbb{B}(\dot{\mathbf{x}})| = \left| \bigcup_{i=1}^n B(\dot{x}_i, \dot{r}) \right|.$$

We denote by  $\mathcal{G}_{\dot{z}}(\dot{z}\mathcal{E})$  the set of Gibbs measures associated to  $\mathcal{E}$  with activity  $\dot{z}$  and inverse temperature  $\dot{z}$ , that is the set of measures solutions of

$$\int_{\mathcal{D}} F(\dot{\mathbf{x}}) \hat{\mu}(d\dot{\mathbf{x}}) = \int_{\mathcal{D}} \frac{1}{\dot{Z}_{\Lambda}(\dot{\mathbf{y}})} \int_{\dot{\mathcal{M}}} F(\dot{\mathbf{y}}_{\Lambda^c} \dot{\mathbf{x}}) \exp(-\dot{z}\mathcal{E}_{\Lambda}(\dot{\mathbf{x}}\dot{\mathbf{y}}_{\Lambda^c})) \mathbf{1}_{\mathcal{D}}(\dot{\mathbf{y}}_{\Lambda^c} \dot{\mathbf{x}}) \pi_{\Lambda}^{\dot{z}}(d\dot{\mathbf{x}}) \hat{\mu}(d\dot{\mathbf{y}}), \quad (4.1)$$

for any  $\Lambda \subset \mathbb{R}^d$  and any positive bounded measurable  $F$  on  $\mathring{\mathcal{M}}$ , where

$$\mathcal{E}_\Lambda(\mathring{\mathbf{x}}) := |\mathbb{B}(\mathring{\mathbf{x}}_\Lambda) \setminus \mathbb{B}(\mathring{\mathbf{x}}_{\Lambda^c})| = \left| \bigcup_{\mathring{x} \in \mathring{\mathbf{x}}_\Lambda} B(\mathring{x}, \mathring{r}) \setminus \bigcup_{\mathring{x} \in \mathring{\mathbf{x}}_{\Lambda^c}} B(\mathring{x}, \mathring{r}) \right|. \quad (4.2)$$

is the conditional energy in  $\Lambda$  for (infinite) configurations of hard spheres  $\mathring{\mathbf{x}} \in \mathring{\mathcal{M}}$ .

**Remark 4.1.** It is worth noting that several choices of conditional energy lead to the same set of Gibbs measures. This is the case, for example, for  $\tilde{\mathcal{E}}_\Lambda(\mathring{\mathbf{x}}) := |\mathbb{B}(\mathring{\mathbf{x}}) \cap \Lambda|$ , since it differs from  $\mathcal{E}_\Lambda$  only for factors that only depend on the external configuration.

Similarly to the Widom–Rowlinson model [WR70], the following correspondence between two-type and one-type models holds, where the intensity  $\dot{z}$  of the particles plays the role of *inverse temperature* for the effective interaction between hard spheres.

**Proposition 4.2.** For any  $\dot{z}, z > 0$ ,

- If  $\mu \in \mathcal{G}_{z, \dot{z}}$  is a Gibbs measure on  $\mathcal{M}$ , then its marginal on  $\mathring{\mathcal{M}}$  is a Gibbs measure  $\dot{\mu}_z \in \mathcal{G}_{\dot{z}}(z\mathcal{E})$ .
- If  $\dot{\mu} \in \mathcal{G}_{\dot{z}}(z\mathcal{E})$  is a Gibbs measure on  $\mathring{\mathcal{M}}$ , then  $\dot{\mu}(d\mathring{\mathbf{x}}) \otimes \pi_{\mathbb{R}^d \setminus \mathbb{B}(\mathring{\mathbf{x}})}^{\dot{z}}(d\mathring{\mathbf{x}}) \in \mathcal{G}_{z, \dot{z}}$  is a Gibbs measure on  $\mathcal{M}$ .

The proof is relatively straightforward, but it is still worthwhile to include here. Indeed, in the Widom–Rowlinson model this result is folklore, and as such a rigorous proof using the DLR formalism is not easily found. Moreover, as far as we are aware, this is the first time this argument is extended to this setting where there is an additional hard-core interaction.

*Proof.* We first show that if  $\dot{\mu}$  satisfies the DLR equations (1.2) for  $\mathcal{E}$ , then  $\mu := \dot{\mu} \otimes \pi_{\mathbb{R}^d \setminus \mathbb{B}(\mathring{\mathbf{x}})}^{\dot{z}}$  satisfies the DLR equations (1.1). For any positive bounded measurable  $F$  on  $\mathcal{D}$ , we have

$$\begin{aligned} \int F(\mathbf{x}) \mu(d\mathbf{x}) &:= \iint F(\mathring{\mathbf{x}}\mathring{\mathbf{x}}) \pi_{\mathbb{R}^d \setminus \mathbb{B}(\mathring{\mathbf{x}})}^{\dot{z}}(d\mathring{\mathbf{x}}) \dot{\mu}(d\mathring{\mathbf{x}}) \\ &= \iiint F(\mathring{\mathbf{x}}'_\Lambda \mathring{\mathbf{x}}'_\Lambda \mathring{\mathbf{x}}_{\Lambda^c} \mathring{\mathbf{x}}_{\Lambda^c}) \frac{e^{-z|\mathbb{B}(\mathring{\mathbf{x}}'_\Lambda) \setminus \mathbb{B}(\mathring{\mathbf{x}}_{\Lambda^c})|}}{Z_\Lambda(\mathring{\mathbf{x}}_{\Lambda^c})} \pi_{\mathbb{B}(\mathring{\mathbf{x}}'_\Lambda \mathring{\mathbf{x}}_{\Lambda^c})^c}^{\dot{z}}(d\mathring{\mathbf{x}}) \mathbb{1}_{\mathcal{D}}(\mathring{\mathbf{x}}'_\Lambda \mathring{\mathbf{x}}_{\Lambda^c}) \pi_\Lambda^{\dot{z}}(d\mathring{\mathbf{x}}'_\Lambda) \dot{\mu}(d\mathring{\mathbf{x}}_{\Lambda^c}). \end{aligned}$$

Writing  $e^{-|\mathbb{B}(\mathring{\mathbf{x}}'_\Lambda) \setminus \mathbb{B}(\mathring{\mathbf{x}}_{\Lambda^c})|} = e^{-|\mathbb{B}(\mathring{\mathbf{x}}'_\Lambda) \cup \mathbb{B}(\mathring{\mathbf{x}}_{\Lambda^c})|} e^{|\mathbb{B}(\mathring{\mathbf{x}}_{\Lambda^c})|}$ , and absorbing the second term into  $Z_\Lambda(\mathbf{x}_{\Lambda^c})$ , we have that this is precisely the Poisson void probability for  $\pi_\Lambda^{\dot{z}} \pi_{\Lambda^c}^{\dot{z}}$ , so that we find

$$\int F(\mathbf{x}) \mu(d\mathbf{x}) = \int \frac{1}{Z_\Lambda(\mathbf{x}_{\Lambda^c})} \left( \iint F(\mathring{\mathbf{x}}'_\Lambda \mathring{\mathbf{x}}_{\Lambda^c}) \mathbb{1}_{\mathcal{D}}(\mathring{\mathbf{x}}'_\Lambda \mathring{\mathbf{x}}_{\Lambda^c}) \pi_\Lambda^{\dot{z}}(d\mathring{\mathbf{x}}'_\Lambda) \pi_\Lambda^{\dot{z}}(\mathring{\mathbf{x}}') \right) \mu(d\mathbf{x}).$$

Conversely, let  $\mu \in \mathcal{G}_{z, \dot{z}}$ . Its marginal on the hard spheres is characterised by integration over test functions  $F$  supported on  $\mathring{\mathcal{M}}$ , that is:  $\int_{\mathring{\mathcal{M}}} F(\mathring{\mathbf{x}}) \dot{\mu}_z(d\mathring{\mathbf{x}}) := \int_{\mathcal{M}} F(\mathring{\mathbf{x}}) \mu(d\mathring{\mathbf{x}}\mathring{\mathbf{x}})$ . The DLR equations for  $\mu$  then yield

$$\begin{aligned} \int F(\mathring{\mathbf{x}}) \mu(d\mathring{\mathbf{x}}\mathring{\mathbf{x}}) &= \int \frac{1}{Z_\Lambda(\mathbf{y})} \int_{\mathring{\mathcal{M}}} \int_{\mathring{\mathcal{M}}} F(\mathring{\mathbf{y}}_{\Lambda^c} \mathring{\mathbf{x}}) \mathbb{1}_{\mathcal{D}}(\mathring{\mathbf{y}}_{\Lambda^c} \mathring{\mathbf{y}}_{\Lambda^c} \mathring{\mathbf{x}}\mathring{\mathbf{x}}) \pi_\Lambda^{\dot{z}}(d\mathring{\mathbf{x}}) \pi_\Lambda^{\dot{z}}(d\mathring{\mathbf{x}}) \mu(d\mathbf{y}) \\ &= \int \frac{1}{Z_\Lambda(\mathbf{y})} \int_{\mathring{\mathcal{M}}} \int_{\mathring{\mathcal{M}}} F(\mathring{\mathbf{y}}_{\Lambda^c} \mathring{\mathbf{x}}) \mathbb{1}_{\mathcal{D}}(\mathbf{y}_{\Lambda^c} \mathring{\mathbf{x}}\mathring{\mathbf{x}}) \pi_\Lambda^{\dot{z}}(d\mathring{\mathbf{x}}) \pi_\Lambda^{\dot{z}}(d\mathring{\mathbf{x}}) \mu(d\mathbf{y}) \\ &= \int \frac{1}{Z_\Lambda(\mathring{\mathbf{y}})} \int_{\mathring{\mathcal{M}}} F(\mathring{\mathbf{y}}_{\Lambda^c} \mathring{\mathbf{x}}) \exp(-z\mathcal{E}_\Lambda(\mathring{\mathbf{x}}\mathring{\mathbf{y}}_{\Lambda^c})) \mathbb{1}_{\mathcal{D}}(\mathring{\mathbf{y}}_{\Lambda^c} \mathring{\mathbf{x}}) \pi_\Lambda^{\dot{z}}(d\mathring{\mathbf{x}}) \mu(d\mathring{\mathbf{y}}\mathring{\mathbf{y}}), \end{aligned}$$



### 4.3 Low-activity regime: no sphere-percolation

As a by-product of the chain estimate of Lemma 3.7, needed to control the dynamics and prove the existence of solutions for  $(\mathcal{S}_\infty)$ , we get the absence of percolation  $\dot{\mu}$ -a.s. for small  $\dot{z}$ . Although our argument is not the classical one in the line of [TMLS97], it still relies on the crucial idea that there is only a very small area in which an additional sphere can be put in order to lengthen a sphere chain, hence a very small, geometrically decreasing, probability to have long chains.

**Proposition 4.3.** We say that two spheres  $\dot{x}_i$  and  $\dot{x}_j$  of a configuration  $\dot{\mathbf{x}}$  *interact* if  $|\dot{x}_i - \dot{x}_j| < 2\dot{r}$ , i.e. if their interaction  $\mathcal{V}_{\text{overlap}}\left(\frac{|\dot{x}_i - \dot{x}_j|}{2\dot{r}}\right)$  does not vanish. Let  $\dot{\mu} \in \mathcal{G}_{\dot{z}}(\dot{z}\mathcal{E})$ . Then, for any  $\dot{z} > 0$  and any  $\dot{z} < (2^d(\dot{r}^d - \dot{r}^d)\mathbf{v}_d)^{-1}$ , there is  $\dot{\mu}$ -a.s. no infinite cluster of interacting spheres.

*Proof.* If a configuration  $\dot{\mathbf{y}}$  has an infinite chain of interacting spheres at distance smaller than  $2\dot{r}$  of one another, there is a radius  $R(\dot{\mathbf{y}})$  such that  $B(0, R)$  intersects this infinite chain for each  $R > R(\dot{\mathbf{y}})$ . Consequently,  $\dot{\mathbf{y}}$  has a chain of interacting spheres from  $B(0, R)$  to  $B(0, 2R)$  which involves at least  $\left\lfloor \frac{R}{2\dot{r}} \right\rfloor$  spheres. We then only have to prove that

$$\dot{\mu}(C_\infty) = 0, \quad \text{where } C_\infty := \left\{ \text{For } R \text{ large enough, } \dot{\mathbf{y}}_{B(0, 2R)} \in \mathcal{B}_{\text{Chain}}\left(R, \left\lfloor \frac{R}{2\dot{r}} \right\rfloor, 2\dot{r}\right) \right\}.$$

This holds as soon as

$$\sum_R \dot{\mu}\left(\dot{\mathbf{y}}_{B(0, 2R)} \in \mathcal{B}_{\text{Chain}}\left(R, \left\lfloor \frac{R}{2\dot{r}} \right\rfloor, 2\dot{r}\right)\right) < +\infty. \quad (4.3)$$

By Proposition 4.2,  $\dot{\mu}$  is the projection on the sphere configuration space  $\dot{\mathcal{M}}$  of the Gibbs measure  $\dot{\mu}(d\dot{\mathbf{x}}) \otimes \pi_{\mathbb{R}^d \setminus \mathbb{B}(\dot{\mathbf{x}})}^{\dot{z}}(d\dot{\mathbf{x}}) \in \mathcal{G}_{\dot{z}, \dot{z}}$  on  $\mathcal{M}$ . Since  $\mathcal{B}_{\text{Chain}}$  events only involve spheres, and  $X^{\mathbf{y}, 2R}(0) = \mathbf{y}_{B(0, 2R)}$  by (2.7) is  $\mu_{2R}^{\mathbf{y}}$ -distributed under  $Q_{2R}^{\mathbf{y}}$ , we have

$$\begin{aligned} \dot{\mu}\left(\dot{\mathbf{y}}_{B(0, 2R)} \in \mathcal{B}_{\text{Chain}}\left(R, \left\lfloor \frac{R}{2\dot{r}} \right\rfloor, 2\dot{r}\right)\right) &= \mu\left(\mathbf{y}_{B(0, 2R)} \in \mathcal{B}_{\text{Chain}}\left(R, \left\lfloor \frac{R}{2\dot{r}} \right\rfloor, 2\dot{r}\right)\right) \\ &= \int_{\mathcal{D}} P\left(X^{\mathbf{y}, 2R}(0) \in \mathcal{B}_{\text{Chain}}\left(R, \left\lfloor \frac{R}{2\dot{r}} \right\rfloor, 2\dot{r}\right)\right) \mu(d\mathbf{y}) \\ &\leq \int_{\mathcal{D}} Q_{2R}^{\mathbf{y}}\left(\mathcal{B}'_{\text{Chain}}\left(1, R, \left\lfloor \frac{R}{2\dot{r}} \right\rfloor, 2\dot{r}\right)\right) \mu(d\mathbf{y}) + \mathbf{d}_{TV}(2R). \end{aligned}$$

Thanks to (3.5) and Lemma 3.7, we obtain

$$\begin{aligned} &\dot{\mu}\left(\dot{\mathbf{y}}_{B(0, 2R)} \in \mathcal{B}_{\text{Chain}}\left(R, \left\lfloor \frac{R}{2\dot{r}} \right\rfloor, 2\dot{r}\right)\right) \\ &\leq \left(\dot{z}((2\dot{r})^d - (2\dot{r})^d)\mathbf{v}_d\right)^{\left\lfloor \frac{R}{2\dot{r}} \right\rfloor} + 2 \int_{\mathcal{D}} \dot{z} \dot{\lambda}_R^{\mathbf{y}}((\dot{B}_R^{\mathbf{y}})^c) + \dot{z} \dot{\lambda}_R^{\mathbf{y}}((\dot{B}_R^{\mathbf{y}})^c) \mu(d\mathbf{y}). \end{aligned}$$

This ensures that (4.3) holds as soon as  $\dot{z} < \frac{1}{((2\dot{r})^d - (2\dot{r})^d)\mathbf{v}_d}$ , completing the proof.  $\square$

**Remark 4.4.** Although the connection between (absence of) percolation and (absence of) phase transition is not automatic – in particular, FKG does not hold and, since this is not a symmetric mixture, a phase transition cannot be proven via symmetry breaking using percolation as in [CCK95, GLM95] – it is still interesting to compare the different regimes.

Let  $\varphi(y) = v_d \overset{\circ}{r}^d - \frac{1}{2} \mathcal{V}_{\text{overlap}}(|y|/(2\overset{\circ}{r})) \geq 0$ . By the classical result of Ruelle [Rue99, Theorem 4.2.3], we know that there is no phase transition if

$$\dot{z} < e^{-1} \left( (2\overset{\circ}{r})^d v_d + \int_{|y| \in (2\overset{\circ}{r}, 2\overset{\circ}{r})} |1 - e^{-\dot{z}\varphi(y)}| dy \right)^{-1}.$$

Comparing with Proposition 4.3, we see (apart from the  $e^{-1}$  factor) an additional  $(2\overset{\circ}{r})^d$  at the denominator, coming from the hard core. It is however worth noting (see [JT20]) that this should be significantly improved by considering the *renormalised activity*  $\dot{z}' = \dot{z} \exp\{-\dot{z}v_d \overset{\circ}{r}^d\}$ .

## 4.4 High-activity regime: towards an optimal packing

We are now interested in the high-density behaviour of the equilibrium measures in  $\mathcal{G}_z(\dot{z}\mathcal{E})$ .

In the case of a finite number of hard spheres, we were able to prove in [FKRZ24] that the reversible probability measure of the corresponding finite-dimensional gradient system for  $n$  hard spheres concentrates around admissible configurations which minimise the energy. Moreover, admissible  $n$ -sphere configurations who realise the minimal energy maximise the contact number.

In the infinite setting, the behaviour is more complicated, as the number of spheres in any volume is random. There is now a competition between the minimisation of the *energy* (achieved either by having less or more densely packed spheres) and that of the *entropy* (achieved by being as close as possible to the completely random Poisson point process); this is the *Gibbs variational principle* [Geo94, Geo95]. In particular then, since the energy is minimised not only by packing a given number of spheres, but also by taking as few of them as possible, only in the limit  $\dot{z} \rightarrow \infty$ , the entropic cost of not putting a lot of points in the system leads to the packing phenomenon. This can be formalised as follows:

**Proposition 4.5.** Let  $\dot{\mu} \in \mathcal{G}_z(\dot{z}\mathcal{E})$ , and  $\rho_z(\dot{z}) = \lim_{n \rightarrow \infty} \frac{1}{|\Lambda_n|} \int |\dot{\mathbf{x}}_{\Lambda_n}| d\dot{\mu}(\dot{\mathbf{x}})$ ,  $\Lambda_n = [-n, n]^d$ , denote its intensity. Then, for any  $\dot{z} > 0$ , as  $\dot{z} \rightarrow \infty$ ,  $\dot{\mu}$  attains the closest packing density, that is:

$$\lim_{\dot{z} \rightarrow \infty} \rho_z(\dot{z}) = \lim_{n \rightarrow \infty} \frac{1}{|\Lambda_n|} N_n =: \rho^*,$$

with  $N_n$  the maximal number of mutually disjoint open balls of radius  $\overset{\circ}{r}$  included in  $\Lambda_n$ .

The proof is a straightforward application of [MMS<sup>+</sup>01, Proposition 2], which states the above result for any regular potential with hard core, as  $\varphi_2$  is (e.g. [Rue99]), using the Gibbs variational principle as main tool.

# A Appendix: Proof of the remaining statements

## A.1 Existence of the penalisation functions

We prove here the existence of the sphere penalisation function  $\psi_R^y$  and the particle penalisation function  $\psi_R^y$  needed in Section 2.1. Let  $\psi$  be a non-increasing  $C^\infty$  function with  $\psi(t) = 1$  if  $t \leq 0$  and

$\psi(t) = 0$  if  $t \geq 1$ , and  $\varphi$  be a non-decreasing  $C^\infty$  function with  $\varphi(t) = 0$  if  $t \leq 0$  and  $\varphi(t) = t + C^{\text{st}}$  if  $t \geq 1$ . Then the two functions

$$\begin{aligned} \psi_R^{\mathbf{y}}(x) &:= 2 \log(R) + \varphi(R^{d+1}(|x| - R)) + \sum_{\substack{\dot{\mathbf{y}} \in \mathbf{y} \\ R \leq |\dot{\mathbf{y}}| \leq R+2\hat{r}}} \psi\left(\frac{|\dot{\mathbf{y}} - x|^2}{4\hat{r}^2}\right) + \sum_{\substack{\dot{\mathbf{y}} \in \mathbf{y} \\ R \leq |\dot{\mathbf{y}}| \leq R+\hat{r}}} \psi\left(\frac{|\dot{\mathbf{y}} - x|^2}{\hat{r}^2}\right) \\ &\quad + \log\left(\#\{\dot{\mathbf{y}} \in \mathbf{y} : R \leq |\dot{\mathbf{y}}| \leq R + 2\hat{r}\}\right) + \log\left(\#\{\dot{\mathbf{y}} \in \mathbf{y} : R \leq |\dot{\mathbf{y}}| \leq R + \hat{r}\}\right), \\ \psi_R^{\mathbf{y}}(x) &:= 2 \log(R) + \varphi(R^{d+1}(|x| - R)) + \sum_{\substack{\dot{\mathbf{y}} \in \mathbf{y} \\ R \leq |\dot{\mathbf{y}}| \leq R+\hat{r}}} \psi\left(\frac{|\dot{\mathbf{y}} - x|^2}{\hat{r}^2}\right) \\ &\quad + \log\left(\#\{\dot{\mathbf{y}} \in \mathbf{y} : R \leq |\dot{\mathbf{y}}| \leq R + \hat{r}\}\right), \end{aligned}$$

with the convention that  $\log(\#\emptyset) + \sum_{\emptyset} \cdots = 0$ , are non-negative functions of class  $C^2$  with bounded derivatives. Moreover,  $\psi_R^{\mathbf{y}}$  is constant on  $\mathring{B}_R^{\mathbf{y}}$ , and  $\psi_R^{\mathbf{y}}$  is constant on  $\mathring{B}_R^{\mathbf{y}}$ . By construction then,

$$\sum_{R=1}^{\infty} \int_{(\mathring{B}_R^{\mathbf{y}})^c} e^{-\psi_R^{\mathbf{y}}(x)} dx < +\infty \quad \text{and} \quad \sum_{R=1}^{\infty} \int_{(\mathring{B}_R^{\mathbf{y}})^c} e^{-\psi_R^{\mathbf{y}}(x)} dx < +\infty.$$

## A.2 Proof of the ball separation and nested inclusion Lemma 3.6

We consider a path  $X : [0, 1] \rightarrow \mathcal{D}$  that belongs neither to  $\mathcal{B}'_{\text{Chain}}(\delta, \alpha, \kappa, \varepsilon)$  nor to  $\mathcal{B}'_{\text{Fast}}(\alpha, \delta, \varepsilon/4)$ , and study it during the time interval  $[a\delta, (a+1)\delta]$ , on balls of radii

$$0 < \rho =: \rho_{1/\delta} < \cdots < \rho_{a+1} < \rho_a := \rho_{a+1} + 2\kappa(2\hat{r} + \varepsilon) < \cdots < \rho_0 := \rho + \frac{2\kappa}{\delta}(2\hat{r} + \varepsilon) \leq \alpha,$$

focusing first on the spheres, then on the particles.

By definition of the index set  $\mathring{\mathcal{J}} := \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon)$ , the spheres of  $X(a\delta)$  whose indices are not in  $\mathring{\mathcal{J}}$  are neither in the ball  $B(0, \rho_a)$  nor in an  $\varepsilon$ -chain connected to it, i.e. not at distance  $(2\hat{r} + \varepsilon)$  of any  $\varepsilon$ -chain connected to  $B(0, \rho_a)$ , that is:

$$\forall j \notin \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \quad |\dot{X}_j(a\delta)| > \rho_a,$$

$$\forall j \notin \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \quad \forall i \in \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \quad |\dot{X}_j(a\delta) - \dot{X}_i(a\delta)| > 2\hat{r} + \varepsilon.$$

$X \notin \mathcal{B}'_{\text{Fast}}(\alpha, \delta, \varepsilon/4)$  implies that no sphere in  $B(0, \alpha)$  moves by more than  $\varepsilon/4$  during the time interval  $[a\delta, (a+1)\delta]$ , hence the separation property:

$$\forall j \notin \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \quad \forall i \in \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \quad \forall t \in [a\delta, (a+1)\delta], \quad |\dot{X}_j(t) - \dot{X}_i(t)| > 2\hat{r} + \frac{\varepsilon}{2}.$$

Since  $X \notin \mathcal{B}'_{\text{Chain}}(\delta, \alpha, \kappa, \varepsilon)$ , the starting configuration  $X(a\delta)$  does not have any  $\varepsilon$ -chain of spheres that intersects  $B(0, \alpha)$  and involves more than  $(\kappa + 1)$  spheres, i.e. is longer than  $\kappa(2\hat{r} + \varepsilon)$ . Consequently, all spheres labelled in  $\mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon)$  start at a distance from the origin of at most:

$$\forall i \in \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \quad |\dot{X}_i(a\delta)| \leq \underbrace{\rho_a}_{\text{radius defining } \mathring{\mathcal{J}}} + \underbrace{\kappa(2\hat{r} + \varepsilon)}_{\text{longest } \varepsilon\text{-chain}}. \quad (\text{A.1})$$

Since they move at most of  $\varepsilon/4$  during the time interval  $[a\delta, (a+1)\delta]$ , we get the localisation property:

$$\forall i \in \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \forall t \in [a\delta, (a+1)\delta], \quad |\dot{X}_i(t)| \leq \rho_a + \kappa(2\mathring{r} + \varepsilon) + \frac{\varepsilon}{4}.$$

Finally, for each  $j \notin \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon)$ , we already have  $|\dot{X}_j(a\delta)| > \rho_a$ , thus  $|\dot{X}_j((a+1)\delta)| > \rho_a - \frac{\varepsilon}{4}$ :

$$\begin{aligned} j \notin \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon) &\implies |\dot{X}_j((a+1)\delta)| > \rho_a - \frac{\varepsilon}{4} > \rho_{a+1} + \kappa(2\mathring{r} + \varepsilon) \\ &\implies j \notin \mathring{\mathcal{J}}(X((a+1)\delta), \rho_{a+1}, \varepsilon), \end{aligned}$$

thanks to (A.1), used for  $a+1$  instead of  $a$ . This proves the first inclusion:

$$\mathring{\mathcal{J}}(X((a+1)\delta), \rho_{a+1}, \varepsilon) \subset \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon).$$

The proof of the analogue properties for the particles is similar. The particles of  $X(a\delta)$  whose indices do not belong to  $\mathring{\mathcal{J}} := \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon)$  are neither in  $B(0, \rho_a)$  nor in the  $\varepsilon$ -neighbourhood of an  $\varepsilon$ -chain connected to it, i.e. not at distance  $\mathring{r} + \varepsilon$  of any  $\varepsilon$ -chain connected to  $B(0, \rho_a)$ :

$$\forall k \notin \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \quad |\dot{X}_k(a\delta)| > \rho_a,$$

$$\forall k \notin \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \forall i \in \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \quad |\dot{X}_k(a\delta) - \dot{X}_i(a\delta)| > \mathring{r} + \varepsilon.$$

Again, no sphere and no particle in  $B(0, \alpha)$  moves by more than  $\varepsilon/4$  during the time interval, hence

$$\forall k \notin \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \forall t \in [a\delta, (a+1)\delta], \quad |\dot{X}_k(t)| > \rho_a - \frac{\varepsilon}{4}, \quad (\text{A.2})$$

$$\forall k \notin \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \forall i \in \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \forall t \in [a\delta, (a+1)\delta], \quad |\dot{X}_k(t) - \dot{X}_i(t)| > \mathring{r} + \frac{\varepsilon}{2}.$$

Particles in  $\mathring{\mathcal{J}}$  stay near the origin as a consequence of (A.1),

$$\forall k \in \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \quad |\dot{X}_k(a\delta)| \leq \underbrace{\rho_a + \kappa(2\mathring{r} + \varepsilon)}_{\text{farthest position of spheres in } \mathring{\mathcal{J}}} + \mathring{r} + \varepsilon,$$

hence

$$\forall k \in \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon), \forall t \in [a\delta, (a+1)\delta], \quad |\dot{X}_k(t)| \leq \rho_a + \kappa(2\mathring{r} + \varepsilon) + \mathring{r} + \frac{5\varepsilon}{4}.$$

Inequality (A.2) for  $t = (a+1)\delta$  yields

$$\begin{aligned} k \notin \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon) &\implies |\dot{X}_k((a+1)\delta)| > \rho_a - \frac{\varepsilon}{4} \geq \rho_{a+1} + \kappa(2\mathring{r} + \varepsilon) + \mathring{r} + \varepsilon \\ &\implies k \notin \mathring{\mathcal{J}}(X((a+1)\delta), \rho_{a+1}, \varepsilon), \end{aligned}$$

thanks to the consequence of (A.1), applied to  $(a+1)$  instead of  $a$ . This proves the second inclusion:

$$\mathring{\mathcal{J}}(X((a+1)\delta), \rho_{a+1}, \varepsilon) \subset \mathring{\mathcal{J}}(X(a\delta), \rho_a, \varepsilon),$$

thus concluding the proof of the lemma.  $\square$



### A.3 Proof of Lemma 3.7 : long sphere chains are improbable

We prove here Lemma 3.7, that is, we find an upper bound for the probability of long sphere chains to form, at some time between 0 and 1, for the mixed penalised process starting from its reversible distribution.

The definition of  $\mathcal{B}'_{\text{Chain}}(\delta, \alpha, \kappa, \varepsilon)$  in Definition 3.1 and the fact that  $X(t)$  is  $\mu_R^y$ -distributed for each  $t$  under  $Q_R^y$  imply, for any time step  $\delta > 0$  :

$$\begin{aligned} Q_R^y\left(\mathcal{B}'_{\text{Chain}}(\delta, \alpha, \kappa, \varepsilon)\right) &= \bigcup_{k=0}^{\lfloor 1/\delta \rfloor} Q_R^y\left(\mathbf{X}(k\delta) \in \mathcal{B}'_{\text{Chain}}(\alpha, \kappa, \varepsilon)\right) \\ &\leq \frac{1}{\delta} \mu_R^y\left(\mathcal{B}_{\text{Chain}}(\alpha, \kappa, \varepsilon)\right). \end{aligned}$$

To complete the proof of Lemma 3.7, we only have to prove the following static estimate:

**Lemma A.1.** For each sphere activity  $\dot{z} > 0$ , any chain length  $\kappa \in \mathbb{N}^*$ , any radius  $\alpha > 0$ , and any leeway  $\varepsilon \in (0, 2\dot{r})$ ,

$$\forall y \in \mathcal{D}, \quad \forall R > 0, \quad \mu_R^y\left(\mathcal{B}_{\text{Chain}}(\alpha, \kappa, \varepsilon)\right) \leq \left(\dot{z} \left((2\dot{r} + \varepsilon)^d - (2\dot{r})^d\right) \nu_d\right)^\kappa.$$

*Proof.* The proof relies on an induction inequality for chains under the measure  $\nu_{R,n,m}^y$ .

We have to carefully manage the difference between labelled and unlabelled sphere configurations. Fix first the sphere indices in the norm, in order to avoid double counting. This is possible because the measure of the set of configurations where several spheres have the same norm is zero. The exchangeability of the measure  $\otimes_{i=1}^n \dot{\lambda}_R^y(d\dot{x}_i)$  ensures that:

$$\begin{aligned} &\nu_{R,n,m}^y\left(\mathcal{B}_{\text{Chain}}(\alpha, \kappa, \varepsilon)\right) \\ &= n! \int_{\mathbb{R}^{nd}} \int_{\mathbb{R}^{md}} \mathbb{1}_{\{|\dot{x}_1| < |\dot{x}_2| < \dots < |\dot{x}_n|\}} \mathbb{1}_{\{(\dot{x}_1, \dots, \dot{x}_n) \in \mathcal{B}_{\text{Chain}}(\alpha, \kappa, \varepsilon)\}} \mathbb{1}_{\mathcal{D}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) \otimes_{k=1}^m \dot{\lambda}_R^y(d\dot{x}_k) \otimes_{i=1}^n \dot{\lambda}_R^y(d\dot{x}_i). \end{aligned}$$

For any finite configuration  $\dot{\mathbf{x}} = (\dot{x}_1, \dots, \dot{x}_n) \in \mathcal{D}$ , the existence of at least one  $\varepsilon$ -chain of length  $\kappa$  starting in  $|B(0, \alpha)|$  is equivalent to the existence of a permutation  $\tau$  on  $\{1, \dots, n\}$  such that  $|\dot{x}_{\tau(1)}| \leq \alpha$  and  $|\dot{x}_{\tau(1)} - \dot{x}_{\tau(2)}| < 2\dot{r} + \varepsilon$ ,  $|\dot{x}_{\tau(2)} - \dot{x}_{\tau(3)}| < 2\dot{r} + \varepsilon$ ,  $\dots$ ,  $|\dot{x}_{\tau(\kappa)} - \dot{x}_{\tau(\kappa+1)}| < 2\dot{r} + \varepsilon$ . Using the inverse permutation, again denoted by  $\tau$ , we can rewrite the expression, and separately integrate with respect to  $\dot{x}_\kappa$  :

$$\begin{aligned} &\nu_{R,n,m}^y\left(\mathcal{B}_{\text{Chain}}(\alpha, \kappa, \varepsilon)\right) \\ &= n! \int_{\mathbb{R}^{nd}} \int_{\mathbb{R}^{md}} \mathbb{1}_{\{|\dot{x}_1| < |\dot{x}_2| < \dots < |\dot{x}_n|\}} \mathbb{1}_{\{\exists \tau: |\dot{x}_{\tau(1)}| \leq \alpha, |\dot{x}_{\tau(1)} - \dot{x}_{\tau(2)}| < 2\dot{r} + \varepsilon, \dots, |\dot{x}_{\tau(\kappa)} - \dot{x}_{\tau(\kappa+1)}| < 2\dot{r} + \varepsilon\}} \\ &\quad \mathbb{1}_{\mathcal{D}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) \otimes_{k=1}^m \dot{\lambda}_R^y(d\dot{x}_k) \otimes_{i=1}^n \dot{\lambda}_R^y(d\dot{x}_i) \\ &= n! \int_{\mathbb{R}^{nd}} \int_{\mathbb{R}^{md}} \mathbb{1}_{\{\exists \tau: |\dot{x}_{\tau(1)}| < |\dot{x}_{\tau(2)}| < \dots < |\dot{x}_{\tau(n)}|\}} \mathbb{1}_{\{|\dot{x}_1| \leq \alpha, |\dot{x}_1 - \dot{x}_2| < 2\dot{r} + \varepsilon, \dots, |\dot{x}_\kappa - \dot{x}_{\kappa+1}| < 2\dot{r} + \varepsilon\}} \\ &\quad \mathbb{1}_{\mathcal{D}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) \otimes_{k=1}^m \dot{\lambda}_R^y(d\dot{x}_k) \otimes_{i=1}^n \dot{\lambda}_R^y(d\dot{x}_i) \\ &= n! \int_{\mathbb{R}^{(n-1)d}} \int_{\mathbb{R}^{md}} \int_{\mathbb{R}^d} \mathbb{1}_{\{|\dot{x}_1| \leq \alpha, |\dot{x}_1 - \dot{x}_2| < 2\dot{r} + \varepsilon, \dots, |\dot{x}_\kappa - \dot{x}_{\kappa+1}| < 2\dot{r} + \varepsilon\}} \\ &\quad \mathbb{1}_{\mathcal{D}}(\dot{\mathbf{x}}\dot{\mathbf{x}}) \dot{\lambda}_R^y(d\dot{x}_{\kappa+1}) \otimes_{k=1}^m \dot{\lambda}_R^y(d\dot{x}_k) \otimes_{i=1, i \neq \kappa+1}^n \dot{\lambda}_R^y(d\dot{x}_i), \quad (\text{A.3}) \end{aligned}$$

since  $\mathbb{1}_{\{\exists \tau: |\dot{x}_{\tau(1)}| < |\dot{x}_{\tau(2)}| < \dots < |\dot{x}_{\tau(n)}|\}}$  is equal to 1 for almost every sphere configuration. Thanks to the inequality

$$\mathbb{1}_{\mathcal{D}}(\dot{x}_1, \dots, \dot{x}_n, \dot{\mathbf{x}}) \leq \mathbb{1}_{\mathcal{D}}(\dot{x}_1, \dots, \dot{x}_\kappa, \dot{x}_{\kappa+2}, \dots, \dot{x}_n, \dot{\mathbf{x}}) \mathbb{1}_{\{|\dot{x}_\kappa - \dot{x}_{\kappa+1}| \geq 2\hat{r}\}},$$

the integral on  $\dot{x}_{\kappa+1}$  can be isolated as

$$\int_{\mathbb{R}^d} \mathbb{1}_{\{2\hat{r} \leq |\dot{x}_\kappa - \dot{x}_{\kappa+1}| < 2\hat{r} + \varepsilon\}} \dot{\lambda}_R^{\mathbf{y}}(d\dot{x}_{\kappa+1}) \leq |B(\dot{x}_\kappa, 2\hat{r} + \varepsilon) \setminus B(\dot{x}_\kappa, 2\hat{r})| = ((2\hat{r} + \varepsilon)^d - (2\hat{r})^d) |B(0, 1)|.$$

We finally get

$$\begin{aligned} & \nu_{R,n,m}^{\mathbf{y}}(\mathcal{B}_{\text{Chain}}(\alpha, \kappa, \varepsilon)) \\ & \leq n! ((2\hat{r} + \varepsilon)^d - (2\hat{r})^d) |B(0, 1)| \\ & \int_{\mathbb{R}^{(n-1)d}} \int_{\mathbb{R}^{md}} \mathbb{1}_{\{|\dot{x}_1| \leq \alpha, |\dot{x}_1 - \dot{x}_2| < 2\hat{r} + \varepsilon, \dots, |\dot{x}_{\kappa-1} - \dot{x}_\kappa| < 2\hat{r} + \varepsilon\}} \\ & \mathbb{1}_{\mathcal{D}}(\dot{\mathbf{x}}) \otimes_{k=1}^m \dot{\lambda}_R^{\mathbf{y}}(d\dot{x}_k) \otimes_{i=1, i \neq \kappa+1}^n \dot{\lambda}_R^{\mathbf{y}}(d\dot{x}_i). \end{aligned}$$

From (A.3), the above integral is equal to  $(n-1)! \nu_{R,n-1,m}^{\mathbf{y}}(\mathcal{B}_{\text{Chain}}(\alpha, \kappa-1, \varepsilon))$ , so that we find

$$\nu_{R,n,m}^{\mathbf{y}}(\mathcal{B}_{\text{Chain}}(\alpha, \kappa, \varepsilon)) \leq n ((2\hat{r} + \varepsilon)^d - (2\hat{r})^d) |B(0, 1)| \nu_{R,n-1,m}^{\mathbf{y}}(\mathcal{B}_{\text{Chain}}(\alpha, \kappa-1, \varepsilon)).$$

Let us now transfer this chain induction inequality to the mixed probability measure  $\mu_R^{\mathbf{y}}$ . According to (2.5), and since configurations with less than  $\kappa+1$  spheres do not have  $\kappa$ -chains:

$$\begin{aligned} & \mu_R^{\mathbf{y}}(\mathcal{B}_{\text{Chain}}(\alpha, \kappa, \varepsilon)) \\ & = \frac{1}{\mathcal{Z}_R^{\mathbf{y}}} \sum_{m=0}^{+\infty} \frac{\dot{z}^m}{m!} \sum_{n=\kappa+1}^{+\infty} \frac{\dot{z}^n}{n!} \nu_{R,n,m}^{\mathbf{y}}(\mathcal{B}_{\text{Chain}}(\alpha, \kappa, \varepsilon)) \\ & \leq ((2\hat{r} + \varepsilon)^d - (2\hat{r})^d) |B(0, 1)| \frac{1}{\mathcal{Z}_R^{\mathbf{y}}} \sum_{m=0}^{+\infty} \frac{\dot{z}^m}{m!} \sum_{n=\kappa+1}^{+\infty} \frac{\dot{z}^n}{n!} n \nu_{R,n-1,m}^{\mathbf{y}}(\mathcal{B}_{\text{Chain}}(\alpha, \kappa-1, \varepsilon)) \\ & \leq ((2\hat{r} + \varepsilon)^d - (2\hat{r})^d) \nu_d \frac{\dot{z}}{\mathcal{Z}_R^{\mathbf{y}}} \sum_{m=0}^{+\infty} \frac{\dot{z}^m}{m!} \sum_{n=\kappa}^{+\infty} \frac{\dot{z}^n}{n!} \nu_{R,n,m}^{\mathbf{y}}(\mathcal{B}_{\text{Chain}}(\alpha, \kappa-1, \varepsilon)) \\ & \leq \dot{z} ((2\hat{r} + \varepsilon)^d - (2\hat{r})^d) \nu_d \mu_R^{\mathbf{y}}(\mathcal{B}_{\text{Chain}}(\alpha, \kappa-1, \varepsilon)). \end{aligned}$$

A straightforward induction argument completes the proof of Lemma A.1. □

#### A.4 Proof of Lemma 3.8 : fast moving spheres or particles are improbable

The oscillation estimates of the penalised processes stated in Lemma 3.8 rely on time reversal techniques. Let us first write the reversible penalised process as a sum of forward and backward Brownian motions. According to  $(\mathcal{S}_R^{\mathbf{y}})$ , the following processes are rescaled Brownian motions for  $1 \leq i, j \leq n$

and  $1 \leq k \leq m$ :

$$\left\{ \begin{array}{l} \dot{W}_i(t) = \dot{X}_i^{\mathbf{y},R,n,m}(t) - \dot{X}_i^{\mathbf{y},R,n,m}(0) + \frac{1}{2} \int_0^t \nabla \psi_R^{\mathbf{y}}(\dot{X}_i^{\mathbf{y},R,n,m}(s)) ds \\ \quad - \sum_{j=1}^n \int_0^t (\dot{X}_i^{\mathbf{y},R,n,m} - \dot{X}_j^{\mathbf{y},R,n,m})(s) dL_{ij}^{\mathbf{y},R,n,m}(s) \\ \quad - \sum_{k=1}^m \int_0^t (\dot{X}_i^{\mathbf{y},R,n,m} - \dot{X}_k^{\mathbf{y},R,n,m})(s) d\ell_{ik}^{\mathbf{y},R,n,m}(s) \\ \dot{W}_k(t) = \dot{X}_k^{\mathbf{y},R,n,m}(t) - \dot{X}_k^{\mathbf{y},R,n,m}(0) + \frac{\sigma^2}{2} \int_0^t \nabla \psi_R^{\mathbf{y}}(\dot{X}_k(s)) ds \\ \quad - \sigma^2 \sum_{i=1}^n \int_0^t (\dot{X}_k^{\mathbf{y},R,n,m} - \dot{X}_i^{\mathbf{y},R,n,m})(s) d\ell_{ik}^{\mathbf{y},R,n,m}(s). \end{array} \right.$$

The forward process  $(\dot{X}_i^{\mathbf{y},R,n,m}(t), \dot{X}_k^{\mathbf{y},R,n,m}(t), L_{ij}^{\mathbf{y},R,n,m}(t), \ell_{ik}^{\mathbf{y},R,n,m}(t))_{t \in [0,1], 1 \leq i, j \leq n, 1 \leq k \leq m}$  has the same distribution as the backward process

$$(\dot{X}_i^{\mathbf{y},R,n,m}(1-t), \dot{X}_k^{\mathbf{y},R,n,m}(1-t), L_{ij}^{\mathbf{y},R,n,m}(1-t), \ell_{ik}^{\mathbf{y},R,n,m}(1-t))_{t \in [0,1], 1 \leq i, j \leq n, 1 \leq k \leq m}.$$

Hence, the following images of the backward process also are rescaled Brownian motions:

$$\left\{ \begin{array}{l} \overleftarrow{W}_i^{\mathbf{y},R,n,m}(t) = \dot{X}_i^{\mathbf{y},R,n,m}(1-t) - \dot{X}_i^{\mathbf{y},R,n,m}(1) + \frac{1}{2} \int_0^t \nabla \psi_R^{\mathbf{y}}(\dot{X}_i^{\mathbf{y},R,n,m}(1-s)) ds \\ \quad - \sum_{j=1}^n \int_0^t (\dot{X}_i^{\mathbf{y},R,n,m} - \dot{X}_j^{\mathbf{y},R,n,m})(1-s) dL_{ij}^{\mathbf{y},R,n,m}(1-s) \\ \quad - \sum_{k=1}^m \int_0^t (\dot{X}_i^{\mathbf{y},R,n,m} - \dot{X}_k^{\mathbf{y},R,n,m})(1-s) d\ell_{ik}^{\mathbf{y},R,n,m}(1-s) \\ \overleftarrow{\dot{W}}_k^{\mathbf{y},R,n,m}(t) = \dot{X}_k^{\mathbf{y},R,n,m}(1-t) - \dot{X}_k^{\mathbf{y},R,n,m}(1) + \frac{\sigma^2}{2} \int_0^t \nabla \psi_R^{\mathbf{y}}(\dot{X}_k(1-s)) ds \\ \quad - \sigma^2 \sum_{i=1}^n \int_0^t (\dot{X}_k^{\mathbf{y},R,n,m} - \dot{X}_i^{\mathbf{y},R,n,m})(1-s) d\ell_{ik}^{\mathbf{y},R,n,m}(1-s). \end{array} \right.$$

As in [FKRZ24], the following relations hold for every  $t \in [0, 1]$ :

$$\begin{aligned} \dot{X}_i^{\mathbf{y},R,n,m}(t) &= \dot{X}_i^{\mathbf{y},R,n,m}(0) + \frac{1}{2} \left( \dot{W}_i(t) + \overleftarrow{W}_i^{\mathbf{y},R,n,m}(1-t) - \overleftarrow{W}_i^{\mathbf{y},R,n,m}(1) \right), \\ \dot{X}_k^{\mathbf{y},R,n,m}(t) &= \dot{X}_k^{\mathbf{y},R,n,m}(0) + \frac{\sigma^2}{2} \left( \dot{W}_k(t) + \overleftarrow{W}_k^{\mathbf{y},R,n,m}(1-t) - \overleftarrow{W}_k^{\mathbf{y},R,n,m}(1) \right). \end{aligned} \tag{A.4}$$

Let us now compute the probability that, among the  $n$  spheres and  $m$  particles in the reversible solution  $(\dot{X}_i^{\mathbf{y},R,n,m}, \dot{X}_k^{\mathbf{y},R,n,m})_{1 \leq i, j \leq n, 1 \leq k \leq m}$  of the finite-dimensional SDE with initial condition distributed according to  $\nu_{R,n,m}^{\mathbf{y}}$ , at least one sphere or particle that enters the  $\alpha$ -neighbourhood of the origin oscillates too much (see Definition 3.2):

$$Q_{R,n,m}^{\mathbf{y}}(\mathcal{B}'_{\text{Fast}}(\alpha, \delta, \varepsilon)) = \int P(\mathbf{X}^{\mathbf{y},R,n,m}(\mathbf{x}, \cdot) \in \mathcal{B}'_{\text{Fast}}(\alpha, \delta, \varepsilon)) \nu_{R,n,m}^{\mathbf{y}}(d\mathbf{x}).$$

The  $n$  spheres (resp.  $m$  particles) are exchangeable, thus

$$\begin{aligned} &Q_{R,n,m}^{\mathbf{y}}(\mathcal{B}'_{\text{Fast}}(\alpha, \delta, \varepsilon)) \\ &\leq n Q_{R,n,m}^{\mathbf{y}}(\dot{X}_1(\cdot) \in \mathcal{B}'_{\text{Fast}}(\alpha, \delta, \varepsilon)) + m Q_{R,n,m}^{\mathbf{y}}(\dot{X}_1(\cdot) \in \mathcal{B}'_{\text{Fast}}(\alpha, \delta, \varepsilon)). \end{aligned} \tag{A.5}$$

We evaluate separately first the sphere term and then the particle term.

According to Definition 3.2, for  $\alpha_0 = 0$  and  $\alpha_j = \alpha + \varepsilon/\delta + j/\delta$  if  $j \in \mathbb{N}^*$ , the event  $\mathring{X}_1(\cdot) \in \mathcal{B}_{\text{Fast}}(\alpha, \delta, \varepsilon)$  is equivalent to:

$$\begin{aligned} & \exists j, j' \in \mathbb{N} \text{ s.t. } \alpha_j \leq |\mathring{X}_1(0)| < \alpha_{j+1} \text{ and } \alpha_{j'} \leq |\mathring{X}_1(1)| < \alpha_{j'+1} \text{ and} \\ & \min_{0 \leq s \leq 1} |\mathring{X}_1(s)| \leq \alpha \text{ and } \sup_{\substack{|t-s| < \delta \\ 0 \leq s, t \leq 1}} |\mathring{X}_1(t) - \mathring{X}_1(s)| > \varepsilon. \end{aligned}$$

For a path to go from the outside of the ball with radius  $\alpha_j$  to the inside of the ball with radius  $\alpha$  in a time duration of less than 1, the distance covered during at least one of the  $1/\delta$  time intervals of length  $\delta$  has to be larger than  $\delta(\alpha_j - \alpha) = \varepsilon + j$ . Hence  $\mathring{X}_1(\cdot) \in \mathcal{B}_{\text{Fast}}(\alpha, \delta, \varepsilon)$  implies:

$$\begin{aligned} & \exists j, j' \in \mathbb{N} \text{ s.t. } \alpha_j \leq |\mathring{X}_1(0)| < \alpha_{j+1} \text{ and } \alpha_{j'} \leq |\mathring{X}_1(1)| < \alpha_{j'+1} \text{ and} \\ & \min_{0 \leq s \leq 1} |\mathring{X}_1(s)| \leq \alpha \text{ and } \sup_{\substack{|t-s| < \delta \\ 0 \leq s, t \leq 1}} |\mathring{X}_1(t) - \mathring{X}_1(s)| > \varepsilon + \max(j, j'). \end{aligned}$$

Using expression (A.4), we get:

$$\begin{aligned} & \sup_{\substack{|t-s| < \delta \\ 0 \leq s, t \leq 1}} |\mathring{X}_1^{\mathbf{y}, R, n, m}(t) - \mathring{X}_1^{\mathbf{y}, R, n, m}(s)| > \eta \\ \implies & \sup_{\substack{|t-s| < \delta \\ 0 \leq s, t \leq 1}} |\mathring{W}_1^{\mathbf{y}, R, n, m}(t) - \mathring{W}_1^{\mathbf{y}, R, n, m}(s)| > \eta \text{ or } \sup_{\substack{|t-s| < \delta \\ 0 \leq s, t \leq 1}} |\mathring{W}_1^{\mathbf{y}, R, n, m}(t) - \mathring{W}_1^{\mathbf{y}, R, n, m}(s)| > \eta. \end{aligned}$$

Thus, thanks to the reversibility of  $Q_{R, n, m}^{\mathbf{y}}$ , we have:

$$\begin{aligned} & Q_{R, n, m}^{\mathbf{y}} \left( \mathring{X}_1^{\mathbf{y}, R, n, m}(\cdot) \in \mathcal{B}_{\text{Fast}}(\alpha, \delta, \varepsilon) \right) \\ & \leq 2 \sum_{j \in \mathbb{N}} Q_{R, n, m}^{\mathbf{y}} \left( \alpha_j \leq |\mathring{X}_1^{\mathbf{y}, R, n, m}(0)| < \alpha_{j+1} \text{ and } \sup_{\substack{|t-s| < \delta \\ 0 \leq s, t \leq 1}} |\mathring{W}_1^{\mathbf{y}, R, n, m}(t) - \mathring{W}_1^{\mathbf{y}, R, n, m}(s)| > \varepsilon + j \right). \end{aligned}$$

We then use the following standard Brownian estimate:

**Lemma A.2.** If  $W$  is a  $d$ -dimensional Brownian motion then, for every  $\varepsilon > 0$  and every  $\delta \in ]0, 1]$ ,

$$P \left( \sup_{\substack{|t-s| < \delta, \\ 0 \leq s, t \leq 1}} |W(t) - W(s)| \geq \varepsilon \right) \leq 4\sqrt{5} \frac{d}{\delta} \exp \left( -\frac{\varepsilon^2}{10 d \delta} \right).$$

The Brownian motion is independent of the starting point in equation  $(S_R^{\mathbf{y}})$ , thus Lemma A.2 implies:

$$\begin{aligned} & Q_{R, n, m}^{\mathbf{y}} \left( \mathring{X}_1^{\mathbf{y}, R, n, m}(\cdot) \in \mathcal{B}_{\text{Fast}}(\alpha, \delta, \varepsilon) \right) \\ & \leq 8\sqrt{5} \frac{d}{\delta} \sum_{j=0}^{+\infty} \nu_{R, n, m}^{\mathbf{y}} \left( \alpha_j \leq |\mathring{x}_1| < \alpha_{j+1} \right) \exp \left( -\frac{(\varepsilon + j)^2}{10 d \delta} \right). \end{aligned}$$

Then (2.3) yields:  $\nu_{R, n, m}^{\mathbf{y}}(\alpha_j \leq |\mathring{x}_1| < \alpha_{j+1}) \leq \nu_{R, n-1, m}^{\mathbf{y}}(\mathcal{D}) |B(0, \alpha_{j+1}) \setminus B(0, \alpha_j)|$ , so that the above quantity is bounded by:

$$\leq 8\sqrt{5} \frac{d}{\delta} \exp \left( -\frac{\varepsilon^2}{10 d \delta} \right) \nu_{R, n-1, m}^{\mathbf{y}}(\mathcal{D}) \sum_{j=0}^{+\infty} e^{-\frac{2\varepsilon j + j^2}{10 d \delta}} |B(0, \alpha_{j+1})|.$$

The above sum is smaller than some constant  $\dot{C}_d$  which only depends on the dimension  $d$ . Putting in  $\dot{C}_d$  the factors depending only on the dimension, we obtain:

$$Q_{R,n,m}^y \left( \dot{X}_1^{y,R,n,m}(\cdot) \in \mathcal{B}_{\text{Fast}}(\alpha, \delta, \varepsilon) \right) \leq \frac{\dot{C}_d}{\delta^{d+1}} \exp\left(-\frac{\varepsilon^2}{10 d \delta}\right) \nu_{R,n-1,m}^y(\mathcal{D}) \alpha^d.$$

We evaluate in the same way the particle term  $Q_{R,n,m}^y \left( \dot{X}_1(\cdot) \in \mathcal{B}_{\text{Fast}}(\alpha, \delta, \varepsilon) \right)$ . The only difference is that the diffusion coefficient is equal to  $\dot{\sigma}$  instead of 1. We obtain again, for  $\varepsilon \leq 1$ ,  $\delta \leq 1$ , and  $\alpha \geq 1$ , and with some constant  $\dot{C}_d$  which depends only on the dimension  $d$ :

$$Q_{R,n,m}^y \left( \dot{X}_1^{y,R,n,m}(\cdot) \in \mathcal{B}_{\text{Fast}}(\alpha, \delta, \varepsilon) \right) \leq \frac{\dot{C}_d}{\delta^{d+1}} \exp\left(-\frac{\varepsilon^2}{10 d \delta \dot{\sigma}^2}\right) \nu_{R,n,m-1}^y(\mathcal{D}) \alpha^d.$$

Thanks to (A.5),

$$\begin{aligned} & Q_{R,n,m}^y(\mathcal{B}'_{\text{Fast}}(\alpha, \delta, \varepsilon)) \\ & \leq \frac{\alpha^d}{\delta^{d+1}} \exp\left(-\frac{\varepsilon^2}{10 d \delta \max(1, \dot{\sigma}^2)}\right) \left( n \dot{C}_d \nu_{R,n-1,m}^y(\mathcal{D}) + m \dot{C}_d \nu_{R,n,m-1}^y(\mathcal{D}) \right). \end{aligned}$$

We can now compute the probability of fast motion under the Poissonian mixture  $Q_R^y$ :

$$\begin{aligned} Q_R^y(\mathcal{B}'_{\text{Fast}}(\alpha, \delta, \varepsilon)) & \leq \frac{\alpha^d}{\delta^{d+1}} \exp\left(-\frac{\varepsilon^2}{10 d \delta \max(1, \dot{\sigma}^2)}\right) \frac{1}{\mathcal{Z}_R^y} \\ & \quad \left( \underbrace{\dot{z} \dot{C}_d \sum_{n=1}^{+\infty} \frac{\dot{z}^{n-1}}{(n-1)!} \sum_{m=0}^{+\infty} \frac{\dot{z}^m}{m!} \nu_{R,n-1,m}^y(\mathcal{D})}_{=\mathcal{Z}_R^y} \right. \\ & \quad \left. + \dot{z} \dot{C}_d \sum_{n=0}^{+\infty} \frac{\dot{z}^n}{n!} \sum_{m=1}^{+\infty} \frac{\dot{z}^{m-1}}{(m-1)!} \nu_{R,n,m-1}^y(\mathcal{D}) \right)_{=\mathcal{Z}_R^y}. \end{aligned}$$

Choosing  $C_{\text{Fast}} := \dot{C}_d \dot{z} + \dot{C}_d \dot{z}$  completes the proof of Lemma 3.8.

## References

- [AO54] S. Asakura and F. Oosawa. On interaction between two bodies immersed in a solution of macro-molecules. *J. Chem. Phys.*, 22(7):1255–1256, 1954.
- [CCK95] J. T. Chayes, L. Chayes, and R. Kotecký. The analysis of the Widom–Rowlinson model by stochastic geometric methods. *Comm. Math. Phys.*, 172(3):551–569, 1995.
- [DI93] P. Dupuis and H. Ishii. SDEs with oblique reflection on nonsmooth domains. *Ann. Probab.*, 21(1):554–580, 1993.
- [DI08] P. Dupuis and H. Ishii. Correction : SDEs with oblique reflection on nonsmooth domains. *Ann. Probab.*, 36(5):1992–1997, 2008.
- [FKRZ24] M. Fradon, J. Kern, S. Roelly, and A. Zass. Diffusion dynamics for an infinite system of two-type spheres and the associated depletion effect. *Stoch. Process. Appl.*, 171:104319, 2024.

- [FR00] M. Fradon and S. Roelly. Infinite dimensional diffusion processes with singular interaction. *Bull. Sci. Math.*, 124(4):287–318, 2000.
- [FR07] M. Fradon and S. Roelly. Infinite system of Brownian balls with interaction: The non-reversible case. *ESAIM Probab. Stat.*, 11:55–79, 2007.
- [Fra13] M. Fradon. Stochastic differential equations on domains defined by multiple constraints. *Electron. Commun. Probab.*, 18(26):1–13, 2013.
- [FRT00] M. Fradon, S. Roelly, and H. Tanemura. An infinite system of Brownian balls with infinite range interaction. *Stoch. Proc. Appl.*, 90:43–66, 2000.
- [Geo79] H.-O. Georgii. *Canonical Gibbs Measures: Some Extensions of de Finetti's Representation Theorem for Interacting Particle Systems*. Lecture Notes in Mathematics. Springer Berlin Heidelberg, 1979.
- [Geo94] H.-O. Georgii. Large deviations and the equivalence of ensembles for Gibbsian particle systems with superstable interaction. *Probab. Theory Relat. Fields.*, 99(2):171–195, 1994.
- [Geo95] H.-O. Georgii. The equivalence of ensembles for classical systems of particles. *J. Stat. Phys.*, 80(5):1341–1378, 1995.
- [GLM95] G. Giacomin, J. L. Lebowitz, and C. Maes. Agreement percolation and phase coexistence in some gibbs systems. *J. Stat. Phys.*, 80(5-6):1379–1403, 1995.
- [JKSZ24] B. Jahnel, J. Köppl, Y. Steenbeck, and A. Zass. The variational principle for a marked Gibbs point process with infinite-range multibody interactions. *Preprint*, arXiv:2408.17170, 2024.
- [JT20] S. Jansen and D. Tsagkarogiannis. Cluster expansions with renormalized activities and applications to colloids. *Ann. Henri Poincaré*, 21(1):45–79, 2020.
- [Lan77] R. Lang. Unendlich-dimensionale Wienerprozesse mit Wechselwirkung. *Z. Wahrsch. Verw. Geb.*, 38:55–72, 1977.
- [LS84] P. L. Lions and A. S. Sznitman. Stochastic differential equations with reflecting boundary conditions. *Commun. Pure Appl. Mathem.*, 37:511–537, 1984.
- [LT11] H. N. W. Lekkerkerker and R. Tuinier. Colloids and the depletion interaction. *Lecture Notes in Physics*, 2011.
- [MMS<sup>+</sup>01] S. Mase, J. Møller, D. Stoyan, R. P. Waagepetersen, and G. Döge. Packing densities and simulated tempering for hard core Gibbs point processes. *Ann. Inst. Statist. Math.*, 53(4):661–680, 2001.
- [Per09] J. Perrin. Mouvement brownien et réalité moléculaire. *Ann. Chim. Phys.*, 15(8):5–114, 1909.
- [Rue99] D. Ruelle. *Statistical Mechanics*. World Scientific and Imperial College Press, 1999.
- [Sai87] Y. Saisho. Stochastic differential equations for multi-dimensional domain with reflecting boundary. *Probab. Theory Rel. Fields*, 74:455–477, 1987.
- [Sko61] A. V. Skorokhod. Stochastic equations for diffusion processes in a bounded region. *Teor. Veroyatn. Primen.*, 6(3):264–274, 1961.
- [Sko62] A. V. Skorokhod. Stochastic equations for diffusion processes in a bounded region. ii. *Teor. Veroyatn. Primen.*, 7(1):3–23, 1962.
- [Tan79] H. Tanaka. Stochastic differential equations with reflecting boundary condition in convex regions. *Hiroshima Math. J.*, 9(1):163 – 177, 1979.

- [Tan96] H. Tanemura. A system of infinitely many mutually reflecting brownian balls. *Probab. Th. Rel. Fields*, 104:399–426, 1996.
- [Tan22] H. Tanemura. *Infinite Particle Systems with Hard-Core and Long-Range Interaction*, pages 511–533. Dirichlet Forms and Related Topics. Springer, 2022.
- [TMLS97] R. H. Schonmann T. M. Liggett and A. M. Stacey. Domination by product measures. *Ann. Probab.*, 25(1):71–95, 1997.
- [Vri76] A. Vrij. Polymers at interfaces and the interactions in colloidal dispersions. *Pure and Appl. Chem.*, 48(4):471–483, 1976.
- [WJL23] R. Wittmann, S. Jansen, and H. Löwen. Geometric criteria for the absence of effective many-body interactions in nonadditive hard particle mixtures. *J. Math Phys.*, 64:103301, 2023.
- [WR70] B. Widom and J. Rowlinson. New model for the study of liquid–vapor phase transitions. *J. Chem. Phys.*, 52(4):1670–1684, 1970.