

Homogenization of a system of multi-species nonlinear diffusion-reaction equations in an $H^{1,p}$ setting

Hari Shankar Mahato

Center of Industrial Mathematics,
Department of Mathematics and Informatics
University of Bremen, Germany

Abstract. In this talk, we consider a system of highly nonlinear multi-species diffusion-reaction equations with homogeneous Neumann boundary condition. All reactions are reversible. For this system, the existence and uniqueness of the weak solution are proved at the micro scale on the interval $[0, T)$ for any $T > 0$. We obtain, global in time, L^∞ - estimates of the solution with the help of a Lyapunov functional. For the existence of the solution, we use Schaefer's fixed point theorem, maximal regularity and Lyapunov type arguments (cf. [3]). In the second part, we upscale the model from the micro to the macro scale by using *two-scale convergence* and *periodic unfolding* (cf. [1], [2]).

References

- 1 G. Allaire, Homogenization and two scale convergence, *SIAM Journal of math. anal.*, Vol. 23, No. 6, pp. 1482-1518, 1992.
- 2 D. Cioranescu, A. Damlamian and G. Griso, Periodic unfolding and homogenization, *C. R. Acad. Sci. Paris, Ser. I 335*, 99104, 2002.
- 3 S. Kräutle, *General multi-species reactive transport problems in porous media: Efficient numerical approaches and existence of global solutions*, Habilitation thesis, Universität Erlangen-Nürnberg, Germany, 2008.